

Research Paper

A real-world 1:1 propensity-matched study revealed unmarried status was independently associated with worse survival for patients with renal clear cell carcinoma

Shi-Long Zhang^{1*}, Hai-Tao Sun^{2,3*}, Zhan-Ming Li¹, Zheng-Yan Zhang¹, Wen-Rong Wang⁴, Xin Wang⁵, Zhi-Ming Wang^{6,7}✉, Li-Shun Wang¹✉

1. Institute of Fudan-Minhang Academic Health System, Minhang Branch, Zhongshan hospital, Fudan University, Shanghai 201199, P.R. China.
2. Shanghai Institute of Medical Imaging, Shanghai 200032, P.R. China
3. Department of Interventional Radiology, Zhongshan Hospital, Fudan University, Shanghai 200032, P.R. China
4. Faculty of Physical Education, Shandong Normal University, Jinan 250014, P.R. China
5. Department of acupuncture and moxibustion, Central Hospital of Shanghai Xuhui District, Shanghai 200031, P.R. China.
6. Department of Medical oncology, Zhongshan Hospital, Fudan University, No. 180 Fenglin Road, Shanghai, 200032, P.R. China
7. Xiamen branch, Zhongshan Hospital, Fudan University, No. 668 Jinhu Road, Xiamen, 361000, P.R. China.

*Shi-Long Zhang and Hai-Tao Sun contribute equally to this paper.

✉ Corresponding authors: Li-Shun Wang, M.D., Ph.D., Institute of Fudan-Minhang Academic Health System, Minhang Branch, Zhongshan hospital, Fudan University, No. 170 Xinsong Road, Minhang, Shanghai, 200032, P.R. China. Tel: 86-021-60267405; E-mail: lishunwang@fudan.edu.cn. Zhi-Ming Wang, M.D., Ph.D., Department of Medical oncology, Zhongshan Hospital, Fudan University, No. 180 Fenglin Road, Shanghai, 200032, P.R. China; Xiamen branch, Zhongshan Hospital, Fudan University, No. 668 Jinhu Road, Xiamen, 361000, P.R. China. Tel: 86-021-64041990; E-mail: wzming@126.com

© Ivyspring International Publisher. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY-NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Received: 2018.11.23; Accepted: 2019.05.07; Published: 2019.06.09

Abstract

Background: Marital status has been reported as an independent prognostic factor for survival in various cancers, but it has been rarely studied in renal clear cell carcinoma (ccRCC). In this study, we aimed to assess the impact of marital status on the survival of ccRCC patients.

Methods: We retrospectively investigated the Surveillance, Epidemiology, and End Results (SEER) database and identified 68599 of ccRCC patients between 1973 and 2015. These patients were divided into married, single, divorced and widowed groups. The survival differences among these groups were assessed by Kaplan-Meier method and log-rank test. Multivariate Cox regression analyses were performed to identify the overall survival (OS) and cancer-specific survival (CSS) independent factors. Furthermore, 1:1 propensity score matching (PSM) analysis was performed to minimize the potential confounding factors.

Results: Of the 68599 ccRCC patients, 44553 (64.95%) patients were married, 7410 (10.80%) were divorced, 10663 (15.54%) were single, and 5973 (8.71%) were widowed. The 5-year OS was 79.0%, 73.8%, 77.3%, and 66.4 % in the married, divorced, single, and widowed groups, respectively ($p = 0.001$) and the corresponding 5-year CSS rates were 85.5%, 83.3%, 80.8%, 76.5%, respectively. Multivariate Cox regression analysis marital status was the independent prognostic factor for OS and CSS. Compared with the married patients, the divorced, single, and widowed patients faced increased higher mortality risks for OS and CSS. In stratified analyses by sex, surgery conditions and cancer stages, those unmarried patients still had worse prognosis. The results were further confirmed in the 1:1 matched group.

Conclusion: Unmarried ccRCC patients experienced worse survival than their married counterparts. Among the unmarried patients, the widowed suffered the highest mortality risks for OS and CSS.

Key words: Renal clear cell carcinoma; marital status; cancer survival; propensity score matching; SEER

Introduction

Renal cell carcinoma (RCC) is a genitourinary malignancy. Its incidence has significantly increased¹, and contributed to approximately 63,990 new cases and approximately 693,000 deaths worldwide in

2017². Among the histological subtypes of RCC, clear cell RCC (ccRCC) is the most common subtype, responsible for 75–80% of all RCC cases. The prognosis of ccRCC is determined by many factors, including age, sex, disease stage, Fuhrman nuclear grade, tumor size, molecular pathogenesis and treatment strategies. Nowadays people have realized that an integrative concept of health and disease should include the interaction of biology, psychology and sociology, which is also known as biological-psychology-social medical model³⁻⁴.

Recently, results from considerable literature have disclosed that that married patients have superior survival compared to the unmarried in various cancers⁵, such as soft tissue sarcoma⁶, liver cancer⁷ and colon cancer⁸. This interesting phenomenon arise much public attentions. It is postulated that married status could contribute to optimistic psychological, enough social support, decent incomes, healthy lifestyle and comfortable living conditions. Similarly, previous studies reported that that marital status was a prognostic factor of survival in kidney cancer patients⁹⁻¹¹. However, all of them merely focus on all kinds of kidney cancer without differentiating pathological subtypes. As we all know, RCC is a highly heterogenous tumor, ccRCC and other subtypes (eg, papillary, chromophobe) have distinct pathologies and biological behaviors, or even different long-term survival. So conducting analysis on the survival of ccRCC and other subtypes separately might be more important and reasonable. Furthermore, previous studies had significant imbalance baseline that married people were more likely to be diagnosed at an earlier stage and receive surgery compared to the unmarried. Thus, the impact of marital status on the survival of ccRCC has not been rigorously investigated.

In this study, we conducted a comprehensive study that involved a large sample ccRCC patients diagnosed between 1973 and 2015 to explore the relationship between marital status and ccRCC survival, as well as the potential underlying mechanisms. Furthermore, we also conducted 1:1 propensity score matching (PSM) analysis, a powerful method to minimize selection bias, to created 1:1 matched cohort with well-balanced baseline characteristics. In addition, we performed Cox proportional hazards regression to explore the impact of marital status on ccRCC patients in the matched cohort.

Patients and Methods

Patient selection

Our study used the SEER database-18 cohort database [Incidence -SEER 18 Regs Research Data +

Hurricane Katrina Impacted Louisiana Cases, Nov 2017 Sub (1973–2015 varying)], released in November 2017, as data source¹². We got access to the SEER database with the ID number 13264-Nov2017. Using the SEER-stat software (SEER*Stat 8.2.1), patients with ICD-O-3 (International Classification of Diseases for Oncology, 3rd edition) site code C64.9 diagnosed between 1973 and 2015 were identified from the SEER database. Patients were included according to the following criteria: (1) their ICD-O-3 morphology code indicated ccRCC; (2) they aged more than 18 years at diagnosis; (3) their marital status were known; (4) they were diagnosed with ccRCC only or more than one primary cancer but ccRCC was the first; (5) their cause of death was known; (6) their survival time were known and greater than 0 month.

Study variables

The sex, age, race, diagnosis year, pathological grade, marital status, AJCC stage, surgery status, median household income, insurance status, cause of death, vital status and survival time from the SEER database. Marital status was categorized as married, divorced, single and widowed. Additionally, marital status was also categorized as married and unmarried (single, divorced and widowed) groups in the 1:1 PSM analysis. Patients were divided into three groups: 18 to 49 years, 50 to 59 years, 60 to 69 years, 70 to 79 years, and ≥ 80 years. Race was classified as white, black, or others (American Indian/AK Native, Asian/Pacific Islander). Diagnosis years were divided into three periods (1973-1993, 1994-2004, and 2005-2015) to eliminate the survival benefits brought by targeted therapy in recent years. County-level median household income was included to represent patients' socioeconomic status and stratified into quintiles for analysis: Quartile 1 (<US \$48700), Quartile 2 (US \$48701-56200), Quartile 3 (US \$56201-66931), and Quartile 4 (>US \$66931). The included patients were furthermore stratified into "insured", "uninsured" and "unknown" groups according to their insurance status. Tumor stage were listed as stage I, stage II, stage III, stage IV according to the American Joint Committee on Cancer staging (AJCC sixth edition).

Statistical analyses

The demographic and clinical data were presented by percentage (%). Differences in baseline characteristics were compared by χ^2 test. The primary endpoints in our study, overall survival (OS) and cancer-specific survival (CSS), were calculated by Kaplan-Meier method and log-rank test was used to detect survival differences. Multivariate Cox regression analysis were performed to identify the independent prognostic factors for ccRCC patients.

To further control for potential baseline confounding factors across groups, we carried out 1:1 PSM analysis based on marital status (married and unmarried group) to re-examine the impact of marital status. In our study, a 1:1 pair matching (without replacement) was conducted through nearest neighbor method with a caliper of 0.1 times the standard deviation of the propensity score¹³. The matching was performed by the MatchIt package in R (version 3.5.1). Standardized differences (SD) were used to examine the balance across baseline covariates before and after matching, and a SD below 0.1 was reliable enough to provide well-balanced covariates after matching¹⁴.

All statistical analyses and figures were generated using the tableone, rms, survival, survminer, ggplot2 and MatchIt packages in R (version 3.5.1), unless otherwise specified. All *p* values were two-sided with statistical significance defined at < 0.05.

Results

Patient baseline characteristics

A total of 68599 eligible patients diagnosed from 1973 to 2015 in the SEER database were included. 43271 (63.08%) were male and 25328 (36.92%) were female, 44553 (64.95%) were married, 7410 (10.80%) were divorced, 10663 (15.54%) were single, and 5973 (8.71%) widowed. Baseline characteristics of ccRCC patients according to marital status were listed in Table 1. Difference of baseline characteristics were noticed significantly in all subgroups. Especially, married group had the highest percentage of male patients (69.2%) and white patients (83.3%), while among unmarried patients, single group had the highest percentage of male patients (63.3%) and widowed group had the highest percentage of female (72.6%). Widowed patients tended to be in age groups of 60-69 (25.8%), 70-79 (39.3%), and ≥ 80 years (26.3%), while the single patients were predominantly in younger age group of 18-49 (33.4%), and more tended to present with smaller tumor sizes (54.7%) and early stages (65.2%), compared to other unmarried groups.

Impact of marital status on overall survival (OS)

The OS of ccRCC patients was calculated by Kaplan-Meier method. The results showed a significant survival difference according to marital status (log-rank test *p* < 0.001) (Figure 1A). The 5-year OS was 79.0% in the married group, 73.8% in the divorced group, 77.3% in the single group, and 66.4% in the widowed group (*p* < 0.001). Univariate analysis identified marital status, age, race, diagnosis year,

histological type, pathological grade, tumor size, AJCC stage, surgery status, median household income and insurance status as significant factors associated with OS (*p* < 0.001). After controlling for above-mentioned factors, multivariate Cox regression analysis showed that, compared to the married (as the reference group), divorced (hazard ratio (HR), 1.33, 95% confidence interval (CI): 1.27-1.40), single (HR, 1.26, 95%CI: 1.20-1.32), and widowed (HR, 1.42, 95%CI: 1.35-1.49) patients had higher death risks for OS (Table 2). In addition, widowed patients had the lowest rate and the highest death risks for OS. Notably, socioeconomic factors including median household income and insurance status were significantly associated with OS in both univariate and multivariate analysis.

Impact of marital status on cancer-specific survival (CSS)

The 5-year CSS rates for married patients, divorced, widowed, single patients were 85.5%, 83.3%, 80.8%, 76.5%, respectively (*p* < 0.001) (Figure 1B, Table 3). By univariate analysis, all the baseline characteristics were found to be associated with CSS among ccRCC patients. In addition, patients with insurance had better CSS than those without (5-year CSS rate 85.2 % vs. 83.2%, *p* < 0.001). When the aforementioned covariates were adjusted in multivariate Cox regression analysis, marital status was confirmed as an independent prognostic factor, with worse CSS among unmarried patients (divorced, HR, 1.16, 95% CI 1.08-1.25, *p* < 0.001; single, HR, 1.09, 95% CI 1.02-1.16, *p* = 0.006; widowed, HR, 1.20, 95% CI 1.13-1.28, *p* < 0.001). However, multivariate Cox regression analysis did not support the association between insurance status and CSS (HR, 1.04, 95% CI 0.96-1.14, *p* = 0.314).

Subgroup analysis stratified by sex, surgery, and cancer stage

Multiple variables including have been identified as prognostic factors for ccRCC mortality, and those variables also been verified independently in our study. Hence, subsequently we divided all ccRCC patients into several subgroups stratified by those variables and investigated their impacts on CSS. The survival curves of the patients within sex, surgery and AJCC stage group were shown in Figure 2, Figure 3 and Figure 4, respectively. Marital status remained an independent prognostic factor in almost all subgroups (*p* < 0.001) (Table 4). Furthermore, in the Cox regression analysis, we also observed several interesting findings: 1) The widowed patients suffered the worst CSS in each sex, surgery and AJCC stage subgroups, and being widowed was associated

with the highest death risks compared to other unmarried status. 2) Being divorced would raise death risks in comparison to being married in males, patients with surgery, and all AJCC stages. 3) No significant survival differences were noticed between the married and single group in the subgroup of female, and stage II stage III.

Survival analysis of ccRCC patients in the 1:1 matched cohort

To eliminate influences of confounding factors across the baseline characteristics and ensure our observations were reliable and stable, we conducted a 1:1 matched cohort analysis through PSM method. After matching, we had 18028 ccRCC patients including 9014 married and another 9014 unmarried patients in our subsequent analysis. Distribution of

the baseline characteristics was well-balanced in the matched cohort (Table 5). In general, an absolute SD < 0.10 indicated a negligible difference across the groups; and the largest SD was 0.03 in the matched cohort.

Even so, unmarried patients persisted to suffer more significant survival disadvantages than married patients in the Kaplan-Meier analysis. The 5-year OS rate in married patients was 75.8% while 5-year OS rate was 68.3% in unmarried group ($p < 0.001$) (Figure 5A). Like OS, the 5-year CSS rate was 84.4% in the married group and 80.7% in unmarried group ($p < 0.001$) (Figure 5B). Despite the basically comparable variables across two groups, we furthermore performed univariate Cox regression to make a more accurate conclusion.

Table 1. Baseline characteristics of ccRCC patients according to marital status in the SEER database

Characteristic	Total (%)	Married (%)	Divorced (%)	Single (%)	Widowed (%)	<i>p</i>
	68599 (100)	44553(64.95)	7410(10.80)	10663(15.54)	5973(8.71)	
Sex						<0.001
Male	43271(63.1)	30817(69.2)	4063(54.8)	6755(63.3)	1636(27.4)	
Female	25328(36.9)	13736(30.8)	3347(45.2)	3908(36.7)	4337(72.6)	
Age						<0.001
18-49	12230(17.8)	7356(16.5)	1236(16.7)	3557(33.4)	81(1.4)	
50-59	16827(24.5)	11062(24.8)	2218(29.9)	3115(29.1)	432(7.2)	
60-69	22019(32.1)	15190(34.1)	2632(35.5)	2653(24.9)	1544(25.8)	
70-79	13322(19.4)	8774(19.7)	1113(15.1)	1085(10.2)	2350(39.3)	
≥80	4201(6.2)	2171(4.9)	211(2.8)	253(2.4)	1566(26.3)	
Race						<0.001
White	55394(80.8)	37131(83.3)	5842(78.8)	7515(70.5)	4906(82.1)	
Black	8030(11.7)	3820(8.6)	1168(15.8)	2350(22.0)	692(11.6)	
Others ¹	5175(7.5)	3602(8.1)	400(5.4)	798(7.5)	375(6.3)	
Diagnosis year						<0.001
1973-1993	15843(23.1)	10490(23.5)	1547(20.9)	1890(17.7)	1916(32.1)	
1994-2004	26157(38.1)	17037(38.3)	2894(39.1)	4079(38.3)	2147(35.9)	
2005-2015	26599(38.8)	17026(38.2)	2969(40.0)	4694(44.0)	1910(32.0)	
Pathological grade						<0.001
Grade I	8446(12.3)	5368(12.0)	893(12.1)	1320(12.4)	865(14.5)	
Grade II	34890(50.9)	22638(50.8)	3821(51.6)	5385(50.5)	3046(51.0)	
Grade III	20046(29.2)	13191(29.6)	2101(28.4)	3154(29.6)	1600(26.8)	
Grade IV	5217(7.6)	3356(7.6)	595(7.9)	804(7.5)	462(7.7)	
AJCC stage						<0.001
Stage I	44110(64.3)	28580(64.1)	4803(64.8)	6952(65.2)	3775(63.2)	
Stage II	6822(9.9)	4402(9.9)	727(9.8)	1129(10.6)	564(9.4)	
Stage III	11165(16.3)	7378(16.6)	1141(15.4)	1587(14.9)	1059(17.8)	
Stage IV	6502(9.5)	4193(9.4)	739(10.0)	995(9.3)	575(9.6)	
Tumor size						<0.001
≤5 cm	37587(54.8)	24464(54.9)	4056(54.7)	5836(54.7)	3231(54.1)	
5-10cm	23197(33.8)	15044(33.8)	2533(34.2)	3487(32.7)	2133(35.7)	
>10 cm	7815(11.4)	5045(11.3)	821(11.1)	1340(12.6)	609(10.2)	
Surgery						<0.001
Performed	66507(96.9)	43459(97.5)	7131(96.2)	10283(96.4)	5634(94.3)	
Not Performed	2092(3.1)	1094(2.5)	279(3.8)	380(3.6)	339(5.7)	
Median household income						<0.001
Quartile 1	19671(28.7)	12515(28.1)	2314(31.2)	2951(27.7)	1891(31.7)	
Quartile 2	15614(22.8)	9958(22.4)	1638(22.1)	2634(24.7)	1384(23.1)	
Quartile 3	16913(24.7)	11096(24.8)	1859(25.1)	2610(24.5)	1348(22.6)	
Quartile 4	16401(23.8)	10984(24.7)	1599(21.6)	2468(23.1)	1350(22.6)	
Insurance status						<0.001
Insured	49926(72.8)	32659(73.3)	5532(74.7)	7832(73.5)	3903(65.3)	
Uninsured	4836(7.1)	2779(6.2)	625(8.4)	976(9.1)	456(7.6)	
Unknown	13837(20.1)	9115(20.5)	1253(16.9)	1855(17.4)	1614(27.1)	

AJCC, the American Joint Committee on Cancer




Figure 1. Kaplan–Meier survival curves according to marital status (married, divorced, widowed, and single) in patients with renal clear cell carcinoma. **A.** Overall survival: $\chi^2=677, p < 0.001$; **B.** Cancer-specific survival: $\chi^2=147, p < 0.001$.

Using married patients as reference, unmarried individuals persisted to be associated with increased risks death for both OS (HR, 1.31, 95% CI 1.24-1.38, $p < 0.001$) and CSS (HR, 1.19, 95% CI 1.11-1.28, $p < 0.001$). These results proved that our analysis was credible and reliable, which meant that the confounding factors were not responsible for the error source.

Table 2. Univariate and multivariate survival analysis for evaluating the impact of marital status on the OS among ccRCC patients

Variables	5-year OS	Univariate analysis		Multivariate analysis		
		Log Rank χ^2	<i>p</i>	HR	95% CI	<i>p</i>
Marital status		677	<0.001			
Married	79.0%			Reference		
Divorced	73.8%			1.33	1.27-1.40	<0.001
Single	77.3%			1.26	1.20-1.32	<0.001
Widowed	66.4%			1.42	1.35-1.49	<0.001
Sex		95.8	<0.001			
Male	75.6%			Reference		
Female	79.4%			0.85	0.82-0.88	0.021
Age		3156	<0.001			
18-49	87.2%			Reference		
50-59	81.7%			1.32	1.24-1.41	<0.001
60-69	77.1%			1.71	1.61-1.82	<0.001
70-79	70.3%			2.55	2.39-2.72	<0.001
≥80	54.2%			4.01	3.73-4.32	<0.001
Race		33.1	<0.001			
White	76.8%			Reference		
Black	76.7%			1.14	1.08-1.19	<0.001
Others	79.9%			0.88	0.82-0.93	<0.001
Diagnosis year		173	<0.001			
1973-1993	73.6%			Reference		
1994-2004	78.5%			0.95	0.91-0.98	0.043
2005-2015	82.5%			0.87	0.82-0.92	<0.001
Pathological grade		6670	<0.001			
Grade I	85.1%			Reference		
Grade II	84.4%			0.97	0.93-1.04	0.477
Grade III	69.9%			1.35	1.27-1.44	<0.001
Grade IV	39.5%			2.26	2.11-2.42	<0.001
Tumor size		5612	<0.001			
≤5 cm	86.7%			Reference		

Variables	5-year OS	Univariate analysis		Multivariate analysis		
		Log Rank χ^2	<i>p</i>	HR	95% CI	<i>p</i>
5-10cm	70.4%			1.47	1.39-1.55	<0.001
>10 cm	50.6%			1.67	1.55-1.79	<0.001
AJCC stage		26079	<0.001			
Stage I	87.5%			Reference		
Stage II	80.9%			1.07	1.01-1.14	<0.001
Stage III	67.0%			1.79	1.71-1.88	<0.001
Stage IV	18.61%			6.62	6.29-6.98	<0.001
Surgery		8574	<0.001			
Performed	78.70%			Reference		
Not Performed	20.35%			0.30	0.28-0.32	<0.001
Median household income		116	<0.001			
Quartile 1	74.9%					
Quartile 2	76.3%			0.92	0.88-0.96	<0.001
Quartile 3	78.3%			0.85	0.81-0.88	<0.001
Quartile 4	78.8%			0.83	0.79-0.87	<0.001
Insurance status		72.5	<0.001			
Insured	78.1%			Reference		
Uninsured	75.4%			1.16	1.07-1.25	<0.001

AJCC, the American Joint Committee on Cancer; OS, overall survival; CI, confidence interval; HR, hazard ratio

Discussion

To our best knowledge, this study is the first and largest study to date investigating the impact of marital status on the survival of ccRCC patients. In this large, population-based study, we firstly used a systematic PSM and subgroup analysis to show that marital status was an independent prognostic factor and contributed to worse OS and CSS for unmarried ccRCC patients. Compared to married patients, unmarried individuals, including divorced, single, and widowed, faced higher mortality risks for OS and CSS irrespective of whether or not the patients were adjusted for age, sex, race, diagnosis year, histological type, pathological grade, AJCC stage, tumor size, surgery condition, median household income or

insurance status. Besides, among the unmarried, widowed patients had the best chance of dying from ccRCC even after adjustment the abovementioned variables. Moreover, in the subgroup analysis, the increased mortality risks were also significant among unmarried patients at each sex, surgery condition, and all cancer stages subgroups. Even in the 1:1 matched cohort with comparable baseline characteristics, unmarried status still contributed to worse survival for ccRCC patients.

Table 3. Univariate and multivariate survival analysis for evaluating the impact of marital status on the CSS among ccRCC patients

Variables	5-year CSS	Univariate analysis		Multivariate analysis		
		Log Rank χ^2	p	HR	95% CI	p
Marital status		147	<0.001			
Married	85.5%			Reference		
Divorced	83.3%			1.16	1.08-1.25	<0.001
Single	80.5%			1.09	1.02-1.16	0.006
Widowed	76.5%			1.20	1.13-1.28	<0.001
Sex		5.3	0.021			
Male	83.8%			Reference		
Female	86.6%			0.96	0.92-0.99	0.011
Age		636	<0.001			
18-49	90.0%			Reference		
50-59	86.5%			1.14	1.05-1.23	<0.001
60-69	84.3%			1.33	1.23-1.43	<0.001
70-79	82.3%			1.67	1.54-1.80	<0.001
≥80	74.8%			2.25	2.04-2.47	<0.001
Race		29.7	<0.001			
White	84.6%			Reference		
Black	86.0%			1.09	1.02-1.17	0.011
Others ¹	85.0%			0.94	0.87-1.01	0.102
Diagnosis year		21	<0.001			
1973-1993	82.9%			Reference		
1994-2004	85.5%			0.94	0.89-1.00	0.0431
2005-2015	87.7%			0.84	0.79-0.90	<0.001
Pathological grade		2085	<0.001			
Grade I	94.2%			Reference		
Grade II	92.4%			1.09	0.99-1.20	0.074
Grade III	77.4%			1.84	1.67-2.02	<0.001
Grade IV	45.0%			3.12	2.81-3.45	<0.001
Tumor size		613	<0.001			
≤5 cm	95.0%			Reference		
5-10cm	78.5%			1.81	1.70-1.93	<0.001
>10 cm	55.2%			2.25	2.09-2.42	<0.001
AJCC stage		5706	<0.001			
Stage I	95.5%			Reference		
Stage II	87.8%			1.59	1.45-1.74	<0.001
Stage III	75.6%			3.32	3.10-3.56	<0.001
Stage IV	21.19%			14.06	13.09-15.10	<0.001
Surgery		4883	<0.001			
Performed	86.5%			Reference		
Not Performed	28.1%			0.26	0.24-0.28	<0.001
Median household income		40.2	<0.001			
Quartile 1	83.9%			Reference		
Quartile 2	84.1%			0.96	0.91-1.02	0.158
Quartile 3	85.3%			0.88	0.83-0.93	<0.001
Quartile 4	85.7%			0.87	0.82-0.92	<0.001
Insurance status		142	<0.001			
Insured	85.2%			Reference		
Uninsured	83.2%			1.04	0.96-1.14	0.314

AJCC, the American Joint Committee on Cancer; CSS, cancer-specific survival; CI, confidence interval; HR, hazard ratio

Table 4. Univariate and multivariate analysis of marital status on the CSS among ccRCC patients according to age, surgery status, and cancer stages

Variables	5-year CSS	Univariate analysis		Multivariate analysis		
		Log rank χ^2	p	HR	95%CI	p
Sex						
Male		86.5	<0.001			
Married	84.5%			Reference		
Divorced	80.3%			1.12	1.04-1.21	<0.001
Single	83.9%			0.96	0.93-1.02	0.265
Widowed	78.3%			1.37	1.29-1.48	<0.001
Female		122	<0.001			
Married	87.8%			Reference		
Divorced	86.9%			1.26	1.16-1.37	<0.001
Single	88.1%			1.05	0.96-1.14	0.247
Widowed	81.6%			2.12	1.99-2.26	<0.001
Surgery						
Performed		35.6	<0.001			
Married	87.3%			Reference		
Divorced	85.5%			1.52	1.29-1.81	<0.001
Single	86.9%			1.49	0.85-2.65	0.165
Widowed	82.8%			1.63	1.46-1.84	<0.001
Not performed		17.5	<0.001			
Married	32.2%			Reference		
Divorced	26.8%			1.16	0.98-1.36	0.066
Single	32.3%			1.02	0.84-1.24	0.834
Widowed	24.2.2%			1.28	1.04-1.71	0.015
AJCC stage						
Stage I		145	<0.001			
Married	96.3%			Reference		
Divorced	95.8%			1.35	1.25-1.47	<0.001
Single	96.2%			1.09	1.02-1.19	<0.001
Widowed	92.9%			2.36	2.21-2.53	<0.001
Stage II		52.3	<0.001			
Married	88.9%			Reference		
Divorced	84.2%			1.51	1.29-1.77	<0.001
Single	89.6%			1.14	0.98-1.33	0.088
Widowed	79.6%			2.12	1.83-2.47	<0.001
Stage III		22.9	<0.001			
Married	77.0%			Reference		
Divorced	72.8%			1.27	1.14-1.42	<0.001
Single	75.2%			1.09	0.99-1.21	0.0784
Widowed	69.1%			1.73	1.57-1.91	<0.001
Stage IV		19.5	<0.001			
Married	22.5%			Reference		
Divorced	18.9%			1.11	1.02-1.22	0.021
Single	19.0%			1.14	1.05-1.23	0.002
Widowed	18.3%			1.26	1.14-1.39	<0.001

AJCC, the American Joint Committee on Cancer; CSS, cancer-specific survival; CI, confidence interval; HR, hazard ratio

More importantly, our study identified several additional associations. Like, ccRCC tended to occur in young adults, more male than female, and male ccRCC patients benefited more from marriage than females. A potential reason for this sex disparity is that male patients might receive more social supports and assistance from their relatives or friends¹⁵. Interestingly, studies have suggested that the socioeconomic status is associated with better prognosis in several cancers¹⁶⁻¹⁸. In our study, this trend was also observed, and the widowed group had the lowest percentage of insurance, which may partly lead to their survival disadvantages. Additionally, it has been pointed that delayed diagnosis could contribute to poor survival among unmarried patients

^{8,19}. However, in our study, incidence of early stage (stage I/II) cancer was higher in the single group (65.2%), divorced (64.8%), and widowed (63.2%) groups, compared with the married (64.1%). Though, the SEER database does not provide information regarding cancer screening results, earlier diagnosis for unmarried people may indicate higher odds for cancer screening. Obviously, for ccRCC patients, delayed diagnosis alone couldn't explain the worse survival.

Our results indicated that unmarried patients experienced a significant survival disadvantage over their unmarried counterparts. Despite unclear mechanisms intrinsic mechanisms behind this

association, we try to analyze several possible reasons. First of all, people in good marital condition may be encouraged by their spouse to regular physical check-up. Moreover, married patients usually has better tolerance and adherence to prescribed treatments due to the medication reminders and assistance for their spouses²⁰, which is crucial to improve the curability rate and survival²¹⁻²³. Secondly, married patients usually possess stronger financial resources such as higher income level, better employment, medical insurance, which ultimately affect the access to early prevention, timely diagnosis and treatment²⁴.




Figure 2. Kaplan–Meier survival curves of cancer-specific survival in patients with renal clear cell carcinoma stratified by sex. **A.** male: $\chi^2=90.9$, $p < 0.001$; **B.** female: $\chi^2= 148$, $p < 0.001$.




Figure 3. Kaplan–Meier survival curves of cancer-specific survival in patients with renal clear cell carcinoma stratified by surgery. **A.** surgery performed: $\chi^2=95.4$, $p < 0.001$; **B.** surgery not performed: $\chi^2= 8.5$, $p = 0.040$.




Figure 4. Kaplan–Meier survival curves of cancer-specific survival in patients with renal clear cell carcinoma stratified by tumor stage at diagnosis. **A.** stage I: $\chi^2=186$, $p < 0.001$; **B.** stage II: $\chi^2=57.8$, $p < 0.001$; **C.** stage III: $\chi^2=27.2$, $p < 0.001$; **D.** stage IV: $\chi^2= 19.8$, $p < 0.001$.




Figure 5. Kaplan–Meier survival curves according to marital status (married, unmarried) in patients with renal clear cell carcinoma. **A.** Overall survival: $\chi^2=121.4$, $p < 0.001$; **B.** Cancer-specific survival: $\chi^2= 66.42$, $p < 0.001$.

Table 5. Patient baseline characteristics before and after PSM

Characteristic	Before matching			<i>p</i> *	After matching			<i>p</i> *
	Married (%)	Unmarried (%)	SD		Married (%)	Unmarried (%)	SD	
Sample size	44553(100)	24046(100)			9014(100)	9014(100)		
Sex				<0.001				0.004
Male	30817(69.17)	12454(51.79)	NA		4805(53.3)	4611(52.2)	NA	
Female	13736(30.83)	11592(48.21)	0.46		4209(46.7)	4403(48.8)	0.02	
Age				<0.001				<0.001
18-49	7356(16.5)	4874(20.3)	NA		88(1.0)	81(0.9)	NA	
50-59	11062(24.8)	5765(24.0)	0.43		2336(25.9)	2463(27.3)	0.01	
60-69	15190(34.1)	6829(28.4)	0.47		4157(46.1)	3847(42.7)	-0.03	
70-79	8774(19.7)	4548(18.9)	0.40		1102(12.2)	1085(12.0)	0.00	
≥80	2171(4.9)	2030(8.4)	0.22		1331(14.8)	1538(17.1)	0.02	
Race				<0.001				0.561
White	37131(83.3)	18263(76.0)	NA		6673(74.0)	6704(74.4)	NA	
Black	3820(8.6)	4210(17.5)	0.28		1733(19.2)	1682(18.7)	-0.01	
Others	3602(8.1)	1573(6.5)	0.27		608(6.7)	628(7.0)	0.00	
Diagnosis year				<0.001				0.01
1973-1993	10490(23.5)	5353(22.3)	NA		10490(23.5)	5353(22.3)	NA	0.964
1994-2004	17037(38.2)	9120(37.9)	0.49		17037(38.2)	9120(37.9)	0.00	
2005-2015	17026(38.2)	9573(39.8)	0.49		17026(38.2)	9573(39.8)	0.00	
Pathological grade				0.002				-0.01
Grade I	5368(12.0)	3078(12.8)	NA		1086(12.0)	1093(12.1)	NA	0.267
Grade II	22638(50.8)	12252(51.0)	0.50		4568(50.7)	4509(50.0)	-0.01	
Grade III	13191(29.6)	6855(28.5)	0.46		2753(30.5)	2738(30.4)	0.00	
Grade IV	3356(7.5)	1861(7.7)	0.26		607(6.7)	674(7.5)	0.01	
AJCC stage								0.495
Stage I	28580(64.1)	15530(64.6)	0.33	0.048	5769(64.0)	5680(63.0)	0.00	
Stage II	4402(9.9)	2420(10.1)	0.30		883(9.8)	886(9.8)	0.00	
Stage III	7378(16.6)	3787(15.7)	0.37		1521(16.9)	1563(17.3)	0.00	
Stage IV	4193(9.4)	2309(9.6)	0.29		841(9.3)	885(9.8)	0.00	
Tumor size				0.626				0.01
≤10 cm	24464(54.9)	13123(54.6)	NA		4890(54.2)	4783(53.1)	NA	0.196
10-20cm	15044(33.8)	8153(33.9)	0.47		3165(35.1)	3214(35.7)	0.01	
>20 cm	5045(11.3)	2770(11.5)	0.32		959(10.6)	1017(11.3)	0.01	
Surgery				<0.001				0.601
Performed	43459(97.54)	23048(95.85)	0.18		8589(95.1)	8574(95.1)	0.00	
Not Performed	1094(2.46)	998(4.15)	0.15		425(4.9)	4404.9)	0.00	
Median household income				<0.001				0.723
Quartile 1	12515(28.1)	7156(29.8)	NA		2984(33.1)	2960(32.8)	NA	
Quartile 2	9958(22.4)	5656(23.5)	0.42		2370(26.3)	2405(26.7)	0.00	
Quartile 3	11096(24.9)	5817(24.2)	0.43		1905(21.1)	1943(21.6)	-0.01	
Quartile 4	10984(24.7)	5417(22.5)	0.43		1755(19.5)	1706(18.9)	-0.01	
Insurance status				<0.001				0.638
Insured	32659(73.3)	17267(71.8)	NA		6374(70.7)	6430(71.3)	NA	
Uninsured	2779(6.2)	2057(8.6)	0.24		887(9.8)	876(9.7)	0.00	
Unknown	9115(20.5)	4722(19.6)	0.40		1753(19.4)	1708(18.9)	0.00	

AJCC, the American Joint Committee on Cancer; SD, standardized difference; NA, not applicable. **p* value for χ^2 test.

Thirdly, emotional support from spouse and other family members also contributed to better prognosis²⁵ compared to unmarried counterparts, unmarried patients, especially the widowed, have to face with higher psychological stress, economic pressures, lack of family support, absence of public assistance policy, leading to psychological imbalance, feel distressful and depressed. Multiple studies have confirmed that depression and stress were strongly associated with carcinoma growth and metastasis²⁶⁻²⁸. In addition, excess stress and depression would dysregulate the immune and endocrine function, induce chronic inflammation and thus result in worse survival²⁹⁻³¹.

In our study, approximately 73% widowed patients were female. Miller et.al have reported that natural killer cells decreased significantly in women whose husbands have died recently²⁸. And

significantly higher proportion of psychological disturbance have been found in widowed patients. It is rather intuitive that lack of social support could seriously damage the function of immune cells, leading immune escape in tumor cells. On the other hand, it has been proposed that widowed patients less tended to receive surgery compared with the married. And undertreatment could have been one explanation for poor survival in widowed patients²². This correlation between marital status and receipt of surgery was independently validated in ccRCC patients in our study. Spouses of these married individuals might encourage them to receive surgery rather than conservative treatment, which could in part account for survival discrepancies.

Our study used a large sample size and sophisticated statistical analysis to investigate in depth the impact of marital status on survival for

ccRCC patients, to ensure reliability. However, some potential limitations cannot be ignored. Firstly, we just simply divided patients into the married and unmarried groups but failed to get more details about their marriage. For example, SEER database do not provide information about marital history, family problems, lifestyle factors, which may serve as potential confounding factors. Secondly, the marital status was recorded only when diagnosed and we might put some patients into the wrong group when their marital status had changed during the follow-up. In addition, for unrecorded marital status (gay, lesbian, bisexual and transgender), we can't rule out that some have regular partner and get social support secretly. Thirdly, as far as the whole population concerned, a predominance of white people was observed in our study. Several studies have showed that racial disparities could lead to various prognoses among ccRCC patients³²⁻³⁴. One final but important point, SEER database does not provide several important clinical information regarding the treatment plan, co-morbid diseases and prognostic biomarkers such as VHL, HIF- α and ALDH2, which had proven to have an effect on RCC patient prognosis³⁵.

Conclusion

Marital status was an independent prognostic factor of survival for ccRCC patients. Unmarried patients faced higher mortality risks for overall and cancer-specific survival, and among these patients, the widowed suffered the highest mortality risks. Unfavorable socioeconomic and psychological status might be responsible for the inferior survival of unmarried patients.

Acknowledgements

The present study was supported by grants from National Natural Science Foundation of China (NO. 81872245 and 81803601).

Competing Interests

The authors have declared that no competing interest exists.

References

- Znaor A, Lortet-Tieulent J, Laversanne M, et al. International variations and trends in renal cell carcinoma incidence and mortality. *Eur Urol*. 2015;67:519-30.
- Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin*. 2017;67:7-30.
- Engel GL. The need for a new medical model: a challenge for biomedicine. *Psychodyn Psychiatry*. 2012;40:377-96.
- Adler RH. Engel's biopsychosocial model is still relevant today. *J Psychosom Res*. 2009;67:607-11.
- Aizer AA, Chen MH, McCarthy EP, et al. Marital status and survival in patients with cancer. *J Clin Oncol*. 2013;31:3869-76.
- Zhang SL, Wang WR, Liu ZJ, et al. Marital status and survival in patients with soft tissue sarcoma: A population-based, propensity-matched study. *Cancer Med*. 2019;8:465-79.
- Wu C, Chen P, Qian JJ, et al. Effect of marital status on the survival of patients with hepatocellular carcinoma treated with surgical resection: an analysis of 13,408 patients in the surveillance, epidemiology, and end results (SEER) database. *Oncotarget*. 2016;7:79442-52.
- Liu Q, Luo D, Cai S, et al. Real-World Implications of Nonbiological Factors with Staging, Prognosis and Clinical Management in Colon Cancer. *Cancers*. 2018;10:263.
- Li Y, Zhu MX, Qi SH. Marital status and survival in patients with renal cell carcinoma. *Medicine*. 2018;97:e0385.
- Miao T, Li Y, Sheng X, et al. Marital status and survival of patients with kidney cancer. *Oncotarget*. 2017;8:86157-67.
- Wang H, Wang L, Kabirov I, et al. Impact of marital status on renal cancer patient survival. *Oncotarget*. 2017;8:70204-13.
- Warren JL, Klabunde CN, Schrag D, et al. Overview of the SEER-Medicare data: content, research applications, and generalizability to the United States elderly population. *Med Care*. 2002;40:lv-3-18.
- Austin PC. The use of propensity score methods with survival or time-to-event outcomes: reporting measures of effect similar to those used in randomized experiments. *Stat Med*. 2014;33:1242-58.
- Austin PC. Balance diagnostics for comparing the distribution of baseline covariates between treatment groups in propensity-score matched samples. *Stat Med*. 2009;28:3083-107.
- Merluzzi J, Dobrev SD. Unequal on top: Gender profiling and the income gap among high earner male and female professionals. *Soc Sci Res*. 2015;53:45-58.
- Mahal A, Mahal B, Nguyen P, et al. Prostate cancer outcomes for men aged younger than 65 years with Medicaid versus private insurance. *Cancer*. 2018;124:752-9.
- Rong X, Yang W, Garzon-Muvdi T, et al. Influence of insurance status on survival of adults with glioblastoma multiforme: A population-based study. *Cancer*. 2016;122:3157-65.
- Ellis L, Canchola A, Spiegel D, et al. Trends in Cancer Survival by Health Insurance Status in California From 1997 to 2014. *JAMA oncology*. 2018;4:317-23.
- Lin D, Gold HT, Schreiber D, et al. Impact of socioeconomic status on survival for patients with anal cancer. *Cancer*. 2018;124:1791-7.
- Haley WE. Family caregivers of elderly patients with cancer: understanding and minimizing the burden of care. *J Support Oncol*. 2003;1:25-9.
- Tammemagi CM, Neslund-Dudas C, Simoff M, et al. Lung carcinoma symptoms--an independent predictor of survival and an important mediator of African-American disparity in survival. *Cancer*. 2004;101:1655-63.
- Ferrante JM, Gonzalez EC, Roetzheim RG, et al. Clinical and demographic predictors of late-stage cervical cancer. *Arch Fam Med*. 2000;9:439-45.
- Nayeri K, Pitaro G, Feldman JG. Marital status and stage at diagnosis in cancer. *N Y State J Med*. 1992;92:8-11.
- Moreno-Smith M, Lutgendorf SK, Sood AK. Impact of stress on cancer metastasis. *Future Oncol*. 2010;6:1863-81.
- Wittmann D. Emotional and sexual health in cancer: partner and relationship issues. *Curr Opin Support Palliat Care*. 2016;10:75-80.
- Janssen LME, Ramsay EE, Logsdon CD, et al. The immune system in cancer metastasis: friend or foe? *J Immunother Cancer*. 2017;5:79.
- Sullivan DR, Forsberg CW, Ganzini L, et al. Depression symptom trends and health domains among lung cancer patients in the CanCORS study. *Lung Cancer*. 2016;100:102-9.
- Miller AH, Raison CL. The role of inflammation in depression: from evolutionary imperative to modern treatment target. *Nat Rev Immunol*. 2016;16:22-34.
- Uchino BN. Social support and health: a review of physiological processes potentially underlying links to disease outcomes. *J Behav Med*. 2006;29:377-87.
- Sapolsky RM, Donnelly TM. Vulnerability to stress-induced tumor growth increases with age in rats: role of glucocorticoids. *Endocrinology*. 1985;117:662-6.
- Miller GE, Cohen S, Ritchey AK. Chronic psychological stress and the regulation of pro-inflammatory cytokines: a glucocorticoid-resistance model. *Health Psychol*. 2002;21:531-41.
- Patel HD, Kates M, Pierorazio PM, et al. Race and sex disparities in the treatment of older patients with T1a renal cell carcinoma: a comorbidity-controlled competing-risks model. *Urol Oncol*. 2014;32:576-83.
- Mafolasire A, Yao X, Nawaf C, et al. Racial disparities in renal cell carcinoma: a single-payer healthcare experience. *Cancer Med*. 2016;5:2101-8.

34. Kates M, Whalen MJ, Badalato GM, et al. The effect of race and gender on the surgical management of the small renal mass. *Urol Oncol.* 2013;31:1794-9.
35. Gao Y, Wu Z, Xie L, et al. VHL deficiency augments anthracycline sensitivity of clear cell renal cell carcinomas by down-regulating ALDH2. *Nat Commun.* 2017;8:15337.