


Validation of Risk Stratification Criteria to Identify Febrile Neonates at Low Risk of Serious Bacterial Infection

Global Pediatric Health
Volume 6: 1–5
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2333794X19845076
journals.sagepub.com/home/gph


Joni E. Rabiner, MD¹ , Maya Capua, MD², Dina Golfeiz, MD³,
Jamie Shoag, MD⁴, and Jeffrey R. Avner, MD⁵

Abstract

Our objective was to evaluate the accuracy of risk stratification criteria for febrile neonates in the emergency department. This was a retrospective study of febrile neonates ≤ 56 days of age. Patients were low risk for serious bacterial infection (SBI) if all test results were within normal ranges. Three hundred thirty-eight patients were enrolled with a mean age of 32 (± 14) days, and 78 (23%) had SBI: 26 (8%) with bacteremia, 48 (14%) with urinary tract infection, 3 (1%) with meningitis, and 11 (3%) with pneumonia. Risk stratification criteria identified 47 (14%) as low risk, 2 of whom had SBI (both with Group B *Streptococcus* bacteremia). The sensitivity was 97.4% (95% confidence interval = 91.0% to 99.7%), and the negative predictive value was 95.7% (95% confidence interval = 84.8% to 98.9%). The risk stratification criteria have high sensitivity and high negative predictive value for identifying infants at low risk for SBI. Care must be taken to assure reliable follow-up.

Keywords

Febrile neonate, serious bacterial infection, risk stratification, low risk criteria

Received October 23, 2018. Received revised March 28, 2019. Accepted for publication March 29, 2019.

Introduction

With an estimated 1.4 million neonatal deaths each year attributed to invasive infections,¹ and up to 10% of infants less than 2 months of age with fever having serious bacterial infection (SBI),² diagnosis and rapid treatment of SBI in febrile infants is critical in the emergency department (ED). Over the past 2 decades, the microbial agents causing SBI in febrile neonates have been changing, with decreasing rates of Group B *Streptococcus* (GBS) and *Listeria monocytogenes* and increasing rates of *Escherichia coli*.^{3–5}

Most evidence-based protocols recommend an evaluation for sepsis including examination of blood, urine, and cerebrospinal fluid (CSF) for all febrile infants 28 days of age or less and for all ill-appearing febrile infants between 29 and 56 days of age to evaluate for SBI, as well as administration of empiric antibiotics. For well-appearing febrile infants between 1 and 2 months of age without a source of infection on physical examination, criteria to identify infants at low risk for SBI have been developed.^{6–8} Utilizing the white blood cell (WBC)

count, band count, urinalysis, CSF cell counts, chest X-ray, and stool smear, these criteria have been found to have good sensitivity (92% to 98%) and negative predictive value (NPV; 99% to 100%) in identifying low-risk infants who can be safely discharged from the ED without antibiotic treatment.^{6,8,9} Most hospitals have established protocols for infants 1 to 2 months of age based on these criteria,¹⁰ and there does not appear to be significant differences in diagnosis or clinical outcomes associated with variations in hospital protocols.^{11,12}

¹New York Presbyterian- Morgan Stanley Children's Hospital / Columbia University Medical Center, New York, NY, USA

²Steven and Alexandra Cohen Children's Medical Center / Northwell Health, New Hyde Park, NY, USA

³Children's Hospital at Montefiore / Albert Einstein College of Medicine, Bronx, NY, USA

⁴Jackson Memorial Hospital / University of Miami, Miami, FL, USA

⁵Maimonides Medical Center, Brooklyn, NY, USA

Corresponding Author:

Joni E. Rabiner, Morgan Stanley Children's Hospital of New York, 3959 Broadway, CHN-1-116, New York, NY 10032, USA.

Email: jer2212@cumc.columbia.edu



However, the epidemiology of SBI in infants is changing, and different organisms elicit different inflammatory responses,¹³ which may change the usefulness of some of the elements of the stratification criteria such as the WBC count.

The purpose of our study was to determine if the risk stratification protocol used at our institution to identify SBI in febrile infants ≤ 56 days of age, based on previous evidence-based guidelines, remains accurate for identifying all cases of SBI in febrile neonates in the setting of the changing epidemiology of neonatal SBI.

Materials and Methods

This was a retrospective study of febrile neonates 56 days of age or younger who presented to 1 of 3 EDs (1 pediatric and 2 general EDs) of a larger medical center system from January 2012 to December 2014. Patients were included in the study if they had a temperature $\geq 38^{\circ}\text{C}$ rectal in the ED or documented within the 24 hours prior to arrival. The medical records of identified patients were reviewed and data were extracted from the electronic medical record utilized in all 3 EDs.

Each patient's documented ED visit was reviewed for history, physical examination, laboratory, and radiological data. Patient disposition (admit to hospital or discharge home) and return to the ED within 3 days of the initial visit was recorded. The final diagnosis for the patient based on ED and/or hospital discharge diagnoses and laboratory results was documented.

Risk Stratification Protocol

The protocol at our institution was for all infants ≤ 28 days and ill-appearing infants 29 to 56 days of age to have blood, urine, and CSF studies and to be admitted for empiric antibiotic treatment. For well-appearing infants between 29 and 56 days of age, we used a stratification management strategy to identify those at low risk for SBI, and it included assessment with complete blood count with manual differential, blood culture, urinalysis, urine culture, CSF cell counts and culture (if indicated), glucose levels, liver function tests, chest X-ray for respiratory symptoms, and stool smear and culture for diarrhea. A patient 29 to 56 days of age with a reliable caregiver and follow-up available could be managed as an outpatient if he/she met the following low-risk criteria: WBC count between 5000 and 15 000/ μL , manual differential of immature: total neutrophil ratio of <0.2 , urinalysis with <10 WBC/high-power field, normal glucose and liver function tests, normal CSF (if obtained), negative chest X-ray (if obtained), and negative fecal leukocytes (if obtained).

Study Definitions

Serious bacterial infection was defined as bacterial growth of a known pathogen with >1000 colony-forming units/mL in blood, urine, CSF, or stool cultures. Contaminated cultures were defined as cultures with bacteria isolated that are not pathogens; a pediatric infectious disease specialist determined whether questionable isolated organisms were true pathogens or contaminants.

In order to evaluate our screening guidelines, patients were considered not low risk or low risk for SBI. Patients were considered not low risk for SBI if they had any laboratory value outside the defined normal ranges, evidence of bacterial infection on physical examination, grossly bloody CSF that was uninterpretable, or incomplete workup at the time of the ED visit. All other patients who had all normal laboratory values were considered low risk for SBI.

Data Analysis

Data were analyzed with SPSS Statistics (IBM, Armonk, NY). Descriptive statistical analyses, including mean and standard deviation (SD), were used for continuous data, and frequencies were used for categorical data. Test performance characteristics, including sensitivity and negative predictive value, were calculated. Two-sample *t* tests were used to compare means for continuous data between 2 groups with a significance of 0.05.

Ethical Approval and Informed Consent

This study was approved by the Institutional Review Board of Montefiore Medical Center (Reference # 002539).

Results

The charts of 338 patients who presented to the ED with fever were analyzed. Two hundred sixty-three (78%) patients were evaluated in the pediatric ED, and 75 (22%) were evaluated in the 2 general EDs. Demographic information and disposition of the study population are included in Table 1.

Seventy-eight patients (23%) were found to have SBI, including 39 with urinary tract infection, 17 with bacteremia, 11 with pneumonia, 9 with urosepsis, and 3 with bacterial meningitis (one patient had *E coli* bacteremia and *E coli* meningitis). Of the 77 positive cultures, *E coli* bacteriuria was the predominate cause of infection, with 39 (51%) cases, followed by GBS bacteremia

Table 1. Demographic Information and Disposition, N = 338.

	Overall (N = 338)	PED (n = 263)	GED (n = 75)
Age, days, mean (SD)	32.3 (13.9)	32.2 (14.1)	32.7 (13.1)
Age ≤28 days, n (%)	143 (42%)	113 (43%)	30 (40%)
Male, n (%)	196 (58%)	152 (58%)	44 (59%)
Preterm, n (%)	30 (9%)	23 (9%)	7 (9%)
Duration of fever, hours, mean (SD)	21.0 (20.4)	21.3 (21.0)	19.8 (18.0)
Ill appearance, n (%)	38 (11%)	37 (14%)	1 (1%)
Admit, n (%)	298 (88%)	229 (87%)	69 (92%)
SBI, n (%)	78 (23%)	57 (22%)	21 (28%)

Abbreviations: PED, pediatric emergency department; GED, general emergency department; SD, standard deviation; SBI, serious bacterial infection.

Table 2. SBI by Source of Infection and Microbiology.

Pathogen	Blood (n = 336)	Urine (n = 324)	CSF (n = 233)
<i>Escherichia coli</i>	8	39	1
GBS	10	3	1
<i>Bacillus cereus</i>	4	0	0
<i>Rothia dentocariosa</i>	1	0	0
<i>Enterococcus gallinarum</i>	1	0	0
<i>Flavobacterium meningosepticum</i>	1	0	0
<i>Enterococcus faecalis</i>	1	5	0
<i>Klebsiella pneumonia</i>	0	3	0
<i>Proteus mirabilis</i>	0	1	0
<i>Staphylococcus aureus</i> (MSSA)	0	0	1
Total	26	51 ^a	3

Abbreviations: SBI, serious bacterial infection; CSF, cerebrospinal fluid; GBS, Group B *Streptococcus*; MSSA, methicillin-sensitive *Staphylococcus aureus*.

^aThree urine cultures were positive for >1 pathogen.

with 9 (12%) cases and *E coli* bacteremia with 8 (10%) cases (Table 2).

The risk stratification results versus SBI and test performance characteristics for all 338 patients are shown in Table 3. Overall, of the 78 cases of SBI in the study population, 76 were correctly identified as not low risk, with a sensitivity of 97.4% (95% confidence interval = 91.0% to 99.7%). Of the 47 infants identified as low risk, 45 did not have SBI, with a NPV of 95.7% (95% confidence interval = 84.8% to 98.9%). All 30 febrile neonates 28 days of age or less with SBI were correctly identified as not low risk.

There were 2 infants identified as low risk that were found to have SBI, and both patients had blood cultures positive for GBS. The first patient was a 47-day-old ex-40-week female with fever for 6 hours prior to arrival, while the second patient was a 50-day-old ex-36-week female with fever for 7 hours prior to arrival, and both infants were recalled, admitted, and treated after blood

cultures became positive. Both patients had no progression of symptoms on reevaluation and an uneventful recovery.

One hundred fourteen patients had incomplete evaluations in the ED. Ninety-two (81%) did not have a manual differential performed and therefore could not have an immature–total neutrophil ratio calculated. Eleven (10%) were 28 days or younger and did not have a lumbar puncture performed. Six (5%) did not have a manual differential and were 28 days or younger and did not have a lumbar puncture. Five (4%) did not have a urinalysis obtained.

The mean WBC count overall was 11.9 (SD = 5.1). The mean WBC count for patients with cultures positive for *E coli* was 14.7 (SD = 5.8), and the mean WBC count for patients with cultures positive for GBS was 11.6 (SD = 5.3). There was no significant difference between the mean WBC count of patients with *E coli* and GBS infection ($P = NS$).

Table 3. Risk Stratification Results Versus SBI and Test Performance Characteristics With 95% Confidence Intervals^a.

	SBI+	SBI-	Total
Not low risk (positive or incomplete screening)	76	215	291
Low risk (negative screening)	2	45	47
Total	78	260	338

Abbreviation: SBI, serious bacterial infection.

^aSensitivity 97.4% (91.0% to 99.7%); specificity 17.3% (12.9% to 22.5%); positive predictive value 26.1% (24.9% to 27.4%); and negative predictive value 95.7% (84.8% to 98.9%).

Discussion

It is critical for emergency clinicians not to miss the diagnosis of SBI in neonates due to the significant potential morbidity and mortality associated with untreated SBI. Up to 10% of febrile infants less than 2 months of age will have SBI, with about 3% having bacteremia or meningitis.² For febrile infants less than 1 month of age, there may be a higher incidence of SBI, with up to 20% of infants 28 days of age or younger having infection.¹¹ In our study, the incidence of SBI was 23% overall, with 30/143 (21%) in infants 28 days of age or younger and 48/195 (25%) in infants 29 to 56 days of age.

Febrile neonates may not be managed according to established standards of care or institutional clinical practice guidelines. In one study, one third of pediatric EDs did not have clinical practice guidelines for the evaluation and management of febrile infants less than 2 months of age.¹⁰ For infants \leq 28 days with fever, recommended management for complete evaluation with blood, urine, and CSF studies is not followed in 28% to 33%.^{11,14} In our study, 114 (34%) febrile infants had incomplete evaluations according to our protocol, due to not having a manual differential performed, a lumbar puncture not performed on a patient under 1 month of age, or a urinalysis not obtained. In the ideal setting, all tests would be obtained; however, in practice, it may not be possible to have every test completed on every patient. At the time of this study, our institution's laboratory did not routinely perform manual differentials on all patients 56 days of age or younger. In addition, in some cases, not enough urine was obtained from a catheterized specimen for a complete urinalysis, and only a culture or urine dipstick was obtained. In this study, we considered patients with incomplete evaluations to be not low risk for SBI.

Based on the low-risk criteria strategies and general consensus for identifying SBI in febrile neonates, for well-appearing febrile infants, a WBC count $<$ 15 000 is required to be considered low risk.¹⁵ However, evidence suggests that WBC is not sufficiently accurate for identification of SBI in children, with low sensitivity for

diagnosis of neonatal sepsis¹⁶ and for SBI.¹⁷ Moreover, different pathogens may cause varying immune and inflammatory responses.¹³ As such, some pathogens may not cause an elevated WBC count or may not elevate the WBC count to the same extent as others, leading to a potentially missed SBI diagnosis based on current screening guidelines. Our study showed a higher WBC for *E coli* infections than for GBS infections that was not statistically significant. However, our 2 missed cases of SBI were both GBS bacteremia. As there were relatively low numbers of GBS and *E coli* infections in our study, more research is needed to investigate this question.

Additionally, the bacteriology of microbial agents causing neonatal SBI in febrile neonates has been changing over the past 2 decades, with decreasing rates of GBS and *Listeria* and increasing rates of *E coli* and other gram-negative infections.^{3-5,16} In accordance with current literature, our study demonstrated that *E coli* is indeed the most common pathogen causing SBI in this age group, causing 47 cases of urinary tract infection, bacteremia, and meningitis. GBS was the second most common pathogen, with 13 positive cultures in blood, urine, and CSF. Both infants with SBI not identified as low risk by our protocol had GBS bacteremia.

The 2 missed cases of GBS bacteremia in our study were infants 47 and 50 days old with fever for 6 to 7 hours prior to arrival to the ED. These were well-appearing infants who had complete workups done in the ED and were discharged home as they were categorized as low risk. Both were recalled, reevaluated promptly without further progression of disease, and treated without further sequelae.

Our sensitivity of 97.4% and NPV of 95.7% are similar to previous studies, with other sources citing sensitivity of 92% to 98% and NPV of 99% to 100%.^{6,8} We believe that our current protocol, established with current evidence, remains accurate to identify neonates at low risk for SBI.

Our study had several limitations. First, this was a retrospective chart review, limited by the data that were recorded in the medical record. However, using objective data from laboratory results minimizes this limitation, allowing for an objective diagnosis of SBI. Another

potential limitation of our study was the relatively small sample size, with a total of 338 patient charts reviewed. However, we included data over 3 years taken from 3 EDs with similar populations of patients, and our incidence of SBI and the sensitivity and NPV of febrile infants at low risk for SBI is similar to that documented in the literature, indicating that our patient population is likely a representative sample.

Conclusions

Escherichia coli was the most common pathogen causing SBI in infants up to 56 days of age, followed by GBS. The current risk stratification criteria used by our institution remain useful as our criteria have both high sensitivity and NPV for identifying infants 29 to 56 days of age at low risk for SBI. Our risk stratification criteria correctly identified all infants less than 28 days of age with SBI; however, 2 infants greater than 28 days of age with SBI were missed, although they both received treatment and recovered uneventfully. Therefore, while the current risk stratification criteria serve a useful tool in the management of febrile infants presenting to the ED, care must be taken to assure reliable follow-up.

Author Contributions

Joni Rabiner contributed to conception and design, data analysis and interpretation, and drafted and critically revised the manuscript.

Maya Capua contributed to conception and design, contributed to acquisition, analysis, and interpretation of data, and drafted the manuscript.

Dina Golfeiz contributed to acquisition of data, analysis, and interpretation of data, and drafted the manuscript.

Jamie Shoag contributed to acquisition of data.

Jeffrey Avner contributed to conception and design, data analysis and interpretation, and critically revised the manuscript.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Joni E. Rabiner  <https://orcid.org/0000-0003-4505-1947>

References

- Shane AL, Stoll BJ. Neonatal sepsis: progress towards improved outcomes. *J Infect*. 2014;68(suppl 1):S24-S32.
- Avner JR. Acute fever. *Pediatr Rev*. 2009;30:5-13.
- Stoll BJ, Hansen NI, Sanchez PJ, et al; Eunice Kennedy Shriver National Institute of Child Health and Human Development Neonatal Research Network. Early onset neonatal sepsis: the burden of group B streptococcal and *E coli* disease continues. *Pediatrics*. 2011;127:817-826.
- Shane AL, Stoll BJ. Recent developments and current issues in the epidemiology, diagnosis, and management of bacterial and fungal neonatal sepsis. *Am J Perinatol*. 2013;30:131-141.
- Greenhow TL, Hung YY, Herz AM, Losada E, Pantell RH. The changing epidemiology of serious bacterial infections in young infants. *Pediatr Infect Dis J*. 2014;33:595-599.
- Baker MD, Bell LM, Avner JR. Outpatient management without antibiotics of fever in selected infants. *N Engl J Med*. 1993;329:1437-1441.
- Baskin MN, O'Rourke EJ, Fleisher GR. Outpatient treatment of febrile infants 28 to 89 days of age with intramuscular administration of ceftriaxone. *J Pediatr*. 1992;120:22-27.
- Jaskiewicz JA, McCarthy CA, Richardson AC, et al. Febrile infants at low risk for serious bacterial infection—an appraisal of the Rochester criteria and implications for management. Febrile Infant Collaborative Study Group. *Pediatrics*. 1994;94:390-396.
- Irwin AD, Wickenden J, Le Doare K, Ladhani S, Sharland M. Supporting decisions to increase the safe discharge of children with febrile illness from the emergency department: a systematic review and meta-analysis. *Arch Dis Child*. 2016;101:259-266.
- Aronson PL, Thurm C, Williams DJ, et al; Febrile Young Infant Research Collaborative. Association of clinical practice guidelines with emergency department management of febrile infants ≤56 days of age. *J Hosp Med*. 2015;10:358-365.
- Aronson PL, Thurm C, Alpern ER, et al; Febrile Young Infant Research Collaborative. Variation in care of the febrile young infant <90 days in US pediatric emergency departments. *Pediatrics*. 2014;134:667-677.
- Chua KP, Neuman MI, McWilliams JM, Aronson PL; Febrile Young Infant Research Collaborative. Association between clinical outcomes and hospital guidelines for cerebrospinal fluid testing in febrile infants aged 29-56 days. *J Pediatr*. 2015;167:1340-1346.
- Abe R, Oda S, Sadahiro T, et al. Gram-negative bacteremia induces greater magnitude of inflammatory response than gram-positive bacteremia. *Crit Care*. 2010;14:R27.
- Jain S, Cheng J, Alpern ER, et al. Management of febrile neonates in US pediatric emergency departments. *Pediatrics*. 2014;133:187-195.
- Meltzer AJ, Powell KR, Avner JR. Fever in infants and children. *Consensus Pediatrics*. 2005;1:1-19.
- Shah BA, Padbury JF. Neonatal sepsis: an old problem with new insights. *Virulence*. 2014;5:170-178.
- De S, Williams GJ, Hayen A, et al. Value of white cell count in predicting serious bacterial infection in febrile children under 5 years of age. *Arch Dis Child*. 2014;99:493-499.