

The potential of P2X7 receptors as a therapeutic target, including inflammation and tumour progression

Geoffrey Burnstock^{1,2,3}  · Gillian E. Knight¹

Received: 6 September 2017 / Accepted: 1 November 2017 / Published online: 21 November 2017
© The Author(s) 2017. This article is an open access publication

Abstract Seven P2X ion channel nucleotide receptor subtypes have been cloned and characterised. P2X7 receptors (P2X7R) are unusual in that there are extra amino acids in the intracellular C terminus. Low concentrations of ATP open cation channels sometimes leading to cell proliferation, whereas high concentrations of ATP open large pores that release inflammatory cytokines and can lead to apoptotic cell death. Since many diseases involve inflammation and immune responses, and the P2X7R regulates inflammation, there has been recent interest in the pathophysiological roles of P2X7R and the potential of P2X7R antagonists to treat a variety of diseases. These include neurodegenerative diseases, psychiatric disorders, epilepsy and a number of diseases of peripheral organs, including the cardiovascular, airways, kidney, liver, bladder, skin and musculoskeletal. The potential of P2X7R drugs to treat tumour progression is discussed.

Keywords Pain · Infection · Cancer · CNS disorders · Cardiovascular · Airways · Diabetes · Kidney · Bladder · Liver · Gut · Immune cells

Introduction

P2X receptors are a family of ionotropic ATP-gated receptors. They are cation-selective channels, equally permeable to Na⁺ and K⁺ and with significant Ca²⁺ permeability. Seven mammalian subunits have been cloned to date (P2X1–7) that form either functional homo- or heterotrimers. When three molecules of ATP bind to a P2X receptor, the pore opens within milliseconds, allowing the cations to flow. P2X7 receptors (P2X7R) differ in that they have extra amino acids in the intracellular C terminus and are bi-functional. The binding of ATP within milliseconds induces the opening of a channel selective for small cations, and within seconds, a larger pore opens, which allows permeation by molecules with a mass of up to 900 Da (leading to the release of inflammatory cytokines and apoptosis) (see [1, 2]).

An important review was published about the development of P2X7R antagonists for anti-inflammatory therapy, including therapeutic potential and recent discoveries [3]. More recent reviews about P2X ion channel receptors and inflammation [1] and the P2X7R as a therapeutic target [4] have been published. This review is focused largely on the recent interest since 2014 about the involvement of P2X7R in inflammation and tumour progression.

P2X7R and inflammation

P2X7R have been implicated in the regulation of inflammation [5] and virtually all immune cell types (both of innate and adaptive immunity, including lymphocytes, macrophages, monocytes, neutrophils, basophils, dendritic cells, eosinophils and mast cells) express P2X7R [6]. They therefore represent significant therapeutic potential. A review was published in 2015 that discussed those P2X7R antagonists that exhibited

✉ Geoffrey Burnstock
g.burnstock@ucl.ac.uk

¹ Autonomic Neuroscience Centre, University College Medical School, Rowland Hill Street, London NW3 2PF, UK

² Department of Pharmacology and Therapeutics, The University of Melbourne, Melbourne, Australia

³ Florey Institute of Neuroscience and Mental Health, Parkville, Melbourne, Australia


promising therapeutic potential for the treatment of inflammatory diseases, pain and cancer [7]. Other reviews have been published that highlight the involvement of P2X7R in neuroinflammation and how medicinal chemists are working to identify centrally penetrant antagonists [8, 9]. For instance, GSK1482160 is a potent P2X7R antagonist with blood–brain barrier penetration, which has the ability to be radiolabelled with ^{11}C . This allows it to be potentially useful as a biomarker of neuroinflammation [10], since P2X7R may mediate the neuroinflammation and cognitive impairment that can result following surgery [11]. Another review explored the role of P2X7R in fibrosis, the pathological outcome of most chronic inflammatory diseases [12].

Activation of P2X7R is associated with immune responses, including allergic inflammation [13]. ATP released from damaged or infected cells causes inflammation by activation of P2X7R and subsequent release of inflammatory cytokines, such as interleukin (IL)-1 β , thereby acting as a danger signal by activating P2X7R on immune cells to increase immune responses [14]. An example of this is seen in enteric epithelial cells in conditions of inflammation of the gut (see [1]; Fig. 1). Pulmonary neutrophils are the inflammatory cells that are initially recruited during lung injury. Antagonism of the P2X7R with AZ106006120, or knockout (KO) of the receptor, reduced neutrophil infiltration and pro-inflammatory cytokine levels in a mouse model of acute lung injury [16]. Human and murine neutrophils express functional P2X7R, which mediate NLRP3 inflammasome-dependent IL-1 β secretion [17]. Nanobodies are small, single-domain antibody fragments and a mouse nanobody, 13A7, was shown to block gating of the P2X7R channel on T cells and macrophages *in vivo* and is proposed as a new drug candidate for inflammatory disorders [18].

P2X7R in inflammation, infection and immunity

Extracellular ATP is an endogenous danger signal that activates inflammatory responses in immune cells. The role of P2X7R in infectious inflammatory diseases has been reviewed [8, 19, 20]. P2X7R are needed for the development of the inflammatory response associated with sepsis in mice [21, 22] and Brilliant Blue G (BBG; a selective antagonist for mouse P2X7R that is blood–brain barrier permeable and safe) was shown to ameliorate sepsis-induced brain damage [23]. Also in mice, P2X7R downstream of caspase-11 play a critical role for pyroptosis and susceptibility to sepsis induced by the non-canonical inflammasome [24].

The use of P2X7R agonists in conjunction with low molecular weight anti-bacterial medicines has been proposed for the treatment of multi-drug-resistant tuberculosis [25]. P2X7R antagonists are potential tools for the treatment of *Clostridium perfringens* type C [26] and *Porphyromonas gingivalis* [27] infections. P2X7 activation was shown to be protective during severe



Enteric epithelial cell
(or enteric neuron, macrophage, etc.)

Fig. 1 Cell type-specific schema of ATP release and action. ATP (red triangles) can be released from the cell cytosol to the extracellular space (dashed red line) via pannexin 1 (Panx1) channels or connexin 43 (Cx43) hemichannels (pictured). Once in the extracellular space, this ATP acts as a paracrine transmitter, as can ATP released from nearby cells that are dead or dying (not shown). Extracellular ATP can activate P2 receptors, such as P2X7 receptors (P2X7R) (pictured), that depolarise the target cell but also activate an inflammatory response in immune cells (dashed gray line) with subsequent release of cytokines such as interleukin (IL)-1 β that can act back at Panx1 and Cx43 to modulate their function (dashed gray lines). Activation of P2X7R also mediates the T cell responses (e.g. Ca^{2+} entry, IL-2 synthesis) and macrophage migration (not shown). (Reproduced from [15], with permission from Frontiers Media S.A. (via Open Access))

Escherichia coli infection in mice [28]. Similarly, ATP release from infected macrophages and subsequent activation of P2X7R are critical for IL-1-dependent host protection from *Bacillus anthracis* [29]. The role of P2X7R and ectonucleotidases in infectious inflammatory diseases has been reviewed [20].

Evidence was presented to show that mouse P2X7R are involved in containing the parasitic protozoan *Toxoplasma gondii* spread *in vivo*, by stimulating inflammation [30]. The dysfunction of P2X7R is likely to contribute to morbidity due to human schistosomiasis, a chronic inflammatory disease [31]. P2X7R-deficient mice were more susceptible to *Leishmania amazonensis* infection than wild type (WT) mice, suggesting that P2X7R play a key role in parasite control by regulating T effector cells and inflammation [32].

P2X7R activation regulates inflammatory responses during acute viral infection in mice [33]. Upregulated expression of P2X7R on peripheral blood mononuclear cells provides anti-viral immunity in patients against hepatitis C virus [34], while activation of P2X7R participates in the exacerbated immune response that occurs during influenza virus infection in mice [35]. P2X7R play a role in control of dengue virus-2 infection and KN62 appeared to have anti-viral and anti-inflammatory actions in infected human monocytes [36]. Purinergic receptors, particularly P2X7, have been identified as key mediators

of human immunodeficiency virus-1 (HIV-1) infection and inflammation [37] and ATP induces rapid release of HIV-1 from virus containing compartments of human macrophages [38]. In HIV-1 infection, abnormalities in neuron-glia interactions result in neuronal damage. HIV-1 Tat induces neuronal apoptosis and augments the expression of P2X7R in astrocytes. Oxidised ATP (oxATP), A-438079 and BBG attenuated Tat-mediated neurotoxicity and may be novel tools for therapeutic management of neuroAIDS [39].

A valuable review discusses the role of purinergic signalling in autoimmunity and highlights a role for P2X7R in systemic lupus erythematosus [40]. P2X7R activation deleted intestinal T cells by apoptosis and suppressed T cell-induced colitis in mice [41]. P2X7R are expressed on peripheral lymphocytes and may influence the immune profile from patients with the indeterminate form of Chagas disease [42].

P2X7R and inflammatory neuropathic pain

The earlier literature about the involvement of P2X7R in neuropathic pain has been reviewed [43].

Genetically determined P2X7R pore formation was shown to regulate variability in chronic pain sensitivity [44]. P2X7R activation by endogenous ATP contributes to the development of inflammatory hyperalgesia. Removal of the P2X7R gene or P2X7R antagonists, such as BBG and oxATP, abolished chronic inflammatory and neuropathic pain in animal models (see review by Alves et al. [45]). Inflammatory pain that occurs in burn patients following dressing changes was relieved by puerarin treatment, which was claimed to act by decreasing the expression levels of P2X7R mRNA and protein in peripheral blood mononuclear cells [46]. P2X7R activation in vivo was involved in the development of central sensitisation in an acute inflammatory pain animal model (see [1]).

P2X7R antagonists were shown to have analgesic activity in a rat model of neuropathic pain. P2X7R expressed by microglia mediate neuropathic pain, which is reduced by antagonists, such as A-740003 and A-438079 (see [47]). Central nervous system (CNS)-penetrant P2X7R antagonists may be beneficial for the treatment of persistent pain by targeting microglia. Resveratrol was shown to be neuroprotective against neuropathic pain mediated by P2X7R expressed on satellite glial cells of the dorsal root ganglia (DRG) [48].

P2X7R and tumour progression

A review covering the early literature about the involvement of purinergic signalling and cancer, including discussion about the roles of the P2X7R in a variety of tumours, is available [49].


P2X7R are a key mediator of inflammation and cancer invasion/metastasis and P2X7R antagonists are potential

anti-metastatic agents [50]. A feature of some cancer cells is the high level of expression of P2X7R, which, depending on the tumour type, can mediate either proliferation or cell death (see [51]). Thus, P2X7R activation may have effects on anti-tumour immunity that are opposed to a direct effect on tumour growth. It was also pointed out in this review that there is increased release of ATP, promoting cancer cell migration and metastasis. High levels of extracellular ATP accumulate in tumour interstitium and high ATP doses inhibit migration of endothelial cells from human breast carcinoma, via the activation of P2X7R [52]. Human pancreatic duct adenocarcinoma cells in vitro express high levels of P2X7R protein and AZ10606120 inhibited cell proliferation [53]. ATP activation of P2X7R triggers immunogenic signalling, which converts dying cancer cells into an effective anti-cancer vaccine [54]. P2X7R activation is linked to elevated expression of inflammation promoting factors, tumour cell migration, increase in $[Ca^{2+}]_i$ and membrane depolarisation in malignant gliomas (see [55]; Fig. 2). Evidence has been presented to suggest that P2X7R antagonists are promising therapeutic tools for the treatment of osteosarcoma [56]. Growth of experimental tumours is strongly inhibited by P2X7R antagonism of both cancer and immune cells (see [57]). A non-functional P2X7R, nfpP2X7R, is expressed on cancer cells and has been proposed as a novel therapeutic target for human cancer [58, 59]. A phase I clinical trial has recently demonstrated that nfpP2X7R-targeted antibodies were a novel, safe and tolerable topical therapy for basal cell carcinoma [60]. P2X7R activation induces apoptosis in acute myeloid leukaemia cells but not in normal hematopoietic stem cells [61].

ACN human neuroblastoma cells express P2X7R, which is a regulator of its growth and angiogenesis, and AZ10606120 and A-740003 reduced ACN-derived tumour growth in nude mice so P2X7R may be a novel therapeutic target for treatment of neuroblastoma [62, 63]. P2X7R are expressed on human malignant glioma and C6 cells [55, 64, 65] and high P2X7R expression has been correlated with progression-free survival and overall survival [66]. P2X7R activation caused release of inflammatory cytokines by macrophages exposed to glioma-conditioned medium that was prevented by A-740003 [67]. A P2X7R agonist, 2'(3')-O-(4-benzoylbenzoyl)-ATP (Bz-ATP), enhanced the temozolomide anti-tumour effect on human cultured glioblastoma stem cells [68].

Autocrine release of ATP and activation of P2X7R influence the metastatic migration of human lung cancer H292 and PC-9 cells [69], and in immunodeficient mice, migration of transplanted HTB177 and HTB183 lung cancer cells was reduced by A-438079 [70]. The role of P2X7R in tumour progression is complex and studies show contradictory effects of P2X7R activation. For instance, an increase in survival of non-small cell lung cancer patients with high P2X7R expression was observed [71], although low P2X7R expression resulted in greater mRNA (miR-21) expression in the non-small

Fig. 2 Altered regulation of calcium channels in brain tumours is part of neoplastic transformation. In the brain, the transformation of a normal cell into a tumour cell might be related to Ca^{2+} oscillations, and the homeostasis misbalance can define the malignant phenotype, which includes uncontrolled proliferation, enhanced migration and invasion and abnormal cell death. The activation of P2X7 receptors (P2X7R) leads to extracellular signal-regulated protein kinases 1 and 2 (ERK1/2), phosphatidylinositol 3-kinase (PI3K) and mitogen-activated protein kinase 1/2 (MEK1/2) activation. High P2X7R functionality and pore activity are linked to apoptosis/necrosis in glioma cells and better progression-free survival. (Reproduced from [55], with permission from the American Society for Pharmacology and Experimental Therapeutics)



cell lung cancer tumours. Thus, the high levels of miR-21 expression in these cancer patients may be a consequence of P2X7R downregulation and resultant promotion of tumour progression. These results agree with the study by Souza and colleagues [72], which showed that defective P2X7R expression following miR-21 activation by a K-Ras mutation, led to reduced tumour-killing activity, resulting in a poorer prognosis. P2X7R are expressed in pancreatic cancer PancTu-1 Luc cells and antagonists, such as AZ10606120, are likely to be effective therapeutic agents [53]. Other antagonists, A-438079 and A-740003, reduced inflammation associated with colitis but increased tumour incidence in a mouse model of colitis-associated cancer [73]. In P2X7R-transfected human embryonic kidney cells and CT26 colon carcinoma, there was enhanced tumorigenesis when the cells were inoculated into either immunodeficient or immunocompetent mice, respectively. It was shown that in tumours derived from B16 mouse melanoma or ACN human neuroblastoma cell

lines, tumour growth was inhibited by oxATP [74]. The anthraquinone, emodin, suppressed the invasiveness of the highly invasive breast cancer cell line MDA-MB-435s, by antagonizing P2X7R [75]. ATP increased $[\text{Ca}^{2+}]_i$ in breast tumour cells and high concentrations produced apoptosis via P2X7R. ATP-mediated activation of the human breast cancer cell line T47D resulted in an increase in cell migration and the development of metastases, suggesting a potential therapeutic role for P2X7R antagonists [76]. P2X7R expression is a prognostic indicator for postoperative cancer-specific survival of patients with clear-cell renal cell carcinoma [77].

Decreased expression of P2X7R is associated with the development of cervical cancer. In women, decreased expression of P2X7R is found in endometrial epithelial pre-cancerous lesions. Activation of P2X7R-dependent apoptosis with BzATP may be a chemotherapeutic approach to prevent cell growth of pre-cancerous and early cancerous epithelial lesions [78]. Clearly, while ATP and P2X7R activation induced

apoptosis of tumour cells in some models, in others it accelerated tumour growth.

It is known that there are many polymorphisms of the P2X7R [79, 80], which, in addition to resulting in a loss of function, may alter the activity of the receptor. Expression of non-functional cytolytic P2X7R was found in all pathology specimens of prostate cancer examined [81]. P2X7R were not expressed in normal tissues from patients with no evidence of cancer, suggesting that the appearance of P2X7R is an early marker of prostate cancer. The expression of P2X7R is increased by hypoxia and hypoxia-driven increase in P2X7R enhances invasion and migration of tumour cells [82]. Ultraviolet (UV) light has been implicated in the genesis of cutaneous cancer, including basal and squamous cell carcinoma as well as melanoma. UV-B irradiation destroys P2X7R, by directly killing epithelial cells and by reducing P2X7R mRNA by degrading the protein, and may contribute to the malignant transformation of keratinocytes [83]. P2X7R were expressed in the necrotic centre of nodular basal cell carcinomas and in apoptotic cells in superficial multifocal and infiltrative cells. ATP caused apoptosis of cultured A431 human squamous cell carcinoma cell line via P2X7R and application of BzATP inhibited the formation of skin papillomas and carcinomas in mice [84]. There was increased expression of P2X7R in patients with superficial spreading melanomas, which were later shown to be functional, and it was suggested that they are a target for melanoma therapy. A low pH environment (like that seen in solid tumours) induced ATP release from B16 melanoma cells to increase proliferation via P2X7R and oxATP inhibited tumour growth in B16 melanoma-bearing mice [85]. In P2X7R KO mice, tumour progression of B16 melanoma was accelerated, showing that P2X7R are critical to support an anti-tumour immune response as well as restricting tumour growth and metastatic diffusion [86]. γ -Irradiation, which causes growth arrest and death of tumour cells, induced P2X7R-dependent ATP release from the B16 melanoma cells. P2X7R KO mice were susceptible to bone cancer pain and had an earlier onset of pain-related behaviours. The majority of human osteosarcomas express P2X7R isoforms A and B and the expression of either isoform is differently coupled to cell growth and activity [87]. P2X7R are involved in cancer-induced bone pain and A839977 was suggested as a useful analgesic tool in a rat model of cancer-induced bone pain [88].

Activation of P2X7R can either promote cellular survival or induce cytotoxicity and depends on the stimulus intensity of ATP to control the ion channel or the P2X7-dependent large pore functions. How these two opposite effects are controlled is not fully understood. Recently, a feedback loop was described showing that sustained activation of P2X7R results in the release of active matrix metalloproteinase 2, which stops ion channel and large pore responses in several different cell types, including macrophages and human tumour cells.

This effect may be an important fine-tuning of P2X7R functions. The authors suggested that P2X7R antagonists could be useful as in treating inflammatory diseases and cancers [89].

As the tumour microenvironment contains very high concentrations of ATP, adenosine is also present in high concentrations, following enzymatic breakdown of ATP by CD39 and CD73. Both ATP and adenosine contribute to immunosuppression or immunostimulation of the host, and stimulation of growth or cytotoxicity of the tumour, depending on the receptors activated. Thus, targeting specific receptor subtypes produces different effects, for example targeting CD73 or A_{2A} receptors reduced immunosuppression and potentially inhibited tumour growth [90]. As has been described above, growth of experimental tumours is also strongly inhibited by targeting P2X7R of cancer and immune cells (see [57]).

P2X7R and diseases of the CNS

The role of P2X7R in diseases related to neuroinflammation and the use of centrally penetrant antagonists has been highlighted [1, 11, 14, 18, 91–94]. Blockade of P2X7R may serve as a therapeutic target in alleviating the degree of inflammation seen in neurodegenerative and neoplastic conditions [95]. Figure 3 is a schematic showing P2X7R-mediated pathways of common disease mechanisms in CNS disorders of different aetiology. P2X7R are upregulated in various disease conditions and stress signals elicit activation, which can lead to excitotoxicity, neuroinflammation, neuronal damage, reactive astrogliosis or neuroplasticity, contributing to disease pathology (see [92]).

Neurodegenerative diseases

Attention has been directed recently towards the use of P2X7R antagonists for the treatment of neurodegenerative diseases, which are often associated with inflammation and damage to both neurons and glia [96–99]. P2X7R antagonists were neuroprotective in an animal model of Alzheimer's disease (AD). P2X7R trigger α -secretase-dependent processing of the amyloid precursor protein to generate β -amyloid peptides, which are present in the amyloid plaques in AD. Inhibition of P2X7R in vivo reduces amyloid plaques through glycogen synthase kinase 3 β and secretases. BBG improved cognition in an animal model of AD and inhibited amyloid- β -induced loss of filopodia and dendrite spines in cultured hippocampal neurons [100]. The control of amyloid plaque formation in vivo by P2X7R was reviewed in 2015 [97] and discussed further in 2017 [101]. It was suggested that P2X7R contribute to Parkinson's disease (PD) pathogenesis through a triple effect on synaptotoxicity, gliosis and neurotoxicity, indicating therapeutic potential of P2X7R antagonists for PD [102–104]. BBG was neuroprotective in an intranigral lipopolysaccharide (LPS) animal model of PD [104]. P2X7R are involved in multiple sclerosis (MS) and in the animal model of




Fig. 3 Common disease mechanism by P2X7 receptor (P2X7R)-mediated pathways in central nervous system (CNS) disorders of different etiology. P2X7R are expressed on nerve terminals, astrocytes and microglia, and they are upregulated in various disease conditions. Stress signals such as hypoxia/ischemia (metabolic limitations), mechanical injury and bacterial or chemical toxins elicit the endogenous activation of P2X7R and lead to a self-amplifying ATP release and to further activation of P2X7R on neighbouring cells. Following the influx of Ca^{2+} through the receptor ion channel complex, P2X7R activation (i) releases glutamate from nerve terminals and astrocytes by both exocytotic and non-exocytotic mechanisms, which may give rise excitotoxicity; (ii) leads to the posttranslational processing of pro-interleukin-1 β (pro-IL-1 β) to the leaderless, mature IL-1 β and to its further release by the NLRP3 inflammasome and that

of other cytokines, which contribute to neuroinflammation; (iii) enhance reactive oxygen species (ROS) production and thereby aggravate protein misfolding and neuronal damage; (iv) leads directly or indirectly to cell death and the following reactive astrogliosis; and (v) directly or indirectly downregulates the production of brain-derived neurotrophic factor (BDNF) and the subsequent neuroplasticity. These key mechanisms could be manifested and contribute to disease pathology in Alzheimer's disease (AD), Parkinson's disease (PD), Huntington's disease (HD), status epilepticus (SE), amyotrophic lateral sclerosis (ALS), multiple sclerosis (MS), stroke, pain and mood disorders in different forms and proportion, depending on the etiology. Abbreviations: GLU, glutamate, ROS, reactive oxygen species. (Reproduced from [92], with permission from Elsevier)

MS, experimental autoimmune encephalomyelitis (see [105]), where increased expression of P2X7R protein was shown in brain homogenates. Genetic variants in P2X7R affect susceptibility to MS [106, 107]. Altered P2X7R level and function in mouse models of Huntington's disease was reported and BBG prevented neuronal apoptosis and reduced body weight loss and motor-coordination deficits [108].

Inflammation features in the pathogenesis of amyotrophic lateral sclerosis (ALS). The potential of antagonism with BBG for the treatment of ALS was proposed, but it was claimed that the treatment was gender-dependent, although this varied in different studies using the SOD1-G93A mouse model of ALS: more effective in males [109], more effective in females [110] and no difference between the sexes [111]. Similarly, the pathogenesis of ALS was shown to differ in different studies. In P2X7R KO-

SOD1-G93A mice, onset of the disease occurred more rapidly than in WT mice, with increased astrogliosis, microgliosis, motoneuron loss, induction of pro-inflammatory markers and activation of mitogen-activated protein kinase (MAPK) pathways [112]. In a later study by this group using the same mouse model of ALS, BBG reduced neuroinflammation at late pre-symptomatic phases of the disease, enhanced motor neuron survival and reduced microgliosis in the lumbar spinal cord [111]. These differences may be due to the timing of the receptor block, complete absence of the receptor in the KO study, as opposed to antagonising the receptor prior to symptoms showing in the latter study. Ca^{2+} abnormalities in ALS exist in both motor neurons and immune cells. Reduction of P2X7R expression on peripheral blood mononuclear cells from ALS patients led to calcium dysregulation, a feature of ALS [112].

Psychiatric disorders

Activation of microglial P2X7R causes inflammation and brain penetrant P2X7R antagonists are being developed since microglial P2X7R are being increasingly recognised as a therapeutic target for the treatment of neurological and psychiatric diseases [113]. A review on this subject is available [114].

There is increased neuronal death via P2X7R and pannexin 1 channels in cultured cerebral cortex from prenatal LPS-exposed mice, which suggested a link to an increased risk of developing neurological disorders, including schizophrenia, in the offspring (see [115]). P2X7R, mediating neuroinflammation via the activity of microglia, may play a role in bipolar disorder and offer therapeutic possibilities (see [116]). Anti-depressant effects of chrysothanol (a natural anthraquinone) via inhibition of P2X7R has been reported in an LPS animal model of depression [117]. Similarly, BBG had anti-inflammatory and anti-depressant effects in mice after LPS administration [118]. Acute restraint stress activates the inflammasome via release of ATP and stimulation of P2X7R in a mouse model of stress-related mood disorders [119] and the antagonist A-804598 was shown to have an impact on the neuroimmune and behavioural consequences of stress in a rat model [120]. The P2X7R-Gln460Arg variant has been associated with mood disorders and co-expression of WT P2X7R with the polymorphic variant Gln460Arg transfected into HEK293 human cells was found to impair receptor function [121].

Neuroprotection from injury

There are reviews that discuss the importance of inflammation and the involvement of purinergic signalling in the responses to brain injury [8] and of the P2X7R in particular [1, 14]. P2X7R antagonists are a promising therapeutic tool for the treatment of cerebral ischaemia (see [122, 123]). BBG reduced delayed neuronal cell death in the hippocampal CA1 region after ischaemia/reperfusion injury and it was found that systemic administration of P2X7R antagonists improved recovery after spinal cord injury. The neuroprotective action of the P2X7R antagonist, GSK1370319A, acts by inhibiting the assembly of the NLRP3 inflammasome in glial cells [124, 125]. ATP enhanced radiation-induced brain injury through microglial activation via P2X7R-mediated release of inflammatory mediators, such as cyclooxygenase-2, tumour necrosis factor- α (TNF- α) and IL-6, and P2X7R antagonists were suggested as a potential strategy for the treatment of patients with radiation-induced brain injury [126]. Activation of P2X7R in the hippocampus following traumatic brain injury was thought to imply cognitive impairment in rats [127], while A-438079 provided neuroprotection toward neurological disorders, including stroke, traumatic brain injury and subarachnoid haemorrhage, as well as preserving blood–brain barrier integrity [128].

Recent papers have recognised a role for P2X7R antagonists for the treatment of epilepsy, including drug-resistant epilepsy [8,

129, 130]. P2X7R antagonism produced lasting reduction in the development of spontaneous seizures and inhibited glial inflammatory responses in mouse models of experimental temporal lobe epilepsy [131] and pentylenetetrazol-induced seizures [132]. However, in pilocarpine-induced status epilepticus in mice, blockade of central P2X7R increased the number of seizures and their severity [133]. P2X7R are considered as a target for the treatment of hypoxic/ischaemic encephalopathy and other causes of neonatal seizures since in A-438079-treated mice, seizure number, EEG power and spiking during hypoxia as well as molecular markers of inflammation and microglia were reduced [130].

Cardiovascular diseases

The long non-coding RNA (lncRNA), NONRATT021972, targeted using a small interfering RNA (siRNA), decreased the upregulation of P2X7R in the superior cervical ganglion and improved cardiac function after myocardial ischaemia in rats [134]. Myocardial ischaemic injury facilitated sympathoexcitatory action via P2X7R and it was suggested that P2X7R antagonists may be useful for the treatment of coronary heart disease as expression of P2X7R is increased during coronary ischaemia-reperfusion [135]. Microglial P2X7R in the rat hypothalamic paraventricular nuclei regulate the sympathoexcitatory responses in acute myocardial infarction and gene knockdown of P2X7R with P2X7-siRNA, or inhibition with BBG, reduced mRNA and protein expression of IL-1 β and TNF- α [136]. P2X7R involvement in dilated cardiomyopathy has been reported in P2X7R KO mice [137]. A-740003 reduced experimental autoimmune myocarditis in a mouse model, suggesting a treatment for clinical myocarditis [138]. P2X7R in the kidney play a role in hypertension and it has been suggested that P2X7R antagonists may have promise as clinical anti-hypertensive agents. In atherosclerotic mice, P2X7R are over-expressed whereas P2X7R KO mice were found to have less plaque formation and decreased leukocyte recruitment following ATP stimulation [139]. P2X7R are pro-thrombotic and genetic KO of the P2X7R gene is protective in a mouse model of carotid artery thrombosis. Vasomotor dysfunction was caused by sub-failure overstretch injury in rat abdominal aorta via activation of P2X7R and this suggested that the use of antagonists for the treatment of vascular stretch injury in humans may be beneficial [140].

Diseases of the airways

Inflammation occurs in most diseases of the airways, including asthma, chronic obstructive pulmonary disease, cystic fibrosis, dyspnea, allergy, infection and injury. P2X7R are a target for therapeutic intervention in lung hypersensitivity reactions associated with chronic inflammatory responses. Extracellular ATP was recognised as a danger signal activating P2X7R in lung

inflammation and fibrosis and P2X7R antagonists were proposed as a novel therapeutic approach in humans to control IL-1 β production and fibrosis in lung injury [141] as well as in silicosis, an occupational lung disease, following a study in a mouse model [142]. Attenuated P2X7R function gives protection from asthma and is age-related, being most effective in young boys. P2X7R have also been implicated in the pathophysiology of allergy-induced lung inflammation. Targeting P2X7R on haematopoietic cells, namely dendritic cells or eosinophils, may be a therapeutic approach for the treatment of allergic asthma. The anti-allergic anti-histamine, oxatamide, is claimed to also act as a P2X7R antagonist in N18TG2 and J774 cells [13]. BBG prevented neurogenic pulmonary oedema after subarachnoid haemorrhage in rats by attenuating inflammation [143]. Cigarette smoke activates P2X7R signalling, which appears to be involved in the pathogenesis of emphysema, and induces ATP and inflammatory cytokine release from neutrophils via P2X7R activation [144]. Polymorphisms of the P2X7R gene are associated with the risk and prognosis of human tuberculosis [145]. Data was presented to support the view that P2X7R antagonists should be used to treat the aggressive forms of tuberculosis [146]. ATP, released by activated macrophages and damaged cells, modulates lung inflammation in pneumonia in cattle. Both epithelial cells and pulmonary microvascular endothelial cells expressed mRNA for P2X7R, as did alveolar macrophages, which, when stimulated, activate the proinflammatory IL-1 to IL-5 cytokine cascade. P2X7R are involved in the pathophysiology of LPS-induced lung injury [147] and there is upregulation of pulmonary P2X7R in both acute and chronic lung injury in mice where P2X7R deletion was shown to be lung protective [148]. Pulmonary neutrophils are the initial inflammatory cells to be recruited during lung injury and are crucial for innate immunity. In contrast, pathological recruitment of neutrophils results in lung injury and P2X7R antagonists were shown to reduce neutrophil infiltration and proinflammatory cytokine levels [16].

Gut disorders

Extracellular nucleotides and their receptors are involved in the pathogenesis of inflammatory bowel disease (IBD), which includes two main forms, namely ulcerative colitis (UC) and Crohn's disease (CD). P2X7R are involved in colonic motor dysfunction associated with bowel inflammation in rats [149] and are over-expressed in gut mucosa of patients with CD [150]. P2X7R KO mice were protected against gut inflammation, while in WT mice, ATP via P2X7R triggers the death of mucosal regulatory T cells [151]. Oestrogen receptor β activation may play a therapeutic role in IBD by downregulation of P2X7R [152]. Reviews discussing the role of P2X7R in IBD are available [15, 153]. UC differentially affects P2X7R-expressing enteric neurons based on their chemical codes [154]. In a later paper, it was shown that trinitrobenzene sulfonic acid-induced UC in rats

affected secretory and vasodilatory neurons, enteric sensory neurons and enteric glia of the submucosal plexus expressing P2X7R [155]. ATP mediates inflammatory responses in dextran sulfate sodium-induced UC in mice via P2X7R signalling and A-438079 down-regulated the production of proinflammatory cytokines and attenuated the colitis [156]. There is increased expression of P2X7R in the inflamed mucosa in CD in humans and animal models, suggesting that P2X7R may be a target for treatment of CD [151, 156, 157]. Purinergic signalling is involved in gastrointestinal motility disorders, such as diarrhoea and constipation. P2X7R activity was enhanced in enteric glia that were isolated from mice with long-term morphine treatment, which can result in colonic inflammation [158].

Diseases of the kidneys

Increased expression of P2X7R in renal hypertension, polycystic kidney disease (PKD) and glomerulonephritis is opening up novel purinergic possibilities for the treatment of kidney failure (see [159, 160]). A-438079 protected against ischaemic acute kidney injury in mice [161]. P2X7R are expressed in collecting duct cysts in the *cpk/cpk* mouse model of congenital PKD, and mRNA and protein increased as the disease developed. ATP inhibits renal cyst growth, via P2X7R and oxATP, and A-438079 reduced cyst formation via MAPK-dependent pathways in a zebrafish model of PKD [162]. P2X7R expression has been demonstrated in both experimental and human glomerulonephritis, which suggests that P2X7R antagonists may have therapeutic potential [163]. P2X7R deficiency attenuated renal injury in experimental glomerulonephritis in mice. BBG attenuated nephritis by inhibiting inflammasome activation in a mouse model of lupus nephritis [164]. There is predominant P2X7R control of glomerular haemodynamics in angiotensin-II hypertension [165]. While P2X7R are only weakly expressed in healthy kidney, they are significantly upregulated in hypertension [166], antagonism of which would reduce interstitial inflammation, prevent interstitial cell death and improve blood pressure control. P2X7R antagonism prevented the development of salt-sensitive hypertension and renal injury in Dahl salt-sensitive rats [167]. P2X7R expression was increased in human kidneys with type 2 diabetes and activation of P2X7R evoked renal inflammation and injury in the high-fat diet model of metabolic syndrome, suggesting that P2X7R antagonists might be useful therapeutically for diabetic nephropathy [168]. P2X7R activation contributes to the high prevalence of kidney disease found in diabetics [160].

Diseases of the lower urinary tract

Localised inhibition of P2X7R at the spinal cord inflammatory injury site in a rat model reduced microglia numbers and improved neurogenic bladder dysfunction [169]. P2X7R,

expressed by macrophages and neutrophils in the bladder sub-mucosa, are increased in cyclophosphamide-induced haemorrhagic cystitis in mice and treatment with A-438079 or genetic ablation of this receptor reduced the tissue levels of IL-1 β and TNF- α and reduced nociceptive behaviour [170]. It was shown that 7 days after unilateral ureteral obstruction in WT mice there was increased expression of P2X7R associated with inflammation and fibrogenic responses in the cortex [171]. It was suggested that there is a potential role for P2X7R antagonists to prevent renal interstitial fibrosis [172].

Diseases of the liver

In mouse models of acetaminophen (APAP)-induced inflammation, liver injury after overdose involved P2X7R activation [173] resulting in hepatic caspase-1 and migration of neutrophils into the liver, suggesting that ATP may play a pivotal role in the development of inflammasomes after APAP overdose [174]. A-438079 was reported to be protective against APAP-induced liver injury, by an effect on metabolic activation and cell death pathways, rather than on involvement of inflammasomes [175]. Blockade of the P2X7R-NLRP3 inflammasome axis is a potential therapeutic target for liver fibrosis [176]. In mice with induced autoimmune hepatitis, activation of P2X7R on natural killer T cells was shown to inhibit naive but stimulate activated cells, resulting in suppression or stimulation of the autoimmune hepatitis [177]. P2X7R-mediated responses participate in infection of human hepatocytes by hepatitis delta virus and hepatitis B virus. P2X7R mediate leptin-induced GLUT4 function in mice stellate cells in non-alcoholic steatohepatitis [178]. P2X7R may be a major component of the purinergic signalling complex in hepatitis C virus-induced liver pathogenesis [34].

Skin diseases

A valuable review discusses the involvement of P2X7R in skin biology and their role in inflammatory skin disorders, including irritant and chronic dermatitis, psoriasis, cancer, graft-versus-host disease as well as wound healing and transplantation [179]. Inflammation of the skin is associated with release of ATP, increase in expression of purinoceptors, particularly the P2X7R, and subsequent release of proinflammatory cytokines from immune cells [180]. P2X7R antagonists reduce skin inflammation. In psoriasis, high P2X7R expression, confined to the cell membrane of the basal layer of the epidermis, plays a role in shaping the inflammatory microenvironment [181], leading to differentiation of Th17 lymphocytes, which are involved in the pathogenesis and potential treatment of psoriasis [182]. Systemic sclerosis fibroblasts express mRNA for several P2R subtypes, including P2X7R,

suggesting a potential therapeutic role for P2X7R antagonists in systemic sclerosis patients. In the initial phase of wound healing, there is acute inflammation. P2X7R on immune cells also mediate killing of intracellular pathogens by inducing apoptosis of host macrophages, chemo-attraction and cell adhesion. P2X7R immunoreactivity was associated with hyperkeratotic areas of the stratum corneum and in nuclei of koilocytes in the suprabasal layers of warts [183]. The nuclei that were positive for P2X7R were shrunken, showing much more intense P2X7R staining. The presence of P2X7R in the nucleus of human papillomavirus-infected cells indicates disruption of the cellular machinery. P2X7R agonists may be used to trigger apoptosis in these virally infected cells. Sensitisation to contact allergens requires activation of the immune system by endogenous danger signals. Mice lacking P2X7R are resistant to contact hypersensitivity. P2X7R-deficient dendritic cells fail to induce sensitisation to contact allergens and do not release IL-1 β in response to ATP, suggesting that P2X7R are crucial for extracellular ATP release in skin in response to contact allergens [184]. Interference with P2X7R signalling may be a promising strategy for the prevention of allergic contact dermatitis. There is a pathogenic role for keratinocyte-derived ATP in irritant dermatitis. The necrosis produced by the chemical irritant croton oil was prevented in mice by pre-treatment with A-438079 [185].

Musculoskeletal diseases

P2X7R proteins were upregulated on dystrophic myoblasts of *mdx* mice (a mouse model of Duchenne muscular dystrophy) and it was suggested that antagonists to these receptors may be of potential therapeutic benefit [186] as treatment with BBG or oxATP slowed the progression of disease in the mouse model [187]. The P2X7R has received the most attention in relation to the treatment of osteoporosis [188]. P2X7R plays an important role in both cortical and cancellous bone mass augmentation. They were shown to mediate stimulation of periosteal and cancellous bone formation and inhibition of cancellous bone resorption during growth [188]. Single nucleotide polymorphisms (SNPs) of the P2X7 gene are associated with fracture risk, decrease in bone mineral density and osteoporosis [189]. P2X7R are also involved in the chain of events leading to the formation of human osteoclasts, and P2X7R antagonists may have therapeutic roles in bone diseases with an increase in osteoclast number, such as Paget's disease [190]. P2X7R antagonists may be useful for the management of osteoporosis and with disorders of remodelling, where there is reduced bone mass [191]. Different polymorphic variants of the P2X7R are associated with high or reduced periprosthetic osteolysis in the long-term complication of total hip arthroplasty due to osteoarthritis [192]. Cross-talk between P2X7R and Wnt/ β -catenin pathways may modulate

osteoblast activity in response to mechanical loading [193]. Treatment with AZD9056, a selective P2X7R antagonist, produced pain-relieving and anti-inflammatory effects in rats with osteoarthritis [194]. ATP, via P2X7R, induced higher levels of IL-1 β in blood samples from rheumatoid arthritis (RA) patients compared to controls. It was suggested that mononuclear cells from these patients were more sensitive to ATP stimulation, perhaps due to genetic polymorphism in the P2X7R gene [195]. P2X7R play a role in the pathogenesis of RA and systemic lupus erythematosus. Human rheumatoid synoviocytes express P2X7R mRNA and protein. Animal models of arthritis have provided evidence for an *in vivo* role for the P2X7R in the progression of inflammatory disease. In P2X7R KO mice, there was a reduced incidence and severity of anti-collagen-induced arthritis symptoms; therefore, targeting the P2X7R may be a promising treatment for RA [196]. Block of P2X7R signalling in the collagen-induced arthritis animal model of RA inhibited peripheral inflammatory tissue destruction. P2X7R antagonists have been explored for the treatment of inflammatory pain in joints [197]. Charcot-Marie-Tooth 1A is a demyelinating hereditary neuropathy and A-438079 improved the clinical phenotype of the disease in a rat model and was recommended for its treatment [198]. A review was published about P2X7R antagonists in rodent models of musculoskeletal and other disorders [199].

P2X7R and diabetes

Type 1 and type 2 diabetes are inflammatory diseases. P2X7R KO in mice prevents streptozotocin (STZ)-induced type 1 diabetes and the levels of proinflammatory mediators (IL-1 β , interferon- γ and nitric oxide) did not increase [200]. P2X7R-pannexin 1 channels impair bone mechanosignalling in osteocytes associated with type 1 diabetes that affects osteoblast function and maintenance of bone health [201]. In STZ-induced diabetic animals, P2X7R located on glucagon-containing α cells in pancreatic islets increase and they migrate centrally to take the place of the missing insulin-containing β cells [202]. OxATP in mice has been proposed as a therapeutic tool to cause immunosuppression and tolerance induction in pancreatic islet transplantation [203, 204]. LncRNA NONRATT021972 siRNA decreases the expression of P2X7 mRNA and protein in DRG, thereby reducing mechanical and thermal hyperalgesia in type 2 diabetic rats [205]. ATP concentrations and P2X7R expression were increased in glial cells in rats with painful diabetic neuropathy [206]. Diabetic sympathetic neuropathy in type 2 diabetic rats was improved by reducing the expression of P2X7R with lncRNAuc.48+ siRNA in superior cervical ganglia [207]. Different P2X7R polymorphisms are associated with pain sensitivity for diabetic neuropathic pain patients, such that patients with the gain-of-function SNP report more severe

pain, whereas those with a loss-of-function SNP report less pain when the pain was tested and scored [208]. P2X7R antagonists may be a useful adjuvant treatment to delay the progression of diabetic nephropathy [209].

Conclusions

It is clear from the recent literature that P2X7R antagonists are potential effective therapeutic agents for the treatment of inflammatory diseases and cancer. There is a problem, however, concerning the many polymorphic forms of the human P2X7R, as care must be taken in identifying those patients that would most benefit by the available antagonists, as different antagonists are not always effective agents at some of the polymorphic types (see [210]). There is an explosion of interest in developing centrally penetrant P2X7R antagonists and their therapeutic explorations in clinical trials [211–213]. The relation between inflammation elicited by P2X7R activation and the immune system needs further exploration. Apart from the potential of the therapeutic use of P2X7R antagonists for CNS disorders (including neurodegenerative diseases, brain injury, psychiatric diseases, epilepsy and neuropathic pain), there is increasing recognition of their use for the treatment of diseases of the heart, lung, gut, kidney, liver and bladder. The reasons for dual roles of P2X7R mediating cell proliferation and apoptotic cell death need further investigation of the mechanisms involved.

The purinergic signalling field is now well established and much known about the physiological roles played by this system. The emphasis now is on the pathophysiology and therapeutic potential of purinergic signalling (see [214, 215]). For example, clopidogrel (trade name Plavix) and ticagrelor (trade name Brilique) are P2Y₁₂ receptor antagonists that inhibit platelet aggregation and are widely used for the treatment of thrombosis and stroke (see [216]).

Characterisation of the P2Y class (see [217]) made it possible to identify specific agonists of the P2Y₂ receptor, which evokes mucus secretion. A new, long-lasting agonist, diquafasol (Diquas), has been developed by Inspire Pharmaceuticals [218]. Diquas was launched by Santen in Japan in 2010, with Inspire Pharmaceuticals.

Through the cloning of the P2X3 ion channel receptor located on nociceptive sensory nerves [219] and later the discovery of the involvement of these channels in signalling pain (see [220]), a new target for pain relief is receiving attention from the pharmaceutical industry. Roche and, more recently, Afferent Pharmaceuticals have developed AF-219, a specific P2X3 antagonist, which is a promising new analgesic, orally bioavailable and stable *in vivo* [221]. AF-219 is currently in phase 2 clinical trials for the treatment of three painful disorders: osteoarthritis; interstitial cystitis/bladder pain syndrome and idiopathic chronic cough. Four phase 1 studies of AF-219 have demonstrated that the compound is safe and well

tolerated and the phase 2 clinical trials were completed earlier this year [222]. Antagonism of P2X3 receptors on primary afferent neurons is a novel analgesic approach that has been pursued by leading pharmaceutical companies for the last 15 years. Over 600 patents are currently held which relate to the P2X3 receptor and pain (<https://patentscope.wipo.int/search/en/result.jsf>). There are currently investigations into the potential therapeutic use of purinergic compounds for osteoporosis, irritable bowel syndrome, atherosclerosis, kidney failure and cancer.

Since P2X7R-mediated inflammation is associated with a wide range of diseases, medicinal chemists in research institutes, universities and drug companies are optimistic that P2X7R antagonists are promising tools for the treatment of inflammatory diseases and more than 70 patents have been filed in the last few years by Glaxo, AstraZeneca, Roche, Janssen and Abbott, to name a few. For instance, Abbot filed a patent from the use of A-438079 in disease models of affective disorders and status epilepticus, while Abbott filed for triazole-based compounds for the CNS disorders and as a therapeutic strategy to treat spinal cord injury and the University Hospitals of Cleveland claimed in vitro and in vivo effects of P2X7R antagonists in epithelial cancer and papilloma (see [213]).

As mentioned above, a major challenge is to understand which polymorphic variations of the P2X7R have relevance to which disease and to fit patients with different polymorphisms to the P2X7R antagonist that would have the greatest efficacy for that variant. For example, the responses of human leukocytes to GSK1370319A were significantly altered, directly related to the SNP genotype, there being a 6.7-fold difference in the inhibition of ATP-stimulated IL-1 β release by GSK1370319A between individuals with the homozygous gain- and loss-of-function genotypes [223]. A correlation was found between some gain-of-function and loss-of-function P2X7R SNPs, transfected into HEK-293 cells, and the expression of the channel protein. A change in both channel and pore function was described for the mutant P2X7R in parallel to the protein levels, although the agonist and antagonist sensitivity was not altered. The presence of the gain-of-function SNP (rs208294 (His155Tyr) and rs1718119 (Ala348Thr)) in female patients with diabetic peripheral neuropathic pain was associated with higher pain intensity scores [208]. Recently, mutations and molecular modelling studies have identified an allosteric binding site forming at the subunit interface at the apex of the receptor, distinct from the ATP-binding pocket, that regulates access of antagonists, such as AZ10606120, to the allosteric site. This allosteric pocket may provide novel targets for P2X7R drug development [224, 225], although the effect of polymorphic variations on the allosteric site has not been elucidated yet.

Also, more needs to be discovered about the relation of inflammatory diseases to the immune system where

lymphocytes, macrophages, monocytes, neutrophils, basophils, dendritic cells, eosinophils and mast cells all express P2X7R (see [6]).

It seems likely that a breakthrough will occur about the use of P2X7R antagonists for a variety of diseases in the future, including, in particular, neurodegenerative diseases and cancer and the description of the crystal structure of mammalian P2X7R can only help in this endeavour [225, 226].

In addition to the therapeutic potential of P2X7R antagonists, P2X3R antagonists look extremely promising for the treatment of chronic cough, visceral pain, bladder disorders and hypertension. Exploration of the use of purinergic drugs for the treatment of obesity is also well worth pursuing.

Funding The writing of this review article was not financially supported.

Compliance with ethical standards

Conflict of interest Geoffrey Burnstock declares that he has no conflict of interest.

Gillian E. Knight declares that she has no conflict of interest.

Ethical approval This review article does not contain any studies with human participants or animals performed by either of the authors.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Burnstock G (2016) P2X ion channel receptors and inflammation. *Purinergic Signalling* 12:59–67
2. Volonté C, Apolloni S, Skaper SD, Burnstock G (2012) P2X7 receptors: channels, pores and more. *CNS & Neurological Disorders-Drug Targets* 11:705–721
3. Mehta N, Kaur M, Singh M, Chand S, Vyas B, Silakari P, Bahia MS, Silakari O (2014) Purinergic receptor P2X₇: a novel target for anti-inflammatory therapy. *Bioorg Med Chem* 22:54–88
4. Burnstock G (2013) Purinergic signalling: pathophysiology and therapeutic potential. *Keio J Med* 62:63–73
5. Di Virgilio F, Dal Ben D, Sarti AC, Giuliani AL, Falzoni S (2017) The P2X₇ receptor in infection and inflammation. *Immunity* 47:15–31
6. Burnstock G, Boeynaems J-M (2014) Purinergic signalling and immune cells. *Purinergic Signal* 10:529–564
7. Baudelet D, Lipka E, Millet R, Ghinet A (2015) Involvement of the P2X₇ purinergic receptor in inflammation: an update of antagonists series since 2009 and their promising therapeutic potential. *Curr Med Chem* 22:713–729
8. Beamer E, Gölöncsér F, Horváth G, Bekö K, Otrokocsi L, Koványi B, Sperlág B (2016) Purinergic mechanisms in neuroinflammation: an update from molecules to behavior. *Neuropharmacology* 104:94–104

9. Rech JC, Bhattacharya A, Letavic MA, Savall BM (2016) The evolution of P2X7 antagonists with a focus on CNS indications. *Bioorg Med Chem Lett* 26:3838–3845
10. Territo PR, Meyer JA, Peters JS, Riley AA, McCarthy BP, Gao M, Wang M, Green MA, Zheng QH, Hutchins GD (2017) Characterization of ¹¹C-GSK1482160 for targeting the P2X7 receptor as a biomarker for neuroinflammation. *J Nucl Med* 58:458–465
11. Zheng B, Lai R, Li J, Zuo Z (2017) Critical role of P2X7 receptors in the neuroinflammation and cognitive dysfunction after surgery. *Brain Behav Immun* 61:365–374
12. Gentile D, Natale M, Lazzarini PE, Capecchi PL, Laghi-Pasini F (2015) The role of P2X₇ receptors in tissue fibrosis: a brief review. *Purinergic Signal* 11:435–440
13. Yoshida K, Ito M, Matsuoka I (2015) P2X7 receptor antagonist activity of the anti-allergic agent oxatomide. *Eur J Pharmacol* 767: 41–51
14. Giuliani AL, Sarti AC, Falzoni S, Di Virgilio F (2017) The P2X7 receptor-interleukin-1 liaison. *Front Pharmacol* 8:123
15. Diezmos EF, Bertrand PP, Liu L (2016) Purinergic signaling in gut inflammation: the role of connexins and pannexins. *Front Neurosci* 10:311
16. Mishra A, Guo Y, Zhang L, More S, Weng T, Chintagari NR, Huang C, Liang Y, Pushparaj S, Gou D, Breshears M, Liu L (2016) A critical role for P2X7 receptor-induced VCAM-1 shedding and neutrophil infiltration during acute lung injury. *J Immunol* 197:2828–2837
17. Karmakar M, Katsnelson MA, Dubyak GR, Pearlman E (2016) Neutrophil P2X7 receptors mediate NLRP3 inflammasome-dependent IL-1 β secretion in response to ATP. *Nat Commun* 7:10555
18. Danquah W, Meyer-Schwesinger C, Rissiek B, Pinto C, Serracant-Prat A, Amadi M, Jacenda D, Knop JH, Hammel A, Bergmann P, Schwarz N, Assunção J, Rothier W, Haag F, Tolosa E, Bannas P, Boué-Grabot E, Magnus T, Laeremans T, Stortelers C, Koch-Nolte F (2016) Nanobodies that block gating of the P2X7 ion channel ameliorate inflammation. *Sci Transl Med* 8:366ra162
19. Di Virgilio F, Vuerich M (2015) Purinergic signaling in the immune system. *Auton Neurosci* 191:117–123
20. Morandini AC, Savio LE, Coutinho-Silva R (2014) The role of P2X7 receptor in infectious inflammatory diseases and the influence of ectonucleotidases. *Biomed J* 37:169–177
21. Csóka B, Németh ZH, Törö G, Idzko M, Zech A, Koscsó B, Spolarics Z, Antonioli L, Cseri K, Erdélyi K, Pacher P, Haskó G (2015) Extracellular ATP protects against sepsis through macrophage P2X7 purinergic receptors by enhancing intracellular bacterial killing. *FASEB J* 29:3626–3637
22. Santana PT, Benjamim CF, Martinez CG, Kurtenbach E, Takiya CM, Coutinho-Silva R (2015) The P2X7 receptor contributes to the development of the exacerbated inflammatory response associated with sepsis. *J Innate Immun* 7:417–427
23. Savio LE, Andrade MG, de Andrade Mello P, Santana PT, Moreira-Souza AC, Kolling J, Longoni A, Feldbrügge L, Wu Y, Wyse AT, Robson SC, Coutinho-Silva R (2016) P2X7 receptor signaling contributes to sepsis-associated brain dysfunction. *Mol Neurobiol*
24. Yang D, He Y, Munoz-Planillo R, Liu Q, Nuñez G (2015) Caspase-11 requires the pannexin-1 channel and the purinergic P2X7 pore to mediate pyroptosis and endotoxin shock. *Immunity* 43:923–932
25. Soares-Bezerra RJ, Pinho RT, Bisaggio RC, Benévolo-de-Andrade TC, Alves LA (2015) The search for new agonists to P2X₇R for clinical use: tuberculosis as a possible target. *Cell Physiol Biochem* 37:409–418
26. Nagahama M, Seike S, Shirai H, Takagishi T, Kobayashi K, Takehara M, Sakurai J (2015) Role of P2X7 receptor in *Clostridium perfringens* beta-toxin-mediated cellular injury. *Biochim Biophys Acta* 1850:2159–2167
27. Almeida-da-Silva CL, Morandini AC, Ulrich H, Ojcius DM, Coutinho-Silva R (2016) Purinergic signaling during *Porphyromonas gingivalis* infection. *Biomed J* 39:251–260
28. Greve AS, Skals M, Fagerberg SK, Tomms W, Ellermann-Eriksen S, Evans RJ, Linkermann A, Praetorius HA (2017) P2X₁, P2X₄, and P2X₇ receptor knock out mice expose differential outcome of sepsis induced by α -haemolysin producing *Escherichia coli*. *Front Cell Infect Microbiol* 7:113
29. Ali SR, Timmer AM, Bilgrami S, Park EJ, Eckmann L, Nizet V, Karin M (2011) Anthrax toxin induces macrophage death by p38 MAPK inhibition but leads to inflammasome activation via ATP leakage. *Immunity* 35:34–44
30. Corrêa G, Almeida Lindenberg C, Moreira-Souza AC, Savio LE, Takiya CM, Marques-da-Silva C, Vommaro RC, Coutinho-Silva R (2017) Inflammatory early events associated to the role of P2X7 receptor in acute murine toxoplasmosis. *Immunobiology* 222:676–683
31. Silva CL (2016) Purinergic signaling in schistosomal infection. *Biomed J* 39:316–325
32. Figliuolo VR, Chaves SP, Savio LE, Thorstenberg ML, Machado Salles É, Takiya CM, D'Império-Lima MR, de Matos Guedes HL, Rossi-Bergmann B, Coutinho-Silva R (2017) The role of the P2X7 receptor in murine cutaneous leishmaniasis: aspects of inflammation and parasite control. *Purinergic Signal* 13:145–152
33. Lee BH, Hwang DM, Palaniyar N, Grinstein S, Philpott DJ, Hu J (2012) Activation of P2X₇ receptor by ATP plays an important role in regulating inflammatory responses during acute viral infection. *PLoS One* 7:e35812
34. Manzoor S, Akhtar U, Naseem S, Khalid M, Mazhar M, Parvaiz F, Khaliq S (2016) Ionotropic purinergic receptors P2X₄ and P2X₇: proviral or antiviral? An insight into P2X receptor signaling and hepatitis C virus infection. *Viral Immunol* 29:401–408
35. Leyva-Grado VH, Ermler ME, Schotsaert M, Gonzalez MG, Gillespie V, Lim JK, Garcia-Sastre A (2017) Contribution of the purinergic receptor P2X₇ to development of lung immunopathology during influenza virus infection. *MBio* 8:e00229–e00217
36. Corrêa G, Lindenberg dAC, Fernandes-Santos C, Gandini M, Petitinga PF, Coutinho-Silva R, Kubelka F (2016) The purinergic receptor P2X₇ role in control of dengue virus-2 infection and cytokine/chemokine production in infected human monocytes. *Immunobiology* 221:794–802
37. Swartz TH, Dubyak GR, Chen BK (2015) Purinergic receptors: key mediators of HIV-1 infection and inflammation. *Front Immunol* 6: 585
38. Graziano F, Desdouits M, Garzetti L, Podini P, Alfano M, Rubartelli A, Furlan R, Benaroch P, Poli G (2015) Extracellular ATP induces the rapid release of HIV-1 from virus containing compartments of human macrophages. *Proc Natl Acad Sci U S A* 112:E3265–E3273
39. Tewari M, Monika VRK, Menon M, Seth P (2015) Astrocytes mediate HIV-1 Tat-induced neuronal damage via ligand-gated ion channel P2X₇R. *J Neurochem* 132:464–476
40. Di Virgilio F, Giuliani AL (2016) Purinergic signalling in autoimmunity: a role for the P2X₇R in systemic lupus erythematosus? *Biomed J* 39:326–338
41. Hashimoto-Hill S, Friesen L, Kim M, Kim CH (2017) Contraction of intestinal effector T cells by retinoic acid-induced purinergic receptor P2X₇. *Mucosal Immunol* 10:912–923
42. Souza VD, Dos Santos JT, Cabral FL, Barbisan F, Azevedo MI, Dias Carli LF, de Avila BS, Dos Santos Jaques JA, Rosa Leal DB (2017) Evaluation of P2X₇ receptor expression in peripheral lymphocytes and immune profile from patients with indeterminate form of Chagas disease. *Microb Pathog* 104:32–38
43. Burnstock G (2013) Purinergic mechanisms and pain—an update. *Eur J Pharmacol* 716:24–40

44. Sorge RE, Trang T, Dorfman R, Smith SB, Beggs S, Ritchie J, Austin JS, Zaykin DV, Vander MH, Costigan M, Herbert TA, Yarkoni-Abitbul M, Tichauer D, Livneh J, Gershon E, Zheng M, Tan K, John SL, Slade GD, Jordan J, Woolf CJ, Peltz G, Maixner W, Diatchenko L, Seltzer Z, Salter MW, Mogil JS (2012) Genetically determined P2X7 receptor pore formation regulates variability in chronic pain sensitivity. *Nat Med* 18:595–599
45. Alves LA, Bezerra RJ, Faria RX, Ferreira LG, da Silva Frutuoso V (2013) Physiological roles and potential therapeutic applications of the P2X7 receptor in inflammation and pain. *Molecules* 18:10953–10972
46. Zhang J, Li X, Gao Y, Guo G, Xu C, Li G, Liu S, Huang A, Tu G, Peng H, Qiu S, Fan B, Zhu Q, Yu S, Zheng C, Liang S (2013) Effects of puerarin on the inflammatory role of burn-related procedural pain mediated by P2X₇ receptors. *Burns* 39:610–618
47. Tsuda M (2017) P2 receptors, microglial cytokines and chemokines, and neuropathic pain. *J Neurosci Res* 95:1319–1329
48. Xie J, Liu S, Wu B, Li G, Rao S, Zou L, Yi Z, Zhang C, Jia T, Zhao S, Schmalzing G, Hausmann R, Nie H, Li G, Liang S (2017) The protective effect of resveratrol in the transmission of neuropathic pain mediated by the P2X7 receptor in the dorsal root ganglia. *Neurochem Int* 103:24–35
49. Burnstock G, Di Virgilio F (2013) Purinergic signalling in cancer. *Purinergic Signalling* 9:491–540
50. Park JH, Williams DR, Lee JH, Lee SD, Lee JH, Ko H, Lee GE, Kim S, Lee JM, Abdelrahman A, Muller CE, Jung DW, Kim YC (2016) Potent suppressive effects of 1-piperidinylimidazole based novel P2X7 receptor antagonists on cancer cell migration and invasion. *J Med Chem* 59:7410–7430
51. Ferrari D, Malavasi F, Antonioli L (2017) A purinergic trail for metastases. *Trends Pharmacol Sci* 38:277–290
52. Avanzato D, Genova T, Fiorio PA, Bernardini M, Bianco S, Bussolati B, Mancardi D, Giraud E, Maione F, Cassoni P, Castellano I, Munaron L (2016) Activation of P2X7 and P2Y11 purinergic receptors inhibits migration and normalizes tumor-derived endothelial cells via cAMP signaling. *Sci Rep* 6:32602
53. Giannuzzo A, Saccomano M, Napp J, Ellegaard M, Alves F, Novak I (2016) Targeting of the P2X7 receptor in pancreatic cancer and stellate cells. *Int J Cancer* 139:2540–2552
54. Garg AD, Krysko DV, Vandenabeele P, Agostinis P (2016) Extracellular ATP and P₂X₇ receptor exert context-specific immunogenic effects after immunogenic cancer cell death. *Cell Death Dis* 7:e2097
55. Morrone FB, Gehring MP, Nicoletti NF (2016) Calcium channels and associated receptors in malignant brain tumor therapy. *Mol Pharmacol* 90:403–409
56. Yadav M, Singh A, Rizvi N, Ali S, Salma S, Kumar V, Hussain SR (2015) The role of P2X7R purinoreceptor in osteosarcoma. *Advances in Modern Oncology Research* 1:88–96
57. Di Virgilio F, Adinolfi E (2017) Extracellular purines, purinergic receptors and tumor growth. *Oncogene* 36:293–303
58. Barden JA, Gilday-Baird A, Teh LC, Rajasekariah G-H, Pedersen J, Christensen NI, Spielman D, Ashley DM (2016) Therapeutic targeting of the cancer-specific cell surface biomarker nP2X₇. *J Clin Cell Immunol* 7:432
59. Barden JA, Yuksel A, Pedersen J, Danieletto S, Delprado W (2014) Non-functional P2X7: a novel and ubiquitous target in human cancer. *J Clin Cell Immunol* 5:327
60. Gilbert SM, Gidley Baird A, Glazer S, Barden JA, Glazer A, Teh LC, King J (2017) A phase I clinical trial demonstrates that nP2X7-targeted antibodies provide a novel, safe and tolerable topical therapy for basal cell carcinoma. *Br J Dermatol* 177:117–124
61. Salvestrini V, Orecchioni S, Talarico G, Reggiani F, Mazzetti C, Bertolini F, Orioli E, Adinolfi E, Di Virgilio F, Pezzi A, Cavo M, Lemoli RM, Curti A (2017) Extracellular ATP induces apoptosis through P2X7R activation in acute myeloid leukemia cells but not in normal hematopoietic stem cells. *Oncotarget* 8:5895–5908
62. Amoroso F, Capece M, Rotondo A, Cangelosi D, Ferracin M, Franceschini A, Raffaghello L, Pistoia V, Varesio L, Adinolfi E (2015) The P2X7 receptor is a key modulator of the PI3K/GSK3 β /VEGF signaling network: evidence in experimental neuroblastoma. *Oncogene* 34:5240–5251
63. Gomez-Villafuertes R, Garcia-Huerta P, Diaz-Hernandez JI, Miras-Portugal MT (2015) PI3K/Akt signaling pathway triggers P2X7 receptor expression as a pro-survival factor of neuroblastoma cells under limiting growth conditions. *Sci Rep* 5:18417
64. Monif M, O'Brien TJ, Drummond KJ, Reid CA, Liubinas SV, Williams DA (2014) P2X7 receptors are a potential novel target for anti-glioma therapies. *Journal of Inflammation* 11:25
65. Fang J, Chen X, Wang S, Xie T, Du X, Liu H, Wang S, Li X, Chen J, Zhang B, Liang H, Yang Y, Zhang W (2015) The expression of P2X₇ receptors in EPCs and their potential role in the targeting of EPCs to brain gliomas. *Cancer Biol Ther* 16:498–510
66. Gehring MP, Kipper F, Nicoletti NF, Sperotto ND, Zanin R, Tamajusuku AS, Flores DG, Meurer L, Roesler R, Filho AB, Lenz G, Campos MM, Morrone FB (2015) P2X7 receptor as predictor gene for glioma radiosensitivity and median survival. *Int J Biochem Cell Biol* 68:92–100
67. Bergamin LS, Braganhol E, Figueiró F, Casali EA, Zanin RF, Sévigny J, Battastini AM (2015) Involvement of purinergic system in the release of cytokines by macrophages exposed to glioma-conditioned medium. *J Cell Biochem* 116:721–729
68. D'Alimonte I, Nargi E, Zuccarini M, Lanuti P, Di Iorio P, Giuliani P, Ricci-Vitiani L, Pallini R, Caciagli F, Ciccarelli R (2015) Potentiation of temozolomide antitumor effect by purine receptor ligands able to restrain the in vitro growth of human glioblastoma stem cells. *Purinergic Signal* 11:331–346
69. Takai E, Tsukimoto M, Harada H, Kojima S (2014) Autocrine signaling via release of ATP and activation of P2X7 receptor influences motile activity of human lung cancer cells. *Purinergic Signal* 10:487–497
70. Schneider G, Glaser T, Lameu C, Abdelbaset-Ismael A, Sellers ZP, Moniuszko M, Ulrich H, Ratajczak MZ (2015) Extracellular nucleotides as novel, underappreciated pro-metastatic factors that stimulate purinergic signaling in human lung cancer cells. *Mol Cancer* 14:201
71. Boldrini L, Giordano M, Ali G, Melfi F, Romano G, Lucchi M, Fontanini G (2015) P2X7 mRNA expression in non-small cell lung cancer: MicroRNA regulation and prognostic value. *Oncol Lett* 9:449–453
72. Souza CO, Santoro GF, Figliuolo VR, Nanini HF, de Souza HS, Castelo-Branco MT, Abalo AA, Paiva MM, Coutinho CM, Coutinho-Silva R (2012) Extracellular ATP induces cell death in human intestinal epithelial cells. *Biochim Biophys Acta* 1820:1867–1878
73. Hofman P, Cherfils-Vicini J, Bazin M, Ilie M, Juhel T, Hébuterne X, Gilson E, Schmid-Alliana A, Boyer O, Adriouch S, Vouret-Craviari V (2015) Genetic and pharmacological inactivation of the purinergic P2RX7 receptor dampens inflammation but increases tumor incidence in a mouse model of colitis-associated cancer. *Cancer Res* 75:835–845
74. Adinolfi E, Raffaghello L, Giuliani AL, Cavazzini L, Capece M, Chiozzi P, Bianchi G, Kroemer G, Pistoia V, Di Virgilio F (2012) Expression of P2X7 receptor increases in vivo tumor growth. *Cancer Res* 72:2957–2969
75. Jelassi B, Anachelin M, Chamouton J, Cayuela ML, Clarysse L, Li J, Gore J, Jiang LH, Roger S (2013) Anthraquinone emodin inhibits human cancer cell invasiveness by antagonizing P2X7 receptors. *Carcinogenesis* 34:1487–1496
76. Xia J, Yu X, Tang L, Li G, He T (2015) P2X7 receptor stimulates breast cancer cell invasion and migration via the AKT pathway. *Oncol Rep* 34:103–110

77. Liu Z, Liu Y, Xu L, An H, Chang Y, Yang Y, Zhang W, Xu J (2015) P2X7 receptor predicts postoperative cancer-specific survival of patients with clear-cell renal cell carcinoma. *Cancer Sci* 106:1224–1231
78. Gorodeski GI (2009) P2X7-mediated chemoprevention of epithelial cancers. *Expert Opin Ther Targets* 13:1313–1332
79. Fuller SJ, Stokes L, SkarRatt KK, BJ G, Wiley JS (2009) Genetics of the P2X7 receptor and human disease. *Purinergic Signal* 5:257–262
80. Shemon AN, Sluyter R, Fernando SL, Clarke AL, Dao-Ung LP, SkarRatt KK, Saunders BM, Tan KS, BJ G, Fuller SJ, Britton WJ, Petrou S, Wiley JS (2006) A Thr357 to Ser polymorphism in homozygous and compound heterozygous subjects causes absent or reduced P2X7 function and impairs ATP-induced mycobacterial killing by macrophages. *J Biol Chem* 281:2079–2086
81. Slater M, Danieleto S, Gidley-Baird A, Teh LC, Barden JA (2004) Early prostate cancer detected using expression of non-functional cytolytic P2X₇ receptors. *Histopathology* 44:206–215
82. Tafani M, Schito L, Pellegrini L, Villanova L, Marfe G, Anwar T, Rosa R, Indelicato M, Fini M, Pucci B, Russo MA (2011) Hypoxia-increased RAGE and P2X7R expression regulates tumor cell invasion through phosphorylation of Erk1/2 and Akt and nuclear translocation of NF- κ B. *Carcinogenesis* 32:1167–1175
83. Ruzsnavszky O, Telek A, Gönczi M, Balogh A, Remenyik E, Csemoch L (2011) UV-B induced alteration in purinergic receptors and signaling on HaCaT keratinocytes. *J Photochem Photobiol B* 105:113–118
84. Fu W, McCormick T, Qi X, Luo L, Zhou L, Li X, Wang BC, Gibbons HE, bdul-Karim FW, Gorodeski GI (2009) Activation of P2X₇-mediated apoptosis inhibits DMBA/TPA-induced formation of skin papillomas and cancer in mice. *BMC Cancer* 9:114
85. Hattori F, Ohshima Y, Seki S, Tsukimoto M, Sato M, Takenouchi T, Suzuki A, Takai E, Kitani H, Harada H, Kojima S (2012) Feasibility study of B16 melanoma therapy using oxidized ATP to target purinergic receptor P2X7. *Eur J Pharmacol* 695:20–26
86. Adinolfi E, Capece M, Franceschini A, Falzoni S, Giuliani AL, Rotondo A, Sarti AC, Bonora M, Syberg S, Corigliano D, Pinton P, Jorgensen NR, Abelli L, Emionite L, Raffaghello L, Pistoia V, Di Virgilio F (2015) Accelerated tumor progression in mice lacking the ATP receptor P2X7. *Cancer Res* 75:635–644
87. Giuliani AL, Colognesi D, Ricco T, Roncato C, Capece M, Amoroso F, Wang QG, De Marchi E, Gartland A, Di Virgilio F, Adinolfi E (2014) Trophic activity of human P2X7 receptor isoforms A and B in osteosarcoma. *PLoS One* 9:e107224
88. Falk S, Schwab SD, Frosig-Jorgensen M, Clausen RP, Dickenson AH, Heegaard AM (2015) P2X7 receptor-mediated analgesia in cancer-induced bone pain. *Neuroscience* 291:93–105
89. Young CNJ, Chira N, Rog J, Al-Khalidi R, Benard M, Galas L, Chan P, Vaudry D, Zablocki K, Gorecki DC (2017) Sustained activation of P2X7 induces MMP-2-evoked cleavage and functional purinoceptor inhibition. *J Mol Cell Biol*. <https://doi.org/10.1093/jmcb/mjx030>
90. Loi S, Pommey S, Haibe-Kains B, Beavis PA, Darcy PK, Smyth MJ, Stagg J (2013) CD73 promotes anthracycline resistance and poor prognosis in triple negative breast cancer. *Proc Natl Acad Sci U S A* 110:11091–11096
91. Burnstock G (2015) Physiopathological roles of P2X receptors in the central nervous system. *Curr Med Chem* 22:819–844
92. Sperlagh B, Illes P (2014) P2X7 receptor: an emerging target in central nervous system diseases. *Trends Pharmacol Sci* 35:537–547
93. Cisneros-Mejorado A, Perez-Samartin A, Gottlieb M, Matute C (2015) ATP signaling in brain: release, excitotoxicity and potential therapeutic targets. *Cell Mol Neurobiol* 35:1–6
94. Pevarello P, Bovolenta S, Tarroni P, Za L, Severi E, Torino D, Vitalone R (2017) P2X7 antagonists for CNS indications: recent patent disclosures. *Pharm Pat Anal* 6:61–76
95. Monif M, Reid CA, Powell KL, Drummond KJ, O'Brien TJ, Williams DA (2016) Interleukin-1 β has trophic effects in microglia and its release is mediated by P2X7R pore. *J Neuroinflammation* 13:173
96. Metzger MW, Walser SM, Aprile-Garcia F, Dedic N, Chen A, Holsboer F, Arzt E, Wurst W, Deussing JM (2017) Genetically dissecting *P2rx7* expression within the central nervous system using conditional humanized mice. *Purinergic Signal* 13:153–170
97. Miras-Portugal MT, Diaz-Hernandez JJ, Gomez-Villafuertes R, Diaz-Hernandez M, Artalejo AR, Gualix J (2015) Role of P2X7 and P2Y₂ receptors on α -secretase-dependent APP processing: control of amyloid plaques formation "in vivo" by P2X7 receptor. *Comput Struct Biotechnol J* 13:176–181
98. Sáez-Orellana F, Godoy PA, Bastidas CY, Silva-Grecchi T, Guzmán L, Aguayo LG, Fuentealba J (2016) ATP leakage induces P2XR activation and contributes to acute synaptic excitotoxicity induced by soluble oligomers of β -amyloid peptide in hippocampal neurons. *Neuropharmacology* 100:116–123
99. Woods LT, Ajit D, Camden JM, Erb L, Weisman GA (2016) Purinergic receptors as potential therapeutic targets in Alzheimer's disease. *Neuropharmacology* 104:169–179
100. Chen X, Hu J, Jiang L, Xu S, Zheng B, Wang C, Zhang J, Wei X, Chang L, Wang Q (2014) Brilliant Blue G improves cognition in an animal model of Alzheimer's disease and inhibits amyloid- β -induced loss of filopodia and dendrite spines in hippocampal neurons. *Neuroscience* 279:94–101
101. Burnstock G (2017) Purinergic signalling and neurological diseases: an update *CNS Neurol Disord Drug Targets* 16:257–265
102. Carmo MR, Menezes AP, Nunes AC, Pliássova A, Rolo AP, Palmeira CM, Cunha RA, Canas PM, Andrade GM (2014) The P2X7 receptor antagonist Brilliant Blue G attenuates contralateral rotations in a rat model of Parkinsonism through a combined control of synaptotoxicity, neurotoxicity and gliosis. *Neuropharmacology* 81:142–152
103. Jörg M, Scammells PJ, Capuano B (2014) The dopamine D₂ and adenosine A_{2A} receptors: past, present and future trends for the treatment of Parkinson's disease. *Curr Med Chem* 21:3188–3210
104. Wang XH, Xie X, Luo XG, Shang H, He ZY (2017) Inhibiting purinergic P2X7 receptors with the antagonist brilliant blue G is neuroprotective in an intranigral lipopolysaccharide animal model of Parkinson's disease. *Mol Med Rep* 15:768–776
105. Burnstock G (2015) Purinergic signalling in neuroregeneration. *Neural Regeneration Research* 10:1919
106. BJ G, Field J, Dutertre S, Ou A, Kilpatrick TJ, Lechner-Scott J, Scott R, Lea R, Taylor BV, Stankovich J, Butzkueven H, Gresle M, Laws SM, Petrou S, Hoffjan S, Akkad DA, Graham CA, Hawkins S, Glaser A, Bedri SK, Hillert J, Matute C, Antiguiedad A, Wiley JS (2015) A rare P2X7 variant Arg307Gln with absent pore formation function protects against neuroinflammation in multiple sclerosis. *Hum Mol Genet* 24:5644–5654
107. Sadovnick AD, BJ G, Traboulsee AL, Bernales CQ, Encarnacion M, Yee IM, Criscuoli MG, Huang X, Ou A, Milligan CJ, Petrou S, Wiley JS, Vilarino-Guell C (2017) Purinergic receptors *P2RX4* and *P2RX7* in familial multiple sclerosis. *Hum Mutat* 38:736–744
108. Díaz-Hernández M, Domenico J, Sánchez-Nogueiro J, Gómez-Villafuertes R, Canals JM, Alberch J, Miras-Portugal MT, Lucas JJ (2009) Altered P2X7-receptor level and function in mouse models of Huntington's disease and therapeutic efficacy of antagonist administration. *FASEB J* 23:1893–1906
109. Cervetto C, Frattaroli D, Maura G, Marcoli M (2013) Motor neuron dysfunction in a mouse model of ALS: gender-dependent effect of P2X7 antagonism. *Toxicology* 311:69–77
110. Bartlett R, Sluyter V, Watson D, Sluyter R, Yerbury JJ (2017) P2X7 antagonism using Brilliant Blue G reduces body weight loss and prolongs survival in female SOD1G93A amyotrophic lateral sclerosis mice. *PeerJ* 5:e3064

111. Apolloni S, Amadio S, Parisi C, Matteucci A, Potenza RL, Armida M, Popoli P, D'Ambrosi N, Volonté C (2014) Spinal cord pathology is ameliorated by P2X7 antagonism in a SOD1-mutant mouse model of amyotrophic lateral sclerosis. *Dis Model Mech* 7:1101–1109
112. Apolloni S, Amadio S, Montilli C, Volonté C, D'Ambrosi N (2013) Ablation of P2X7 receptor exacerbates gliosis and motoneuron death in the SOD1-G93A mouse model of amyotrophic lateral sclerosis. *Hum Mol Genet* 22:4102–4116
113. Bhattacharya A, Biber K (2016) The microglial ATP-gated ion channel P2X7 as a CNS drug target. *Glia* 64:1772–1787
114. Krügel U (2016) Purinergic receptors in psychiatric disorders. *Neuropharmacology* 104:212–225
115. Avendano BC, Montero TD, Chávez CE, von Bernhardt R, Orellana JA (2015) Prenatal exposure to inflammatory conditions increases Cx43 and Panx1 unopposed channel opening and activation of astrocytes in the offspring effect on neuronal survival. *Glia* 63:2058–2072
116. Barron ML, Werry EL, McGregor IS, Kassiou M (2014) P2X7 in bipolar and depressive disorders. In: Weiss N, Koschak A (eds) *Pathologies of calcium channels*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp 635–661
117. Zhang K, Liu J, You X, Kong P, Song Y, Cao L, Yang S, Wang W, Fu Q, Ma Z (2016) P2X7 as a new target for chrysophanol to treat lipopolysaccharide-induced depression in mice. *Neurosci Lett* 613:60–65
118. Ma M, Ren Q, Zhang JC, Hashimoto K (2014) Effects of brilliant blue G on serum tumor necrosis factor- α levels and depression-like behavior in mice after lipopolysaccharide administration. *Clin Psychopharmacol Neurosci* 12:31–36
119. Iwata M, Ota KT, Li XY, Sakae F, Li N, Duthel S, Banasr M, Duric V, Yamanashi T, Kaneko K, Rasmussen K, Glasebrook A, Koester A, Song D, Jones KA, Zorn S, Smagin G, Duman RS (2016) Psychological stress activates the inflammasome via release of adenosine triphosphate and stimulation of the purinergic type 2X7 receptor. *Biol Psychiatry* 80:12–22
120. Catanzaro JM, Hueston CM, Deak MM, Deak T (2014) The impact of the P2X₇ receptor antagonist A-804598 on neuroimmune and behavioral consequences of stress. *Behav Pharmacol* 25:582–598
121. Aprile-Garcia F, Metzger MW, Paez-Pereda M, Stadler H, Acuña M, Liberman AC, Senin SA, Gerez J, Hoijman E, Refojo D, Mitkovski M, Panhuysen M, Stühmer W, Holsboer F, Deussing JM, Arzt E (2016) Co-Expression of wild-type P2X₇R with Gln460Arg variant alters receptor function. *PLoS One* 11:e0151862
122. Cisneros-Mejorado A, Gottlieb M, Cavaliere F, Magnus T, Koch-Nolte F, Scemes E, Pérez-Samartín A, Matute C (2015) Blockade of P2X7 receptors or pannexin-1 channels similarly attenuates postischemic damage. *J Cereb Blood Flow Metab* 35:843–850
123. Mahi N, Kumar A, Jaggi AS, Singh N, Dhawan R (2015) Possible role of pannexin 1/P₂X₇ purinoceptor in neuroprotective mechanism of ischemic postconditioning in mice. *J Surg Res* 196:190–199
124. Yue N, Huang H, Zhu X, Han Q, Wang Y, Li B, Liu Q, Wu G, Zhang Y, Yu J (2017) Activation of P2X7 receptor and NLRP3 inflammasome assembly in hippocampal glial cells mediates chronic stress-induced depressive-like behaviors. *J Neuroinflammation* 14:102
125. Murphy N, Cowley TR, Richardson JC, Virley D, Upton N, Walter D, Lynch MA (2012) The neuroprotective effect of a specific P2X₇ receptor antagonist derives from its ability to inhibit assembly of the NLRP3 inflammasome in glial cells. *Brain Pathol* 22:295–306
126. Xu P, Xu Y, Hu B, Wang J, Pan R, Murugan M, LJ W, Tang Y (2015) Extracellular ATP enhances radiation-induced brain injury through microglial activation and paracrine signaling via P2X7 receptor. *Brain Behav Immun* 50:87–100
127. Sun L, Gao J, Zhao M, Cui J, Li Y, Yang X, Jing X, Wu Z (2015) A novel cognitive impairment mechanism that astrocytic p-connexin 43 promotes neuronal autophagy via activation of P2X₇R and down-regulation of GLT-1 expression in the hippocampus following traumatic brain injury in rats. *Behav Brain Res* 291:315–324
128. Zhao H, Zhang X, Dai Z, Feng Y, Li Q, Zhang JH, Liu X, Chen Y, Feng H (2016) P2X7 receptor suppression preserves blood-brain barrier through inhibiting RhoA activation after experimental intracerebral hemorrhage in rats. *Sci Rep* 6:23286
129. Cieslak M, Wojtczak A, Komoszynski M (2017) Role of the purinergic signaling in epilepsy. *Pharmacol Rep* 69:130–138
130. Rodriguez-Alvarez N, Jimenez-Mateos EM, Engel T, Quinlan S, Reschke CR, Conroy RM, Bhattacharya A, Boylan GB, Henshall DC (2017) Effects of P2X7 receptor antagonists on hypoxia-induced neonatal seizures in mice. *Neuropharmacology* 116:351–363
131. Jimenez-Pacheco A, Diaz-Hernandez M, Arribas-Blázquez M, Sanz-Rodríguez A, Olivos-Oré LA, Artalejo AR, Alves M, Letavic M, Miras-Portugal MT, Conroy RM, Delanty N, Farrell MA, O'Brien DF, Bhattacharya A, Engel T, Henshall DC (2016) Transient P2X7 receptor antagonism produces lasting reductions in spontaneous seizures and gliosis in experimental temporal lobe epilepsy. *J Neurosci* 36:5920–5932
132. Fischer W, Franke H, Krügel U, Müller H, Dinkel K, Lord B, Letavic MA, Henshall DC, Engel T (2016) Critical evaluation of P2X7 receptor antagonists in selected seizure models. *PLoS One* 11:e0156468
133. Rozmer K, Gao P, Araujo MG, Khan MT, Liu J, Rong W, Tang Y, Franke H, Krügel U, Fernandes MJ, Illes P (2016) Pilocarpine-induced status epilepticus increases the sensitivity of P2X7 and P2Y1 receptors to nucleotides at neural progenitor cells of the juvenile rodent hippocampus. *Cereb Cortex* 27:3568–3585
134. Tu G, Zou L, Liu S, Wu B, Lv Q, Wang S, Xue Y, Zhang C, Yi Z, Zhang X, Li G, Liang S (2016) Long noncoding NONRATT021972 siRNA normalized abnormal sympathetic activity mediated by the upregulation of P2X7 receptor in superior cervical ganglia after myocardial ischemia. *Purinergic Signal* 12:521–535
135. Granado M, Amor S, Montoya JJ, Monge L, Fernández N, García-Villalón AL (2015) Altered expression of P2Y2 and P2X7 purinergic receptors in the isolated rat heart mediates ischemia-reperfusion injury. *Vascul Pharmacol* 73:96–103
136. Du D, Jiang M, Liu M, Wang J, Xia C, Guan R, Shen L, Ji Y, Zhu D (2015) Microglial P2X₇ receptor in the hypothalamic paraventricular nuclei contributes to sympathoexcitatory responses in acute myocardial infarction rat. *Neurosci Lett* 587:22–28
137. Martinez CG, Zamith-Miranda D, da Silva MG, Ribeiro KC, Brandão IT, Silva CL, Diaz BL, Bellio M, Persechini PM, Kurtenbach E (2015) P2x7 purinergic signaling in dilated cardiomyopathy induced by auto-immunity against muscarinic M2 receptors: autoantibody levels, heart functionality and cytokine expression. *Sci Rep* 5:16940
138. Zempo H, Sugita Y, Ogawa M, Watanabe R, Suzuki J, Isobe M (2015) A P2X7 receptor antagonist attenuates experimental autoimmune myocarditis via suppressed myocardial CD4⁺ T and macrophage infiltration and NADPH oxidase 2/4 expression in mice. *Heart Vessels* 30:527–533
139. Stachon P, Heidenreich A, Merz J, Hoppe N, Bode C, Idzko M, Zirlik A (2016) Deficiency of purinergic receptor P2X7 reduces atherosclerosis in mice. *Circulation* 134(Suppl_1):A16502
140. Luo W, Guth CM, Jolayemi O, Duvall CL, Brophy CM, Cheung-Flynn J (2016) Subfailure overstretch injury leads to reversible functional impairment and purinergic P2X7 receptor activation in intact vascular tissue. *Front Bioeng Biotechnol* 4:75
141. Gicquel T, Le Dare B, Boichot E, Lagente V (2017) Purinergic receptors: new targets for the treatment of gout and fibrosis. *Fundam Clin Pharmacol* 31:136–146

142. Monção-Ribeiro LC, Faffe DS, Santana PT, Vieira FS, da Graca CL, Marques-da-Silva C, Machado MN, Caruso-Neves C, Zin WA, Borojevic R, Takiya CM, Coutinho-Silva R (2014) P2X7 receptor modulates inflammatory and functional pulmonary changes induced by silica. *PLoS One* 9:e110185
143. Chen S, Zhu Z, Klebe D, Bian H, Krafft PR, Tang J, Zhang J, Zhang JH (2014) Role of P2X purinoceptor 7 in neurogenic pulmonary edema after subarachnoid hemorrhage in rats. *PLoS One* 9:e89042
144. Lucattelli M, Cicko S, Muller T, Lommatzsch M, De Cunto G, Cardini S, Sundas W, Grimm M, Zeiser R, Durk T, Zissel G, Sorichter S, Ferrari D, Di Virgilio F, Virchow JC, Lungarella G, Idzko M (2011) P2X₇ receptor signaling in the pathogenesis of smoke-induced lung inflammation and emphysema. *Am J Respir Cell Mol Biol* 44:423–429
145. Zheng X, Li T, Chen Y, Pan H, Zhang Z, Dai Y, Wang J (2017) Genetic polymorphisms of the P2X7 gene associated with susceptibility to and prognosis of pulmonary tuberculosis. *Infect Genet Evol* 53:24–29
146. Amaral EP, Ribeiro SC, Lanes VR, Almeida FM, de Andrade MR, Bomfim CC, Salles EM, Bortoluci KR, Coutinho-Silva R, Hirata MH, Alvarez JM, Lasunskiaia EB, D'Império-Lima MR (2014) Pulmonary infection with hypervirulent *Mycobacteria* reveals a crucial role for the P2X7 receptor in aggressive forms of tuberculosis. *PLoS Pathog* 10:e1004188
147. Moncao-Ribeiro LC, Cagido VR, Lima-Murad G, Santana PT, Riva DR, Borojevic R, Zin WA, Cavalcante MC, Rica I, Brando-Lima AC, Takiya CM, Faffe DS, Coutinho-Silva R (2011) Lipopolysaccharide-induced lung injury: role of P2X7 receptor. *Respir Physiol Neurobiol* 179:314–325
148. Hafner S, Wagner K, Weber S, Gröger M, Wepler M, McCook O, Scheuerle A, Stahl B, Huber-Lang M, Jung B, Calzia E, Georgieff M, Möller P, Frick M, Radermacher P, Wagner F (2017) Role of the purinergic receptor P2XR4 after blunt chest trauma in cigarette smoke-exposed mice. *Shock* 47:193–199
149. Antonioli L, Giron MC, Colucci R, Pellegrini C, Sacco D, Caputi V, Orso G, Tuccori M, Scarpignato C, Blandizzi C, Fornai M (2014) Involvement of the P2X7 purinergic receptor in colonic motor dysfunction associated with bowel inflammation in rats. *PLoS One* 9:e116253
150. Neves AR, Castelo-Branco MT, Figliuolo VR, Bernardazzi C, Buongusto F, Yoshimoto A, Nanini HF, Coutinho CM, Carneiro AJ, Coutinho-Silva R, de Souza HS (2014) Overexpression of ATP-activated P2X7 receptors in the intestinal mucosa is implicated in the pathogenesis of Crohn's disease. *Inflamm Bowel Dis* 20:444–457
151. Figliuolo VR, Savio LEB, Safya H, Nanini H, Bernardazzi C, Abalo A, de Souza HSP, Kanellopoulos J, Bobe P, Coutinho C, Coutinho-Silva R (2017) P2X7 receptor promotes intestinal inflammation in chemically induced colitis and triggers death of mucosal regulatory T cells. *Biochim Biophys Acta* 1863:1183–1194
152. Ma B, Jiang Q, Li W, Li Z (2016) Estrogen receptor beta (ER β) activation plays a therapeutic role in murine models of inflammatory bowel disease (IBD) via inhibiting P2X3 and P2X7 receptors. *The FASEB Journal* 30:1023–1023
153. Kurashima Y, Kiyono H, Kunisawa J (2015) Pathophysiological role of extracellular purinergic mediators in the control of intestinal inflammation. *Mediators Inflamm* 2015:427125
154. da Silva MV, Marosti AR, Mendes CE, Palombit K, Castelucci P (2015) Differential effects of experimental ulcerative colitis on P2X7 receptor expression in enteric neurons. *Histochem Cell Biol* 143:171–184
155. da Silva MV, Marosti AR, Mendes CE, Palombit K, Castelucci P (2017) Submucosal neurons and enteric glial cells expressing the P2X7 receptor in rat experimental colitis. *Acta Histochem* 119:481–494
156. Wan P, Liu X, Xiong Y, Ren Y, Chen J, Lu N, Guo Y, Bai A (2016) Extracellular ATP mediates inflammatory responses in colitis via P2 x 7 receptor signaling. *Sci Rep* 6:19108
157. Eser A, Colombel JF, Rutgeerts P, Vermeire S, Vogelsang H, Braddock M, Persson T, Reinisch W (2015) Safety and efficacy of an oral inhibitor of the purinergic receptor P2X7 in adult patients with moderately to severely active Crohn's disease: a randomized placebo-controlled, double-blind, phase IIa study. *Inflamm Bowel Dis* 21:2247–2253
158. Bhavsar S, Gade A, Kang M, Hauser KF, Dewey WL, Akbarali HI (2017) Connexin-purinergic signaling in enteric glia mediates the prolonged effect of morphine on constipation. *FASEB J* 31:2649–2660
159. Burnstock G, Evans L, Bailey M (2014) Purinergic signalling in the kidney in health and disease. *Purinergic Signalling* 10:71–101
160. Menzies RI, Tam FW, Unwin RJ, Bailey MA (2017) Purinergic signaling in kidney disease. *Kidney Int* 91:315–323
161. Yan Y, Bai J, Zhou X, Tang J, Jiang C, Tolbert E, Bayliss G, Gong R, Zhao TC, Zhuang S (2015) P2X7 receptor inhibition protects against ischemic acute kidney injury in mice. *Am J Physiol Cell Physiol* 308:C463–C472
162. Chang MY, JK L, Tian YC, Chen YC, Hung CC, Huang YH, Chen YH, MS W, Yang CW, Cheng YC (2011) Inhibition of the P2X7 receptor reduces cystogenesis in PKD. *J Am Soc Nephrol* 22:1696–1706
163. Turner CM, Tam F, Lai P-C, Tarzi RM, Burnstock G, Pusey CD, Cook HT, Unwin RJ (2007) Increased expression of the pro-apoptotic ATP-sensitive P2X₇ receptor in experimental and human glomerulonephritis. *Nephrology Dialysis Transplantation* 22:386–395
164. Zhao J, Wang H, Dai C, Wang H, Zhang H, Huang Y, Wang S, Gaskin F, Yang N, SM F (2013) P2X7 blockade attenuates lupus nephritis by inhibiting NLRP3/ASC/caspase-1 activation. *Arthritis Rheum* 65:3176–3185
165. Franco M, Bautista-Perez R, Cano-Martinez A, Pacheco U, Santamaria J, Del Valle-Mondragon L, Perez-Mendez O, Navar LG (2017) Physiopathological implications of P2X1 and P2X7 receptors in regulation of glomerular hemodynamics in angiotensin II-induced hypertension. *Am J Physiol Renal Physiol*:ajprenal.00663.02016
166. Menzies RI, Unwin RJ, Bailey MA (2015) Renal P2 receptors and hypertension. *Acta Physiol (Oxf)* 213:232–241
167. Ji X, Naito Y, Hirokawa G, Weng H, Hiura Y, Takahashi R, Iwai N (2012) P2X₇ receptor antagonism attenuates the hypertension and renal injury in Dahl salt-sensitive rats. *Hypertens Res* 35:173–179
168. Solini A, Menini S, Rossi C, Ricci C, Santini E, Blasetti Fantauzzi C, Iacobini C, Pugliese G (2013) The purinergic 2X7 receptor participates in renal inflammation and injury induced by high-fat diet: possible role of NLRP3 inflammasome activation. *J Pathol* 231:342–353
169. Munoz A, Yazdi IK, Tang X, Rivera C, Taghipour N, Grossman RG, Boone TB, Tasciotti E (2017) Localized inhibition of P2X7R at the spinal cord injury site improves neurogenic bladder dysfunction by decreasing urothelial P2X3R expression in rats. *Life Sci* 171:60–67
170. Martins JP, Silva RB, Coutinho-Silva R, Takiya CM, Battastini AM, Morrone FB, Campos MM (2012) The role of P2X7 purinergic receptors in inflammatory and nociceptive changes accompanying cyclophosphamide-induced haemorrhagic cystitis in mice. *Br J Pharmacol* 165:183–196
171. Gonçalves RG, Gabrich L, Rosário A Jr, Takiya CM, Ferreira ML, Chiarini LB, Persechini PM, Coutinho-Silva R, Leite M Jr (2006) The role of purinergic P2X₇ receptors in the inflammation and fibrosis of unilateral ureteral obstruction in mice. *Kidney Int* 70:1599–1606
172. Xiao H, Si LY, Liu W, Li N, Meng G, Yang N, Chen X, Zhou YG, Shen HY (2013) The effects of adenosine A_{2A} receptor knockout on renal interstitial fibrosis in a mouse model of unilateral ureteral obstruction. *Acta Histochem* 115:315–319

173. Amaral SS, Oliveira AG, Marques PE, Quintão JL, Pires DA, Resende RR, Sousa BR, Melgaco JG, Pinto MA, Russo RC, Gomes AK, Andrade LM, Zanin RF, Pereira RV, Bonorino C, Soriani FM, Lima CX, Cara DC, Teixeira MM, Leite MF, Menezes GB (2013) Altered responsiveness to extracellular ATP enhances acetaminophen hepatotoxicity. *Cell Commun Signal* 11:10
174. Hoque R, Sohail MA, Salhanick S, Malik AF, Ghani A, Robson SC, Mehal WZ (2012) P2X7 receptor-mediated purinergic signaling promotes liver injury in acetaminophen hepatotoxicity in mice. *Am J Physiol Gastrointest Liver Physiol* 302:G1171–G1179
175. Xie Y, Williams CD, McGill MR, Lebofsky M, Ramachandran A, Jaeschke H (2013) Purinergic receptor antagonist A438079 protects against acetaminophen-induced liver injury by inhibiting P450 isoenzymes, not by inflammasome activation. *Toxicol Sci* 131:325–335
176. Jiang S, Zhang Y, Zheng JH, Li X, Yao YL, YL W, Song SZ, Sun P, Nan JX, Lian LH (2017) Potentiation of hepatic stellate cell activation by extracellular ATP is dependent on P2X7R-mediated NLRP3 inflammasome activation. *Pharmacol Res* 117:82–93
177. Kawamura H, Aswad F, Minagawa M, Govindarajan S, Dennert G (2006) P2X7 receptors regulate NKT cells in autoimmune hepatitis. *J Immunol* 176:2152–2160
178. Chandrashekar V, Das S, Seth RK, Dattaroy D, Alhasson F, Michelotti G, Nagarkatti M, Nagarkatti P, Diehl AM, Chatterjee S (2016) Purinergic receptor X7 mediates leptin induced GLUT4 function in stellate cells in nonalcoholic steatohepatitis. *Biochim Biophys Acta* 1862:32–45
179. Geraghty NJ, Watson D, Adhikary SR, Sluyter R (2017) P2X7 receptor in skin biology and diseases. *World J Dermatol* 5:72–83
180. Inoue K, Hosoi J, Denda M (2007) Extracellular ATP has stimulatory effects on the expression and release of IL-6 via purinergic receptors in normal human epidermal keratinocytes. *J Invest Dermatol* 127:362–371
181. Lioi AB, Ferrari BM, Dubyak GR, Weinberg A, Siegf SF (2015) Human β defensin-3 increases CD86 expression on monocytes by activating the ATP-gated channel P2X7. *J Immunol* 195:4438–4445
182. Killeen ME, Ferris L, Kupetsky EA, Falo L Jr, Mathers AR (2013) Signaling through purinergic receptors for ATP induces human cutaneous innate and adaptive Th17 responses: implications in the pathogenesis of psoriasis. *J Immunol* 190:4324–4336
183. Greig AVH, Cuthill S, Linge C, Clayton E, Burnstock G (2006) P2X₅ and P2X₇ receptors in human warts and CIN 612 organotypic raft cultures of human papillomavirus infected keratinocytes. *Purinergic Signalling* 2:509–515
184. Weber FC, Esser PR, Müller T, Ganesan J, Pellegratti P, Simon MM, Zeiser R, Idzko M, Jakob T, Martin SF (2010) Lack of the purinergic receptor P2X₇ results in resistance to contact hypersensitivity. *J Exp Med* 207:2609–2619
185. Zanin RF, da Silva GL, Erig T, Sperotto ND, Leite CE, Coutinho-Silva R, Batastini AM, Morrone FB (2015) Decrease of serum adenosine nucleotide hydrolysis in an irritant contact dermatitis mice model: potential P2X7R involvement. *Mol Cell Biochem* 404:221–228
186. Gazzo E, Baldassari S, Assereto S, Fruscione F, Pistorio A, Panicucci C, Volpi S, Perruzza L, Fiorillo C, Minetti C, Traggiai E, Grassi F, Bruno C (2015) Enhancement of muscle T regulatory cells and improvement of muscular dystrophic process in *mdx* mice by blockade of extracellular ATP/P2X axis. *Am J Pathol* 185:3349–3360
187. Sinadinos A, Young CN, Al-Khalidi R, Teti A, Kalinski P, Mohamad S, Floriot L, Henry T, Tozzi G, Jiang T, Wurtz O, Lefebvre A, Shugay M, Tong J, Vaudry D, Arkle S, doRego JC, Górecki DC (2015) P2RX7 purinoceptor: a therapeutic target for ameliorating the symptoms of duchenne muscular dystrophy. *PLoS Med* 12:e1001888
188. Kvist TM, Schwarz P, Jørgensen NR The P2X7 receptor: a key player in immune-mediated bone loss? *Scic World J* 2014, 2014:10
189. Varley I, Greeves JP, Sale C, Friedman E, Moran DS, Yanovich R, Wilson PJ, Gartland A, Hughes DC, Stellingwerff T, Ranson C, Fraser WD, Gallagher JA (2016) Functional polymorphisms in the P2X7 receptor gene are associated with stress fracture injury. *Purinergic Signalling* 12:103–113
190. Agrawal A, Buckley KA, Bowers K, Furber M, Gallagher JA, Gartland A (2010) The effects of P2X7 receptor antagonists on the formation and function of human osteoclasts in vitro. *Purinergic Signal* 6:307–315
191. Jørgensen NR, Boeynaems JM, Di Virgilio F (2011) European meeting "P2 receptors: new targets for the treatment of osteoporosis". *Purinergic Signal* 7:275–276
192. Mrazek F, Gallo J, Stahelova A, Petrek M (2010) Functional variants of the *P2RX7* gene, aseptic osteolysis, and revision of the total hip arthroplasty: a preliminary study. *Hum Immunol* 71:201–205
193. Grol MW, Brooks PJ, Pereverzev A, Dixon SJ (2016) P2X7 nucleotide receptor signaling potentiates the Wnt/ β -catenin pathway in cells of the osteoblast lineage. *Purinergic Signal* 12:509–520
194. Hu H, Yang B, Li Y, Zhang S, Li Z (2016) Blocking of the P2X7 receptor inhibits the activation of the MMP-13 and NF- κ B pathways in the cartilage tissue of rats with osteoarthritis. *Int J Mol Med* 38:1922–1932
195. Portales-Cervantes L, Niño-Moreno P, Salgado-Bustamante M, García-Hernández MH, Baranda-Candido L, Reynaga-Hernández E, Barajas-López C, González-Amaro R, Portales-Pérez DP (2012) The His155Tyr (489C>T) single nucleotide polymorphism of *P2RX7* gene confers an enhanced function of P2X7 receptor in immune cells from patients with rheumatoid arthritis. *Cell Immunol* 276:168–175
196. Alqallaf SM, Evans BA, Kidd EJ (2009) Atypical P2X receptor pharmacology in two human osteoblast-like cell lines. *Br J Pharmacol* 156:1124–1135
197. Beswick PJ, Billinton A, Chambers LJ, Dean DK, Fonfria E, Gleave RJ, Medhurst SJ, Michel AD, Moses AP, Patel S, Roman SA, Roomans S, Senger S, Stevens AJ, Walter DS (2010) Structure-activity relationships and in vivo activity of (1*H*-pyrazol-4-yl)acetamide antagonists of the P2X₇ receptor. *Bioorg Med Chem Lett* 20:4653–4656
198. Sociali G, Visigalli D, Prukop T, Cervellini I, Mannino E, Venturi C, Bruzzone S, Sereda MW, Schenone A (2016) Tolerability and efficacy study of P2X7 inhibition in experimental Charcot-Marie-Tooth type 1A (CMT1A) neuropathy. *Neurobiol Dis* 95:145–157
199. Bartlett R, Stokes L, Sluyter R (2014) The P2X7 receptor channel: recent developments and the use of P2X7 antagonists in models of disease. *Pharmacol Rev* 66:638–675
200. Vieira FS, Nanini HF, Takiya CM, Coutinho-Silva R (2016) P2X7 receptor knockout prevents streptozotocin-induced type 1 diabetes in mice. *Mol Cell Endocrinol* 419:148–157
201. Seref-Ferlenguez Z, Maung S, Schaffler MB, Spray DC, Suadcani SO, Thi MM (2016) P2X7R-Panx1 complex impairs bone mechanosignaling under high glucose levels associated with type-1 diabetes. *PLoS One* 11:e0155107
202. Coutinho-Silva R, Parsons M, Robson T, Lincoln J, Burnstock G (2003) P2X and P2Y purinoceptor expression in pancreas from streptozotocin-diabetic rats. *Molecular and Cellular Endocrinology* 204:141–154
203. Vergani A, Fotino C, D'Addio F, Tezza S, Podetta M, Gatti F, Chin M, Bassi R, Molano RD, Corradi D, Gatti R, Ferrero ME, Secchi A, Grassi F, Ricordi C, Sayegh MH, Maffi P, Pileggi A, Fiorina P (2013) Effect of the purinergic inhibitor oxidized ATP in a model of islet allograft rejection. *Diabetes* 62:1665–1675
204. Aikin RA (2013) Targeting purinergic receptors in islet transplantation. *Diabetes* 62:1394–1395
205. Liu S, Zou L, Xie J, Xie W, Wen S, Xie Q, Gao Y, Li G, Zhang C, Xu C, Xu H, Wu B, Lv Q, Zhang X, Wang S, Xue Y, Liang S (2016) LncRNA NONRATT021972 siRNA regulates neuropathic

- pain behaviors in type 2 diabetic rats through the P2X₇ receptor in dorsal root ganglia. *Mol Brain* 9:44
206. Liu W, Ao Q, Guo Q, He W, Peng L, Jiang J, Hu X (2017) miR-9 mediates CALHM1-activated ATP-P2X₇R signal in painful diabetic neuropathy rats. *Mol Neurobiol* 54:922–929
 207. Wu B, Zhang C, Zou L, Ma Y, Huang K, Lv Q, Zhang X, Wang S, Xue Y, Yi Z, Jia T, Zhao S, Liu S, Xu H, Li G, Liang S (2016) LncRNA uc.48+ siRNA improved diabetic sympathetic neuropathy in type 2 diabetic rats mediated by P2X₇ receptor in SCG. *Auton Neurosci* 197:14–18
 208. Ursu D, Ebert P, Langron E, Ruble C, Munsie L, Zou W, Fijal B, Qian YW, McNearney TA, Mogg A, Grubisha O, Merchant K, Sher E (2014) Gain and loss of function of P2X₇ receptors: mechanisms, pharmacology and relevance to diabetic neuropathic pain. *Mol Pain* 10:37
 209. Rodrigues AM, Bergamaschi CT, Fernandes MJ, Paredes-Gamero EJ, Buri MV, Ferreira AT, Araujo SR, Punaro GR, Maciel FR, Nogueira GB, Higa EM (2014) P2X₇ receptor in the kidneys of diabetic rats submitted to aerobic training or to N-acetylcysteine supplementation [corrected]. *PLoS One* 9:e97452
 210. Caseley EA, Muench SP, Roger S, Mao HJ, Baldwin SA, Jiang LH (2014) Non-synonymous single nucleotide polymorphisms in the P2X receptor genes: association with diseases, impact on receptor functions and potential use as diagnosis biomarkers. *Int J Mol Sci* 15:13344–13371
 211. Barniol-Xicota M, Kwak SH, Lee SD, Caseley E, Valverde E, Jiang LH, Kim YC, Vázquez S (2017) Escape from adamantane: scaffold optimization of novel P2X₇ antagonists featuring complex polycycles. *Bioorg Med Chem Lett* 27:759–763
 212. Müller CE (2015) Medicinal chemistry of P2X receptors: allosteric modulators. *Curr Med Chem* 22:929–941
 213. Park JH, Kim YC (2017) P2X₇ receptor antagonists: a patent review (2010–2015). *Expert Opin Ther Pat* 27:257–267
 214. Burnstock G (2006) Pathophysiology and therapeutic potential of purinergic signaling. *Pharmacological Reviews* 58:58–86
 215. Burnstock G (2007) Physiology and pathophysiology of purinergic neurotransmission. *Physiol Rev* 87:659–797
 216. Zhang Y, Zhang S, Ding Z (2017) Role of P2Y₁₂ receptor in thrombosis. In: Islam MS (ed) *Thrombosis and embolism: from research to clinical practice: volume 1*. Springer International Publishing, Cham, pp 307–324. https://doi.org/10.1007/5584_2016_123
 217. Ralevic V, Burnstock G (1998) Receptors for purines and pyrimidines. *Pharmacol Rev* 50:413–492
 218. Matsumoto Y, Ohashi Y, Watanabe H, Tsubota K (2012) Efficacy and safety of diquafosol ophthalmic solution in patients with dry eye syndrome: a Japanese phase 2 clinical trial. *Ophthalmology* 119:1954–1960
 219. Chen CC, Akopian AN, Sivilotti L, Colquhoun D, Burnstock G, Wood JN (1995) A P2X purinoceptor expressed by a subset of sensory neurons. *Nature* 377:428–431
 220. Burnstock G (2016) Purinergic receptors and pain—an update. *Frontiers in Medicinal Chemistry* 9:3–55
 221. Gever JR, Rothschild S, Henningsen R, Martin R, Hackos D, Panicker S, Rubas W, Oglesby I, Dillon MP, Milla ME, Burnstock G, Ford APDW (2010) AF-353, a novel, potent orally bioavailable P2X₃/P2X_{2/3} receptor antagonist. *Br J Pharmacol* 160:1387–1398
 222. Abdulqawi R, Dockry R, Holt K, Layton G, McCarthy BG, Ford AP, Smith JA (2015) P2X₃ receptor antagonist (AF-219) in refractory chronic cough: a randomised, double-blind, placebo-controlled phase 2 study. *Lancet* 385:1198–1205
 223. McHugh SM, Roman S, Davis B, Koch A, Pickett AM, Richardson JC, Miller SR, Wetten S, Cox CJ, Karpe F, Todd JA, Bullmore ET (2012) Effects of genetic variation in the P2RX₇ gene on pharmacodynamics of a P2X₇ receptor antagonist: a prospective genotyping approach. *Br J Clin Pharmacol* 74:376–380
 224. Allsopp RC, Dayl S, Schmid R, Evans RJ (2017) Unique residues in the ATP gated human P2X₇ receptor define a novel allosteric binding pocket for the selective antagonist AZ10606120. *Sci Rep* 7:725
 225. Karasawa A, Kawate T (2016) Structural basis for subtype-specific inhibition of the P2X₇ receptor. *Elife* 5:e22153
 226. Kasuya G, Yamaura T, Ma XB, Nakamura R, Takemoto M, Nagumo H, Tanaka E, Dohmae N, Nakane T, Yu Y, Ishitani R, Matsuzaki O, Hattori M, Nureki O (2017) Structural insights into the competitive inhibition of the ATP-gated P2X receptor channel. *Nat Commun* 8:876