

Article

Antimicrobial Activity of Extracts from the *Humiria balsamifera* (Aubl)

Edelson de J. S. Dias ^{1,*}, Antônio J. Cantanhede Filho ¹, Fernando J. C. Carneiro ¹, Cláudia Q. da Rocha ², Luís Cláudio N. da Silva ³, Joice C. B. Santos ³, Thayná F. Barros ³ and Deivid M. Santos ³

¹ Chemistry Graduate Program, Federal Institute of Education, Science and Technology of Maranhão, Avenida Getúlio Vargas 04, Monte Castelo, São Luís 65030-005, MA, Brazil; prof.antoniofilho@ifma.edu.br (A.J.C.F.); fernandocarneiro@ifma.edu.br (F.J.C.C.)

² Chemistry Graduate Program, Federal University of Maranhão, Av. dos Portugueses, 1966—Vila Bacanga, São Luís 65080-805, MA, Brazil; rocha.claudia@ufma.br

³ Graduate Program in Microbial Biology, CEUMA University, Rua Josué Montello, 1—Renascença II, São Luís 65075-120, MA, Brazil; luiscln.silva@ceuma.br (L.C.N.d.S.); joic.cast@hotmail.com (J.C.B.S.); thata.fernandesbarros@hotmail.com (T.F.B.); deivid.martinss98@gmail.com (D.M.S.)

* Correspondence: edelson.sa@acad.ifma.edu.br; Tel.: +55-016-99419-2922

Abstract: *Humiria balsamifera* (Aubl), commonly known as “mirim”, is a plant of the Humiriaceae family, which consists of 39 species divided between eight genera: *Duckesia*, *Endopleura*, *Humiria*, *Humiriastrum*, *Hylocara*, *Sacoglottis*, *Schistostemon*, and *Vantenea*. This study aimed to characterize *H. balsamifera* extracts by LC-MS/MS and evaluate their antimicrobial potential through in vitro and in vivo assays. The leaves and stem bark of *H. balsamifera* were collected and dried at room temperature and then ground in a knife mill. The extracts were prepared with organic solvents in order to increase the polarity index (hexane, ethyl acetate, and methanol). The antimicrobial effects of these extracts were evaluated against the following bacterial strains: *Escherichia coli* ATCC 25922, *Listeria monocytogenes* ATCC 15313, *Salmonella enterica* Typhimurium ATCC 14028, and *Staphylococcus aureus* ATCC 6538. The best activity was observed in the ethyl acetate (EALE = 780 µg/mL), methanol (MLE = 780 µg/mL), and hexane (HLE = 1560 µg/mL) leaf extracts against *S. aureus*. Considering the results for both antimicrobial and antibiofilm activities, the EALE extract was chosen to proceed to the infection assays, which used *Tenebrio molitor* larvae. The EALE treatment was able to extend the average lifespan of the larvae (6.5 days) in comparison to *S. aureus*-infected larvae (1 day). Next, the samples were characterized by High-Performance Liquid Chromatography coupled to a mass spectrometer, allowing the identification of 11 substances, including seven flavonoids, substances whose antimicrobial activity is already well-reported in the literature. The number of bioactive compounds found in the chemical composition of *H. balsamifera* emphasizes its significance in both traditional medicine and scientific research that studies new treatments based on substances from the Brazilian flora.

Keywords: Humiriaceae; *Humiria*; antimicrobial; microorganisms; flavonoids



Citation: Dias, E.d.J.S.; Cantanhede Filho, A.J.; Carneiro, F.J.C.; da Rocha, C.Q.; da Silva, L.C.N.; Santos, J.C.B.; Barros, T.F.; Santos, D.M. Antimicrobial Activity of Extracts from the *Humiria balsamifera* (Aubl). *Plants* **2021**, *10*, 1479. <https://doi.org/10.3390/plants10071479>

Academic Editors: Yun-Soo Seo and Joong-Sun Kim

Received: 9 June 2021

Accepted: 12 July 2021

Published: 19 July 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Microorganisms are naturally well-spread out in the environment, and they can easily reach surfaces people come into contact with, including food products, whether at the harvest, slaughter, processing, or even packaging. Once in contact with the food, they start their growth process by consuming nutrients and causing the product to deteriorate [1–3].

Bacteria, fungi, viruses, and protozoa are the main microorganisms responsible for food contamination, infecting humans through the consumption of beef, fish, poultry, eggs, unhygienic fruits and fresh produce, causing a variety of diseases [4,5]. The World Health Organization estimates that one in 10 people worldwide become ill after consuming

contaminated food and about 420,000 people die each year, resulting in the loss of 33 million healthy life years (DALYs) [2,6].

Bacteria represent an added concern for health and food safety organizations, especially those able to grow at low temperatures and resist a wide range of temperature variations [7,8]. Bacterial pathogens such as *Escherichia coli*, *Salmonella enterica*, *Listeria monocytogenes*, and *Staphylococcus aureus*, among others, are responsible for several global foodborne outbreaks and cause life-threatening illnesses such as diarrheal diseases [5,9–12]. Another problem in fighting bacteria is their ability to develop resistance to conventional antimicrobials. These pathogens can use various strategies to inhibit the effects of antimicrobials, such as the production of inactivating enzymes, reduction of outer membrane permeability, efflux system, and blocking or altering the target site of antibiotics, further motivating the research focused on finding alternative ways to combat them [13,14].

In the search for new effective substances against resistant pathogens, several secondary metabolites from plants and endophytic microorganisms have shown promise [15,16]. Most of the drugs used in general today were developed based on ethnopharmacological knowledge [17–19], indicating that the chemistry of natural products is a big ally in the development of therapeutic agents [20,21].

The plant species *Humiria balsamifera* (Aubl), popularly known as “mirim”, presents interesting biological activities. The literature reports, most of all, anti-inflammatory [22,23], antimalarial [24], antioxidant [25,26], and antifungal activity [27], highlighting the therapeutic potential of this plant. Some substances isolated from this species so far have already been reported as well, such as bergenin, arjunolic acid, friedelin, lupeol, phytol, caryophyllene oxide, epoxide humulene, and trans-isolongifolanone, among others [24]. However, the antibacterial and antibiofilm activities of its derived products have not been extensively examined. Thus, this work aims to characterize and evaluate the effectiveness of *H. balsamifera* extracts in terms of the antimicrobial and antibiofilm activities against foodborne pathogens (*Escherichia coli* ATCC 25922, *Listeria monocytogenes* ATCC 15313, *Salmonella enterica* Typhimurium ATCC 14028, and *Staphylococcus aureus* ATCC 6538). The *in vivo* antimicrobial action of the most active extract was analyzed using a method based on the infection of *Tenebrio molitor* larvae.

2. Results

2.1. Antimicrobial Activity Evaluation

The antimicrobial activity of *Humiria balsamifera* (Aubl) leaf and stem bark extracts was evaluated by the determination of their minimum inhibitory concentrations (MIC) against four foodborne bacteria species: *E. coli*, *L. monocytogenes*, *S. enterica* Typhimurium, and *S. aureus* (Table 1).

Table 1. Determination of the minimum inhibitory concentrations (MIC) of the leaf and stem bark extracts of *Humiria balsamifera* (Aubl).

Bacteria Species	HSBE	EASBE	MSBE	HLE	EALE	MLE
<i>E. coli</i> ATCC 25922	>12,500	>12,500	>12,500	>12,500	>12,500	>12,500
<i>L. monocytogenes</i> ATCC 6538	>12,500	>12,500	>12,500	>12,500	3120	3120
<i>S. aureus</i> ATCC 6538	>12,500	>12,500	>12,500	1560	780	780
<i>S. enterica</i> Typhimurium ATCC 14028	>12,500	>12,500	>12,500	>12,500	>12,500	>12,500

Footnote: HSBE = hexane stem bark extract, EASBE = ethyl acetate stem bark extract, MSBE = methanol stem bark extract, HLE = hexane leaf extract, EALE = ethyl acetate leaf extract, and MLE = methanol leaf extract. MIC values are expressed in µg/mL.

Stem bark extracts did not exhibit antimicrobial action at any of the concentrations tested (MIC > 1250 µg/mL). However, the leaf extracts successfully inhibited *S. aureus*, with MIC = 780 µg/mL (EALE and MLE) and 1560 µg/mL (HLE). The EALE and the MLE also inhibited *L. monocytogenes* (MIC = 3120 µg/mL). The leaf extracts presented no action against the Gram-negative bacteria tested in this study.

2.2. Evaluation of the Antibiofilm Activity of *Humiria balsamifera* (Aubl) Extracts

Since the leaf extracts of *Humiria balsamifera* presented better inhibition results against *S. aureus*, their antibiofilm action at subinhibitory concentrations ($0.5 \times \text{MIC}$, $0.25 \times \text{MIC}$, $0.125 \times \text{MIC}$, and $0.0625 \times \text{MIC}$) was also evaluated (Figure 1). The EALE and the HLE reduced biofilm production by *S. aureus* by nearly 25% at concentrations higher than $390 \mu\text{g/mL}$ (Figure 1A,B). The MLE did not exhibit significant antibiofilm activity (Figure 1C). Considering the results for the antimicrobial and antibiofilm activities, the EALE was chosen to proceed to the in vivo tests, using *Tenebrio molitor* larvae.




Figure 1. Inhibition of the biofilm formation in *Staphylococcus aureus* by *Humiria balsamifera* (Aubl) extracts. (A) HLE = hexane leaf extract, (B) EALE = ethyl acetate leaf extract, and (C) MLE = methanol leaf extract. (*) Significant differences ($p < 0.05$) in relation to untreated biofilm (C+).

2.3. Evaluation of the In Vivo Activity of the Ethyl Acetate Leaf Extract of *Humiria balsamifera* (Aubl)

To evaluate the antimicrobial efficacy of the ethyl acetate leaf extract (EALE), the selected method was an alternative infection model based on the *S. aureus* ability to infect *T. molitor* larvae (Figure 2). The group infected with a lethal dose of *S. aureus* with no treatment presented an average lifespan of 1 day. In contrast, the uninfected larvae inoculated with the extract or its vehicle did not show a decrease in their lifespan. The EALE treatment was able to extend the average lifespan of the larvae (6.5 days), and by the end of the evaluation period, 50% of all larvae in this group were still alive.




Figure 2. Effect of the *Humiria balsamifera* (Aubl) EALE treatment on the lifespan of *Tenebrio molitor* larvae during infection by *S. aureus* ATCC 6538. The larvae were treated with EALE at a dose of 3.12 mg/kg. Negative control groups injected with PBS (phosphate-buffered saline) were also included.

2.4. Chemical Characterization of *Humiria balsamifera* (Aubl) Leaf and Stem Bark Extracts

Analyses of the leaf and stem bark extracts of *Humiria balsamifera* (Aubl) by HPLC-ESI-IT/MS in negative-ion mode identified 11 molecular ions (Tables 2–4). Their structures were proposed (Figure 3) based on the fragments originated from the molecular ion by multi-stage mass spectrometry (MSⁿ). The mass spectrometry ionization source was the electrospray (ESI). The ESI source may not have ionized the compounds like steroids and triterpenes. It was possible to identify only phenolic compounds in the extracts. From the 11 identified substances, seven were flavonoids (galocatechin, kaempferol 3-neohesperidoside, sophoricoside, quercetin 3-arabinoside, quercetin-*O*-rhamnoside, kaempferol-dirhamnoside, and vitexin-dirhamnoside); three were coumarins (bergenin and two derivatives: galloylbergenin and hydroxybenzoyl bergenin); and one was an oligosaccharide (maltotetraose).

Table 2. Identification of the substances present in the ethyl acetate stem bark extract of *Humiria balsamifera* (Aubl).

RT (min)	[M-H]	MSn	Proposed Substance	Reference
1.97	665	664, 305	Maltotetraose	[28]
2.85	327	249, 234, 207	Bergenin	[29]
3.58	479	327, 249, 234, 207	Galloylbergenin	[29]
4.87	463	327, 249, 234, 207	Hydroxybenzoyl bergenin	[29]

Table 3. Identification of the substances present in the *Humiria balsamifera* (Aubl) methanol stem bark extract.

RT (min)	[M-H]	MSn	Proposed Substance	Reference
3.01	305	179, 165	Galocatechin	[30]
3.81	327	234, 207, 192	Bergenin	[29]
6.35	593	431, 411	Kaempferol 3-neohesperidoside	[31]

Table 4. Identification of the substances present in the *Humiria balsamifera* (Aubl) ethyl acetate leaf extract.

RT (min)	[M-H]	MSn	Proposed Substance	Reference
2.83	327	249, 234, 207, 192	Bergenin	[29]
5.27	431	311, 283, 341	Sophoricoside	[32]
6.09	433	300, 271, 243, 227	Quercetin 3-arabinoside	[33]
6.46	447	300, 271, 243	Quercetin- <i>O</i> -rhamnoside	[33]
7.17	577	431, 413, 341, 300	Kaempferol-dirhamnoside	[33]
8.38	561	431, 293	Vitexin-dirhamnoside	[32]




Figure 3. Chemical structures of the substances identified in the extracts of *Humiria balsamifera* (Aubl) by HPLC-ESI-IT/MS using the MassBank Spectral DataBase. Maltotetraose (1); Bergenin (2); Galloylbergenin (3); Hydroxybenzoyl bergenin (4); Gallic acid (5); Kaempferol 3-O-neohesperidoside (6); Sophoric acid (7); Quercetin 3-O-arabinoside (8); Quercetin-3-O-rhamnoside (9); Kaempferol-3-O-dirhamnoside (10); and Vitexin-3-O-dirhamnoside (11).

Maltotetraose (1) presented a molecular ion of $m/z = 665$, with a fragment of $m/z = 664$ after the loss of a proton. In the third stage of the fragmentation process, the loss of 341 Da as $C_{12}H_{21}O_{11}$ and 18 Da as a water molecule produced a fragment of $m/z = 305$.

Bergenin (2) originated four ionic fragments: from an initial loss of 60 Da as $C_2H_4O_2$, and 18 Da as a water molecule, resulted the fragment $C_{12}H_{10}O_6^-$ ($m/z = 249$); the second one was produced from the loss of 93 Da as $C_2H_4O_2$, a methyl group, and a water molecule, resulting in the ion $C_{11}H_7O_6^-$ ($m/z = 234$); the third fragment $C_{10}H_8O_5^-$ ($m/z = 207$) was the result of the loss of 120 Da as $C_4H_8O_4$; and finally, the loss of 135 Da as $C_4H_8O_4$ and a methyl group led to the fourth fragment, $C_9H_5O_5^-$ ($m/z = 192$).

All these ions were also present in the spectra of the bergenin derivatives, plus one other fragment, observed in both spectra.

The fragment $C_{14}H_{15}O_9^-$ ($m/z = 327$) was observed in both spectra—galloylbergenin (3) and hydroxybenzoyl bergenin (4). For galloylbergenin, this fragment was a result of the loss of 152 Da as the galloyl group. For hydroxybenzoyl bergenin, this fragment resulted from the elimination of the hydroxybenzoyl group (136 Da).

Gallic acid (5) produced two fragment ions: $C_9H_8O_4^-$ ($m/z = 179$), resulting from the loss of 126 Da as $C_6H_6O_3$, and $C_8H_8O_4$ ($m/z = 165$) from the loss of 140 Da as $C_7H_8O_3$.

Kaempferol 3-O-neohesperidoside (6) also produced only two fragment ions: $C_{21}H_{19}O_{10}^-$ ($m/z = 431$), generated by the loss of 162 Da as $C_6H_{10}O_5$, and $C_{21}H_{15}O_9^-$ ($m/z = 411$), originated by the ion $C_{21}H_{19}O_{10}^-$, after a water loss. Bergenin (2) was also identified.

Sophoric acid (7) originated three ions: $C_{21}H_{19}O_{10}^-$ ($m/z = 431$), as a result of the loss of 90 Da as $C_3H_6O_3$, $C_{17}H_{11}O_6^-$ ($m/z = 311$) from the loss of 120 Da as $C_4H_8O_4^-$, and from $C_{17}H_{11}O_6^-$, the third fragment, $C_{16}H_{11}O_5^-$ ($m/z = 283$), was formed as a result of the loss of 28 Da as a carbon monoxide molecule.

In the quercetin 3-arabinoside spectrum (8), four fragment ions were observed: $C_{15}H_9O_7^-$ ($m/z = 300$) from the loss of 133 Da as $C_5H_9O_4$, $C_{14}H_7O_6^-$ ($m/z = 271$) from the loss of 162 Da as $C_6H_{10}O_5$, and the fragments $C_{13}H_7O_5^-$ ($m/z = 243$) and $C_{13}H_7O_4^-$ ($m/z = 227$), resulting from the loss of a carbon monoxide and dioxide, respectively.

The fragmentation of quercetin-*O*-rhamnoside (9) were similar to what was observed for quercetin 3-arabinoside. The loss of a rhamnose molecule, $C_6H_{11}O_4$ ($m/z = 147$ Da), led to the fragment ion $C_{15}H_9O_7^-$ ($m/z = 300$). Then, the loss of carbon monoxide produced the fragments $C_{14}H_7O_6^-$ ($m/z = 271$) and $C_{13}H_7O_5^-$ ($m/z = 243$).

The fragmentation of kaempferol-dirhamnoside produced four ions. The first one, $C_{21}H_{19}O_{10}^-$ ($m/z = 431$), was generated by the loss of a rhamnose molecule ($C_6H_{11}O_4$, $m/z = 147$ Da). From the $C_{21}H_{19}O_{10}^-$ fragment ($m/z = 431$), the loss of a water molecule produced the second fragment ion, $C_{21}H_{17}O_9^-$ ($m/z = 413$), which, in turn, generated the fragment $C_{18}H_{13}O_7^-$ ($m/z = 341$) by the loss of 72 Da as $C_3H_4O_2$. Finally, the loss of 41 Da as C_2OH produced the fragment $C_{16}H_{12}O_6^-$ ($m/z = 300$).

The fragmentation of vitexin-dirhamnoside resulted in two ionic products: $C_{21}H_{17}O_9^-$ ($m/z = 413$), generated by the loss of a rhamnose molecule ($C_6H_{11}O_4$, 147 Da), and $C_{17}H_9O_5^-$ ($m/z = 293$), produced by the loss of $C_4H_6O_3$ ($m/z = 102$), followed by a water molecule.

3. Discussion

This research aimed to characterize and evaluate the antimicrobial potential of the extracts of *Humiria balsamifera* (Aubl), also known as “mirim”. This species belongs to the Humiriaceae family, and its tea is used in many Brazilian regions for its anti-inflammatory action, especially for treating uterine inflammation [22,23].

Analyses of the leaf extracts by HPLC-ESI-MS and FIA-ESI-IT/MS led to the identification of 11 substances, 10 of which had not yet been reported for this species: seven flavonoids (galocatechin, kaempferol 3-neohesperidoside, sophoricoside, quercetin 3-arabinoside, quercetin-*O*-rhamnoside, kaempferol-dirhamnoside, and vitexin-dirhamnoside); one oligosaccharide (maltotetraose), bergenin; and two derivatives (galloylbergenin and hydroxybenzoyl bergenin).

The substances present in the chemical composition of *H. balsamifera* tell a lot about the species. Flavonoids, according to the characterization presented in this study, are the most abundant class of compounds. These substances exhibit high bioactive potential and present anti-ulcer, antioxidant, anti-inflammatory, anti-allergic, antitumor, antiviral, antifungal, and antidiabetic activities [34–37].

Antimicrobial tests with flavonoids have received increasing attention in recent years, since these compounds are synthesized by plants in response to various types of stress, including microbial infections [38–40]. Researchers are also interested in how flavonoids are able to exhibit antibacterial activity through mechanisms different from conventional drugs, hindering the development of microbial resistance [41,42].

During this study, the *H. balsamifera* extracts were subjected to three tests for the evaluation of their antimicrobial potential against different bacteria strains. By the end of the in vitro tests, the extracts which presented the best results were selected for the in vivo anti-infective assay using *T. molitor* larvae. Our results showed that the *H. balsamifera* ethyl acetate leaf extract (EALE) showed efficacy against *S. aureus*, one of the most resistant pathogens in existence, in all three tests (MIC, antibiofilm potential, and the tests in vivo). The efficacy of this extract is believed to be due to the flavonoids present in its composition. Flavonoids are well-known in the literature, as other polyphenols, to be able to inhibit microbial growth through several mechanisms, such as the inhibition of ATP synthesis in the electron transport chain, inhibition of nucleic acid synthesis, inhibition of the efflux pump, inhibition of biofilm formation, inhibition of virulence factors, inhibition of quorum sensing, membrane disruption, inhibitors of bacterial toxins, and inhibition of cell envelope synthesis [41,43,44].

Bergenin, identified in the three extracts analyzed, has already been reported in *H. balsamifera* [24] and other two species from the Humiriaceae family: *Endopleura uchi* and *Sacoglottis gabonensis* [45,46]. This isocoumarin and its derivatives, such as the identified flavonoids, can be directly related to the antimicrobial activity of the extracts against *S. aureus*. A recent study showed that six synthetic derivatives of bergenin obtained by Williamson synthesis inhibited *S. aureus* growth, especially 8,10-dihexyl-bergenin and 8,10-didecyl-bergenin, which presented the most promising MIC value: 3.12 µg/mL [47].

Besides antimicrobial activity, bergenin has also been pointed out as one of the main substances responsible for antimalarial [24], anti-inflammatory [48,49], antinociceptive [50], anxiolytic [51], and antioxidant activities [52,53].

4. Materials and Methods

4.1. Botanical Material

The leaves and stem bark of *Humiria balsamifera* (Aubl) were collected during the rainy season in Contrato Village, located in the municipality of Morros—MA (2°55′14.8″ S 43°55′38.8″ W). The material was identified at the Rosa Mochel Herbarium (SLUI) of the State University of Maranhão—UEMA, where an exsiccate was deposited under registration number 4769.

4.2. Preparation of the Extracts

The leaves were dried at room temperature, while the stem barks were dried at 40–50 °C in an oven for 24 h; after which, they were ground in a knife mill separately. The crude extracts were prepared from the ground materials by cold percolation, and each extraction was performed twice for each solvent within a period of 5 days, following the sequence hexane, ethyl acetate, and methanol. The extracts obtained were filtered and concentrated on a rotary evaporator [54].

4.3. Test Microorganisms

The microbial strains used in this work were obtained from the Microbial Culture Collection of CEUMA University. It used the following strains: *Escherichia coli* ATCC 25922, *Listeria monocytogenes* ATCC 15313, *Salmonella enterica* Typhimurium ATCC 14028, and *Staphylococcus aureus* ATCC 6538.

4.4. Minimum Inhibitory Concentration (MIC)

The antimicrobial activity of the extracts was evaluated by the determination of the minimum inhibitory concentration (MIC). The MIC was determined by the broth microdilution method. Sterile 96-well plates were prepared with 150 µL of Müeller Hinton broth (MHB) and 50 µL of the extract following the serial dilution method. After the dilutions (1250 µg/mL–2.0 µg/mL for the stem bark extracts and 6250 µg/mL–1.0 µg/mL for the leaf extracts), 10 µL of the microbial suspension were added to the plates until a 0.5 McFarland standard was reached, and they were then incubated in a lab oven at 37 °C for 24 h. The test was performed in duplicate. Once the incubation period ended, 20 µL of resazurin (Sigma-Aldrich, St. Louis, MO, USA; 0.03%) were added, and the readings were executed after 40 min of incubation at 37 °C. Alterations in color, from blue to pink, were considered an indication of microbial growth. The MIC was defined as the lowest concentration able to prevent microbial growth [55].

4.5. Antibiofilm Test

To evaluate the antibiofilm activity, a sample of 10 µL of *S. aureus* suspension (prepared as described in the MIC section) was mixed with 140 µL of MHB and 50 µL of EACH to reach subinhibitory concentrations (0.5 × MIC, 0.25 × MIC, 0.125 × MIC, and 0.0625 × MIC). After 24 h of incubation at 37 °C, the formed biofilm was fixed with methanol (P.A.), stained with violet crystal (0.1%), and washed with ethanol (P.A.). The biofilm mass was

measured using a spectrophotometer at 550 nm. The absorbances of the wells that each received only the culture medium and bacterial solution were used as the positive control.

4.6. Infection Model Using *Tenebrio molitor* Larvae

The assessment of the antimicrobial effect *in vivo* used larvae of the insect *T. molitor* (Tenebrionidae). Larvae of approximately 100 mg were randomized into groups with a minimum of 10 individuals. Before inoculation, the cuticles were cleaned with 70% alcohol. An aliquot of 10 μ L of the microbial suspension (1.0×10^{11} CFU/mL) was injected in the membrane region between the penultimate and antepenultimate rings of the larvae, which were then incubated at 37 °C. After 2 h, the larvae groups received 10 μ L of each extract at different concentrations. Viability was assessed daily by the absence of movement. Larvae inoculated with the microorganism and treated with PBS were used as the negative control, while the noninfected larvae were selected as the positive control. Death of all larvae or transition into pulp form in the experimental group determined the end of the experiment [55].

4.7. Extracts Characterization by HPLC-ESI-MS and FIA-ESI-IT/MS

For the HPLC-ESI-IT/MS/MS and FIA-ESI-IT/MSⁿ analyses, a clean-up step was performed to remove any contaminants; the solution was purified by solid-phase extraction (SPE) using Phenomenex Strata C18 cartridges (500 mg of stationary phase) that were previously activated with 5 mL of MeOH and equilibrated with 5 mL of MeOH:H₂O (1:1, *v/v*). The compounds were eluted from the cartridges using 1 mL of MeOH:H₂O (1:1, *v/v*) with a final volume of 5 mL. The samples were then filtered using a 0.22- μ m PTFE filter and dried. The extract was diluted to 10 μ g/mL in the HPLC solvent, and then, aliquots of 20 μ L were injected directly into the LC-ESI-IT/MS [56].

The analysis was carried out on an online LC-ESI-IT-MS in a LCQ Fleet Ion Trap Mass Spectrometer, Thermo Scientific® (Waltham, MA, EUA). A Kinetex® (Torrance, CA, USA) C18 LC column (2.1 \times 100 mm, 100 Å, and 5 μ m) was used to separate the components. The analysis was executed using water with formic acid 0.1% (A) and acetonitrile + formic acid 0.1% (B), with formic acid 0.1% added in gradient boosting, going from 10% to 100% in 6 min with a flow rate of 0.4 mL/min. The sample was injected into the HPLC system, where it was analyzed online by ESI-MS in the negative ion mode with a UV detector. Mass spectrometry was performed in an LCQ Fleet Ion Trap LC/MSⁿ—Thermo Scientific® (Waltham, MA, EUA) [56].

A FIA-ESI-IT-MSⁿ flow injection analysis was performed in an LTQ XL™ linear ion trap mass spectrometer with an ESI ion source (electrospray ionization) in negative mode (Thermo, San Jose, CA, USA), using a stainless-steel capillary tube at 280 °C, 5.00 kV, capillary voltage of −90 V, −100 V tubular lens, and a flow rate of 5 μ L min^{−1}. A full-scan analysis was conducted at 100–1000 *m/z*. Multiple-stage fragmentations by electrospray ionization/multi-stage mass spectrometry (ESI-MSⁿ) were performed using collision-induced dissociation (CID) by helium for ionic activation. The first step was a full-scan MS to acquire data from the ions at the selected *m/z*. The second step was a tandem mass spectrometry (MS/MS) in data-dependent acquisition mode on [M−H][−] molecules of the compounds of interest at a collision energy of 30% and activation time of 30 ms. The product ions were subjected to further fragmentation under the same conditions, until no more fragments were observed. The identification of different compounds in the chromatographic profile of the hydroalcoholic extract was carried out by comparing their retention times and UV spectra with the literature data [56].

4.8. Statistical Analysis

All tests were performed in at least two independent assays in quadruplicate. The statistical analyses were performed using the software GraphPad Prism version 5.01 (GraphPad Software Inc., La Jolla, CA, USA). Data were analyzed by two-way analysis of variance

(ANOVA), followed by Tukey's test. A p -value of < 0.05 was considered statistically significant.

5. Conclusions

The number of bioactive compounds found in the chemical composition of *H. balsamifera* emphasized its significance in both traditional medicine and scientific research of studies with new treatments based on substances from the Brazilian flora. This highlighted the importance of this study, since the analyses by HPLC-ESI-MS and FIA-ESI-IT/MS identified 11 substances, 10 of which had not yet been reported for *H. balsamifera*, improving the literature regarding its composition. Additionally, the antimicrobial activity of the ethyl acetate leaf extract (EALE) against *S. aureus* was, for the first time, described using three different strategies (MIC, antibiofilm activity, and tests in vivo), highlighting all the potential of this plant against one of the most resistant bacteria of the present day, which encourages further studies on natural bioactive metabolites that can be isolated from *H. balsamifera*.

Author Contributions: E.d.J.S.D.: methodology, validation, investigation, formal analysis, writing original draft; J.C.B.S.: methodology, investigation; T.F.B.: methodology, investigation; D.M.S.: methodology, investigation; F.J.C.C.: conceptualization, resources, writing review & editing, project administration; L.C.N.d.S.: conceptualization, investigation, resources, writing review & editing, project administration; C.Q.d.R.: conceptualization, investigation, resources, writing review & editing, project administration, funding acquisition; A.J.C.F.: conceptualization, investigation, resources, writing review & editing, project administration, funding acquisition. All authors have read and agreed to the published version of the manuscript.

Funding: The authors would like to thank the Foundation for Research and Scientific Development of Maranhão (FAPEMA) for the research grants (processes numbers: BM QUOTA IFMA 2017/BM-02144, BIC-07392/16, and UNIVERSAL-00998/16) and the Federal Institute of Education, Science and Technology of Maranhão (IFMA).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Gonelimali, F.D.; Lin, J.; Miao, W.; Xuan, J.; Charles, F.; Chen, M.; Hatab, S.R. Antimicrobial Properties and Mechanism of Action of Some Plant Extracts Against Food Pathogens and Spoilage Microorganisms. *Front. Microbiol.* **2018**, *9*, 1–9. [CrossRef]
2. Franz, C.M.A.P.; Den Besten, H.M.W.; Böhnlein, C.; Gareis, M.; Zwietering, M.H.; Fusco, V. Microbial food safety in the 21st century: Emerging challenges and foodborne pathogenic bacteria. *Trends Food Sci. Technol.* **2018**, *81*, 155–158. [CrossRef]
3. Parlapani, F.F.; Mallouchos, A.; Haroutounian, S.A.; Bozaris, I.S. Volatile organic compounds of microbial and non-microbial origin produced on model fish substrate un-inoculated and inoculated with gilt-head sea bream spoilage bacteria. *LWT Food Sci. Technol.* **2018**, *78*, 54–62. [CrossRef]
4. Lianou, A.; Panagou, E.Z.; Nychas, G.-J.E. Microbiological spoilage of foods and beverages. In *The Stability and Shelf Life of Food*, 2nd ed.; Subramaniam, P., Ed.; Woodhead Publishing: Cambridge, UK, 2016; pp. 3–42. [CrossRef]
5. Nahar, S.; Jeong, H.L.; Kim, Y.; Ha, A.J.; Roy, P.K.; Park, S.H.; Ashrafudoulla, M.; Mizan, M.F.R.; Ha, S. Inhibitory effects of Flavourzyme on biofilm formation, quorum sensing, and virulence genes of foodborne pathogens *Salmonella Typhimurium* and *Escherichia coli*. *Food Res. Int.* **2021**, *147*, 110461. [CrossRef]
6. WHO (World Health Organisation). WHO Estimates of the Global Burden of Foodborne Diseases Foodborne Diseases Burden Epidemiology Reference Group 2007–2015. 2015. Available online: <https://www.who.int/en/news-room/fact-sheets/detail/food-safety> (accessed on 1 June 2021).
7. Moyer, C.L.; Morita, R.Y. Psychrophiles and psychrotrophs. In *Encyclopedia of Life Sciences*; John Wiley & Sons Ltd.: Chichester, UK, 2007; pp. 1–6. [CrossRef]
8. Wei, Q.; Wang, X.; Sun, D.; Pu, H. Rapid detection and control of psychrotrophic microorganisms in cold storage foods: A review. *Trends Food Sci. Technol.* **2019**, *86*, 453–464. [CrossRef]
9. Kirk, M.D.; Ângulo, F.J.; Havelaar, A.H.; Black, R.E. Diarrhoeal disease in children due to contaminated food. *Bull. World Health Organ.* **2017**, *95*, 233–234. [CrossRef] [PubMed]

10. Anderson, C.J.; Kendall, M.M. *Salmonella typhimurium* Strategies for Host Adaptation. *Front. Microbiol.* **2017**, *8*, 1983. [CrossRef] [PubMed]
11. Neri, D.; Antoci, S.; Iannetti, L.; Ciorba, A.B.; D'Aurelio, R.; Del Matto, I.; Di Leonardo, M.; Giovannini, A.; Prencipe, V.A.; Pomilio, F.; et al. EU and US control measures on *Listeria monocytogenes* and *Salmonella* spp. in certain ready-to-eat meat products: An equivalence study. *Food Control* **2019**, *96*, 98–103. [CrossRef]
12. Shahid, A.H.; Nazir, K.H.M.; El Zowalaty, M.E.; Kabir, A.; Sarker, S.A.; Siddique, M.P.; Ashour, H.M. Molecular detection of vancomycin resistance and methicillin resistance in *Staphylococcus aureus* from food processing environments. *One Health* **2021**, *100276*, 1–14. [CrossRef]
13. Bello, A.; Dingle, T.C. What's That Resistance Mechanism? Understanding Genetic Determinants of Gram-Negative Bacterial Resistance. *Clin. Microbiol. Newsl.* **2018**, *40*, 165–174. [CrossRef]
14. Holmes, A.H.; Moore, L.S.P.; Sundsfjord, A.; Steinbakk, M.; Regmi, S.; Karkey, A.; Guerin, P.J.; Piddock, L.J.V. Understanding the mechanisms and drivers of antimicrobial resistance. *Lancet* **2016**, *387*, 176–187. [CrossRef]
15. Newman, D.J.; Cragg, G.M. Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J. Nat. Prod.* **2020**, *83*, 770–803. [CrossRef] [PubMed]
16. Chandra, H.; Bishnoi, P.; Yadav, A.; Patni, B.; Mishra, A.P.; Nautiyal, A.R. Antimicrobial Resistance and the Alternative Resources with Special Emphasis on Plant-Based Antimicrobials—A Review. *Plants* **2017**, *6*, 16. [CrossRef]
17. Da Silva, L.C.N.; Da Silva, M.V.; Correia, M.T.S. New Frontiers in the Search of Antimicrobials Agents from Natural Products. *Front. Microbiol.* **2017**, *8*, 210. [CrossRef] [PubMed]
18. Valli, M.; Bolzani, V.S. Natural Products: Perspectives and Challenges for use of Brazilian Plant Species in the Bioeconomy. *An. Acad. Bras. Cienc.* **2019**, *91*, e20190208. [CrossRef] [PubMed]
19. Rodrigues, T.; Reker, D.; Schneider, P.; Schneider, G. Counting on natural products for drug design. *Nat. Chem.* **2016**, *8*, 531–541. [CrossRef]
20. Imai, Y.; Meyer, K.J.; Iinishi, A. A new antibiotic selectively kills Gram-negative pathogens. *Nature* **2019**, *576*, 459–464. [CrossRef] [PubMed]
21. Melander, R.J.; Basak, A.K.; Melander, C. Natural products as inspiration for the development of bacterial antibiofilm agents. *Nat. Prod. Rep.* **2020**, *11*. [CrossRef]
22. Carneiro, D.B.; Barboza, M.S.L.; Menezes, M.P. Plantas nativas úteis na Vila dos Pescadores da Reserva Extrativista Marinha Caeté-Taperaçu, Pará, Brasil. *Acta Botânica Bras.* **2010**, *24*, 1027–1033. [CrossRef]
23. Santos, M.G.; Fevereiro, P.C.A.; Reis, G.L.; Barcelos, J.I. Recursos vegetais da restinga de Carapebus, Rio de Janeiro, Brasil. *J. Neotrop. Biol.* **2009**, *6*, 35–54. [CrossRef]
24. Silva, T.B.C.; Alves, V.L.; Mendonça, L.V.H.; Conserva, L.M.; Rocha, E.M.M.; Andrade, E.H.A.; Lemos, R.P.L. Chemical Constituents and Preliminary Antimalarial Activity of *Humiria balsamifera*. *Pharm. Biol.* **2004**, *42*, 94–97. [CrossRef]
25. Paz, S.A.; Marques, G.E.C. Formulação de Hidratante com Extratos de *Humiria balsamifera* (Aubl A. St. Hil) com Ação Antioxidante. In *Cadernos de Agroecologia; Anais do XI Congresso Brasileiro de Agroecologia; ABA—Associação Brasileira de Agroecologia: Rio de Janeiro, Brasil, 2020; Volume 15, p. 2*. Available online: <http://cadernos.aba-agroecologia.org.br/index.php/cadernos/issue/view/4> (accessed on 1 May 2021).
26. Dutra, A.R.; Sutilli, F.K.; Muzitano, M.F.; Guimarães, D.O.; Souza, R.O.; Leal, I.C. Microwave-assisted extraction of metabolites from *Humiria balsamifera* leaves: A rapid and efficient methodology for antioxidant constituents obtainment. *Planta Med.* **2013**, *79*. [CrossRef]
27. Lima, C.N.S.; Farias, F.A.C.; Marques, G.E.C. Atividade Antimicrobiana in vitro dos Extratos de *Humiria balsamifera* (Aubl) A. St. Hil. Frente a Cepas Bacterianas e Fúngicas. In *Mirim (Humiria balsamifera (Aubl.): Riqueza da Biodiversidade Maranhense*, 1st ed.; Luís, S., Ed.; IFMA: São Luís, Brazil, 2019; Volume 1, pp. 99–107. Available online: <https://editora.ifma.edu.br/livraria-edifma-2019/> (accessed on 2 May 2021).
28. Tüting, W.; Adden, R.; Mischnick, P. Fragmentation pattern of regioselectively O-methylated maltooligosaccharides in electrospray ionisation-mass spectrometry/collision induced dissociation. *Int. J. Mass Spectrom.* **2004**, *232*, 107–115. [CrossRef]
29. Song, H.; Wang, J.; Zhang, R.; Liu, X.; Yuan, G.; Wei, C.; Zhao, W.; Li, R.; Wang, B.; Guo, R. In vivo metabolism study of bergenin in rats by HPLC-QTOF mass spectrometry. *Biomed. Chromatogr.* **2013**, *27*, 1398–1405. [CrossRef] [PubMed]
30. Callemien, D.; Collin, S. Use of RP-HPLC-ESI (–)—MS/MS to Differentiate Various Proanthocyanidin Isomers in Lager Beer Extracts. *J. Am. Soc. Brew. Chem.* **2008**, *66*, 109–115. [CrossRef]
31. March, R.E.; Miao, X. A fragmentation study of kaempferol using electrospray quadrupole time-of-flight mass spectrometry at high mass resolution. *Int. J. Mass Spectrom.* **2004**, *231*, 157–167. [CrossRef]
32. Ablajan, K. A study of characteristic fragmentation of isoflavonoids by using negative ion ESI-MSn. *Int. J. Mass Spectrom.* **2011**, *46*, 77–84. [CrossRef] [PubMed]
33. Ćirić, A.; Prosen, H.; Jelikić-Stankov, M.; Đurđević, P. Evaluation of matrix effect in determination of some bioflavonoids in food samples by LC–MS/MS method. *Talanta* **2012**, *99*, 780–790. [CrossRef]
34. Havsteen, B.H. The biochemistry and medical significance of the flavonoids. *Pharmacol. Ther.* **2002**, *96*, 67–202. [CrossRef]
35. Ng, K.R.; Lyu, X.; Mark, R.; Chen, W.N. Antimicrobial and antioxidant activities of phenolic metabolites from flavonoid-producing yeast: Potential as natural food preservatives. *Food Chem.* **2019**, *270*, 123–129. [CrossRef]
36. Lalani, S.; Poh, C.L. Flavonoids as Antiviral Agents for Enterovirus A71 (EV-A71). *Viruses* **2020**, *12*, 184. [CrossRef]

37. Sun, H.; Wang, D.; Song, X. Natural Prenylchalconaringenins and Prenylnaringenins as Antidiabetic Agents: α -Glucosidase and α -Amylase Inhibition and in Vivo Antihyperglycemic and Antihyperlipidemic Effects. *J. Agric. Food Chem.* **2017**, *65*, 1574–1581. [[CrossRef](#)]
38. Samy, R.P.; Gopalakrishnakone, P. Therapeutic Potential of Plants as Anti-microbials for Drug Discovery. *Evid. Based Complementary Altern. Med.* **2010**, *7*, 283–294. [[CrossRef](#)]
39. Buer, C.S.; Imin, N.; Djordjevic, M.A. Flavonoids: New roles for old molecules. *J. Integr. Plant Biol.* **2010**, *52*, 98–111. [[CrossRef](#)]
40. Petrucci, E.; Braidot, E.; Zancani, M.; Peresson, C.; Bertolini, A.; Patui, S.; Vianello, A. Plant flavonoids-biosynthesis, transport and involvement in stress responses. *Int. J. Mol. Med. Sci.* **2013**, *14*, 14950–14973. [[CrossRef](#)]
41. Biharee, A.; Sharma, A.; Kumar, A.; Jaitak, V. Antimicrobial flavonoids as a potential substitute for overcoming antimicrobial resistance. *Fitoterapia* **2020**, *146*, 104720. [[CrossRef](#)] [[PubMed](#)]
42. Cushnie, T.P.T.; Lamb, A.J. Antimicrobial activity of flavonoids. *Int. J. Antimicrob. Agents* **2005**, *26*, 343–356. [[CrossRef](#)] [[PubMed](#)]
43. Solnier, J.; Martin, L.; Bhakta, S.; Bucar, F. Flavonoids as Novel Efflux Pump Inhibitors and Antimicrobials Against Both Environmental and Pathogenic Intracellular Mycobacterial Species. *Molecules* **2020**, *25*, 734. [[CrossRef](#)]
44. Araya-Cloutier, C.; Vincken, J.; Ederen, R.V.; Den Besten, H.M.W.; Gruppen, H. Rapid membrane permeabilization of *Listeria monocytogenes* and *Escherichia coli* induced by antibacterial prenylated phenolic compounds from legumes. *Food Chem.* **2018**, *240*, 147–155. [[CrossRef](#)]
45. Silva, S.L.; Oliveira, V.G.; Yano, T.; Nunomura, R.C.S. Atividade antimicrobiana da bergenia de *Endopleura uchi* (Huber) Cuatrec. *Acta Amaz.* **2009**, *39*, 187–191. [[CrossRef](#)]
46. Maduka, H.C.C.; Okoye, Z.S.C.; Eje, A. The influence of *Sacoglottis gabonensis* stem bark extract and its isolate bergenin, Nigerian alcoholic beverage additives, on the metabolic and haematological side effects of 2,4-dinitrophenyl hydrazine-induced tissue damage. *Vasc. Pharmacol.* **2003**, *39*, 317–324. [[CrossRef](#)]
47. Silva Neto, O.C.; Teodoro, M.T.F.; Nascimento, B.O. Bergenin of *Peltophorum dubium* (Fabaceae) Roots and Its Bioactive Semi-Synthetic Derivatives. *J. Braz. Chem. Soc.* **2020**, *31*, 2644–2650. [[CrossRef](#)]
48. Nunomura, R.C.S.; Oliveira, V.G.; Silva, S.L.; Nunomura, S.M. Characterization of bergenin in *Endopleura uchi* bark and its anti-inflammatory activity. *J. Braz. Chem. Soc.* **2009**, *20*, 1060–1064. [[CrossRef](#)]
49. Shah, M.R.; Arfan, M.; Amin, H. Synthesis of new bergenin derivatives as potent inhibitors of inflammatory mediators NO and TNF- α . *Bioorganic Med. Chem. Lett.* **2012**, *22*, 2744–2747. [[CrossRef](#)]
50. Oliveira, C.M.; Nonao, F.B.; Lima, F.O. Antinociceptive Properties of Bergenin. *J. Nat. Prod.* **2011**, *74*, 2062–2068. [[CrossRef](#)] [[PubMed](#)]
51. Singh, J.; Kumar, A.; Sharma, A. Antianxiety activity guided isolation and characterization of bergenin from *Caesalpinia digyna* Rottler roots. *J. Ethnopharmacol.* **2017**, *195*, 182–187. [[CrossRef](#)] [[PubMed](#)]
52. Uddin, G.; Sadat, A.; Siddiqui, B.S. Comparative antioxidant and antiplasmodial activities of 11-O-galloylbergenin and bergenin isolated from *Bergenia ligulata*. *Trop. Biomed.* **2014**, *31*, 143–148. [[CrossRef](#)]
53. Muniz, M.P.; Nunomura, S.M.; Lima, E.S.; Lima, A.S.; Almeida, P.D.O.; Nunomura, R.C.S. Quantification of bergenin, antioxidant activity and nitric oxide inhibition from bark, leaf and twig of *Endopleura uchi*. *Química Nova* **2020**, *43*, 413–418. [[CrossRef](#)]
54. Cantanhede Filho, A.J.; Santos, L.S.; Guilhon, G.M.S.P.; Zoghbi, M.G.B.; Ports, P.S.; Rodrigues, I.C.S. Triterpenoides, Fenólicos e Efeito Fitotóxico de *Eugenia flavescens* DC (Myrtaceae). *Química Nova* **2017**, *40*, 252–259. [[CrossRef](#)]
55. Da Fonseca Amorim, E.A.; Castro, E.J.M.; da Souza, S.V.; Alves, M.S.; Dias, L.R.L.; Melo, M.H.F.; de Miranda, R.D.C.M. Antimicrobial Potential of *Streptomyces ansochromogenes* (PB3) Isolated from a Plant Native to the Amazon Against *Pseudomonas aeruginosa*. *Front. Microbiol.* **2020**, *11*. [[CrossRef](#)]
56. Serra, M.B.; Barroso, W.A.; Rocha, C.; Furtado, P.G.R.; Borges, A.C.R.; Silva, S.N.; Tangerina, M.M.P.; Nascimento, J.R.; Vilegas, W.; Alves, A.C.; et al. Chemical Characterization and Wound Healing Property of *Jacaranda decurrens* Cham. (Bignoniaceae): An Experimental Study Based on Molecular Mechanisms. Evidence-Based. *Complementary Altern. Med.* **2020**. [[CrossRef](#)] [[PubMed](#)]