



POSTER PRESENTATION

Open Access

# The impact of HLA-Cw\*12:02 on control of HIV-1 infection

M Koyanagi<sup>1\*</sup>, K Honda<sup>1</sup>, T Chikata<sup>1</sup>, T Akahoshi<sup>1</sup>, H Murakoshi<sup>1</sup>, H Gatanaga<sup>2</sup>, S Oka<sup>2</sup>, M Takiguchi<sup>1</sup>

From AIDS Vaccine 2012

Boston, MA, USA. 9-12 September 2012

## Background

Previous studies have demonstrated that higher HLA-C expression, which is determined through a single nucleotide polymorphism 35 kb upstream and the variation within the 3' untranslated region of the HLA-C locus, associate with slow progression of HIV-1-infected disease. Although HLA-C plays important roles in presenting antigens to CTLs or a ligand for inhibitory killer cell Ig-like receptors (KIR), the role of HLA-C-restricted CTL and NK cells in the control of HIV-1 is still unclear. Our recent study of chronically HIV-1 infected Japanese cohort showed that the HLA-B\*52:01-Cw\*12:02 haplotype was significantly associated with lower viral load. In this study, we investigated whether HLA-Cw\*12:02-restricted CTLs or NK cells via KIR have a significant impact on viraemic control.

## Methods

We sequenced Pol, Gag and Nef from 400 chronically HIV-1 clade B-infected treatment-naïve Japanese individuals and then analyzed amino acid polymorphisms associated with HLA-B\*52:01-Cw\*12:02 haplotype using Fisher's exact test. Next we performed intracellular IFN $\gamma$  staining or IFN $\gamma$  ELISPOT assay to detect CTL responses to the peptides including those polymorphisms.

## Results

We found 9 amino acid polymorphisms significantly associated with HLA-B\*52:01-Cw\*12:02 haplotype ( $p < 0.002$   $q < 0.2$ ). By using ICC assay, we identified 2 Cw\*12:02-restricted CTL epitopes and 4 B\*52:01-restricted ones. Four Cw\*12:02-restricted CTL epitopes including previously reported ones were analyzed to investigate the effect of Cw\*12:02-restricted CTLs on control of HIV-1. No significant correlation between the responses to these

Cw\*12:02-restricted epitopes and viral load was found in chronically HIV-1 infected Cw\*12:02 positive Japanese individuals.

## Conclusion

Those results showed that HLA-Cw\*12:02-restricted CTLs have no effect on control of HIV-1 and suggested that HLA-B\*52:01-restricted CTLs or Cw\*12:02-restricted NK cells control HIV-1 viraemia in Japanese cohort.

## Author details

<sup>1</sup>Center for AIDS Research, Kumamoto University, Kumamoto, Japan. <sup>2</sup>AIDS Clinical Center, National Center for Global Health and Medicine, Tokyo, Japan.

Published: 13 September 2012

doi:10.1186/1742-4690-9-S2-P257

Cite this article as: Koyanagi et al.: The impact of HLA-Cw\*12:02 on control of HIV-1 infection. *Retrovirology* 2012 **9**(Suppl 2):P257.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
www.biomedcentral.com/submit



<sup>1</sup>Center for AIDS Research, Kumamoto University, Kumamoto, Japan  
Full list of author information is available at the end of the article