

SCIENTIFIC REPORTS



OPEN

High-dimensional entanglement certification

Zixin Huang^{1,2}, Lorenzo Maccone³, Akib Karim^{1,2}, Chiara Macchiavello³, Robert J. Chapman^{1,2} & Alberto Peruzzo^{1,2}

Received: 04 January 2016

Accepted: 23 May 2016

Published: 17 June 2016

Quantum entanglement is the ability of joint quantum systems to possess global properties (correlation among systems) even when subsystems have no definite individual property. Whilst the 2-dimensional (qubit) case is well-understood, currently, tools to characterise entanglement in high dimensions are limited. We experimentally demonstrate a new procedure for entanglement certification that is suitable for large systems, based entirely on information-theoretics. It scales more efficiently than Bell's inequality and entanglement witness. The method we developed works for arbitrarily large system dimension d and employs only two local measurements of complementary properties. This procedure can also certify whether the system is maximally entangled. We illustrate the protocol for families of bipartite states of qudits with dimension up to 32 composed of polarisation-entangled photon pairs.

As the dimension of investigated systems increases, it becomes more complicated to demonstrate their quantum effects^{1–3}. Indeed, a full characterization (quantum tomography) becomes practically impossible already for systems of rather small dimensionality^{4,5}. It is therefore important to explore new avenues to prove the presence of such effects, e.g. entanglement, for arbitrary dimensions. High-dimensional entangled states offer a larger code space, attracting interests for quantum key distribution⁶, teleportation⁷ and security-enhanced quantum cryptography^{8,9}.

In this context, proving that one has achieved entanglement (entanglement certification) and detecting entanglement are different primitives. Indeed, entanglement detection methods^{10–14} must be as sensitive as possible and must be able to detect the largest possible class of entangled states. Often such methods are inapplicable to large system dimensions or scale poorly with increasing dimension as they entail increasingly complicated measurements and data analysis. Entanglement detection such as witness operators, for instance, typically requires a number of local measurement settings that scales linearly in d^{15} . In contrast, entanglement certification refers to the fact that one has simply to prove that the system is entangled. To do entanglement certification, one can optimize the method to the specific entangled state that one is producing. It must fulfill different requirements: it must be as robust and simple as possible. In other words a good method for entanglement detection can also work for entanglement certification, but not vice versa.

Here, we present an entanglement certification procedure that is extremely simple to implement (the measurement of two local observables^{16,17} suffices), is compatible with current state-of-the-art experimental techniques, and can be easily scaled up to arbitrary dimension. In addition to certifying the production of entangled states, our procedure can also certify the production of maximal entanglement. To prove its simplicity, we present an experimental test that uses entangled systems with dimension up to $d = 32$, constructed by suitably grouping couples of entangled photon pairs. In the general case, for a two-qudit experiment with arbitrary d , two measures would still be sufficient to implement our method, but not sufficient for tomography or other entanglement detection methods. To certify the presence of entanglement, one has only to calculate the (classical) correlations among the measurement outcomes of the two observables, for example, through their mutual information. If such correlations are larger than some threshold, the state is guaranteed to be entangled (Fig. 1). If they attain their maximum value, the state must be maximally entangled.

If A and C are complementary properties, the knowledge of A gives no information on C and vice versa. This happens whenever $|\langle a|c\rangle|^2 = 1/d$, for all $|a\rangle$ and $|c\rangle$ eigenstates of the observables A and C , where d is the Hilbert space dimension. This is equivalent to two sets of mutually unbiased bases (MUB's). Consider the two-qubit maximally entangled state $|\Phi^+\rangle \propto |00\rangle + |11\rangle$; it clearly has maximal correlation among results for the observables

¹Quantum Photonics Laboratory, School of Engineering, RMIT University, Melbourne, Victoria 3000, Australia.

²School of Physics, University of Sydney, NSW 2006, Australia. ³Dip. Fisica and INFN Sez. Pavia, University of Pavia, via Bassi 6, I-27100 Pavia, Italy. Correspondence and requests for materials should be addressed to A.P. (email: alberto.peruzzo@rmit.edu.au)




Figure 1. Illustration of the procedure. (a) Measure two complementary observables: AB and CD , where A, C are observables for system 1 and C, D for system 2. Then calculate the mutual information I_{AB} and I_{CD} of their outcomes. (b) If the value of the sum $I_{AB} + I_{CD}$ is larger than $\log_2(d)$ bits (d being the system dimension), then the two systems are certified to be entangled. If the value of the sum is $2 \log_2(d)$ bits, they are certified to be maximally entangled.

with eigenstates $|0\rangle, |1\rangle$: both qubits have the same value. This state has also maximal correlation among results for a complementary observable with eigenvalues $|\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$, since it can be written as $|\Phi^+\rangle \propto |++\rangle + |--\rangle$. Thus, the mutual information between measurement outcomes on the two qubits is one bit per observable, summing to two bits. If one has only separable states, the sum cannot be larger than one: for example, the classically correlated state $\rho \propto |00\rangle\langle 00| + |11\rangle\langle 11|$ has one bit of mutual information for the outcomes of the first observable, but zero bits for the outcomes of the second.

Starting from the theoretical suggestions of¹⁸, we extend the method from the qubit ($d=2$) case to arbitrary high dimensions. The proposed mechanism to certify entanglement thus uses the following procedure:

1. Identify two bipartite complementary observables AB and CD , where A, C are system 1 observables and B, D are system 2 observables (Fig. 1a).
2. Measure the statistics of the outcomes of the two observables: local measurements on the two systems suffice.
3. These measurements return the joint probabilities $p(a_o, b_o)$ of obtaining outcome a_o on system 1 for A and b_o on system 2 for B , and $p(c_o, d_o)$ of obtaining outcome c_o on system 1 and d_o on system 2. Use these results to calculate the mutual information I_{AB} among measurement results for AB and I_{CD} among results for CD , with

$$I_{AB} \equiv \sum_{ab} p(a, b) \log_2 \frac{p(a, b)}{\sum_a p(a, b) \sum_b p(a, b)}. \tag{1}$$

4. Certification: if $I_{AB} + I_{CD} > \log_2 d$, the two systems are entangled; if $I_{AB} + I_{CD} = 2 \log_2 d$, the two systems are maximally entangled.

The proof of these statements, based on Maassen and Uffink's entropic uncertainty relation¹⁹ is given in¹⁸ (see also^{20–22}).

In regards to the choice of observables, for pure states, the obvious choice would be the Schmidt bases and their respective Fourier bases. Whilst for mixed states, one can diagonalise the density matrix, identify the eigenvector with the largest weight and use the Schmidt basis (and its respective Fourier bases). Alternatively, one can choose the bases that diagonalises the reduced density matrices. While there is no guarantee that these choices will allow one to implement the procedure, they are the ones that may uncover the most correlations.

This method is simple to implement for systems of any dimensionality: it only entails independent measurements of two local observables on the two systems. Moreover, it is robust since, although one can optimize the choice of observables to maximize the sum of mutual information, the systems are guaranteed to be entangled if the above conditions are satisfied for *any* couple of complementary observables. It is interesting to note that, coherently performing sequential complementary measurements on the same system may generate the entanglement itself²³.

To date, the most prominent way of producing higher dimensional entangled systems is via the orbital angular momentum degree of freedom of a photon^{6,24,25}; schemes to produce three-level entangled states in trapped ions have also been proposed^{26,27}. Our method, however, works for any d -dimensional system as long as the appropriate measurements can be performed.

Based on the theoretical work by Collins *et al.*, other recent experiments^{25,28} have studied higher dimensional entangled system via generalised Bell's inequalities, where the correlations between the two measurement settings on each qudit system are studied. The violation of Bell's inequality is usually discussed in terms of quantum non-locality. In this context, the framework of hidden variable theories is in general specified by the 4 measurement settings, each with d outcomes, therefore needing d^4 joint probabilities²⁹ to describe the system globally. In this case^{25,28,29}, examine two settings on each system, and how the observables on one system are correlated with the two observables on the second system, requiring a total of $2d \times 2d$ joint outcomes. On the other hand, our method is based on entropic relations, requiring only two measurement settings, and is completely specified by $2d^2$ probabilities. Similarly to²⁸, we used ensembles of individual entangled photon pairs to construct a higher dimensional state. All the subsystems that compose this state do not exist at the same time, but that is irrelevant to our current aims.

To demonstrate the method in practice, we performed an experiment using high-dimensional entangled systems. We certify the presence of entanglement for various families of states, by measuring pairs of complementary observables. These families are obtained by appropriately grouping couples of polarisation entangled photonic qubits, mapping qudits onto qubits:

$$\frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} |j\rangle |j\rangle = \frac{1}{\sqrt{d}} (|00\rangle + |11\rangle)^{\otimes n}, \quad (2)$$

where this mapping is obtained by expressing j in binary notation and rearranging the qubits in such a way that the ones placed in odd positions in the tensor products on the right hand side are assigned to system 1 while the ones in even positions to system 2. For example, a $d=4$ maximally entangled state can be expressed as $(|00\rangle + |11\rangle + |22\rangle + |33\rangle)/2 \equiv (|0000\rangle + |0101\rangle + |1010\rangle + |1111\rangle)/2 \equiv (|0\rangle_\alpha |0\rangle_\beta + |1\rangle_\alpha |1\rangle_\beta) \otimes (|0\rangle_\gamma |0\rangle_\delta + |1\rangle_\gamma |1\rangle_\delta)/2$, where qubits labeled α and γ belong to system 1, while β and δ to system 2. It would also be extremely interesting to show that one could coherently manipulate the n entangled pairs locally, though that is not necessary to implement our method (which is also one of the main strengths of the method). An analogous mapping applies to the mixed states:

$$\frac{1}{d} \sum_j |j\rangle \langle j| \otimes |j\rangle \langle j| = \frac{1}{d} (|00\rangle \langle 00| + |11\rangle \langle 11|)^{\otimes n}. \quad (3)$$

Regarding the necessary measurements, the observable AB corresponds to σ_z , namely the computational basis $\{|0\rangle, |1\rangle\}$ for each qubit. As to the observable CD , we will consider two possibilities: the Fourier basis and the σ_x basis. The Fourier basis is defined as

$$|f_j\rangle \equiv \frac{1}{\sqrt{d}} \sum_{k=0}^{d-1} \omega^{kj} |k\rangle \Rightarrow |k\rangle = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} \omega^{-kj} |f_j\rangle, \quad (4)$$

with $\omega \equiv \exp(2\pi i/d)$. It can be expressed as tensor products of single-qubit states (see Supplementary Material).

The Fourier basis in arbitrary dimension identifies an observable complementary to the computational basis. However, in the case considered here, there are complementary bases that are simpler to access experimentally, namely the bases where one measures σ_x or σ_y on each qubit. We consider σ_x here. The σ_x basis is given by $|c_k\rangle$ ($k = 0, \dots, d-1$) obtained by expressing the binary digits of k in the $\{+, -\}$ basis, i.e.

$$|c_k\rangle = [(|0\rangle + (-1)^{\gamma_1} |1\rangle)(|0\rangle + (-1)^{\gamma_2} |1\rangle) \dots] / \sqrt{2^n}, \quad (5)$$

where γ_ℓ are the bits of the number k . The σ_x basis is complementary to the computational basis, since $|\langle j|c_k\rangle|^2 = 1/2^n = 1/d$ for all j, k .

We tested our entanglement certification procedure on several families of states. These particular states that we consider are arbitrarily chosen for their simplicity. For the specific examples, the entanglement of the d -dimensional entangled pair is exactly the same as the one obtained from consecutive entangled two-photon states. This happens only when d is a power of two, though the method works for any d . The first is

$$\rho_c(p) = \frac{p}{d} \sum_{j'j} |j'j\rangle \langle j'j| + \frac{1-p}{d} \sum_j |jj\rangle \langle jj|, \quad (6)$$

which mixes the maximally entangled state of equation (2) with the state in equation (3) with classical correlation only on the computational basis, the above state is always entangled for $p \neq 0$.

The experimental test of this prediction is presented in Fig. 2, red circles (the markers are the data points and the connecting lines are the expected curves fitted to density matrices from quantum state tomography).




Figure 2. Experimental test of the method for d -dimensional systems. The plots refer to the experimentally determined sum of mutual information $I_{AB} + I_{CD}$ for ρ_c of equation (6) (red circles), ρ_w of equation (7) (green triangles), ρ_a of equation (8) (blue stars when CD is the σ_x observable, and purple diamonds when CD is the Fourier basis). The points that fall above the entanglement threshold (orange dashed line) refer to states that are certified to be entangled. Taking into account the confidence interval of the measurement outcome, the state was maximally entangled if the maximum sum $I_{AB} + I_{CD} = 2 \log_2 d$ is achieved. The bands are a Monte Carlo simulation, taking into account realistic experimental conditions (e.g. phase inaccuracy due to the wave plates used in state tuning); for $d = 32$ they are narrower than the fit. The error bars represent bounds for 2 standard deviations (95% confidence), for most of the data points the error bars are smaller than the markers.

The second family of states we consider are the Werner states: a mixture of a maximally entangled and a maximally mixed state,

$$\rho_w = \frac{p}{d} \sum_{jj'} |jj\rangle \langle jj'| + \frac{1-p}{d^2} I, \tag{7}$$

where I is the identity $d^2 \times d^2$ matrix. These states are entangled for $p > 1/(d+1)$ (e.g.³⁰). Their mutual information is experimentally determined as the green triangles in Fig. 2.

The third family of states is

$$\rho_a = \frac{p}{d} \sum_{jj'} |jj\rangle \langle jj'| + (1-p) \left(\frac{|+-\rangle \langle +-| + |-+\rangle \langle -+|}{2} \right)^{\otimes n}, \tag{8}$$

where $|\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$. The experimental results are plotted in Fig. 2 as blue stars (when the CD observable is the σ_x basis $|c_k\rangle$) and as purple diamonds (when CD is the Fourier basis $|f_j\rangle$). Measurements in the Fourier basis and in the σ_x basis are performed for $d=4$ and 8 only, due to the large number of projections required for higher dimensions where temporal phase instability would significantly affect the measurements. This state shows the difference between the complementary bases: its form implies that correlations are greater in the σ_x than the Fourier basis.

The entanglement certification method can be modified by using a different measure of correlations. Indeed, instead of the mutual information, one can also measure correlations with the Pearson coefficient

$$C_{AB} \equiv (\langle AB \rangle - \langle A \rangle \langle B \rangle) / \sigma_A \sigma_B, \tag{9}$$

where $\langle X \rangle$ is the expectation value of X and σ_X^2 its variance. It measures the linear correlation between the outcomes of two observables A and B , and takes values $-1 \leq C_{AB} \leq 1$. It is conjectured¹⁸ that if $C_{AB} + C_{CD} > 1$, then the state is entangled. Moreover, it is known that if $C_{AB} + C_{CD} = 2$, the state is maximally entangled. This may be helpful in high-noise scenarios, since it allows for the certification of a larger class of entangled states. The Pearson correlation coefficient seems to be more robust against imbalanced probabilities or decoherence in the experiment (see Fig. 3), and numerical simulations suggest that it is more effective in detecting entanglement¹⁸. So, in addition to the sum of mutual information $I_{AB} + I_{CD}$, we can use $C_{AB} + C_{CD}$ as an alternative way of certifying entanglement (modulo a conjecture).

We consider the two-qubit states

$$\rho_b = p |\Phi^+\rangle \langle \Phi^+| + (1-p) |\Phi^-\rangle \langle \Phi^-|, \tag{10}$$

$$\rho_s = p \left(\frac{|00\rangle \langle 00| + |11\rangle \langle 11|}{2} \right) + (1-p) \left(\frac{|+-\rangle \langle +-| + |-+\rangle \langle -+|}{2} \right), \tag{11}$$

and the Werner state ρ_w of Eq. (7) with $d=2$. In (10), $|\Phi^\pm\rangle \equiv (|00\rangle \pm |11\rangle) / \sqrt{2}$. The state ρ_b is entangled for $p \neq 1/2$, whereas ρ_s is always separable and has zero discord only for $p=0, 1$. For two-qubit states, the σ_x basis $|c_k\rangle$ coincides with the Fourier basis $|f_j\rangle$, so the two possible observables CD we used above coincide here.




Figure 3. Experimental comparison between mutual information (above) and Pearson coefficient (below). The insets show the theoretical curves. The data points refer to ρ_b of equation (10) (red circles), ρ_s of equation (11) (blue diamonds), ρ_w of equation (7) (green triangles) for $d=2$. The states whose data points are strictly above the entanglement threshold ($I_{AB} + I_{CD} = 1$ for the upper graph and $C_{AB} + C_{CD} = 1$ for the lower graph, conjectured) are certified to be entangled. In the lower graph this threshold overlaps with the fit for ρ_s , which is then not entangled, as expected. The markers represent the data points, the lines represent a fit from reconstructed density matrices, and the bands are a Monte Carlo simulation, taking into account realistic experimental conditions (eg. phase inaccuracy due to the wave plates used in state tuning). The error bars represent bounds for 2 standard deviations. It is evident that the experimental results are in good agreement with the theoretical values.

However, we preferred working with mutual information since the fact that the Pearson coefficient can be used to certify entanglement is still a conjecture, whereas it has been rigorously proved for the sum of mutual information.

Whilst it is true that the standard entanglement witness

$$W = I - \sigma_z \otimes \sigma_z - \sigma_x \otimes \sigma_x \quad (12)$$

out-performs our method for entanglement detection for the ($d=2$ case) Werner state (detects entanglement at $p \geq 0.5$), this is not true for arbitrary dimensions in general.

Finally, also comparing this against Bell's inequalities using the Werner state, our method using mutual information with two complementary observables (computational and σ_x basis), for $d=2$, the threshold of $\log_2(d)$ is surpassed at $p \approx 0.78$. However, if we increase this to three observables (indeed one can, and in this case we consider the σ_y basis), the entanglement is detected at $p \approx 0.65$ (see supplementary material of¹⁸), which outperforms Bell's inequalities at $1/\sqrt{2}^{11}$. Also, in the limit of large d , our method (see Eq. (S7) in the Supplementary Material) also out-performs the threshold in Bell's inequality violation, given in²⁹.




Figure 4. Scheme for polarisation entangled source. Type-I SPDC source is used to generate pairs of horizontally polarised single photons. HWP_1 rotates $|HH\rangle$ to $|++\rangle$, $|+\rangle = \frac{1}{\sqrt{2}}(H + V)$. The top arm experiences a polarisation phase shift, and the state of the bottom arm is purified and matched to the top arm by the PBS rotated by $\pi/4$ radians, which transmits $|+\rangle$. The photons are collected into polarisation maintaining fibres (PMF) and are incident on the two input faces of a PBS, which transmits H and reflects V. This prepares the state $\frac{1}{\sqrt{2}}(HV + e^{i\theta}VH)$ when measuring in the coincidence basis. Standard tomography set up projects the state onto different bases for measurements. See text for further details.

In summary, we have presented an entanglement certification method that is suitable for large dimensional systems. To our knowledge, there are no such methods that are as efficient to implement on a 32-dimensional system such as the one we experimentally study here. We have shown that entangled states are more correlated on the outcomes of complementary observables. One may think that this property is shared by other types of quantum correlations such as quantum discord³¹ but, surprisingly, this is not the case, at least when the correlations are gauged through mutual information. Indeed, the separable state ρ_s has highest $I_{AB} + I_{CD}$ for $p = 0, 1$ (Fig. 3), the values for which its discord is null. In contrast, $I_{AB} + I_{CD}$ is lowest for $p = 1/2$ which is where such state has highest discord. This unexpected property is lost when the correlation is gauged with the Pearson coefficient, as $C_{AB} + C_{CD} = 1$ for all values of p .

Methods

In order to generate polarisation-entangled photon pairs, a set up similar to³² and identical to³³ is used. We use the polarisation encoding where $|H\rangle \equiv |0\rangle$ and $|V\rangle \equiv |1\rangle$. The set up is shown in Fig. 4. A type-I nonlinear crystal (BiBO) is pumped with a vertically polarised cw laser (404 nm) at 80 mW, generating pairs of horizontally polarised single photons (808 nm). The photons are spectrally filtered using 3 nm top-hat filters centred on 808 nm. The first half way plate HWP_1 (optics axis set to $\pi/8$ radians) rotates the output photon pair from $|HH\rangle$ to $|++\rangle$, where $|+\rangle = \frac{1}{\sqrt{2}}(H + V)$. The polarisation in the top arm is matched and purified to that of the bottom arm using a polarisation beam splitter (PBS) rotated by $\pi/4$ radians, which transmits $|+\rangle$.

The unitary operation realised by the sequence of wave plates $QWP_1 + HWP_2(\theta) + QWP_2$, as a function θ of the HWP_2 angle, applies a phase to photon in V state, effectively that of a phase gate: $\begin{pmatrix} 1 & 0 \\ 0 & e^{i4\theta} \end{pmatrix}$. Denoting the input photon in the top (bottom) arm with subscript 1 (2), the two-photon state collected into the polarisation mainting fibre (PMF) is therefore $\frac{1}{\sqrt{2}}(|H\rangle_1 + e^{i\theta}|V\rangle_1) \otimes \frac{1}{\sqrt{2}}(|H\rangle_2 + |V\rangle_2)$.

We use a silicon avalanche single photon counter and measure in coincidence with a time window of 2.5 ns. At the fibre coupled PBS, H is transmitted and V is reflected. After the fibre PBS crystal, the top (bottom) arm fibre contains the components $(V_1$ and $H_2)$ (H_1 and $e^{i\theta}V_2$). However, the design of the fibre PBS is such that the output coupler flips the polarisation state in the bottom arm. Without this flip, the registered coincidence will be either $e^{i\theta}|V\rangle_1|V\rangle_2$ or $|H\rangle_1|H\rangle_2$. After the flip the detector will register either $e^{i\theta}|V\rangle_1|H\rangle_2$ or $|H\rangle_2|V\rangle_1$, and these two components are made spatially indistinguishable by adjusting the position of the fibre, giving the post-selected state $\frac{1}{\sqrt{2}}(|HV\rangle + e^{-i\theta}|VH\rangle)$. HWP_3 and HWP_4 apply the appropriate bit flip operations on each single qubit. $QWP_3 + HWP_5$ and $QWP_4 + HWP_6$ along with the final two PBSs project each photon onto any polarisation state.

The polarisation maintaining fibres have a beat length (H and V delayed with respect to each other by 2π) of 24 mm; the coherence length of the photon pairs is $250 \mu\text{m}$ (extracted from a Hong-Ou-Mandel dip of 99% visibility), which means that if uncompensated, 1.0 m of fibre will spatially separate the H and V components of a photon out of coherence, turning a Bell state $\frac{1}{\sqrt{2}}(|HH\rangle + |VV\rangle)$ into the classically correlated mixture

$\left(\frac{|\langle HH \rangle| + |\langle VV \rangle|}{2}\right)$. When producing a Bell state, this spatial decoherence was compensated by crossing the slow axis of the PMF and tuning the stage position of the fibre (and vice versa when the mixed state is required).

References

1. Arndt, M. & Hornberger, K. Testing the limits of quantum mechanical superpositions. *Nat. Phys.* **10**, 271–277 (2014).
2. Sciarrino, F. Quantum optics: Micro meets macro. *Nat. Phys.* **9**, 529–529 (2013).
3. Friedman, J. R., Patel, V., Chen, W., Tolpygo, S. & Lukens, J. E. Quantum superposition of distinct macroscopic states. *Nat.* **406**, 43–46 (2000).
4. Gühne, O. *et al.* Detection of entanglement with few local measurements. *Phys. Rev. A* **66**, 062305 (2002).
5. Artiles, L., Gill, R. *et al.* An invitation to quantum tomography. *J. Roy. Statist. Soc. Ser. B* **67**, 109–134 (2005).
6. Mair, A., Vaziri, A., Weihs, G. & Zeilinger, A. Entanglement of the orbital angular momentum states of photons. *Nat.* **412**, 313–316 (2001).
7. Bennett, C. H. & Wiesner, S. J. Communication via one- and two-particle operators on Einstein-Podolsky-Rosen states. *Phys. Rev. Lett.* **69**, 2881–2884 (1992).
8. Bruß, D. & Macchiavello, C. Optimal eavesdropping in cryptography with three-dimensional quantum states. *Phys. Rev. Lett.* **88**, 127901 (2002).
9. Durt, T., Kaszlikowski, D., Chen, J.-L. & Kwek, L. C. Security of quantum key distributions with entangled qudits. *Phys. Rev. A* **69**, 032313 (2004).
10. Bruß, D. Characterizing entanglement. *J. Math. Phys.* **43**, 4237–4251 (2002).
11. Horodecki, R., Horodecki, P., Horodecki, M. & Horodecki, K. Quantum entanglement. *Rev. Mod. Phys.* **81**, 865–942 (2009).
12. Gühne, O. & Tóth, G. Entanglement detection. *Phys. Rep.* **474**, 1–75 (2009).
13. Rudnicki Ł., Horodecki P., and Życzkowski K. Collective Uncertainty Entanglement Test. *Phys. Rev. Lett.* **107**, 150502 (2011).
14. Rudnicki Ł., Puchała Z., Horodecki P., and Życzkowski K. Collectibility for mixed quantum states. *Phys. Rev. A* **86**, 062329 (2012).
15. Gühne, O. *et al.* Experimental detection of entanglement via witness operators and local measurements. *J. Modern Opt.* **50**, 1079–1102 (2003).
16. Kothe, C. & Björk, G. Entanglement quantification through local observable correlations. *Phys. Rev. A* **75**, 012336 (2007).
17. Abascal, I. S. & Björk, G. Bipartite entanglement measure based on covariance. *Phys. Rev. A* **75**, 062317 (2007).
18. Maccone, L., Bruß, D. & Macchiavello, C. Complementarity and correlations. *Phys. Rev. Lett.* **114**, 130401 (2015).
19. Maassen, H. & Uffink, J. B. M. Generalized entropic uncertainty relations. *Phys. Rev. Lett.* **60**, 1103–1106 (1988).
20. Schneeloch, J., Broadbent, C. J. & Howell, J. C. Uncertainty relation for mutual information. *Phys. Rev. A* **90**, 062119 (2014).
21. Coles, P. J. & Piani, M. Improved entropic uncertainty relations and information exclusion relations. *Phys. Rev. A* **89**, 022112 (2014).
22. Coles, P. J., Colbeck, R., Yu, L. & Żwolak, M. Uncertainty relations from simple entropic properties. *Phys. Rev. Lett.* **108**, 210405 (2012).
23. Coles, P. J. & Piani, M. Complementary sequential measurements generate entanglement. *Phys. Rev. A* **89**, 010302 (2014).
24. Fickler, R. *et al.* Quantum entanglement of high angular momenta. *Sci.* **338**, 640–643 (2012).
25. Dada, A. C., Leach, J., Buller, G. S., Padgett, M. J. & Andersson, E. Experimental high-dimensional two-photon entanglement and violations of generalized Bell inequalities. *Nat. Phys.* **7**, 677–680 (2011).
26. Zheng, S.-B. Generation of entangled states for many multilevel atoms in a thermal cavity and ions in thermal motion. *Phys. Rev. A* **68**, 035801 (2003).
27. Ye, S.-Y., Zhong, Z.-R. & Zheng, S.-B. Deterministic generation of three-dimensional entanglement for two atoms separately trapped in two optical cavities. *Phys. Rev. A* **77**, 014303 (2008).
28. Lo, H.-P. *et al.* Experimental violation of Bell inequalities for multi-dimensional systems. *arXiv preprint arXiv:1501.06429* (2015).
29. Collins, D., Gisin, N., Linden, N., Massar, S. & Popescu, S. Bell inequalities for arbitrarily high-dimensional systems. *Phys. Rev. Lett.* **88**, 040404 (2002).
30. Kendon, V. M., Życzkowski, K. & Munro, W. J. Bounds on entanglement in qudit subsystems. *Phys. Rev. A* **66**, 062310 (2002).
31. Ollivier, H. & Zurek, W. H. Quantum discord: A measure of the quantumness of correlations. *Phys. Rev. Lett.* **88**, 017901 (2001).
32. Matthews, J. C. *et al.* Observing fermionic statistics with photons in arbitrary processes. *Sci. Rep.* **3**, 1539 (2013).
33. Chapman, R. J. *et al.* Experimental perfect state transfer of an entangled photonic qubit. *Nat. Commun.* **7**:11339 doi: 10.1038/ncomms11339 (2016).

Acknowledgements

A.P. acknowledges an Australian Research Council Discovery Early Career Researcher Award under project number DE140101700 and an RMIT University Vice-Chancellor’s Senior Research Fellowship.

Author Contributions

Z.H., L.M., C.M. and A.P. designed the experiment. Z.H. performed the experiment and analysed the data. All authors contributed to writing the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Huang, Z. *et al.* High-dimensional entanglement certification. *Sci. Rep.* **6**, 27637; doi: 10.1038/srep27637 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article’s Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>