



Olfactory Performance among Hospital Residents

Gustavo Lara Rezende^{1,2,3} Max Sarmet⁴ Gabrielle Everton Sousa² Felipe Carneiro Krier²
Diderot Rodrigues Parreira² Selma Aparecida Kuckelhaus³

¹Otorhinolaryngology Department, Hospital de Base do Distrito Federal (HBDF), Brasília, DF, Brazil

²Faculdade de Medicina, Universidade Católica de Brasília (UCB), Taguatinga, Brasília, DF, Brazil

³Morphology Department, Faculdade de Medicina, Universidade de Brasília (UnB), Brasília, DF, Brazil

⁴Hospital de Apoio de Brasília (HAB), Brasília, DF, Brazil

Address for correspondence Gustavo Lara Rezende, MD, PhD, SEPS 709/909, CMJA, sala 223 A, Brasília, DF, 70390-095, Brazil (e-mail: glararezende@gmail.com).

Int Arch Otorhinolaryngol 2022;26(1):e32–e37.

Abstract

Introduction Smell plays an important role in the maintenance of health and quality of life of the general population. Health workers with olfactory impairment may not be able to help diagnose certain diseases, and subsequently increase the risk of hazardous events and mortality among those affected. ‘Odor learning’ requires repeated experiences with different smells to develop a discriminatory ability, and this is a process that takes years. Because of that, physicians of certain medical specialties have better odor detection than others.

Objective To study the olfactory performance and associated factors of otorhinolaryngology residents compared with residents of different medical specialties in a representative sample of a tertiary hospital.

Methods The University of Pennsylvania Smell Identification Test (UPSIT) was used to compare olfactory performance. Clinical and epidemiological data were collected among 42 hospital residents.

Results Otorhinolaryngology residents presented an average UPSIT score of 35.0, and the other residents, a score of 32.8 ($p = 0.02$) Of all the residents, 40.5% showed some grade of olfactory impairment. Half of the females students in the first year of residency showed olfactory dysfunction. The multivariate analyses found age ($p = 0.03$; 95% confidence interval for $\beta = 0.33$) to be an independent predictor of the UPSIT score.

Conclusion The present study demonstrated that otorhinolaryngology residents have greater olfactory capacity compared with other residents. Future studies should explore the relevant factors of olfactory impairment and its impact on quality of life in this population.

Keywords

- ▶ olfactory test
- ▶ dysosmia
- ▶ residents
- ▶ health workers

Introduction

Smell plays an important role in the maintenance of health and quality of life of the general population. Health workers with olfactory impairment may not be able to help diagnose

certain diseases,^{1–3} and subsequently increase the risk of hazardous events and mortality among those affected.⁴

Little research has been conducted on the prevalence of olfactory disorders among health workers due to the

received

July 5, 2020

accepted

September 27, 2020

published online

February 19, 2021

DOI <https://doi.org/>

10.1055/s-0040-1719122.

ISSN 1809-9777.

© 2021. Fundação Otorrinolaringologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

difficulty of standardising tests and their costs. Cross-sectional studies have demonstrated the prevalence of olfactory disorders in ~ 19% of individuals in the general population.⁵ Among health professionals, there is still no data on the use of this essential work tool during the physical examination of patients and its important role in detecting diseases.⁶ 'Odor learning' requires repeated experiences with different smells to develop a discriminatory ability, and this is a process that takes years.⁴ Because of that, physicians of certain medical specialities would be able to have better odor detection than others.

In addition to the importance of smell for health professionals in the diagnosis of diseases in general, the determination of the olfaction status is an important tool of occupational medicine and public health planning.⁷ The present study employed a highly reliable and standardized 40-item odor identification test⁸ to study the olfactory performance and associated factors of otorhinolaryngology residents compared with residents of different medical specialities in a representative sample of a tertiary hospital.

Methods

Subject Enrollment

The study was performed with subjects who were recruited at a tertiary hospital to test their olfactory capacity using the University of Pennsylvania Smell Identification Test (UPSIT) (► Fig. 1). All subjects attended a residency program. The following epidemiological data were collected: age, sex, ethnic group, monthly household income, presence of nasal diseases, presence of rhinitis, smoking behavior, continuous use of medications, and comorbidities.

Individuals with an acute respiratory infection or a history of head trauma were excluded to avoid participants with odor impairment secondary to other etiologies. The initial study population consisted of 50 subjects. Residents from all

units of the hospital were invited to participate voluntarily. The present study was approved by the institutional Ethics Committee for Analysis of Research Projects of the Clinical Board.

Residency Programs

The individuals were divided into two groups according to the specialty of the residency program: otorhinolaryngology or other specialties. They were also divided into groups according to the year of residency (R1 = first year in the program; R2 = second year in the program etc.) and type of residency program (clinical or surgical).

Monthly Household Income

We also divided the residents into four income-based groups, analogous to the categories previously used in Fornazieri et al.,⁹ which were based on the Brazilian classification of socioeconomic classes.

Ethnicity

Ethnicity was classified according to the definitions of the Brazilian Institute of Geography and Statistics (Instituto Brasileiro de Geografia e Estatística, IBGE, in the Portuguese acronym).⁹ The subjects of the present study were self-defined as White, Black, and mixed-race Brazilians.

Olfactory Evaluation

All participants completed the UPSIT, a widely used forced-choice standardized test of olfactory function.⁸ This test consisted of four booklets of ten odorants each. The stimuli were embedded into microcapsules present in brown strips at the bottom of each page. The examiner directed the subjects to scrape the strip with a pencil, which released the odorant. The subjects then answered a multiple-choice question to describe the smell as they perceived it. Based on the test scores, each subject's olfactory function could be




Fig. 1 Key elements of the study.

classified into the normosmia, microsmia (mild, moderate, and severe), and anosmia categories. The scores ranged from 0 to 40. The UPSIT was originally developed for North American English-speaking groups, but it has since been modified for other cultures and translated into several other languages. The Brazilian-Portuguese version of the scale was used in the present study. This version was developed through a series of experiments to adapt the test to Brazilian populations.⁹

Statistical Analysis

Data were analyzed using the Statistical Package for the Social Sciences (SPSS, IBM Corp., Armonk, NY, US) software, version 24. Descriptive statistics were performed using epidemiological data. The UPSIT scores were compared among the groups using a two-sample *t*-test for the parametric analyses and the Wilcoxon-Mann-Whitney test for the non-parametric analyses. The relationships among the variables were assessed using the Pearson correlation. The Chi-squared test was used to assess the associations among frequencies within the categorical variables. A series of hierarchical multiple linear regressions was used to study the independent influence of the years of residency on the UPSIT score. The same analysis was performed using the following variables: age, ethnicity, presence of comorbidities, and smoking status.

Quality Assessment

A Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist of cross-sectional studies¹⁰ was used as a guide to ensure the quality of the data presentation.

Results

Study Population and Demographics

Out of 50 potential subjects, 42 were included in the study. A total of eight subjects were excluded because they had acute respiratory infections. Out of all the included subjects, 47.6%

were male ($n = 20$), and 52.4% were female ($n = 22$) (►Table 1). The median age of the subjects was 27 years (range: 23 to 35 years). With regard to monthly household income, 38.1% ($n = 16$) were in Class B, and 61.9% ($n = 26$) were in Class A. Regarding ethnicity, 73.8% of the subjects were self-defined as White Brazilian ($n = 31$), 2.4%, as Black Brazilian ($n = 1$), and 23.8%, as mixed-race Brazilian ($n = 10$). The UPSIT results were correlated with age ($r = 0.33$; $p = 0.03$) and the number of continuous-use medications ($r = 0.35$; $p = 0.02$), but not with monthly household income.

Residency Programs

There was an equal number of subjects in the 2 residency groups (otorhinolaryngology residents: $n = 21$; other residents: $n = 21$). The group of other residents consisted of residents of: anaesthesiology (19.0%, $n = 4$), infectology (4.8%, $n = 1$), gastroenterology (4.8%, $n = 1$), medical clinic (9.5%, $n = 2$), nephrology (14.3%, $n = 3$), neurology (4.8%, $n = 1$), neurosurgery (4.8%, $n = 1$), oral and maxillofacial surgery (4.8%, $n = 1$), physiatry (9.5%, $n = 2$), and physical therapy (23.8%, $n = 5$). The residents were also divided into either a surgical (64.3%, $n = 27$) or clinical (35.7%, $n = 15$) group.

Comorbidities and Medication Use

Regarding comorbidities, 16.7% ($n = 7$) reported having a disease or condition. The most common diseases and conditions were hypothyroidism and asthma, but attention deficit hyperactivity disorder, melanoma, and shoulder dislocation were also reported. With regard to nasal diseases and conditions, 40.5% reported a previous diagnosis of nasal disease ($n = 17$), and, out of this group, 26.2% ($n = 11$) specifically reported rhinitis (►Table 1). In total, 40% of the participants reported continuous use of medications ($n = 17$). Considering the whole sample, we observed mainly the continuous use of contraceptives (26.8%, $n = 11$) and antidepressants (11.9%, $n = 5$) (venlafaxine, desvenlafaxine, escitalopram). A minority of subjects reported continuous use of topical corticosteroids (budesonide, mometasone, fluticasone), attention deficit hyperactivity disorder

Table 1 Demographic and clinical characteristics of the study population

	Otorhinolaryngology residents	Other residents	<i>p</i> -value
N	21	21	
Age (years)	27.9 ± 2.5	27.7 ± 2.7	0.70 ^a
University of Pennsylvania Smell Identification Test (UPSIT) score	35.0 ± 2.5	32.8 ± 3.9	0.02 ^a
Gender no. female (%)	11 (52.4)	11 (52.4)	1.00 ^b
Comorbidity (%)	19.0	14.3	0.68 ^b
Number of chronic use medication per individual, median	1	0	0.25 ^a
Nasal disease (%)	38.1	42.9	0.75 ^b
Rhinitis (%)	19.0	33.3	0.33 ^b
Continuous use of medication (%)	52.4	33.3	0.36 ^b
Smoking (%)	4.8	19.0	0.15 ^b

Notes: ^aTwo-sample *t*-test; ^bChi-squared test.

Table 2 University of Pennsylvania Smell Identification Test (UPSIT) scores regarding residency

Variables	N	Age*	UPSIT score*	Standard deviation	Median	Minimum	Maximum	p-value
Residency program								0.02 ^a
Otorhinolaryngology	21	28	35.0	2.5	35	29	40	
Others	21	27	32.8	3.9	34	26	39	
Year of residency								0.72 ^b
R1	18	27	33.8	3.7	34	26	40	
R2	13	28	33.3	3.5	34	27	38	
R3	10	29	34.9	3.2	35	27	39	
R4	1	28	34.0	–	–	–	–	
Residency type								0.23 ^b
Clinical	15	27	32.8	4.2	34	26	39	
Surgical	27	28	34.5	2.8	35	29	40	

Notes: *average; ^atwo-sample *t*-test for the UPSIT scores; ^bWilcoxon-Mann-Whitney test for the UPSIT scores.

drugs (methylphenidate and lisdexamfetamine), thyroid drugs (levothyroxine), systemic corticosteroids (prednisone), and pain medication (trometamol). There was no statistically significant difference in the UPSIT scores regarding the type of drug.

Olfactory Evaluation

With regard to olfactory function, 59.5% of the sample had normosmia, and 40.5% showed some grade of olfactory impairment. We observed no cases of severe microsmia or anosmia. The median overall UPSIT score was of 35 (range: 26–40). The otorhinolaryngology residents presented an average UPSIT score of 35.0, and the other residents, of 32.8 ($p = 0.02$) (►Table 2). The UPSIT scores according to the residency program, type and year of residency are presented in ►Table 2, and, according to the demographics, in ►Table 3. The UPSIT results were not found to be correlated with year of residency ($r = 0.86$; $p = 0.58$). ►Table 4 shows the values and associations between olfactory function and gender, specialty of the residency program, and year of residency. We found no differences in UPSIT results between groups in terms of gender, ethnicity, household income, specialty of the residency program, year of residency, comorbidities, presence of nasal diseases, presence of rhinitis, and smoking status. The multivariate analyses found age ($r = 0.32$, $p = 0.03$; 95% confidence interval for $\beta = 0.33$) to be an independent predictor of the UPSIT score.

Discussion

The results of the 40-item olfaction identification test demonstrated that otorhinolaryngology residents have greater olfactory capacity compared with residents of other specialties. Although small differences in the UPSIT scores were observed between these two groups (35.0 versus 32.8, $p = 0.02$), they were clinically significant. According to the UPSIT Administration Manual, normosmia is defined when UPSIT scores range from 34 to 40 points (males) and 35 to 40 points (females). Mild microsmia ranged from 30 to 33 points (males) and from 31 to

34 points (females). Thus, the average UPSIT score of otorhinolaryngology residents was classified as normosmia, whereas the other group had mild microsmia.⁸

We believe that the awareness of otorhinolaryngology residents regarding the prevention of nasal inflammatory and infectious processes should be associated with these results.^{11,12} In addition, we believe that easy access to free samples of topical nasal corticosteroids and the common practice of washing the nose with saline solution were significant factors that subjects forgot to mention during the interview. These factors would have prevented any inflammatory disease of the olfactory epithelium¹² and the common epistaxis caused by the dry climate of our geographic area.¹³ Moreover, these factors can also be considered as specific and involuntary olfactory training, which models the use of smell among wine and coffee experts.¹⁴ Future studies that examine the role of individual practices and training in the prevention of olfaction disorders should be performed.

With regard to gender, in our sample, women had better olfactory performance than men (34.2 versus 33.6 respectively). It is well known that, among the general population, women have greater olfactory and gustatory sensitivity than men,^{15,16} and, in a recent meta-analysis¹⁷ that assessed the effect of gender on odor identification, a superior performance was observed among women, but only among adults between 18 and 50 years of age. However, we observed a greater prevalence of dysosmia among females than among males (50% versus 30% respectively). The literature shows that the complaint of dysosmia is more frequent in women than in men.¹⁵

The percentage of smokers in the otorhinolaryngology group was much lower (4.8%) than that of the other group (19%). Although the differences in the UPSIT scores between smokers and non-smokers were not statistically significant ($p = 0.67$), we believe that this environmental factor could result in lower UPSIT scores, according to the available medical literature.^{8,9} Only 11.9% of our sample were smokers, but, unfortunately, we did not control the tobacco load of the participants.

In a previous study¹⁸ on the cross-cultural adaptation of the UPSIT for use in the Brazilian population ($n = 49$; mean

Table 3 University of Pennsylvania Smell Identification Test (UPSIT) scores according to demographics

Variables	N	Mean score	Standard deviation	Median	Minimum	Maximum	p-value
Entire sample	42	33.9	3.4	35	26	40	–
Gender							0.54 ^a
Male	20	33.6	3.5	35	26	39	
Female	22	34.2	3.5		27	40	
Ethnicity							0.76 ^b
White Brazilian	31	34.1	3.3	35	27	39	
Black Brazilian	1	–	–	–	–	–	
Mixed Brazilian	10	33.2	4.2	34	26	40	
Monthly household income							0.80 ^b
Group 3	16	33.9	2.7	34	29	38	
Group 4	26	33.9	3.9	35	26	40	
Comorbidity							0.46 ^b
Yes	7	34.5	4.0	35	29	39	
No	35	33.8	3.4	34	26	40	
Nasal disease							0.43 ^b
Yes	17	34.3	3.5	35	27	39	
No	25	33.6	3.5	34	26	40	
Rhinitis							0.26 ^b
Yes	11	33.0	3.6	34	27	39	
No	31	34.2	3.4	35	26	40	
Smoking							0.67 ^b
Yes	5	34.6	3.8	35	29	39	
No	37	33.8	3.4	35	26	40	

Notes: ^aTwo-sample t-test; ^bWilcoxon-Mann-Whitney test.

Table 4 Percentage of each group divided by clinical classification of olfactory deficit

Degree of olfactory loss	Normosmia	Mild microsmia	Moderate microsmia	p-value
Gender				0.21 ^a
Male	70.0	10.0	20.0	
Female	50.0	31.8	18.2	
Residency group				0.18 ^a
Otorhinolaryngology residents	66.7	23.8	9.5	
Other residents	52.4	19.0	28.6	
Year of residency				0.51 ^a
R1	50.0	33.3	16.7	
R2	53.8	23.1	23.1	
R3	80.0	–	20.0	
R4	100.0	–	–	

Note: ^aChi-squared test.

age: 30 years), which used the same translated version of the test, the investigators observed a mean score of 35 points. This mean score was higher than the one observed in the present study, suggesting that residents of health programs had a worse sense of smell than the general population. Work-related factors impacting the lives of the residents,

such as depression and excessive workload, may influence the well-being of these individuals,¹⁹ and could be related to lower UPSIT scores.^{20,21} Sleep deprivation also affects the activity of the insula and piriform cortex, a brain region responsible for odor processing.²² Moreover, the greater olfactory capacity observed in senior residents in

comparison with first-year residents can be explained by the odor learning process that has been confirmed in multiple psychosocial studies⁴ and has served as a foundational tent for clinical olfactory training as we know today.⁴

In the sample of the present study, half of the female participants and half of the students in the first year of residency presented with olfactory dysfunction. The differences observed may also be due to the fact that menstrual cycles are associated with higher levels of stress among first-year residents.¹⁹ Instead of findings such as the one demonstrating that brain activation was consistently lower in females than in males, using functional magnetic resonance imaging,²² we still have more questions than answers regarding this complex understanding of the functional properties of the human olfactory system and neuroendocrine factors.²³

Fornazieri et al.¹⁸ showed that socioeconomic status and schooling influence olfactory performance. Unfortunately, we were unable to confirm this hypothesis, because our sample consisted only of subjects with high socioeconomic status and level of schooling. We aimed for a sufficiently large sample size to be able to generalize our results to the general population of residents. Our sample size can be externally validated, especially when compared with the UPSIT cultural adaptation study performed by Fornazieri et al.¹⁸ with 49 participants.

Conclusion

The present study demonstrated that otorhinolaryngology residents have greater olfactory capacity compared with other residents. Future studies should explore the relevant factors of olfactory impairment and their impact on the quality of life of this population.

Conflict of Interests

The authors have no conflict of interests to declare.

Acknowledgements

The authors would like to thank the Otolaryngologist residents Shaadya R Said and Aline Almeida Liberato for their contribution in some data collection.

References

- Fitzgerald FT, Tierney LM Jr. The bedside Sherlock Holmes. *West J Med* 1982;137(02):169–175
- Nagappan PG, Subramaniam S, Wang DY. Olfaction as a soldier—a review of the physiology and its present and future use in the military. *Mil Med Res* 2017;4:9
- Smith DV. Assessment of patients with taste and smell disorders. *Acta Otolaryngol Suppl* 1988;458:129–133
- Turner JH. Olfactory training: what is the evidence? [published online ahead of print, 2020 Aug 10] *Int Forum Allergy Rhinol* 2020. Doi: 10.1002/alr.22681
- Brämerson A, Johansson L, Ek L, Nordin S, Bende M. Prevalence of olfactory dysfunction: the skövde population-based study. *Laryngoscope* 2004;114(04):733–737
- Sugiura M, Aiba T, Mori J, Nakai Y. An epidemiological study of postviral olfactory disorder. *Acta Otolaryngol Suppl* 1998;538:191–196
- Haxel BR, Nisius A, Fruth K, Mann WJ, Muttray A. [Deficits in medical counseling in olfactory dysfunction]. *HNO* 2012;60(05):432–438
- Doty RL, Shaman P, Dann M. Development of the University of Pennsylvania Smell Identification Test: a standardized micro-encapsulated test of olfactory function. *Physiol Behav* 1984;32(03):489–502
- Fornazieri MA, Doty RL, Santos CA, Pinna FdeR, Bezerra TF, Voegels RL. A new cultural adaptation of the University of Pennsylvania Smell Identification Test. *Clinics (São Paulo)* 2013;68(01):65–68
- Vandenbroucke JP, von Elm E, Altman DG, STROBE Initiative. et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): explanation and elaboration. *Int J Surg* 2014;12(12):1500–1524
- Kaper NM, Aarts MCJ, van Benthem PPG, van der Heijden GJMG. Otolaryngologists adhere to evidence-based guidelines for chronic rhinosinusitis. *Eur Arch Otorhinolaryngol* 2019;276(04):1101–1108
- Fowler J, Chin CJ, Massoud E. Rhinitis medicamentosa: a nationwide survey of Canadian otolaryngologists. *J Otolaryngol Head Neck Surg* 2019;48(01):70
- Rezende GL, Granjeiro RC, Furtado PL, Pinheiro GB, Nakanishi M. Is dry climate related to hospital admission for epistaxis? *Int Arch Otorhinolaryngol* 2009;13:172–177
- Croijmans I, Majid A. Not All Flavor Expertise Is Equal: The Language of Wine and Coffee Experts. *PLoS One* 2016;11(06):e0155845
- Deems DA, Doty RL, Settle RG, et al. Smell and taste disorders, a study of 750 patients from the University of Pennsylvania Smell and Taste Center. *Arch Otolaryngol Head Neck Surg* 1991;117(05):519–528
- Doty RL. Olfactory dysfunction and its measurement in the clinic and workplace. *Int Arch Occup Environ Health* 2006;79(04):268–282
- Wang X, Zhang C, Xia X, Yang Y, Zhou C. Effect of gender on odor identification at different life stages: a meta-analysis. *Rhinology* 2019;57(05):322–330
- Fornazieri MA, Doty RL, Bezerra TFP, et al. Relationship of socioeconomic status to olfactory function. *Physiol Behav* 2019;198:84–89
- Rezende GL, Mello MS, Granjeiro RC, Nakanishi M, Oliveira CA. The quality of life among Otorhinolaryngology residents in Distrito Federal (Brazil). *Braz J Otorhinolaryngol* 2011;77(04):466–472
- Hur K, Choi JS, Zheng M, Shen J, Wrobel B. Association of alterations in smell and taste with depression in older adults. *Laryngoscope Investig Otolaryngol* 2018;3(02):94–99
- Kohli P, Soler ZM, Nguyen SA, Muus JS, Schlosser RJ. The Association Between Olfaction and Depression: A Systematic Review. *Chem Senses* 2016;41(06):479–486
- Bhutani S, Howard JD, Reynolds R, Zee PC, Gottfried J, Kahnt T. Olfactory connectivity mediates sleep-dependent food choices in humans. *eLife* 2019;8:e49053
- Doty RL, Cameron EL. Sex differences and reproductive hormone influences on human odor perception. *Physiol Behav* 2009;97(02):213–228