



Common Contaminants in Next-Generation Sequencing That Hinder Discovery of Low-Abundance Microbes

Martin Laurence¹, Christos Hatzis², Douglas E. Brash^{3*}

1 Shipshaw Labs, Montreal, Quebec, Canada, **2** Yale Comprehensive Cancer Center, Yale School of Medicine, New Haven, Connecticut, United States of America, **3** Department of Therapeutic Radiology, Yale School of Medicine, New Haven, Connecticut, United States of America

Abstract

Unbiased high-throughput sequencing of whole metagenome shotgun DNA libraries is a promising new approach to identifying microbes in clinical specimens, which, unlike other techniques, is not limited to known sequences. Unlike most sequencing applications, it is highly sensitive to laboratory contaminants as these will appear to originate from the clinical specimens. To assess the extent and diversity of sequence contaminants, we aligned 57 “1000 Genomes Project” sequencing runs from six centers against the four largest NCBI BLAST databases, detecting reads of diverse contaminant species in all runs and identifying the most common of these contaminant genera (*Bradyrhizobium*) in assembled genomes from the NCBI Genome database. Many of these microorganisms have been reported as contaminants of ultrapure water systems. Studies aiming to identify novel microbes in clinical specimens will greatly benefit from not only preventive measures such as extensive UV irradiation of water and cross-validation using independent techniques, but also a concerted effort to sequence the complete genomes of common contaminants so that they may be subtracted computationally.

Citation: Laurence M, Hatzis C, Brash DE (2014) Common Contaminants in Next-Generation Sequencing That Hinder Discovery of Low-Abundance Microbes. PLoS ONE 9(5): e97876. doi:10.1371/journal.pone.0097876

Editor: Tom Gilbert, Natural History Museum of Denmark, Denmark

Received: March 4, 2014; **Accepted:** April 25, 2014; **Published:** May 16, 2014

Copyright: © 2014 Laurence et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: DEB is supported by NIH Skin Cancer SPORE grant 5P50CA121974, CH was supported by Yale University, and ML was supported by Shipshaw Labs Inc. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. No patents, products in development or marketed products are related to the findings reported in this paper. These affiliations do not alter the authors adherence to PLOS ONE Policies on sharing data and materials. The Leif Microbiome Analyzer is a commercial product of Shipshaw labs.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: douglas.brash@yale.edu

Introduction

Systematic pathogen discovery based on unbiased high-throughput sequencing [1] was first used in 2008 to detect two novel viruses by pyrosequencing clinical specimens. The first study attributed 2 fragments out of 395,734 (0.0005%) to a novel polyomavirus [2]; the second study attributed 14 fragments out of 103,632 (0.0135%) to a novel arenavirus [3]. Recent improvements in sequencing technology have rendered this method much more sensitive for detecting low-abundance pathogens and other medically important microbes in clinical specimens. For example, PathSeq [4], a bioinformatics toolkit designed to detect novel sequences, was recently used to identify a novel *Bradyrhizobium* species in clinical specimens using whole metagenome shotgun DNA libraries and the Illumina sequencing platform, which can currently produce up to 186 million read pairs per lane [5]. A novel infectious species representing only 0.000001% of the total DNA in a clinical specimen can theoretically be detected using a single Illumina HiSeq 2000 lane, since at least one read pair out of 186 million would originate from the novel microbe in 84% of runs.

Unbiased high-throughput sequencing has been suggested as a way of detecting etiological microbes in cancer tissue [6], an approach we consider promising for prostate cancer [7,8]. To facilitate such studies we constructed the Leif Microbiome Analyzer, a bioinformatics tool similar to PathSeq which was designed to eliminate the need for cluster computing typically required by NCBI blastn based tools such as PathSeq—even when

aligning against the largest NCBI BLAST databases. A basic assumption in calculating the sensitivity of such approaches is that one microbial read is informative [6]. However, while testing the Leif Microbiome Analyzer by examining two clinical samples for reads not aligning to the human genome, we encountered many reads from diverse species not known to be part of the human microbiome, suggesting the presence of contamination; members of the *Bradyrhizobium* genus were particularly prominent. If this situation arises commonly, it would be cost-prohibitive to screen candidate non-aligning reads using polymerase chain reaction (PCR) on the original specimens. For example, 1000 novel microbe reads would entail 1000 confirmatory PCR reactions on the original tissue sample; at \$100 each, including custom primers and optimization, the \$100,000 validation cost would greatly exceed the \$2500 cost of one Illumina HiSeq 2500 lane and would be prone to error. As the cost of sequencing continues to decline faster than the cost of PCR, this overhead is expected to worsen. In the case of novel contaminant microbes whose genome has not been completely sequenced, the 1000 confirmatory PCR reaction figure cannot be reduced by sampling only some reads of each species, as it is not possible to know which reads arose from the same species.

Various types of contamination in sequencing runs have been reported before [9,10], as well as a few mitigation techniques [11–13] which are not suitable for the discovery of novel microbes. We therefore inquired whether contamination was restricted to our libraries or sequencing center, or is a general property of next-generation sequencing workflows.

Materials and Methods

Sample Selection

Initial sequence analysis was performed on two human prostate samples which are not reported here but which motivated the larger studies of publicly available genomic sequences from cultures expected to contain cells of a single species. To obtain sequences of healthy human cells generated by next-generation sequencing workflows which closely resembled our own samples and runs, we searched the NCBI Sequence Read Archive database for “1000 genomes project AND hiseq AND CEU”. The search was restricted to Illumina HiSeq because the Leif Microbiome Analyzer is optimized for this platform; CEU are genomes of Northern European ancestry. Paired-end whole genome shotgun sequencing runs done for the 1000 Genomes Project [14] used DNA extracted from cultured Epstein-Barr virus immortalized B-lymphocytes drawn from blood of healthy individuals. A full description of specimens can be found at www.ncbi.nlm.nih.gov/sra for each of the runs used. This search gave 44 runs, some of which were too small (<18 M read pairs), had reads that were too short (<2×100), or crashed when converting to FASTQ format using the NCBI tool fastq-dump. Of the 44 runs, 23 passed these criteria. This initial search resulted in only Baylor College of Medicine and Broad Institute runs, so we then widened the search to runs from other sequencing centers. An SRA search for “hapmap AND hiseq AND CEU” resulted in 1298 runs. From these, runs from other sequencing centers were chosen randomly for a total of 57 runs. These include sequencing centers from the US, UK, and Germany. Non-human next-generation sequencing runs were also analyzed by searching the NCBI Sequence Read Archive database for “*Candida albicans* AND HiSeq AND WGS”; this search resulted in 198 runs, 142 of which were suitable for analysis (read pairs formatted as 2×100, >2M read pairs and *Candida albicans* species). All 142 runs were done at the Broad Institute.

Sequence Alignment

For all nucleotide sequence alignments in this study, we used the Leif Microbiome Analyzer version 0.7.3 (www.shipshaw.com/leif), a short-read alignment tool that uses an algorithm similar to NCBI blastn and is optimized for paired-end reads produced by the Illumina platform. Reference sequences were obtained from the four largest NCBI BLAST databases (“nt”, “human_genomic”, “other_genomic” and “wgs”) downloaded in FASTA format on April 1st 2014 from ftp.ncbi.nlm.nih.gov/blast/db/FASTA. These databases contain taxonomically labeled Genbank nucleotide sequences of organisms deemed BLAST-worthy by NCBI.

We screened NCBI BLAST eukaryotic sequences for possible *Bradyrhizobium* contamination by first randomly sampling *Bradyrhizobium* sequences found in the NCBI BLAST databases (simulating an Illumina sequencing run producing 200,000 2×100 read pairs), then aligning these read pairs against all eukaryotic sequences in the NCBI BLAST databases. Positive genomes involved sequencing centers from the US, Canada, and China. This screening technique is not exhaustive since only a random sampling of *Bradyrhizobium* sequences was used; aligning against all *Bradyrhizobium* sequences (or all bacterial sequences) is beyond the means of an academic research project. The list of commands required to replicate this analysis are provided in Text S1.

Publicly available Illumina HiSeq 2000 whole genome runs from the 1000 Genomes Project and *Candida albicans* runs were downloaded from the NCBI Sequence Read Archive database. Read pairs from these runs were aligned against the four largest NCBI BLAST databases, and contamination candidates were

identified as described in the next section. The list of commands required to replicate this analysis are provided in Text S2 and S3.

Identification of Contamination Candidates

The Leif Microbiome Analyzer performed the following ten steps to identify contamination candidates in 1000 Genome Project runs. 1) Bases whose quality letter was ≤ ‘%’ were deemed incorrectly called (‘N’). 2) Read pairs which contained many bases deemed to be incorrectly called were tagged as “low-quality” and discarded. Remaining read pairs satisfied the following criteria: ≥ 90% of bases in each read were deemed correctly called and ≥ 50% of the 32 base substrings in each read contained only bases deemed correctly called. 3) Read pairs which contained a 32 base substring matching exactly with human, EBV or phage sequences were discarded. 4) Read pairs which were nearly identical (bases 5 to 64 were compared and deemed nearly identical if ≥57 bases matched) were deemed to be clones produced as an artifact of library preparation and discarded. 5) The DUST algorithm [15] was used to mask bases which were deemed part of a low entropy sequence. 6) Read pairs which contained too many bases masked by DUST were tagged as “low-entropy” and discarded. Remaining read pairs satisfied the following criterion: ≥50% of the 32 base substrings in each read contained only unmasked bases. 7) Read pairs which contained an identical or reverse complement unmasked 32 base substring were merged into a “contig-like group”, and were deemed to have originated from the same template DNA strand pair. 8) A few read pairs from each “contig-like group” were randomly sampled for alignment; the number of sampled read pairs in each “contig-like group” was determined by the equation $1 + \text{ceiling}(\log_2(\text{number_of_read_pairs_in_group}))$. 9) Sampled read pairs were aligned to all sequences in the NCBI “nt”, “human_genomic”, “other_genomic” and “wgs” databases by Leif’s qblast command. Reads (mates) in each read pair were aligned both independently (single alignment) and together (dual alignment, which supports fragments of up to 1024 nucleotides in length). Single alignment was used to produce all results reported here except Text S4, S5, S6 and S7; though single alignment considers mates independently, mates are always assigned to the same “contig-like group”, so the link between mates is not completely lost. 10) “Contig-like groups” whose sampled read pairs aligned with primate, EBV or phage sequences were discarded. A “contig-like group” was deemed to have aligned to primate, EBV or phage sequences if ≥50% of sampled reads had ≥70% homology to primate, EBV or phage sequences present in the NCBI BLAST databases. For example, if a “contig-like group” contained three read pairs, then three out of the six reads must have ≥70% homology to primate, EBV or phage sequences to be discarded during this step.

The consensus set produced by Leif reports the taxonomic node which encompasses all sequences aligning at a given homology percentage for each sampled read individually, and also merged for all sampled reads in a “contig-like group”. For example, in a “contig-like group” containing a single read pair, if read 1 has 93% homology to *Streptococcus mitis* and read 2 has 89% homology to *Enterococcus faecalis*, then the consensus taxonomic node for this “contig-like group” would be: 100%–94% = none, 93%–90% *Streptococcus mitis* and 89%–0% Lactobacillales—since Lactobacillales is the narrowest taxonomic node which contains both *Streptococcus mitis* and *Enterococcus faecalis*. Homology tests described below use the merged consensus set of each “contig-like group” to assign taxonomic classification to all read pairs within this group. “Contig-like groups” with a <90% homology to all database sequences were not given a taxonomic classification and were instead reported as “Low homology”: sampled read pairs in these

groups either originated from a novel species/strain or contained sequence errors. “Contig-like groups” with $\geq 90\%$ homology to any NCBI BLAST database sequence were deemed to be known contaminants. These were split into three broad categories: “Eukaryote”, “Prokaryote” and “Viral”. The “Dual homology” category contains “contig-like groups” with a high homology to two or more broad categories; since these are non-specific matches, they should not be reported as either “Eukaryote”, “Prokaryote” or “Viral”. Finally, the “Prokaryote” category was split into five parts. Many “contig-like groups” aligned specifically to ultrapure water system contaminants reported by Kulakov et al. [16] (genera *Bradyrhizobium*, *Rhizobium/Agrobacterium*, *Sphingomonas*, *Burkholderia*, *Ralstonia*, *Pseudomonas*, *Stenotrophomonas*, *Flavobacterium*) and to Enterobacteriaceae (probably to *Escherichia coli*, although the alignments were not sufficiently specific to exclude other Enterobacteriaceae species). Since *Bradyrhizobium* and *Bradyrhizobium sp. DFCI-1* were highly prevalent, they were reported in separate columns. For a “contig-like group” to be reported under a taxonomic name, all alignments to other taxonomic names were required to have a significantly lower homology (a 5% homology margin was chosen to generate Figure 1). For example, if a “contig-like group” had 93% homology to a eukaryote and 89% to a prokaryote (4% homology margin), its read pairs are reported under the taxonomic node “cellular organisms”, which is placed in the “Dual homology” category in Figure 1.

Candida albicans runs were analyzed similarly to human 1000 Genomes Project runs with the following differences: 1) the low quality bases were identified using $\leq \#$ (rather than $\leq \%$); 2) reads aligning to any sequence from the genus *Candida* were discarded (rather than reads aligning with any sequence from primates or EBV).

Results

The presence of significant levels of *Bradyrhizobium* genus sequence in our two clinical samples led us to examine, as a negative control, reads from two human 1000 Genomes Project runs which did not align to the human genome. Manual review of Leif’s qblast results for these two runs (SRR768303 and SRR385759) aligned against the four largest NCBI BLAST databases revealed that about one third of non-human reads matched specifically with *Bradyrhizobium* sequences, especially *Bradyrhizobium sp. DFCI-1* (Text S4). In addition, some reads matched specifically to exotic species such as the Tibetan antelope, *Pantholops hodgsonii*. Further investigation of *Pantholops hodgsonii* nucleotide sequences in Genbank revealed that regions of this genome align very well with *Bradyrhizobium* sequences. The presence of *Bradyrhizobium* sequences in our two clinical specimens, in two randomly selected 1000 Genomes Project runs and in the assembled genome of *Pantholops hodgsonii* suggests that this bacterium is a common contaminant in next-generation sequencing workflows.

Randomly selected *Bradyrhizobium* sequences were then aligned against eukaryotic sequences in the NCBI BLAST databases, revealing many additional species which match specifically with *Bradyrhizobium* sequences (Table 1). It appears that *Bradyrhizobium* contamination may have been present in the sequencing runs used to assemble some genomes, such as the *Pantholops hodgsonii* genome, and were incorporated into assembled genomes in the NCBI Genome database. The problem is therefore not limited to searches for low-abundance microbes: contamination can go unrecognized in *de novo* assembled genomes, due to low coverage or inadequate curation. The seven Genbank entries listed in

Table 1 suggest that *Bradyrhizobium* contamination is not limited to a single sequencing center, technology or eukaryotic species.

To compare a large number of whole genome shotgun sequencing libraries processed at different sequencing centers by a standardized protocol and not expected to contain etiologic microbes or be complicated by issues of handling of pathology samples, we used the Leif Microbiome Analyzer to identify non-aligning reads from 57 Illumina HiSeq 2000 runs performed by various sequencing centers on 1000 Genomes Project samples. The results are shown in Figure 1. Known contaminant sequences (defined here as read pairs in a human sample which match specifically with NCBI BLAST sequences other than primate, EBV, and phage) were present in all runs, varying from 0.000007% to 0.015% of total read pairs with a median of 0.0003%. Low homology sequences (defined here as read pairs which did not match with sequences in the NCBI BLAST databases, usually due to either a high number of sequencing errors or to the presence of novel contaminant strains/species in the run) are listed in a separate column and are not counted as known contaminant sequences—although some may well originate from novel contaminants. Eukaryotic DNA contamination was common, typically aligning to the genus *Bos*, which may be originating from fetal calf serum used in cell culture media. *Bradyrhizobium* contamination was found in 25 out of 57 runs. This particular contaminant varied from center to center (Figure 1). The highest levels of *Bradyrhizobium* were found in runs from center BCM, where 19 runs out of 30 were contaminated, reaching levels as high as 0.003% of reads (Figure 1). Some runs from centers SC, BI, and MPIMG contained a few reads which matched specifically with *Bradyrhizobium sp. DFCI-1* (Figure 1 and Text S5, S6 and S7). No *Bradyrhizobium* read pairs were found in two runs submitted by Illumina or four runs submitted by WUGSC. However, runs from these centers did show contamination from other organisms. Other species commonly encountered included genera *Rhizobium/Agrobacterium*, *Sphingomonas*, *Burkholderia*, *Ralstonia*, *Pseudomonas*, *Stenotrophomonas*, *Flavobacterium* (reported together in column “Ultrapure water system contaminants – Other” of Figure 1); sequence alignments to all species are reported in Spreadsheet S1. These 57 sequencing runs indicate that contamination is widespread and *Bradyrhizobium sp. DFCI-1* is a prominent contaminant, but contamination levels are highly variable between runs—even runs from the same sequencing center.

Alignment results from 142 *Candida albicans* sequencing runs performed using Illumina HiSeq 2000 are reported in Table S1 and in Spreadsheet S2. *Bradyrhizobium* contamination can be found in 136 out of 142 runs, reaching levels as high as 0.0126% of reads. Ultrapure water system contaminants were detected in all runs.

Discussion

Many different microbe discovery techniques have been developed over the last 150 years [1]; the advent of next-generation sequencing technology has enabled molecular detection techniques which allow the discovery of fastidious or unculturable microbes in samples containing mixed flora such as the human skin [17]. These new techniques revolutionized our understanding of the human microbiome, revealing many previously unknown species in clinical specimens from healthy individuals.

The most common technique in use today for microbiome surveys is based on selective amplification of small regions of microbial DNA using consensus PCR primers prior to high-throughput sequencing. This technique has three major

Center	Run	Primate, EBV, phage or discarded due to low quality/ duplicate/ low entropy	All read pairs								
			Contamination candidates								Low homology (<90%)
			Known contaminants (high homology - ≥90%)								
			Eukaryote	Prokaryote			Enterobacteriaceae	Others & dual homology	Viral	Dual homology	
Ultrapure water system contaminants		Others & dual homology									
<i>Bradyrhizobium</i>		Others & dual homology									
<i>DFCI-1</i>											
BCM	SRR385754	50112167	9	438	433	431	0	172	2	61	543
BCM	SRR385755	43233137	1	227	179	234	0	88	3	23	278
BCM	SRR385758	48859661	12	0	0	1	6	32	0	66	61
BCM	SRR385759	79360966	9	1153	1126	1514	0	517	6	190	1736
BCM	SRR385761	52277896	8	0	0	2	4	14	0	84	209
BCM	SRR385762	103886262	164	0	0	1	1	127	0	5	278
BCM	SRR385763	53807301	9	321	282	363	0	120	1	53	411
BCM	SRR385764	46462143	6	149	145	169	0	65	0	20	277
BCM	SRR385765	43367799	18	625	623	792	171	298	4	4015	1055
BCM	SRR385767	63753008	44	0	0	3	0	27	0	131	747
BCM	SRR385768	46642864	28	0	0	2	1	18	0	62	470
BCM	SRR385769	92608130	21	0	0	22	0	222	0	112	361
BCM	SRR385770	36818041	7	1	4	6	0	2	0	1	4
BCM	SRR385772	60254401	9	7	10	4	16	21	1	460	23
BCM	SRR385773	52061042	0	30	76	18	0	10	7	16	38
BCM	SRR385774	42439279	2	28	41	14	14	22	1	372	39
BCM	SRR385776	42169205	3	35	61	27	0	11	0	3	50
BCM	SRR385777	48462601	4	34	54	24	0	24	0	0	43
BCM	SRR393988	58072454	1	145	102	156	0	57	1	14	211
BCM	SRR393989	49568344	1	67	65	63	0	35	0	13	102
BCM	SRR393990	51017564	34	0	0	4	0	34	0	7	561
BCM	SRR393993	48369009	82	0	0	5	13	0	0	90	67
BCM	SRR393994	56927949	25	2	0	2	0	26	0	5	436
BCM	SRR400037	47847795	16	0	0	21	8	87	0	153	92
BCM	SRR741366	74950213	22	56	40	1	15	36	0	267	56
BCM	SRR768303	69790675	9	18	19	0	2	27	0	23	12
BCM	SRR768304	51864089	15	12	11	2	8	15	2	88	14
BCM	SRR768309	80919926	13	24	23	4	5	18	0	127	24
BI	SRR067576	18153122	21	0	0	15	35	7	0	639	280
BI	SRR067577	46251353	37	0	0	0	0	5	0	0	40
BI	SRR067578	46490939	49	0	0	0	0	4	0	1	53
BI	SRR067579	46015083	27	0	0	0	0	6	0	1	51
BI	SRR068130	125872967	19	0	2	1	0	139	0	106	47
BI	SRR075005	29392952	7	0	1	1	0	43	0	44	17
BI	SRR075006	62188209	496	5	5	0	0	3	0	5	78
ILLUM	ERR091571	211437693	95	0	0	2	1	58	0	6	64
ILLUM	ERR091575	205185449	5	0	0	42	1	21	0	20	103
MPIMG	ERR233225	104620504	0	0	0	4	1	1	0	1	8
MPIMG	ERR233227	106377916	638	0	0	2	5	7	1	62	27
MPIMG	ERR233301	119475893	28	0	0	0	0	3	0	3	18
MPIMG	ERR233302	111852187	63	0	0	54	0	154	0	10	219
MPIMG	ERR234321	108573882	725	0	0	1	0	0	2	4	14
MPIMG	ERR234322	113157820	473	0	0	1	1	12	0	5	30
MPIMG	ERR234323	114214873	13	0	0	0	0	17	0	1	9
MPIMG	ERR234324	118549617	1235	0	0	0	0	2	0	0	21
MPIMG	ERR234325	111531797	386	0	0	0	0	4	0	0	32
MPIMG	ERR234327	112766941	530	0	0	1	7	12	0	16	75
MPIMG	ERR234328	114857252	2388	0	0	1	1	2	0	0	128
MPIMG	ERR234329	111235200	188	0	0	0	0	4	0	2	67
MPIMG	ERR239333	116722187	892	0	0	3	0	3	0	9	50
MPIMG	ERR239334	105779573	577	0	1	0	0	10	0	8	362
SC	ERR050082	42242519	89	0	1	7	0	20	0	6	2432
SC	ERR050083	66388415	61	1	0	4	0	18	0	1422	1146
WUGSC	SRR211275	43009432	50	0	0	0	0	93	2	4	59
WUGSC	SRR211278	55193260	23	0	0	0	1	42	0	7	39
WUGSC	SRR407429	41694818	6	0	0	1	1	23	4	27	13
WUGSC	SRR407508	42586738	240	0	0	1	2	87	2	42	33

Figure 1. The contents of non-aligning reads from 57 human whole genome sequencing runs. Categories are defined in Methods. Sequencing center acronyms in this table are: Baylor College of Medicine (BCM), the Broad Institute (BI), Illumina (ILLUM), the Max Planck Institute for Molecular Genetics (MPIMG), the Sanger Center (SC), and Washington University Genome Sequencing Center (WUGSC). Runs are sorted alphabetically by center, then by SRA number which are assigned successively over time. Units are read pairs.
doi:10.1371/journal.pone.0097876.g001

advantages over unbiased high-throughput sequencing: 1) most human DNA is eliminated prior to sequencing, which reduces sequencing costs for equal assay sensitivity; 2) only a single region of the genome is amplified, so aligning to all known sequences is not required, reducing alignment time approximately one hundred thousand fold; 3) only a single region of the genome is amplified, allowing the abundance of each novel species/sequence to be quantified by counting identical reads, rather than by doing a custom quantitative PCR (qPCR) reaction for each novel sequence. However, this technique has one major drawback: some medically important species' DNA cannot be amplified by consensus PCR primers, so it only paints a partial picture of the microbiome. Despite these three disadvantages of unbiased high-throughput sequencing compared to the conventional method listed above, this new technique will likely enable a second major breakthrough in microbiome analysis, revealing medically important species whose DNA is not amplified by typically used consensus PCR primers, as is the case for *Pneumocystis jirovecii*, *Encephalitozoon hellem* and many other species.

Unbiased high-throughput sequencing can be done using either DNA or RNA. Library preparation for DNA is much simpler and faster than for RNA. For clinical applications or to discover novel microbes during an outbreak, this is a key consideration—especially with the introduction of real-time portable sequencers such as Oxford Nanopore's MinION, whose complete workflow is very short. RNA sequencing was used in the two first unbiased high-throughput sequencing studies which discovered novel viruses in human specimens [2,3]. It has two major advantages over DNA sequencing: 1) some types of viruses do not contain DNA (such as RNA viruses and retroviral virions), thus would be missed; 2) the protein product can easily be deduced from mRNA and aligned—and since proteins are typically better conserved than nucleotide sequences, protein alignment may reveal more accurate taxonomic information. For example, protein homology to known viruses allowed Palacios et al. [3] to identify unknown mRNA sequences as likely originating from an arenavirus, whereas nucleotide homology gave ambiguous results (see Table 2 of their article); this approach rendered their experiment immune to bacterial contamination and works well for studies focusing on viruses.

It is unclear whether DNA or RNA sequencing will typically yield a higher microbial/human read ratio, and thus be more sensitive: this property is obviously microbial species dependent. Furthermore, the minimum microbial/human read ratio which results in adverse health outcomes (such as acute illness, chronic inflammation or increased cancer risk) is not known, and is very likely species dependent as well. Feng et al. found a viral/human RNA read ratio of 0.0005% and associated this novel infectious agent with Merkel cell carcinoma [2]; Palacios et al. found a viral/human RNA read ratio of 0.0135%, and this novel virus caused febrile illness resulting in the death of four patients [3]. It is not possible to establish a lower bound of microbial/human read ratio at which health outcomes would be unaffected based on currently available data.

In our prostate microbiome study and in the present study, we chose whole metagenome shotgun sequencing and Illumina's HiSeq platform as the simplest, cheapest (at equal sensitivity) and most reliable unbiased high-throughput sequencing protocol. We used the Leif Microbiome Analyzer in order to minimize costs;

without this tool, the bioinformatics cost for this study would have been about fifty times higher, putting it well beyond our means (see Table S2 and S3). Aside from the presence of contaminants described here, this approach performed remarkably well, detecting many unknown DNA sequences which are novel species candidates. However, the qPCR step required to accurately assess the abundance of these microbes in clinical specimens and link them to prostate cancer proved too costly and uncertain for our project. Before proceeding to the very expensive and labor intensive qPCR step, we chose to study publicly available next-generation sequencing runs in order to better understand the scope and impact of common contaminants such as *Bradyrhizobium* species.

Bradyrhizobium sp. *DFCI-1* was first discovered in cord colitis syndrome specimens [5], suggesting that it may be medically important. Another novel *Bradyrhizobium* species has been reported in a blood culture from a patient with a poorly defined illness [18]. Prior to the cord colitis syndrome study, *Bradyrhizobium* species were not believed to infect humans. *Bradyrhizobium* species are known to be common contaminants in industrial ultrapure water systems, as a consequence of their predilection for nitrogen-flushed water [16]. Other microbes reported in ultrapure water systems and found here include genera *Rhizobium*/*Agrobacterium*, *Sphingomonas*, *Burkholderia*, *Ralstonia*, *Pseudomonas*, *Stenotrophomonas* and *Flavobacterium*. The presence of *Bradyrhizobium* sequences in several Genbank entries of eukaryotic species (Table 1), in sequencing runs from human specimens (Figure 1) and from *Candida albicans* specimens (Table S1) strongly suggests that it is a ubiquitous laboratory contaminant, although the exact source of contamination remains unclear: it could have been inserted during cell culture, DNA extraction, library preparation or sequencing.

Conclusion

Novel sequences originating from clinical specimens are of great interest, but they and novel laboratory contaminants both appear in the “Low homology” column in Figure 1. It is very difficult to distinguish microbes of interest from novel contaminants by looking at individual reads. The presence of contamination from organisms that have not been fully sequenced and included in NCBI BLAST databases is particularly problematic when considering projects that aim to identify novel microbes present at low prevalence in clinical specimens, as it is possible that this contamination would obscure detection of such microbes. This problem could be addressed either by eliminating the source of contamination in the laboratory or at the reagent supplier, or by fully sequencing the genomes of contaminants known to be present in these sequencing runs, allowing them to be eliminated during data analysis. A UVC dose of 8000 J/m², four times the dose specified by the US EPA for making surface water sources safe for drinking, will introduce a sufficient density of cyclobutane pyrimidine dimers to prevent ~99.99% of 100 nucleotide long DNA fragments from being PCR-amplified or sequenced. The numbers of contaminant reads encountered in Figure 1 and Table S1 suggests that this level of inactivation may be necessary.

We advocate a directed effort to sequence contaminant genomes, as this method is well suited to detecting *Bradyrhizobium* and other contaminants that may already be present in Genbank

Table 1. Genbank entries which appear to be contaminated.

Genbank sequence labeled as a eukaryote	Accession	Contig N50	Taxonomic label	Sequencing center	Sequencing platform
CCTCGACGTAGCGGATGACATCCTCAAGCTTGCG CTCGCGAGGGCAGCTTCGCCAGATCCCAAAGCT CGATGTCACACAGATAGGATGTCTGCGCGACGAAAT CCTCGACATCAGCGAAACCTTCTCCCGCAAGCGTGG TCCGATCCGATCACGCGGCTCGACCGATCCAATGCTTT CGATCCCGAGACGAAGTTGAC	AGTT01257760.1 (737 to 938)	18,674	<i>Panthalops hodgsoni</i>	Beijing Genomics Institute	Illumina Genome Analyser
	APJD01000040.1 (17553 to 17753)	329,545	<i>Bradyrhizobium sp. OHSU_III</i>	Unknown	Illumina MiSeq
ATCGGGAAGCGTCCGGGCTGGTAGCCGGGATTGACAT AGGTGACGTCCGGCATGCCGTGCGCGCCATGTCGTA ATGATCGAAGGCCTTCCGAGCTGCTGGGCCGAAACA CCTTGCCGGTATGGTGCCGCGGAATCCTTGTTACC GCAGCCACCCAGTCTTCCAGCGACTTCTGCAGCGGATGC GAGGCGGGCACCC	AUYS01000696.1 (1109 to 1310)	Unknown	<i>Melampsora pinitorqua</i>	Canada's Michael Smith Genome Sciences Centre	Illumina HiSeq 2000
	NC_017082.1 (1001905 to 1002106)	Complete	<i>Bradyrhizobium sp. 523321</i>	Unknown	454 GS FLX Titanium
TTGATCCGGGATTGTCCGCGGGCGCAGCGGGCCCA GGCACTCCGCGGCCAGGATGAAGCGGGTGAGGCGCG CAGCGGATCTGCCGGCCGCCGAAATCGAGCGTTTGCA GATTGCCCTCGCGGGCAGCGCTGCACCGAGAACTCCA GGATCGCGGCCACCATCGCGCTGAGCAGATCGGCGCGC TCGTCGGGGTGTC	AHJH01004981.1 (282 to 483)	84,429	<i>Hammondia hammondi</i>	J. Craig Venter Institute	454 GS FLX Titanium and Illumina
	AMFB01000038.1 (11173 to 11974)	141,525	<i>Bradyrhizobium sp. DFCl-1</i>	Unknown	Illumina HiSeq 2000
TCGGTTACGAATCCGATCGGCTGGTTGTCGCTCAAAC GGCCGTGGTGTCCCGTCTTCAGTGGTCTCGGCGTTCG ACTGGCGGGTTCGTCATTCTCTGTCGGTGGTGTCTT TTCGTGCTGCATAAGACGGCGCTGACGCTT	AADB02296177.1 (1 to 150)	Unknown	<i>Homo sapiens</i>	Celera	PE Biosystems ABI Prism
	AMFB01000035.1 (33174 to 33323)	141,525	<i>Bradyrhizobium sp. DFCl-1</i>	Unknown	Illumina HiSeq 2000
CGAGCCGCTCGGCGGCAGCGGGATATCGTCCTTGGAGA GTGCCATGCGTGCTACACAAAATGGATGCCGGGAAATGA TTAACATGTTAAGTATTCCCGCAAATGAATCAGCGTAT GTTGCGGTGCAAAGGTTGCGGCGGCGGATTAGGTGATGG CGCGAAGAAATTGTGCAACGAGACAGGGCAGGAGCAGGG TGACGACACATCGTCGCGACGTGGCCCGATGCGCGAATTC TCACGTTGCTGCCGATGTCGCTGCTCCGTGCGCGCAGG CGGTGATGCGGCAGTTTCGTCCTCGTTGCGCAATCACGG GCTGACCGAACAGCAAT	ADNL01003140.1 (1 to 337)	685	<i>Astrammia rara</i>	Unknown	454 GS FLX
	AMFB01000042.1 (23424 to 23760)	141,525	<i>Bradyrhizobium sp. DFCl-1</i>	Unknown	Illumina HiSeq 2000
AGACGCACTCGTTCGCGGAAGGATCGACCCAGCTCAAGAA CGGCCAGAGCTTCATCCTGGATTCCGACAAGACGCCGGGC GACAACAGCCGCTCCAGCTTCCGCATCCGGAAATCCTCG CCGCGCTCCGCCCGGCCACGCGCTGCTGCTGACGACGG CAAGGTGCGGCTGATCGCCGAGGAGACCTCGCCGAGCGC GCCGTGACGCGCTGCTGATCGGCGGCAAGATGTCGGACC GCAAAGGCGTCAACCTGCCGACACCGATCTTCCCGTGTG CGCGATGACGCCAAGGACCGTCCGACTCGAAGCTGCG CTGCCGAGGGAATCGACTGGGTGGCGCTGCTCCTCGTACA GCGGGCCGAGGACGTGATCGAGGCCAAGAAGATGATCCG CGGCCGCGTGGCTGATGGCCAAGATCGAGAAGCCTCAG GCGATGACCGGCTCGCCGACATCATCGATGCGGCGGACG CGC	AKIRO1004699.1 (1 to 482)	35,272	<i>Toxoplasma gondii</i>	J. Craig Venter Institute	454 GS FLX Titanium and Illumina
	AMFB01000031.1 (95879 to 96354)	141,525	<i>Bradyrhizobium sp. DFCl-1</i>	Unknown	Illumina HiSeq 2000

Table 1. Cont.

AAGCGGCGCTGCATGTATTGCGCCGAGGAATTTGCCGAGTG CGTCGGCCGCAAGTTCCGGAAGGGTCATCACGGCATTGAC CTCCGAGATCGGGTGGCGAGCTTACGGCCTGCCAGCATTT CAGGGTAGTCCCAGGATGTGACATAATCGTTGCGGCCGGA CCGAAGCGGCTGGACCGTAGTGACCCGTAATAATGCGGTG TGGGGTGGGAGCTGGTGGATGGGTGTCGATCTTTGAACG TCAAAGGCTTGAGCGAGCTGGATCAAACGGCCCCGGTCTG GATGGTCAATCTGATGCGATTTCGCCAGCGGTGCGCTCGAC GGCGACGGTTCGGGTTGGGATGCCTATCTGCGCTACAGCG CGCTCACTGTCCCATGATCAAGGCCGACGCGGGGCTAC GGGCTGCTCTGGACCGCAACGCCGAGACGGTCCGCGCTCG GCGAGCCGGACGGCCAGCGTTGGG	ADAS01038761.1 (1 to 464)	9,773	<i>Puccinia triticina</i>	Broad Institute	454; ABI
	AMFB01000048.1 (5318 to 5776)	141,525	<i>Bradyrhizobium sp.</i> <i>DFCI-1</i>	Unknown	Illumina HiSeq 2000

Seven sequences which match specifically with both the *Bradyrhizobium* genus and a Genbank entry of a eukaryote. Two hundred thousand randomly selected *Bradyrhizobium* sequences were aligned against eukaryotes in the NCBI BLAST databases: this search is therefore indicative rather than exhaustive. *Bradyrhizobium* contamination inserted prior to *de novo* assembly of the eukaryote genome appears to have caused this double match, as it is not likely that different parts of the *Bradyrhizobium* genome would be conserved in select eukaryotes.
doi:10.1371/journal.pone.0097876.t001

entries. Moreover, until an effective strategy is found to eliminate contaminants from sequencing runs, novel sequences detected in such runs should be scrutinized by qPCR analysis or otherwise to ensure they originate from clinical specimens rather than from the laboratory. Comparing the number of novel organism cells per human cell calculated by using the read counts versus by qPCR or in situ hybridization could highlight discrepancies. Contamination notwithstanding, novel microbe identification based on unbiased high-throughput sequencing is very promising in the study of idiopathic disease, especially as sequencing technology continues to improve.

Supporting Information

Table S1 The contents of non-aligning reads from 142 *Candida albicans* whole genome sequencing runs.
(DOC)

Table S2 Runtime and cost of Leif Microbiome Analyzer for 57 human runs.
(DOC)

Table S3 Runtime and cost of Leif Microbiome Analyzer for 142 *Candida albicans* runs.
(DOC)

Text S1 Batch file commands to analyze *Bradyrhizobium* contaminants in NCBI BLAST databases using the Leif Microbiome Analyzer.
(DOC)

Text S2 Batch file commands to analyze contaminants in 57 publicly available 1000 Genomes Project runs using the Leif Microbiome Analyzer.
(DOC)

Text S3 Batch file commands to analyze contaminants in 142 publicly available *Candida albicans* runs using the Leif Microbiome Analyzer.
(DOC)

References

- Lipkin WI (2010) Microbe hunting. *Microbiol Mol Biol Rev* 74: 363–377.
- Feng H, Shuda M, Chang Y, Moore PS (2008) Clonal integration of a polyomavirus in human Merkel cell carcinoma. *Science* 319: 1096–1100.

Text S4 Example of a specific alignment to *Bradyrhizobium sp. DFCI-1* from an Illumina HiSeq 2000 run at the Baylor College of Medicine.
(DOC)

Text S5 Example of a specific alignment to *Bradyrhizobium sp. DFCI-1* from an Illumina HiSeq 2000 run at the Sanger Center.
(DOC)

Text S6 Example of a very specific alignment to *Bradyrhizobium sp. DFCI-1* from an Illumina HiSeq 2000 run at the Broad Institute.
(DOC)

Text S7 Example of a less specific alignment to *Bradyrhizobium sp. DFCI-1* from an Illumina HiSeq 2000 run at the Max Planck Institute for Molecular Genetics.
(DOC)

Spreadsheet S1 Maximum taxonomic resolution of non-aligning reads from 57 human whole genome sequencing runs.
(XLS)

Spreadsheet S2 Maximum taxonomic resolution of non-aligning reads from 142 *Candida albicans* whole genome sequencing runs.
(XLS)

Acknowledgments

D.E.B. is supported by NIH Skin Cancer SPORE grant 5P50CA121974.

Author Contributions

Conceived and designed the experiments: ML DEB. Performed the experiments: ML. Analyzed the data: ML CH. Wrote the paper: ML CH DEB.

4. Kostic AD, Ojesina AI, Pedamallu CS, Jung J, Verhaak RG, et al. (2011) PathSeq: software to identify or discover microbes by deep sequencing of human tissue. *Nat Biotechnol* 29: 393–396.
5. Bhatt AS, Freeman SS, Herrera AF, Pedamallu CS, Gevers D, et al. (2013) Sequence-based discovery of *Bradyrhizobium enterica* in cord colitis syndrome. *N Engl J Med* 369: 517–528.
6. Duncan CG, Leary RJ, Lin JC, Cummins J, Di C, et al. (2009) Identification of microbial DNA in human cancer. *BMC Med Genom* 2: 22.
7. Stott-Miller M, Wright JL, Stanford JL (2013) MSMB gene variant alters the association between prostate cancer and number of sexual partners. *Prostate* 73: 1803–1809.
8. Sutcliffe S, De Marzo AM, Sfanos KS, Laurence M (2014) MSMB variation and prostate cancer risk: Clues towards a possible fungal etiology. *Prostate* 74: 569–578.
9. Naccache SN, Greninger AL, Lee D, Coffey LL, Phan T, et al. (2013) The Perils of Pathogen Discovery: Origin of a Novel Parvovirus-Like Hybrid Genome Traced to Nucleic Acid Extraction Spin Columns. *J Virol* 87: 11966–11977.
10. Percudani R (2013) A Microbial Metagenome (*Leucobacter* sp.) in Caenorhabditis Whole Genome sequences. *Bioinf Biol Insights* 7: 55.
11. Schmieder R, Edwards R (2011) Fast identification and removal of sequence contamination from genomic and metagenomic datasets. *PLoS One* 6: e17288.
12. Cibulskis K, McKenna A, Fennell T, Banks E, DePristo M, et al. (2011) ContEst: estimating cross-contamination of human samples in next-generation sequencing data. *Bioinformatics* 27: 2601–2602.
13. Jun G, Flickinger M, Hetrick Kurt N, Romm Jane M, Doheny Kimberly F, et al. (2012) Detecting and Estimating Contamination of Human DNA Samples in Sequencing and Array-Based Genotype Data. *Am J Hum Genet* 91: 839–848.
14. Abecasis GR, Altshuler D, Auton A, Brooks LD, Durbin RM, et al. (2010) A map of human genome variation from population-scale sequencing. *Nature* 467: 1061–1073.
15. Morgulis A, Gertz EM, Schäffer AA, Agarwala R (2006) WindowMasker: window-based masker for sequenced genomes. *Bioinformatics* 22: 134–141.
16. Kulakov LA, McAlister MB, Ogden KL, Larkin MJ, O'Hanlon JF (2002) Analysis of bacteria contaminating ultrapure water in industrial systems. *Appl Environ Microbiol* 68: 1548–1555.
17. Findley K, Oh J, Yang J, Conlan S, Deming C, et al. (2013) Topographic diversity of fungal and bacterial communities in human skin. *Nature* 498: 367–370.
18. Lo S-C, Hung G-C, Li B, Lei H, Li T, et al. (2013) Isolation of Novel *Afipia* septicemia and Identification of Previously Unknown Bacteria *Bradyrhizobium* sp. OHSU_III from Blood of Patients with Poorly Defined Illnesses. *PLoS One* 8: e76142.