

Research Article

Depression among Adolescents of Rural Nepal: A Community-Based Study

Prayas Gautam,¹ Maginsh Dahal ,² Harishchandra Ghimire,³ Sakuntala Chapagain,⁴ Kushalata Baral ,⁵ Rohit Acharya,⁶ Sudip Khanal ,⁵ and Anu Neupane⁷

¹School of Public Health, Chitwan Medical College, Chitwan, Nepal

²Department of Public Health, Asian College for Advance Studies, Purbanchal University, Satdobato, Lalitpur, Nepal

³Department of Community Medicine and Public Health, Chitwan Medical College, Chitwan, Nepal

⁴Department of Nursing, Chitwan Medical College, Chitwan, Nepal

⁵Department of Public Health, Nobel College, Pokhara University, Sinamangal, Kathmandu, Nepal

⁶Population Services International/Nepal, Western Regional Hub, Nepalgunj, Nepal

⁷Department of Public Health, Little Buddha College of Health Science, Minbhavan, Kathmandu, Nepal

Correspondence should be addressed to Maginsh Dahal; maginsh@gmail.com

Received 23 June 2020; Revised 25 January 2021; Accepted 29 January 2021; Published 5 February 2021

Academic Editor: Janusz K. Rybakowski

Copyright © 2021 Prayas Gautam et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This community-based study is aimed at finding out the prevalence of depression, and its correlates among adolescents of rural Nepal. Data were collected among adolescents after multistage stratified proportionate random sampling by using the Patient Health Questionnaire (PHQ-9) to assess the depression among adolescent. Chi-square test and logistic regression model were applied. The prevalence of depression was found to be 27%. Adolescents who were not satisfied with their academic performance were 2.4 times more likely to have the risk of depression (AOR = 2.417, CI: 1.097-5.326). Likewise, tobacco users were almost fourteen times (AOR = 13.858, CI: 2.090-91.90), who intended to harm themselves were two times (AOR = 2.323, CI: 1.078-5.005), sleep deprivation were fifteen folds (AOR = 14.911, CI: 7.376-30.145), often scolded by their parents was almost three times (AOR = 2.931, CI: 1.195-7.436), and having poor relationship with friends were 2.4 times more likely (AOR = 2.371, CI: 1.078-5.215) of having depression. Sleep deprivation has a long-term negative impact on health leading depression.

1. Introduction

Adolescence aged 10 to 19 years is a critical period for the achievement of social emotional capabilities which would be the foundation for future life, health, and wellbeing. Helpful and supportive environments in the family, schools, and communities empower adolescents to maintain good mental health and wellbeing, but adolescents are also in the period of vulnerability for the onset of mental health conditions [1]. Depression is a common mental disorder and one of the main causes of disability globally. Depression is the ninth leading cause of illness and disability in all adolescents (WHO). Depression is a foremost contributor to the global burden of disease and often starts at a young age (WHO).

Globally, an estimated 300 million people are affected by depression [2]. Evidences show that depression is considered a major risk factor for increasing premature death among adolescent [3]. Adolescent depressive symptoms are emerging as a public health problem, mostly in developed countries as well as in developing countries [2]. Depression often starts in young age and has negative consequences in adolescent's health [4]. The highest prevalence of adolescent suicide is evidenced in Southeast Asia and Eastern Europe [3]. It can be persistent, comprehensively impairing adolescent's ability to function at work or school and to cope with daily life [5, 6]. Literatures have revealed that sleeping disorder may cause psychiatric disorders and psychological stress leading to acute and chronic depression [7]. Sleep deprivation in

adolescence period is a risk for depression and suicide and increased substance abuse [8].

Mental health problems are estimated to affect 10-20% of children and adolescent worldwide, accounting 15-30% of disability-adjusted life years (DALYs) lost during first three decades of life [9]. Poor mental health can have negative effects on health and development of adolescents. It is associated with several health and social outcomes such as higher alcohol, tobacco and drug use, adolescent pregnancy, school dropout, and involvement in criminal behaviour (WHO).

Mental health problems in young people are not only a major public health challenge but are also a development issue in low- and middle-income countries. Recognizing and addressing depression of young people help them function better socially, academically, and vocationally and develop into well-adjusted and productive adults is prime [9]. Various researches explained that sleep deprivation among adolescents is one of the major contributor for depression [10, 11].

Evidence on the burden of depression among adolescents in Nepal is not well measured due to the lack of adequate research [5, 6]. In Nepal, mental health services receive insignificant attention at all levels of society, from the government to the general public. This is despite studies showing that mental health problems account for a significant proportion of the global burden of disease [12].

This study will help us to shed light on the present situation of adolescent depression. Thus, we conducted the study to find out the prevalence of depression and its associated factors among adolescents of Nepal.

2. Methods

A descriptive cross-sectional study was conducted among the adolescent students of age 15-19 years currently studying in grade 11 and 12 of rural Nepal.

2.1. Sample Size and Sampling Technique. The sample size was determined by using the prevalence of 21.2% (prevalence was obtained from a study conducted in Nepal among undergraduate students) [13].

The sample size was determined by using a formula.

Sample size for finite population (n) = $n_0 / (1 + (n_0 - 1) / N)$ [14].

Then, the sample size for infinite population (n_0) = $z^2 pq / d^2$ [14].

Where n is the minimum required sample size. z is the critical value on standard normal distribution at 95% confidence interval (1.96). p is the prevalence of self-medication (34.4%). d is the margin of error desired around p to be estimated (5%/0.05). $n = 1.962 * 0.344 * (1 - 0.656) / 0.05^2$. $n = 347$.

Additional 7% sample was added for nonresponse. Thus the final sample size was 371.

Multistage stratified proportionate random sampling was done among 7 public and 9 private schools. The total number of students in public was 3197 and private was 3342. Schools were selected in equal number, i.e., three schools from private and three schools from the public as the total number of students in both public and private

was almost equal. The numbers of schools were selected by using probability proportionate to size (PPS). Then, in the second stage, the sample size from each school was determined proportionately, and then in the final stage, further stratification was done in class 11 and 12 and from each faculty (science, management, and education). The sample frame was prepared from the school attendance. The required sample of 371 was selected by random number table through excel. Adolescent students aged between 15-19 years who were present during the day of data collection were included in this study.

2.2. Data Collection Tools. Semistructured, self-administrated questionnaire was used for data collection. The questionnaire included sociodemographic information, behavioural, and psychological characteristics of students and Patient Health Questionnaire (PHQ-9) tool for assessing the depressive symptoms. The Patient Health Questionnaire (PHQ-9) is a self-administered version for the assessment of depression. PHQ-9 is a nine-item scale, in which score for each item of PHQ-9 ranges from 0 (not at all) to 3 (almost every day) based on the frequency of symptoms experienced. PHQ-9 score ranges from 0 to 27, in which a greater score reflects more depressive symptoms. A PHQ score ≥ 10 has a sensitivity of 88% and specificity of 88% and was used to measure the depressive symptoms [15].

2.3. Data Collection Procedure and Technique. The questionnaire was in Nepali language and piloted in 50 individuals before use in the survey. Translated version of PHQ-9 used in this study has undergone internal validity test using Chronbach's alpha, and the value was 0.75. Participants were briefed about the study objectives, and written parental consent was taken for the respondent age below 18 years. Data were collected in a separate classroom. Participants were briefed about the techniques of filling the questionnaire. Seating arrangements of the students were made properly in such a way that chances of peeking each other's answers were as low as possible.

2.4. Statistical Analysis. Data collected went through coding, editing, entry, and rechecking and were done by researcher's each day. Data entry and analysis were done using Epi-Data 3.1 and IBM SPSS version 20. Descriptive analysis was used to describe background characteristics and the prevalence of depression among adolescents. Data were reported in the form of proportion, means, and standard deviation. Chi-square test was used to test the difference between the categorical variables. Variables that were found statistically significant ($p < 0.05$) during bivariate analysis were checked for multicollinearity and then further analyzed using logistic regression model in multivariate analysis. Adjusted odds ratio with 95 percent CI and p value was calculated. Hosmer and Lemeshow test was used to test the goodness-of-fit for the regression model. Predictability of regression and coefficient of determination (Nagelkerke R square) for the equation were calculated. Multicollinearity test has been done to find out the possibility of multicollinearity among independent variables. To identify whether there exists serious

TABLE 1: Demographic and economic characteristics of respondents (*n* = 371).

| Characteristics | <i>n</i> (%) |
|--------------------------------------|-------------------|
| Age mean age ± SD (years) | 17.39 (SD ± 0.92) |
| Gender | |
| Male | 189 (50.9) |
| Ethnicity | |
| Brahmin/Chhetri | 219 (59) |
| Religion | |
| Hindu | 331 (89.2) |
| Place of residence of the respondent | |
| Urban | 292 (79.3) |
| Type of family (<i>n</i> = 370) | |
| Nuclear | 252 (68.1) |
| Family size | |
| ≤5 | 235 (63.5) |
| Parent living status | |
| Living together | 342 (92.2) |
| Monthly income of family (in NRs) | |
| ≤20,000 | 141 (38) |
| >20,000—≤25,000 | 76 (20.5) |
| >25,000—≤30,000 | 81 (21.8) |
| >30,000 | 73 (19.7) |

collinearity problem, variance inflation factor (VIF) has been estimated which is reciprocal to tolerance.

3. Results

Table 1 portrays the demographical and economical characteristics of 371 respondents. Mean age (in years) of the respondents was 17.39 (SD ± 0.92). Almost three-fifths of the participants (59%) were from advantaged caste group (Brahmin/Chhetri). 89.2 percent of the students were following Hindu religion, and almost four-fifth (79.3%) were receding in urban area. Majority (88.1%) of the students live in nuclear family, and more than three-fifth (63.5%) have a family size of up to five. Majority (92.2%) of the parents were living together. More than one third (38%) of the family monthly income was up to twenty thousand.

Table 2 presents characteristics of respondent’s parents. One third (33.91%) of respondent’s father have studied more than basic level but only less than one fifth (18.41%) of respondent’s mother have studied more than eighth grade. Most of the student’s parents have agriculture (41.0%) as major occupation. Almost two-fifth (39.37%) of the respondent’s fathers had alcohol but majority (95.6%) of respondent’s mother never had alcohol.

Table 3 represents the educational characteristics of the students. 53.6 percent of the respondents were from grade 11, almost equal students were enrolled in data collection in public and private institution. More than one third (39.6%) were studying in management faculty. Four fifth (80.6%) of the students currently studying their respective faculty were

TABLE 2: Distribution of characteristics of family of respondents’ (*n* = 371).

| Characteristics | <i>n</i> (%) |
|---|--------------|
| Educational status of father (<i>n</i> = 348) | |
| ≤8 | 230 (66.1) |
| >8 | 118 (33.9) |
| Educational status of mother (<i>n</i> = 364) | |
| ≤8 | 297 (81.6) |
| >8 | 67 (18.4) |
| Occupational status of father (<i>n</i> = 346) | |
| Government job | 62 (17.9) |
| Private job | 47 (13.5) |
| Business | 61 (17.6) |
| Agriculture | 142 (41.0) |
| Others* | 34 (9.8) |
| Occupational status of mother (<i>n</i> = 362) | |
| Government job | 16 (4.4) |
| Private job | 34 (9.3) |
| Business | 45 (12.4) |
| Agriculture | 209 (57.7) |
| Others** | 58 (16.0) |
| Drinking status of father (<i>n</i> = 348) | |
| No | 211 (60.6) |
| Drinking status of mother (<i>n</i> = 364) | |
| No | 348 (95.6) |

*Includes foreign employment and labour work. **Includes housewife, labour work, and foreign job.

because of their own preference and more than three quarters (78.1%) were passed in their previous exams. About two-third (63.6%) of the students was not satisfied with their academic performance.

Table 4 represents the behavioural characteristics of respondents. Among 371 students, majority (94.1%) of the students were nontobacco users, and 94.3% of students have never had alcohol. More than three-fifths of the students occasionally do physical exercise. 56.3% of the respondents participated in extracurricular activities conducted by their schools. Almost half of the respondents used the internet up to two hours daily, and more than one-fifth of the respondents used the internet for more than four hours of internet. More than half (54.5%) of the respondents used a minimum of 1 hour of internet before going to bed.

Table 5 portrays the psychological factors of respondents. More than three quarters (78.7%) report they have had conflict in their families. Great majority (92.5%) of the parents never scold their children. Almost half (49.3%) of the students had a good relationship with friends, and about one third (30.5%) students had a good relationship with teachers as well. More than a quarter (28.9%) students have either girlfriend or boyfriend. More than one third (38.8%) of the respondents felt lonely. Less than one-tenth (7.6%) of the students have left their home without informing their parents regardless of the frequency of running away from home.

TABLE 3: Educational characteristics of the respondents ($n = 371$).

| Characteristics | n (%) |
|---|------------|
| Grade | |
| 11 | 209 (56.3) |
| 12 | 162 (43.7) |
| Type of school | |
| Public | 183 (49.3) |
| Private | 188 (50.7) |
| Faculty of the respondent | |
| Science | 112 (30.2) |
| Management | 147 (39.6) |
| Education | 112 (30.2) |
| Reason for selecting the currently studying faculty | |
| Own decision | 299 (80.6) |
| Others* | 72 (19.4) |
| Achievement in last exam ($n = 370$) | |
| Passed | 290 (78.1) |
| Fail | 81 (21.9) |
| Satisfied with academic performance | |
| Yes | 236 (63.6) |
| No | 135 (36.4) |

*Includes failing to qualify in other faculty, family pressure, and friends pressure.

More than one fifth (21.6%) of the students had tried to hurt themselves irrespective to the frequency of hurting themselves. Students frequently share their feeling more with their close friends than with family.

Figure 1 depicts the prevalence of depression among adolescents. Respondents had a mean PHQ-9 score of 7.35 ($SD \pm 4.04$). More than one quarter (27%) of the respondents were depressed.

Table 6 shows the multicollinearity by the VIF factor. None of them have tolerance < 0.1 and $VIF > 10$. There was no problem of collinearity among independent variables as the highest VIF was 1.765.

The results from multivariable linear regression models of the relationship of students' characteristics with sleep deprivation, internet addiction, and depressive symptoms are tabled in Table 7. Those respondents who were not satisfied with their academic performance were 2.4 times more likely to have risk of depression than those students who were satisfied with their academic performance (AOR = 2.417, CI: 1.097-5.326). Tobacco users were almost fourteen times more likely to be depressed than their counterparts (AOR = 13.858, CI: 2.090-91.90). Those students who intended to harm themselves were two times more likely to develop depression (AOR = 2.323, CI: 1.078-5.005). Those students who have sleep deprivation were fifteen folds in risk developing depression than students without sleep deprivation (AOR = 14.911, CI: 7.376-30.145). Those students who were often scolded by their parents were almost three times more likely to be depressed than those students whose parent never scolds (AOR = 2.931, CI: 1.195-7.436). Similarly, those students who have poor relationship with friends were 2.4 times more

TABLE 4: Behavioural characteristics of respondents ($n = 371$).

| Characteristics | n (%) |
|---|------------|
| Tobacco users | |
| No | 349 (94.1) |
| Alcohol users | |
| No | 350 (94.3) |
| Do physical exercise | |
| Regularly | 46 (12.4) |
| Frequently | 37 (10) |
| Occasionally | 237 (63.9) |
| Rarely | 31 (8.4) |
| Never | 20 (5.4) |
| Participate in extracurricular activities | |
| Yes | 209 (56.3) |
| Daily time spent on internet (hours) | |
| Never | 46 (12.4) |
| $>0-\leq 2$ | 151 (46.5) |
| $>2-\leq 4$ | 132 (40.6) |
| >4 | 42 (12.9) |
| Daily time spent on internet (hours) before going bed ($n = 325$) | |
| Never use | 22 (6.8) |
| $>0-\leq 1$ | 177 (54.5) |
| >1 | 126 (38.7) |

likely to be in risk of depression than those students who have a good relationship with their friends (AOR = 2.371, CI: 1.078-5.215).

4. Discussion

We found that mean family size was 5.51 (± 2.12 SD) which was higher than National 4.6 [16]. In this study, the prevalence of depression was found to be 27%. Prevalence of depression among adolescent students as reported from existing literature varies from (6.4% to 52.9%) [13, 17-22]. A study with similar setting in Bangladesh among secondary school children found the prevalence was 25%, which was similar to our research findings [20]. Prevalence was found a little bit lower in a study conducted by Bhandari et al. [13]. However, Guo et al. found the prevalence very low compared to our findings, i.e., 6.4% [18]. The difference may be due to difference in sample size in this study was very low compared to that study, and the tools used in to measure depression different research were different. Study conducted by Sandal et al. in India reported prevalence was found comparatively higher than ours which was 36.1, and study conducted in Iran also reported a higher prevalence than ours [17, 22]. Malik et al. showed the highest prevalence of depression among adolescent students of Hariyana, India, which was 52.9%, and another study conducted in Bihar, India have also similar prevalence, 49.2% [19, 21].

In this study, very few sociodemographic variables have a significant association with depression. Age, sex, ethnicity, family type, family size, parent marital status, monthly

TABLE 5: Description of psychological factors of respondents ($n = 371$).

| Characteristics | Percentage |
|-------------------------------------|------------|
| Conflict in family | |
| Yes | 293 (78.7) |
| Family members scold you | |
| No | 343 (92.5) |
| Relationship with friends | |
| Good | 183 (49.3) |
| Average | 128 (34.5) |
| Poor | 60 (16.2) |
| Relationship with teacher | |
| Good | 113 (30.5) |
| Average | 169 (45.6) |
| Poor | 89 (24.0) |
| Do you have girlfriend or boyfriend | |
| No | 265 (71.4) |
| Felt lonely | |
| No | 227 (61.2) |
| Run away from home | |
| No | 343 (92.5) |
| Hurt yourself | |
| No | 291 (78.4) |
| Sharing with parents ($n = 370$) | |
| Frequently | 78 (21.1) |
| Occasionally | 258 (69.7) |
| Rarely | 23 (6.2) |
| Never | 11 (3.0) |
| Sharing with friends | |
| Frequently | 150 (40.4) |
| Occasionally | 181 (48.8) |
| Rarely | 30 (8.1) |
| Never | 10 (2.7) |

income occupation, occupation of father, and occupation of mother were found insignificant with depression among students. However, sex was found to be significantly associated with depression with female more at risk of developing depression in studies conducted in Brazil and Malaysia [23, 24]. Similar to our findings, age was found insignificant in the study conducted by Singh et al. in Chandigarh North India [25]. Educational status of parents and drinking status of parents were found to be associated with depression in this study. Those students whose parents were educated more than basic levels were more likely to develop depression than those students whose parents were less educated. The finding could be attributed to the fact that educated parents have higher expectation from children. Apart from that, they often try to indulge their children more toward academic activities and often much of their time of enjoyment or recreational activities get reduced. So when compared with educated parents who are often involved in a job or other activities, the less educated parents are more available to their children to




FIGURE 1: Prevalence of depression among respondents ($n = 371$).

TABLE 6: Test of multicollinearity of different independent variables related to depression.

| Model (constant) | Tolerance | VIF |
|---|-----------|-------|
| Grade of the student | 0.951 | 1.052 |
| Reason for selecting currently studying faculty | 0.823 | 1.215 |
| Achievement in last exam | 0.607 | 1.649 |
| Satisfied with academic performance | 0.567 | 1.765 |
| Tobacco use | 0.620 | 1.613 |
| Family members scold you often | 0.836 | 1.196 |
| Relationship with teacher | 0.697 | 1.435 |
| Relationship with friends | 0.725 | 1.379 |
| Run away from home | 0.727 | 1.376 |
| Hurt yourself | 0.796 | 1.256 |
| Time spend on internet | 0.606 | 1.650 |
| Spend time on internet before going to bed | 0.605 | 1.654 |
| Sleep deprivation | 0.869 | 1.151 |
| Drinking status of father | 0.845 | 1.183 |
| Drinking status of mother | 0.923 | 1.084 |
| Education status of father | 0.655 | 1.527 |
| Education status of mother | 0.644 | 1.553 |

share their feelings and thoughts, thereby reducing the risk of depression. However, a study conducted in Iran does not show any significant association between parents' level of education and depression among children [17]. In contrary with this findings, a study conducted in Malaysia, those students whose parents are less educated were more likely to be depressed [26].

In this study grade, reason for selecting the faculty, achievement in previous exam, and satisfaction with academic performance was found to be statistically significant with depression, whereas the type of school and faculty of students were not found significant with depression of students. This study revealed that those students studying in higher grades, i.e., 12 standard were 1.7 times more likely to have risk of depression than their counterparts (OR = 1.676, CI: 1.056-2.659), and a study conducted by Singh et al. in India shows similar findings, where highest grade students

TABLE 7: Multivariable linear regression of factors associated with depression.

| Characteristics | UOR (95% CI) | AOR (95% CI) | p value |
|---|-----------------------|----------------------|---------|
| Grade (11) | 1.676 (1.056-2.659) | 1.675 (0.881-3.187) | 0.116 |
| Reason for selecting current faculty (own decision) | 3.159 (1.8466-5.406) | 1.712 (0.778-3.769) | 0.182 |
| Achievements in last exam (pass) | 2.675 (1.590-4.499) | 1.121 (0.460-2.735) | 0.801 |
| Satisfied with academic performance (yes) | 3.920 (2.424-6.338) | 2.417 (1.097-5.326) | 0.029* |
| Tobacco use (yes) | 6.655 (2.626-16.866) | 13.858 (2.090-91.90) | 0.006* |
| Run away from home (yes) | 4.841 (2.180-10.749) | 1.075 (0.337-3.424) | 0.903 |
| Tried to hurt yourself (yes) | 2.902 (1.724-4.884) | 2.323 (1.078-5.005) | 0.031* |
| Time spend on internet (never) | | | |
| >0-≤2 | 1.381 (0.563-2.673) | 1.480 (0.256-8.558) | 0.662 |
| >2-≤4 | 2.252 (0.926-5.476) | 0.658 (0.092-4.698) | 0.677 |
| >4 | 8.193 (2.974-22.571) | 1.167 (0.142-9.594) | 0.886 |
| Time spend on internet before going to bed (never) | | | |
| ≤1 | 1.232 (0.576-2.673) | 1.322 (0.283-6.181) | 0.723 |
| >1 | 5.167 (2.396-10.887) | 4.679 (0.851-25.728) | 0.761 |
| Sleep deprivation (yes) | 15.041 (8.397-26.939) | 14.911 (7.376-30.14) | 0.0001* |
| Education status of father (≤8) | 1.671 (1.026-2.723) | 1.775 (0.935-3.366) | 0.079 |
| Education status of mother (≤8) | 1.860 (1.061-3.262) | 1.313 (0.621-2.776) | 0.475 |
| Drinking status of father (no) | 1.935 (1.198-3.125) | 1.643 (0.963-2.773) | 0.069 |
| Drinking status of mother (no) | 3.799 (1.374) | 3.056 (0.986-9.527) | 0.054 |
| Family members scold you (no) | 5.729 (2.544-12.903) | 2.931 (1.195-7.436) | 0.024* |
| Relationship with friend (good) | | | |
| Poor | 3.122 (1.677-5.812) | 2.371 (1.078-5.215) | 0.032* |
| Average | 1.436 (0.847-2.345) | 1.416 (0.762-2.636) | 0.271 |
| Relationship with teachers (good) | | | |
| Poor | 3.264 (1.733-6.149) | 1.445 (0.649-3.217) | 0.367 |
| Average | 1.403 (0.778-2.532) | 2.047 (0.519-2.114) | 0.897 |

Characteristics mentioned in brackets resembles reference category. UOR: unadjusted odds ratio; AOR: adjusted odds ratio; *Include statistically significant; Cox & Snell $R^2 = 0.365$, Nagelkerke $R^2 = 0.531$, and Hosmer and Lemeshow = 0.154.

were 1.6 times more likely to be depressed than those students from lowest grade (OR = 1.659, CI: 0.228-12.099) [25]. In this study, those students who do not have their own decision for selecting currently studying faculty were more likely to have depressive symptoms than those students who selected their subject with their own decision. The findings could be attributed to the fact that those students who selected their career as a peer, their interest could feel less pressure in studies, could do well in their studies, and could be more satisfied with their academic life, thereby reducing stress and risk of depression. In a similar study in India, those students whose parents choose their career without their children decision were more likely to be depressed than their counterparts [25]. Similarly, those students who failed in their previous exams were more likely to be at risk of depression than those students who have passed their previous exam. Study conducted by Bhandari et al. also shows the similar findings [13]. Satisfaction with academic performance was also associated with depression among adolescent students. Those students who were not satisfied with academic performance were more than two times more likely to be depressed than those students who were satisfied with their

academic performance (OR = 2.417, CI: 1.097-5.326). Similar findings were reported in a study conducted in India (OR = 5.089, CI: 11.180-21.957) [25].

Tobacco use and alcohol use were found to be statistically significant with depression among adolescent students, whereas physical activity and participation in extracurricular activities were not found significant with depression. But a study conducted in Korea revealed that those adolescents with no physical activity were found to be a risk factor for depression [27]. In these findings, tobacco use increases the risk of developing depression by almost 7 folds (OR = 6.655, CI: 2.626-16.866), and a study conducted among adolescent of Brazil also showed similar significant with depression (OR = 2.23, CI: 1.12-4.43) [24]. Similarly, alcohol-user students were 2.6 times more likely to be depressed than their counterparts (OR = 2.626, 1.079-0.639). Likewise, significant association was found in a study conducted in Brazil, where alcohol users were on more risk of developing depression (OR = 2.22, CI: 1.21-4.07).

Conflict in family, having a girlfriend or boyfriend, and feeling lonely were found to have no relationship with developing risk of depression among adolescents. The findings

could attribute to the fact that having a girlfriend or boyfriend could help in sharing thoughts and feelings with each other, thereby reducing the risk of depression. Scolding by family members often was significantly associated with depression among adolescent students in this study. The alike findings were reported from a study conducted in India [25]. Relationship with a teacher and relationship with a friend were also a risk factor for developing depression among adolescents according to our findings. However, a study conducted among Chinese adolescents found no significant association between relationship with teacher and risk factor for depression [18]. Having poor relationship with classmates was a risk factor for depression (AOR = 2.371, CI: 1.078-5.215), and alike findings were reported in a study conducted in China (AOR = 1.60, CI: 1.14-2.25) [18]. In this study, run away from home without informing their parents and tried to hurt them were also significantly associated with depression. Study conducted among Chinese students reported significant association between run away from home with depression but no significant association between tried to hurt themselves on purpose and risk of developing depression [18].

Having smartphone and use of internet were not found to be significant with risk of developing depression but more use of internet on a daily basis and use of the internet before falling asleep was found to be statistically associated with developing risk of depression among students. Those students who used the internet for more than an hour before falling asleep were two times more likely to develop depression than those who never used the internet before sleeping. Students who used the internet for more than four hours daily were most likely to be depressed than those who never use the internet (OR = 3.412, CI: 1.419-8.205). Likewise, longer daily time use of internet was analyzed as a risk factor for depression in a study conducted by Lim et al. in Korea. Those students using the internet more than three hours a day were two times more likely to have increased risk of depression (OR = 2.235, CI: 1.078-4.637) [27]. Similarly, another study conducted in India shows the use of social sites was associated with developing risk of developing depression [25].

Sleep deprivation was statistically significant with depression among students in this study. Evidences from other studies clearly show sleep deprivation and depression are strongly interrelated. Those students who have sleep deprivation have a strong effect on the risk of developing depression [10, 24, 28–30]. The odds of increasing risk of developing depression almost increase by five folds in a student with sleep deprivation according to study conducted by Roberts et al. in Texas [10]. In this study, those students lacking proper sleep or having sleep deprivation were fifteen times more likely to have depression than those students with normal sleep (OR = 15.041, CI: 8.397-26.939). Similar findings were reported in Brazil where the odds of developing risk of depression increase by ten times among poor sleepers (OR = 10.440, CI: 1.400-46.070) [24]. The findings of our study provide the baseline information for the policymakers so that they could not deny the existence of problem in Nepalese community. Our finding shows that there might be a possible association between the various factors and depres-

sion among the adolescents but it cannot establish the causal relationships. We suggest the broader spectrum of similar kind of study in large population on interventional setting for the clear picture of the real situation.

5. Conclusion

We can conclude that the problem of depression among adolescents is like an iceberg tip in Nepalese context. This study reflects the clear picture in a small arena and cannot be generalized but can show the policymakers and stakeholders that the problems exist in our context. Improved sleep quality will likely to benefit Nepalese school students in their mental health status, daily activities, and academic performance.

6. Limitations

Since the study is conducted among adolescent and in a small area, it cannot be generalized to the whole population all over Nepal. School dropout adolescent and those adolescents who were currently not studying were excluded from the sample.

Abbreviations

| | |
|--------|---|
| AOR: | Adjusted odds ratio |
| BMC: | Bio med central |
| CI: | Confidence interval |
| CMC: | Chitwan Medical College |
| DALY: | Disability-adjusted life years |
| DAS: | Depression and anxiety scale |
| DPHO: | District Public Health Office |
| IOM: | Institute of Medicine |
| OR: | Odds ratio |
| PHQ-9: | Patient Health Questionnaire-9 |
| PPS: | Probability proportionate to size |
| PSQI: | Pittsburgh Sleep Quality Index |
| PSS: | Perceived Stress Scale |
| SD: | Standard deviation |
| SPSS: | Statistical Package for Social Sciences |
| VIF: | Variance inflation factor |
| WHO: | World Health Organisation |

Data Availability

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Additional Points

Code Availability. The software application or custom codes used in the research are available from the corresponding author (MD) upon reasonable request.

Ethical Approval

Ethical approval was taken from the Institutional Review Board at Chitwan Medical College. Before starting the data collection, permission was taken from concerned authorities. Permission was obtained from the respective schools as well.

The objectives were clearly shared among the participants, and the participation was completely voluntary. Written parental written consent was taken from the parents whose children were below 18 years through the help of students. Then after with the student, written consent was obtained.

Consent

The informed consent and objectives were verbally shared with each respondent, and written consent was taken. Participation in this study was fully voluntary. Confidentiality of students was assured and maintained by the coding system. The authors declare that any person named as coauthor of the contribution is aware of the fact and has agreed to be named.

Conflicts of Interest

The authors declare they have no competing interests.

Authors' Contributions

PG and MD were the principle investigators. KB and MD did the development of the paper. RA, MN, and AN did the article review. MD, HG, and SC did the writing of the manuscript. SK did the data analysis. All authors revised the draft critically and have approved the final text. Prayas Gautam and Maginsh Dahal considered as combined first author. Kushalata Baral and Rohit Acharya contributed equally to this work.

References

- [1] WHO, *Adolescent Mental Health*, World Health Organisation, 2018.
- [2] World Health Organization, R O F S A, *Mental health status of adolescents in South-East Asia: evidence for action*, World Health Organization, 2017.
- [3] A. B. McLoughlin, M. S. Gould, and K. M. Malone, "Global trends in teenage suicide: 2003–2014," *QJM: An International Journal of Medicine*, vol. 108, no. 10, pp. 765–780, 2015.
- [4] X.-L. Jiang, X.-Y. Zheng, J. Yang et al., "A systematic review of studies on the prevalence of insomnia in university students," *Public Health*, vol. 129, no. 12, pp. 1579–1584, 2015.
- [5] K. Baral, M. Dahal, G. Khadka, and A. Adhikari, "Prevalence of depressive symptoms and its associated factors among male prisoners at Dillibazar prison, Kathmandu, Nepal," *European Journal of Medical Sciences*, vol. 2, no. 1, pp. 30–36, 2020.
- [6] K. Baral, M. Dahal, and S. Pradhan, "Knowledge regarding Alzheimer's disease among college students of Kathmandu, Nepal," *International Journal of Alzheimer's Disease*, vol. 2020, pp. 1–6, 2020.
- [7] H. M. Abdulghani, N. A. Alrowais, N. S. Bin-Saad, N. M. al-Subaie, A. M. Haji, and A. I. Alhaqwi, "Sleep disorder among medical students: relationship to their academic performance," *Medical Teacher*, vol. 34, no. sup1, pp. S37–S41, 2012.
- [8] A. Winsler, A. Deutsch, R. D. Vorona, P. A. Payne, and M. Szklo-Coxe, "Sleepless in Fairfax: the difference one more hour of sleep can make for teen hopelessness, suicidal ideation, and substance use," *Journal of Youth and Adolescence*, vol. 44, no. 2, pp. 362–378, 2015.
- [9] M. N. Cook, J. Peterson, and C. Sheldon, "Adolescent depression: an update and guide to clinical decision making," *Psychiatry (Edgmont)*, vol. 6, no. 9, p. 17, 2009.
- [10] R. E. Roberts and H. T. Duong, "The prospective association between sleep deprivation and depression among adolescents," *Sleep*, vol. 37, no. 2, pp. 239–244, 2014.
- [11] WHO, *WHO technical meeting on sleep and health*, vol. 2018, 2004.
- [12] P. Simkhada, E. Van Teijlingen, and S. Marahatta, "Mental health services in Nepal: is it too late?," *Journal of Manmohan Memorial Institute of Health Sciences*, vol. 1, no. 4, pp. 1–2, 2015.
- [13] P. M. Bhandari, D. Neupane, S. Rijal, K. Thapa, S. R. Mishra, and A. K. Poudyal, "Sleep quality, internet addiction and depressive symptoms among undergraduate students in Nepal," *BMC Psychiatry*, vol. 17, no. 1, pp. 106–106, 2017.
- [14] G. D. Israel, *Determining Sample Size*, Florida Cooperative Extension Service/Institute of Food and Agricultural Sciences (IFAS)/ University of Florida, 1992.
- [15] K. Kroenke, R. L. Spitzer, and J. B. Williams, "The phq-9," *Journal of General Internal Medicine*, vol. 16, no. 9, pp. 606–613, 2001.
- [16] Central Bureau of Statistics-National Planning Commission Secretariat, G o N, *Nepal-Annual Household Survey 2015-2016*, Government of Nepal, 2018.
- [17] A. Daryanavard, "Prevalence of depression among high school students and its relation to family structure," *American Journal of Applied Sciences*, vol. 8, no. 1, pp. 39–44, 2011.
- [18] L. Guo, J. Deng, Y. He et al., "Prevalence and correlates of sleep disturbance and depressive symptoms among Chinese adolescents: a cross-sectional survey study," *BMJ Open*, vol. 4, no. 7, article e005517, 2014.
- [19] K. K. Jha, S. K. Singh, S. K. Nirala, C. Kumar, P. Kumar, and N. Aggrawal, "Prevalence of depression among school-going adolescents in an urban area of Bihar, India," *Indian Journal of Psychological Medicine*, vol. 39, no. 3, pp. 287–292, 2017.
- [20] A. Khan, R. Ahmed, and N. W. Burton, "Prevalence and correlates of depressive symptoms in secondary school children in Dhaka city, Bangladesh," *Ethnicity & Health*, vol. 25, no. 1, pp. 34–46, 2020.
- [21] M. Malik, P. Khanna, R. Rohilla, B. Mehta, and A. Goyal, "Prevalence of depression among school going adolescents in an urban area of Haryana, India," *International Journal of Community Medicine And Public Health*, vol. 2, no. 4, pp. 624–626, 2015.
- [22] R. K. Sandal, N. K. Goel, M. K. Sharma, R. K. Bakshi, N. Singh, and D. Kumar, "Prevalence of depression, anxiety and stress among school going adolescent in Chandigarh," *Journal of family medicine and primary care*, vol. 6, no. 2, pp. 405–410, 2017.
- [23] M. Ramli, S. Adlina, A. Suthahar et al., "Depression among secondary school students: a comparison between urban and rural populations in a Malaysian community," *Hong Kong Journal of Psychiatry*, vol. 18, no. 2, p. 39, 2008.
- [24] L. Zinn-Souza, R. Nagai, L. Teixeira et al., "Factors associated with depression symptoms in high school students in São Paulo, Brazil," *Revista de Saúde Pública*, vol. 42, no. 1, pp. 34–40, 2008.
- [25] M. M. Singh, M. Gupta, and S. Grover, "Prevalence & factors associated with depression among schoolgoing adolescents in Chandigarh, North India," *The Indian Journal of Medical Research*, vol. 146, no. 2, pp. 205–215, 2017.

- [26] J. Kaur, S. M. Cheong, B. Mahadir Naidu et al., "Prevalence and correlates of depression among adolescents in Malaysia," *Asia Pacific Journal of Public Health*, vol. 26, 5_suppl, pp. 53S–62S, 2014.
- [27] C. H. Lim, E. J. Kim, J. H. Kim, J. S. Lee, Y. Lee, and S. H. Park, "The correlation of depression with internet use and body image in Korean adolescents," *Korean Journal of Pediatrics*, vol. 60, no. 1, pp. 17–23, 2017.
- [28] M. A. Al-Abri, "Sleep deprivation and depression: a bi-directional association," *Sultan Qaboos University Medical Journal*, vol. 15, no. 1, pp. e4–e6, 2015.
- [29] A. Postans and A. Pidgeon, "Sleep Quality and Mindfulness as Predictors of Depression, Anxiety and Stress," *International Journal of Psychology and Behavioral Sciences*, vol. 6, no. 2, pp. 71–75, 2016.
- [30] S. T. Wiebe, J. Cassoff, and R. Gruber, "Sleep patterns and the risk for unipolar depression: a review," *Nature and Science of Sleep*, vol. 4, p. 63, 2012.