

Reproducibility of fluid-phase measurements in PBS-treated sputum supernatant of healthy and stable COPD subjects

This article was published in the following Dove Medical Press journal:
International Journal of COPD

Fengyan Wang^{1,*}
Zhenyu Liang^{1,*}
Yuqiong Yang¹
Luqian Zhou¹
Lili Guan¹
Weiliang Wu¹
Mei Jiang¹
Weijuan Shi¹
Kuimiao Deng¹
Jianhua Chen²
Rongchang Chen¹

¹State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, Guangzhou Institute of Respiratory Health, First Affiliated Hospital of Guangzhou Medical University, Guangzhou 510120, People's Republic of China; ²Respiratory Medicine Department, Guangzhou Panyu Central Hospital, Guangzhou 511400, People's Republic of China

*These authors contributed equally to this work

Correspondence: Rongchang Chen
State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, Guangzhou Institute of Respiratory Health, First Affiliated Hospital of Guangzhou Medical University, No 151, Yanjiang Road, Guangzhou 510120, People's Republic of China
Tel/fax +86 20 8306 2882
Email chenrc@vip.163.com

Purpose: The purpose of this study was to investigate the reproducibility of fluid-phase measurements in PBS-treated sputum supernatant, processed using the two-step method, of healthy and stable COPD individuals.

Methods: Nine healthy subjects and 23 stable COPD patients provided sputum twice within 6 days. A two-step sputum processing method was used to obtain PBS-treated supernatant and sputum cells. Soluble protein markers and IgG and IgM autoantibody profiles in PBS supernatant were analyzed using customized microarrays. Repeatability of measurements was assessed by paired-sample testing and an intraclass correlation coefficient, then graphically reported by Bland–Altman plot.

Results: There was no significant difference between the repeated detection of 8/10 types of soluble protein markers, all 13 types of IgG autoantibodies, and 12/13 types of corresponding IgM autoantibodies in PBS supernatant. The repeatability of measurements in PBS supernatant was substantial to very good for interleukin 6 (IL6), IL8, IL13, IL10, IL33, vascular endothelial growth factor, soluble receptor for advanced glycation end-products, and tumor necrosis factor- α ; for IgG autoantibodies against aggrecan, centromere protein B (CENP-B), collagen II, collagen IV, cytochrome C, elastin, heat shock protein 47 (HSP47), HSP70, and La/Sjögren syndrome type B antigen; for IgM autoantibodies against CENP-B, collagen I, collagen II, collagen IV, cytokeratin 18, and HSP70; and for sputum neutrophils, macrophages and eosinophils count. Bland–Altman plots suggested good consistency within repeated measurements. Stable COPD patients differed from healthy subjects in the proportion of neutrophils and eosinophils; relative fluorescence intensity of anti-cytochrome C IgG, anti-aggrecan IgM, and anti-cytochrome C IgM. There was a significant positive correlation for stable COPD patients between sputum anti-collagen II IgG and post-bronchodilator FEV₁ %.

Conclusion: We confirmed fluid-phase measurements in PBS-treated sputum supernatant by high-throughput techniques with good repeatability. We demonstrated the presence of IgG and IgM autoantibodies to multiple antigens in the airways of COPD patients.

Keywords: chronic obstructive pulmonary disease, soluble protein markers, autoantibody, reproducibility, sputum

Introduction

Sputum examination is increasingly used as a noninvasive tool to assess airway disease, especially asthma and COPD. The measurement of fluid-phase markers in sputum can help to explore mechanisms of airway disease progression and patient response to drugs.¹ The detection of cytokine levels in induced sputum (eg, interleukin 6 [IL6], IL8, tumor necrosis factor- α [TNF α], and vascular endothelial growth factor [VEGF])

contributed to investigations of the effect of cigarette smoke on chronic airway inflammation, an important pathological feature of COPD.^{2,3} In 2003, a largely unexplored role was suggested for the immune system in the pathogenesis of COPD.⁴ Recent findings indicate that autoantibody profiles may be used to determine COPD risk and provide a diagnostic and prognostic tool.⁵ The detection of soluble protein markers and autoantibodies in sputum may provide a basis for exploring local immune subtypes of COPD. However, there have been no available studies of sputum autoantibody profiles in COPD patients.

As noted in a recent review, the production, preparation, and dispersal of sputum all can affect the levels of fluid-phase markers in sputum.¹⁶ In the traditional sputum processing method, dithiothreitol (DTT) is used to break the disulfide bonds in mucin molecules, thereby allowing the release of cells; this is more effective in dispersing cells than PBS.⁷ However, as a strong reducing agent, DTT may decrease the biological activity of cytokines, many of which rely on disulfide bonds to provide a stable structure for bioactivity.¹ DTT may also disrupt the disulfide bonds present in the capture antibodies of immunoassays, resulting in reduced assay sensitivity.^{1,8} Bafadhel et al proposed a “two-step” sputum processing method for supernatant and cytospin preparation that yielded increased recovery of mediators in PBS-treated sputum, compared with DTT-treated sputum.⁹ This method has been widely used in supernatant purification, as it involves initial PBS processing for PBS supernatant, followed by DTT to obtain cells.^{10–12} This avoids interference of DTT on soluble mediator analysis from sputum supernatant. However, the reproducibility of fluid-phase measurements in sputum treated with PBS and without DTT has not yet been investigated.

In this study, we investigated the reproducibility of fluid-phase measurements in PBS supernatant, by recruiting stable COPD and healthy subjects for comparing high-throughput measurements of soluble protein markers and autoantibody profiles of sputum obtained within a 6-day period.

Materials and methods

Study design

This was a cross-sectional study examining soluble protein markers and autoantibodies in the sputum supernatant of healthy subjects and stable COPD patients. The study was conducted in accordance with the Declaration of Helsinki and approved by The Ethics Committee of The First Affiliated Hospital of Guangzhou Medical University (No 201722).

The clinical trial was registered with www.ClinicalTrials.gov (NCT 03240315).

Subjects

The study population comprised nine healthy subjects and 23 stable COPD patients. COPD was identified on the basis of the GOLD guidelines. Medications for COPD patients were unchanged during the study. Exclusion criteria for COPD patients were a history of significant inflammatory disease, a history of other lung disease, recent exacerbation of COPD treated with antibiotics or steroid, active cancer under treatment, enrollment in a long-term blinded drug study. Healthy subjects were free from significant disease as determined by history, physical examination, and screening investigations. They had post-bronchodilator FEV₁/FVC ratio >70%. Signed and dated written informed consent was obtained from all subjects prior to participation. All consenting subjects provided sputum twice within 6 days. Baseline conditions were required to be constant (diet, time of collection, therapies, and absence of respiratory tract infection).

Sputum processing

Sputum was processed within 2 hours, in accordance with an adaptation of previous methodology.¹³ Selected and weighed sputum specimens were placed in an Eppendorf tube and incubated with an 8× volume (sputum weight) of PBS, placed on ice on a bench rocker for 15 minutes, rocked for 15 seconds every 5 minutes during ice bath. The mixture was centrifuged at 1,000 ×g for 10 minutes by a low-temperature centrifuge at 4°C. The suspension was filtered through a 0.22-μm bacterial filter to remove bacteria, then a 4× volume of the supernatant was aspirated and stored in frozen storage tubes at –80°C for subsequent assays; this was identified as the PBS supernatant. The remainder of the solute was treated with a 2× volume of 0.3% DTT for cytospin preparation. Adjusting for the specimen size, the PBS supernatant of all subjects was examined for soluble protein markers; the PBS supernatant of nine healthy subjects and 19 stabilized COPD patients was used for autoantibody detection.

Microarray analysis of sputum soluble protein markers

The levels of B-cell activation factor (BAFF), interferon-γ (INFγ), IL4, IL5, IL6, IL8, IL10, IL13, IL17, IL33, IL1β, TNFα, leptin, transforming growth factor-β (TGFβ), soluble receptor for advanced glycation end-products (sRAGE), and VEGF in sputum supernatant samples were determined in quadruplicate, using a customized microarray (Human

Cytokine Antibody Microarray slides; RayBiotech, Inc., Norcross, GA, USA).

Microarray analysis of sputum autoantibody profiles

The levels of IgG autoantibodies and IgM autoantibodies against elastin, aggrecan, amyloid, centromere protein B (CENP-B), collagen I, collagen II, collagen IV, cytochrome C, cytokeratin 18, heat shock protein 47 (HSP47), HSP70, La/Sjögren syndrome type B antigen (SS-B), and vimentin in sputum supernatant samples were determined in quadruplicate, using a customized microarray (Human Array-COPD Microarray slides; RayBiotech, Inc., Norcross, GA, USA).

Data analysis

We conducted statistical analysis using SPSS 19.0, MedCalc 15.8, and Origin 2017. The demographic characteristics of subjects were presented by descriptive statistics. Categorical variables were expressed in integers; differences between groups were analyzed by chi-squared tests. Parametric and nonparametric data were presented as mean and median, respectively; while differences between groups were performed by independent-samples *t*-test and Mann–Whitney *U* test, respectively.

A paired-sample test was used to compare variables between day 1 and day 2 to determine whether the overall mean effects at different times were identical. Comparisons between repeated measurements were performed using a paired *t*-test if the difference between 2 days ($d = \text{day 1} - \text{day 2}$) was normally or logarithmically normally distributed, and using a Wilcoxon matched-pair signed-rank test if d was non-normally distributed. The repeatability of measurements was tested by intraclass correlation coefficient (ICC) and graphically depicted by Bland–Altman plotting.^{14–16} The ICC version was defined by two-way random effects model, using average and consistency measurement type with inter-item correlations. According to Landis and Koch, the degree of repeatability between 0.01 and 0.20 was deemed minor, 0.21–0.40 was fair, 0.41–0.60 was moderate, 0.61–0.80 was substantial, and 0.81–1.00 was very good.¹⁷ The bias between repeated measurements was estimated as the mean difference between 2 days (\bar{d}). The variation of \bar{d} is described by the SD of the difference (SD). In Bland–Altman plotting, the limits of agreement (LoA) were estimated by 95% CI of \bar{d} ($\bar{d} - 1.96 \text{ SD}$, $\bar{d} + 1.96 \text{ SD}$); adjusting for sampling error, the 95% CI of the upper and lower bounds of LoA (LOA CI) was also shown in the Bland–Altman plots. Outliers with *z*-scores of >3 or <-3 were removed to avoid distorting SD boundaries.

The correlation between variables was examined by Spearman rank correlation coefficient; correlation (*r*) values above 0.50 were considered relevant if they were significant at the 0.05 level. Between-group comparisons and correlations were examined for sputum obtained on day 1.

Results

The demographic and clinical characteristics of all subjects are shown in Table 1. There were no significant differences in age, sex, number of smokers, or smoking history (pack-years) between healthy and stable COPD groups. More than 50% of the data were undetectable for BAFF, IFN- γ , IL4, IL17, leptin, and TGF β ; thus, these markers were removed. All measurement results are provided in Table S1 and basic scatter plots with the pair-wise comparisons of the two measurements are provided in Figures S1–S4.

Reproducibility of soluble protein marker detections in sputum

Paired *t*-tests and Wilcoxon matched-pair signed-rank tests showed no significant differences between the repeated

Table 1 Demographic and clinical characteristics of subjects

Characteristics	Healthy (n=9)	Stable COPD (n=23)	P-value
Age, years ^a	68 (62–74)	67 (64–71)	0.857
Sex, male	7	23	0.128 ^c
Smoking, n	5	21	0.068 ^c
Former smoking, n	4	13	0.825 ^c
Active smoking, n	1	8	0.367 ^c
Smoking, pack-years ^b	3 (0–45)	30 (15–40)	0.213
pre-FEV ₁ , %pred ^a	92 (85–99)	46 (38–54)	0.000
post-FEV ₁ , %pred ^a	96 (87–104)	51 (43–59)	0.000
pre-FEV ₁ /FVC, % ^a	83 (80–86)	48 (44–52)	0.000
post-FEV ₁ /FVC, % ^a	85 (81–88)	49 (45–54)	0.000
GOLD I–IV, n			NA
I	NA	2	
II	NA	9	
III	NA	8	
IV	NA	4	
Therapies, n			NA
LAMA	NA	7	
LABA	NA	15	
ICS	NA	15	

Notes: ^aData with normal distribution were expressed as mean and 95% CI and analyzed by independent-samples *t*-test. ^bData with non-normal distribution were expressed as median and interquartile range (IQR) and analyzed by Mann–Whitney *U* test. ^cCategorical variables were expressed in number and analyzed by chi-squared tests. **Abbreviations:** GOLD, Global Initiative for Chronic Obstructive Lung Disease; NA, not applicable; LAMA, long-acting muscarinic antagonists; LABA, long-acting beta-agonists; ICS, inhaled corticosteroid.

Table 2 Paired-samples test results

Soluble protein markers	P-value	Autoantibodies	P-IgG	P-IgM	Sputum cell counts	P-value
IL1 β	0.036	Aggrecan	0.665	0.117	Neutrophils (%)	0.235
IL5	0.129	Amyloid	0.318	0.600	Macrophages (%)	0.400
IL6	0.396	CENP-B	0.951	0.014	Eosinophils (%)	0.722
IL8	0.072	Collagen I	0.977	0.301	Lymphocytes (%)	0.345
IL10	0.213	Collagen II	0.859	0.954		
IL13	0.152	Collagen IV	0.269	0.520		
IL33	0.511	Cytochrome C	0.982	0.631		
sRAGE	0.588	Cytokeratin 18	0.517	0.939		
TNF α	0.039	Elastin	0.106	0.754		
VEGF	0.112	HSP47	0.914	0.078		
		HSP70	0.214	0.238		
		SS-B	0.387	0.532		
		Vimentin	0.178	0.829		

Notes: Day 1 – day 2 data with normal distribution or logarithmic normal distribution were analyzed by paired t-test. Non-normally distributed day 1 – day 2 data were analyzed by Wilcoxon test.

Abbreviations: CENP-B, centromere protein B; HSP, heat shock protein; IL, interleukin; sRAGE, the soluble receptor for advanced glycation end-products; SS-B, La/Sjögren syndrome type B antigen; TNF α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor.

detection of IL5, IL6, IL8, IL10, IL13, IL33, sRAGE, and VEGF; differences were detected for IL1 β ($P=0.036$) and TNF α ($P=0.039$) (Table 2; Figure S1). The repeatability of soluble protein markers in PBS supernatant shown by ICC was very good for IL10, IL33, and VEGF; substantial for IL6, IL8, IL13, sRAGE, and TNF α ; moderate for IL5; and fair for IL1 β (Table 3). For IL5, IL8, IL13, and VEGF in PBS supernatant, all data points were within the LoA CI in Bland–Altman plots (Figure 1). There was only 1/32 (3.1%) data point outside the LoA CI for IL1 β , IL6, IL10, IL33,

and TNF α ; 2/32 (6.2%) data points were outside the LoA CI for sRAGE (Figure 1).

Reproducibility of autoantibody detections in sputum

There were no differences between repeated detections for all 13 types of IgG autoantibodies (aggrecan, amyloid, CENP-B, collagen I, collagen II, collagen IV, cytochrome C, cytokeratin 18, elastin, HSP47, HSP70, SS-B, and vimentin) and for 12 types of corresponding IgM autoantibodies;

Table 3 Intraclass correlation coefficient (ICC) assessment results

Soluble protein markers	ICC	IgG	ICC-IgG	ICC-IgM	Sputum cell counts	ICC
IL1 β	0.292	Aggrecan	0.924	0.073	Neutrophils (%)	0.861
IL5	0.603	Amyloid	0.070	0.061	Macrophages (%)	0.871
IL6	0.783	CENP-B	0.863	0.958	Eosinophils (%)	0.840
IL8	0.770	Collagen I	0.333	0.754	Lymphocytes (%)	0.413
IL10	0.831	Collagen II	0.882	0.936		
IL13	0.774	Collagen IV	0.742	0.732		
IL33	0.991	Cytochrome C	0.793	0.533		
sRAGE	0.711	Cytokeratin 18	0.318	0.826		
TNF α	0.698	Elastin	0.661	0.306		
VEGF	0.903	HSP47	0.834	0.062		
		HSP70	0.830	0.862		
		SS-B	0.923	0.598		
		Vimentin	0.514	0.462		

Abbreviations: ICC, intraclass correlation coefficient; CENP-B, centromere protein B; HSP, heat shock protein; IL, interleukin; sRAGE, the soluble receptor for advanced glycation end-products; SS-B, La/Sjögren syndrome type B antigen; TNF α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor.




Figure 1 Bland–Altman plots of soluble protein markers in sputum supernatant. Bland–Altman plots of (A) IL1β; (B) IL5; (C) IL6; (D) IL8; (E) IL10; (F) IL13; (G) IL33; (H) sRAGE; (I) TNFα; and (J) VEGF.

Abbreviations: IL, interleukin; sRAGE, soluble receptor for advanced glycation end-products; TNFα, tumor necrosis factor-α; VEGF, vascular endothelial growth factor.

a difference was detected for CENP-B IgM ($P=0.014$) in sputum supernatant (Table 2; Figures S2 and S3).

The repeatability shown by ICC was very good for IgG autoantibodies against aggrecan, CENP-B, collagen II,

HSP47, HSP70, and SS-B; substantial for IgGs against cytochrome C, collagen IV, and elastin; moderate for anti-vimentin IgG; fair for anti-collagen I and anti-cytokeratin 18 IgGs; and minor for IgG against amyloid (Table 3).

All data points were within the LoA CI in Bland–Altman plots for IgGs against amyloid, collagen I, cytochrome C, cytokeratin 18, HSP70, and vimentin (Figure 2). There was only 1/28 (3.6%) data point outside the LOA CI for the remaining seven types of IgGs (Figure 2).

The repeatability of IgM autoantibody in PBS supernatant, shown by ICC, was very good for IgMs against CENP-B, collagen II, cytokeratin 18, and HSP70; substantial for IgMs against collagen I and collagen IV; moderate for anti-cytochrome C, anti-SS-B, and anti-vimentin IgMs; fair for anti-elastin IgM; and minor for IgMs against aggrecan, amyloid, and HSP47 (Table 3). All data points were within the LoA CI in Bland–Altman plots for IgMs against amyloid

and collagen I (Figure 3). There was only 1/28 (3.6%) data point outside the LOA CI for the remaining eleven types of IgM (Figure 3).

All Bland–Altman plots for fluid phase in PBS supernatant showed that the solid line representing the mean difference was close to the dotted line representing a mean difference of 0. In the assessment of the distribution of data points in Bland–Altman plots as described above, there was a good consistency in repeated measurement.

Repeatability of sputum cell counts

There were no differences between repeated detection for differential cell counts in sputum (Table 2; Figure S4).




Figure 2 (Continued)




Figure 2 Bland–Altman plots of IgG autoantibodies in sputum supernatant. Bland–Altman plots of (A) anti-aggreccan IgG; (B) anti-amyloid IgG; (C) anti-CENP-B IgG; (D) anti-collagen I IgG; (E) anti-collagen II IgG; (F) anti-collagen IV IgG; (G) anti-cytochrome C IgG; (H) anti-cytokeratin 18 IgG; (I) anti-elastin IgG; (J) anti-HSP47 IgG; (K) anti-HSP70 IgG; (L) anti-SS-B IgG; and (M) anti-vimentin IgG.

Abbreviations: CENP-B, centromere protein B; HSP47, heat shock protein 47; HSP70, heat shock protein 70; SS-B, La/Sjögren syndrome type B antigen.

The reproducibility of sputum cell counts was very good for neutrophils (0.86), macrophages (0.87), and eosinophils (0.84); it was moderate for lymphocytes (0.40) (Table 3). All data points were within the LoA CI in Bland–Altman plots for macrophages and lymphocytes (Figure 4). There was only 1/32 (3.1%) data point outside the LoA CI for neutrophils; 2/32 (6.2%) data points were outside the LoA CI for eosinophils (Figure 4).

Comparison of measurements in healthy subjects and stable COPD

Compared with healthy subjects, stable COPD patients showed trends for higher relative fluorescence intensity of anti-cytochrome C IgG (665.9 vs 902.0, $P=0.008$) and anti-aggreccan IgM (0.0 vs 148.5, $P=0.006$) in sputum; they also showed trends for higher neutrophil percentage (53.3% vs 77.4%, $P=0.001$) and eosinophil percentage (1.6% vs 3.6%, $P=0.004$) in sputum (Table S2). Anti-cytochrome C IgM (11,815.6 vs 8,710.0, $P=0.036$) and macrophages (51.2% vs 15.1%, $P=0.001$) were lower in sputum of patients with COPD than in sputum of healthy subjects (Table S2).

Correlations between measurements in stable COPD patients

Correlation analysis suggested a positive correlation for stable COPD patients between sputum anti-collagen II IgG and post-bronchodilator FEV₁% ($r=0.670$, $P=0.002$), as well as between sputum anti-collagen II IgG and sputum lymphocytes ($r=0.649$, $P=0.003$); there were positive correlations between anti-aggreccan IgG and eosinophils in sputum ($r=0.574$, $P=0.010$), between anti-collagen IV IgM and lymphocytes in sputum ($r=0.610$, $P=0.006$), and between post-bronchodilator FEV₁% and sputum macrophages ($r=0.519$, $P=0.002$); there was a negative correlation between anti-cytochrome C IgG and sputum eosinophils ($r=-0.639$, $P=0.003$).

Discussion

In this study, we treated selected fresh sputum with PBS, followed by low-temperature centrifugation to obtain supernatant; we then investigated soluble protein markers and IgG and IgM autoantibody profiles of these PBS supernatants by microarray analysis. We found that the measurements

obtained twice within 6 days were highly reproducible in healthy subjects and stable COPD patients. Notably, this is the first assessment of the repeatability of fluid-phase measurements in PBS-treated sputum supernatant with high-throughput method. This study confirmed the presence of autoantibodies in the airways of healthy subjects and COPD patients, which can be detected by microarrays in the analyses of sputum supernatant.

The “two-step” sputum processing method allows investigators to obtain supernatant by PBS processing and cell lysis with DTT; notably, it avoids the destruction of soluble media by DTT. In this study, the reproducibility of fluid-phase measurements in PBS-supernatant and cell proportion from sputum treated with “two-step” processing method was

good. These results suggested the validity of this method, which may be useful in clinical analyses.

ICC and Bland–Altman plots are commonly used statistical methods for consistency testing. The paired *t*-test can only indicate whether the average difference between the two measurements is equal to 0, according to the current data it cannot fully reflect the consistency between these measurements. In addition, the *P*-value is affected by the degree of freedom; thus, it is not appropriate to use the paired *t*-test instead of the consistency test. The result of ICC is susceptible to the presence of the outliers, while the Bland–Altman consistency analysis method is relatively more robust. For the above reasons, a paired-samples test approach, including ICC and Bland–Altman plots, was used to evaluate repeatability




Figure 3 (Continued)




Figure 3 Bland–Altman plots of IgM autoantibodies in sputum supernatant. Bland–Altman plots of (A) anti-aggrecan IgM; (B) anti-amyloid IgM; (C) anti-CENP-B IgM; (D) anti-collagen I IgM; (E) anti-collagen II IgM; (F) anti-collagen IV IgM; (G) anti-cytochrome C IgM; (H) anti-cytokeratin 18 IgM; (I) anti-elasticin IgM; (J) anti-HSP47 IgM; (K) anti-HSP70 IgM; (L) anti-SS-B IgM; and (M) anti-vimentin IgM.

Abbreviations: CENP-B, centromere protein B; HSP47, heat shock protein 47; HSP70, heat shock protein 70; SS-B, La/Sjögren syndrome type B antigen.

in this study. We evaluated the reproducibility of 10 soluble protein markers that often serve as biomarkers for chronic inflammatory airway disease. Most of these soluble protein markers show good consistency when measured twice within 6 days. Although 96.9% of data points were inside the LoA CI for IL1 β and TNF α in Bland–Altman plots, there was a significant difference between the repeated measurements. The ICC was 0.292 for IL1 β and 0.698 for TNF α , indicating low to medium reproducibility, which may be because of fluctuations that occur under physiological conditions.

We tested the detection reproducibility for IgG and IgM autoantibody profiles by microarray analysis and confirmed that these autoantibodies are detectable in PBS-treated supernatant with a good reproducibility; IgG was clearly detectable. ICC indicated that results regarding IgM autoantibodies generally tended to be less reproducible than those regarding IgG autoantibodies for aggrecan, cytochrome C, elastin, HSP47, and SS-B. This might be because IgM is often produced in early stages of humoral immunity, is maintained for a short time, and then cleared rapidly. A wide variety of natural autoantibodies (NAAs) of the IgM, IgG, and IgA

classes are present in normal individuals.¹⁸ NAAs play an important role in maintaining the immunological homeostasis by participating in immune regulation, repertoire selection, and resistance to infections. Natural IgM antibodies are required in early effective control of bacterial and viral infections and serve to prime the ensuing IgG response.¹⁹ Notably, autoantibodies restrain the harmful consequences of inflammatory autoimmune conditions.²⁰ In this study, we confirmed the existence of NAAs in normal human airway, as well as in the airway of COPD patients. The role of autoimmune pathology in the development and progression of COPD is increasingly appreciated. Packard et al demonstrated that COPD patients produce autoantibodies in serum which is correlated with the disease phenotype.⁵ A case series identified antibody deficiency as a potentially treatable risk factor for frequent COPD exacerbations.²¹ Collagen V-mediated autoimmunity was increased in smoking controls and may potentially contribute to the pathogenesis of COPD.²² Here, we have revealed the presence of IgG and IgM autoantibodies to multiple antigens, including collagen, in the airways of COPD patients. Stable COPD patients differed from healthy




Figure 4 Bland–Altman plots of sputum cell counts. Bland–Altman plots of (A) sputum neutrophils; (B) sputum macrophages; (C) sputum eosinophils; and (D) sputum lymphocytes.

subjects in anti-cytochrome C IgG and anti-aggrecan IgM and anti-cytochrome C IgM. We also found a significant positive correlation for stable COPD patients between sputum anti-collagen II IgG and post-bronchodilator FEV₁%, as well as between anti-aggrecan IgG and eosinophils; and a negative correlation between anti-cytochrome C IgG and sputum eosinophils. Further studies with larger samples are warranted to explore the role of these autoantibodies in the development of COPD.

Comparison of sputum measurements in healthy subjects and COPD patients showed significant differences in the proportion of neutrophils and eosinophils, the levels of anti-cytochrome C IgG and anti-aggrecan IgM and anti-cytochrome C IgM. The comparison and the correlation between markers in sputum and the clinical parameters suggested the validity of this method. Biomarker distributions

are often complex because of heterogeneities due to various disease stages and endotypes. In particular, COPD is a highly heterogeneous disease. Therefore, we believe that overlapping distributions between healthy subjects and COPD patients are logical and acceptable.

There are some limitations in this study. First, for BAFF, IFN- γ , IL4, IL17, leptin, and TGF β , >50% of the data were undetectable. The levels of these protein markers may indeed be low in the airway secretions of healthy and COPD individuals. However, this phenomenon may be related to the reduction in soluble protein marker concentrations after ninefold dilution in sputum processing. To clarify the cause, further studies with high-speed centrifugation of the original sputum, in order to obtain supernatant for detection of soluble protein markers, can be conducted. Second, the sample size was small, although it was comparable to that of a classic

study with similar design.⁶ In the future, we will expand the sample size and evaluate the reproducibility in COPD patients and healthy subjects.

Conclusion

The current study confirms the high repeatability of fluid-phase measurements in PBS supernatant, using the two-step sputum processing method. Autoantibodies of IgG and IgM of multiple antigens were found in the airways of COPD patients and are detected by using a high-throughput protein chip for analyses of sputum supernatant. These results demonstrated the potential use of the two-step sputum processing method as a noninvasive tool for the study of biomarkers, airway inflammation, and localized immunity in COPD patients.

Data sharing statement

The authors intend to share individual deidentified participant data, including demographics, detailed clinical information, and cell and fluid-phase measurements in sputum. The data will be accessible by emailing the authors after the manuscript is published. Subjects' identity information will not be available for privacy reasons.

Acknowledgments

We thank Ryan Chastain-Gross, PhD, from Liwen Bianji, Edanz Group China (www.liwenbianji.cn/ac), for editing the English text of a draft of this manuscript. This study was supported by National Key R&D Program of China (2017YFC1310600, 2017YFC1310601), the Guangzhou Healthcare Collaborative Innovation Major Project (201604020012), Medical Scientific Research Foundation of Guangdong Province (C2017050), and Postdoctoral start-up funding of Guangzhou (191710).

Author contributions

All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Kelly MM, Keatings V, Leigh R, et al. Analysis of fluid-phase mediators. *Eur Respir J Suppl.* 2002;37:24s–39s.
- Hacievliyagil SS, Mutlu LC, Temel I. Airway inflammatory markers in chronic obstructive pulmonary disease patients and healthy smokers. *Niger J Clin Pract.* 2013;16(1):76–81. doi:10.4103/1119-3077.106771
- Rovina N, Papapetropoulos A, Kollintza A, et al. Vascular endothelial growth factor: an angiogenic factor reflecting airway inflammation in healthy smokers and in patients with bronchitis type of chronic obstructive pulmonary disease? *Respir Res.* 2007;8:53. doi:10.1186/1465-9921-8-91
- Agusti M, Donaldson C. Hypothesis: does COPD have an autoimmune component? *Thorax.* 2003;58(10):832. doi:10.1136/thorax.58.10.832
- Packard TA, Li QZ, Cosgrove GP, Bowler RP, Cambier JC. COPD is associated with production of autoantibodies to a broad spectrum of self-antigens, correlative with disease phenotype. *Immunol Res.* 2013;55(1–3):48–57. doi:10.1007/s12026-012-8347-x
- Pizzichini E, Pizzichini MM, Efthimiadis A, et al. Indices of airway inflammation in induced sputum: reproducibility and validity of cell and fluid-phase measurements. *Am J Respir Crit Care Med.* 1996; 154(2 Pt 1):308–317. doi:10.1164/ajrccm.154.2.8756799
- Efthimiadis A, Spanevello A, Hamid Q, et al. Methods of sputum processing for cell counts, immunocytochemistry and in situ hybridisation. *Eur Respir J Suppl.* 2002;37:19s–23s.
- Grebski E, Peterson C, Medici TC. Effect of physical and chemical methods of homogenization on inflammatory mediators in sputum of asthma patients. *Chest.* 2001;119(5):1521–1525.
- Bafadhel M, McCormick M, Saha S, et al. Profiling of sputum inflammatory mediators in asthma and chronic obstructive pulmonary disease. *Respiration.* 2012;83(1):36–44. doi:10.1159/000330667
- Hinks TSC, Brown T, Lau LCK, et al. Multidimensional endotyping in patients with severe asthma reveals inflammatory heterogeneity in matrix metalloproteinases and chitinase 3–like protein 1. *J Allergy Clin Immunol.* 2016;138(1):61–75. doi:10.1016/j.jaci.2015.11.020
- Ghebre MA, Desai D, Singapuri A, et al. Sputum inflammatory mediators are increased in aspergillus fumigatus culture-positive asthmatics. *Allergy Asthma Immun.* 2017;9(2):177. doi:10.4168/aa.2017.9.2.177
- Kolsum U, Donaldson GC, Singh R, et al. Blood and sputum eosinophils in COPD; relationship with bacterial load. *Respir Res.* 2017;18(1):88. doi:10.1186/s12931-017-0570-5
- Pizzichini E, Pizzichini MM, Efthimiadis A, et al. Measurement of inflammatory indices in induced sputum: effects of selection of sputum to minimize salivary contamination. *Eur Respir J.* 1996;9(6): 1174–1180.
- Bartko JJ. The intraclass correlation coefficient as a measure of reliability. *Psychol Rep.* 1966;19(1):3–11. doi:10.2466/pr0.1966.19.1.3
- Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet.* 1986;1(8476): 307–310.
- Bland JM, Altman DG. Measuring agreement in method comparison studies. *Stat Methods Med Res.* 1999;8(2):135–160. doi:10.1177/ 096228029900800204
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics.* 1977;33(1):159–174.
- Lacroix-Desmazes L, Kaveri SV, Mouthon L, et al. Self-reactive antibodies (natural autoantibodies) in healthy individuals. *J Immunol Methods.* 1998;216(1–2):117–137.
- Gronwall C, Silverman GJ. Natural IgM: beneficial autoantibodies for the control of inflammatory and autoimmune disease. *J Clin Immunol.* 2014;34(Suppl 1):S12–S21. doi:10.1007/s10875-014-0025-4
- Zohar Y, Wildbaum G, Karin N. Beneficial autoimmunity participates in the regulation of rheumatoid arthritis. *Front Biosci.* 2006;11: 368–379.
- McCullagh BN, Comellas AP, Ballas ZK, et al. Antibody deficiency in patients with frequent exacerbations of Chronic Obstructive Pulmonary Disease (COPD). *PLoS One.* 2017;12(2):e172437. doi:10.1371/journal.pone.0172437
- Rinaldi M, Lehouck A, Heulens N, et al. Anti-elastin B-cell and T-cell immunity in patients with chronic obstructive pulmonary disease. *Thorax.* 2012;67(8):694–700. doi:10.1136/thoraxjnl-2011-200690

Supplementary materials

Table S1 All measurement results

Soluble protein markers	Day 1	Day 2	P-value
IL1 β ^a	46.5 (27.5, 65.5)	63.0 (39.7, 86.3)	0.036
IL5 ^a	6.4 (3.5, 9.2)	11.8 (4.6, 18.9)	0.129
IL6 ^a	352.0 (208.2, 495.8)	337.4 (186.3, 488.5)	0.396
IL8 ^a	1,551.1 (1,076.5, 2,025.7)	1,648.1 (1,344.2, 1,951.9)	0.072
IL10 ^b	0.4 (0.0, 1.8)	0.9 (0.3, 1.7)	0.213
IL13 ^b	0.5 (0.2, 1.0)	0.8 (0.3, 1.4)	0.152
IL33 ^b	1.6 (0.0, 10.1)	4.1 (0.0, 13.1)	0.511
sRAGE ^b	406.3 (39.3, 1,244.3)	274.6 (77.5, 1,238.4)	0.588
TNF α ^a	85.8 (18.8, 152.7)	109.4 (59.6, 159.2)	0.039
VEGF ^a	2,584.4 (2,159.1, 3,009.7)	2,869.7 (2,402.8, 3,336.6)	0.112
IgG autoantibodies			
Aggrecan ^b	568.3 (501.5, 707.6)	556.8 (462.9, 698.9)	0.665
Amyloid ^a	89.2 (66.9, 111.5)	114.2 (67.9, 160.5)	0.318
CENP-B ^b	3,071.7 (2,490.1, 4,090.3)	3,042.8 (2,820.1, 4,063.2)	0.951
Collagen I ^b	278.1 (223.6, 390.4)	278.7 (207.4, 333.7)	0.977
Collagen II ^b	301.2 (232.0, 638.0)	311.3 (198.8, 765.1)	0.859
Collagen IV ^b	116.9 (46.8, 344.4)	130.5 (56.1, 464.1)	0.269
Cytochrome C ^b	795.4 (637.4, 989.6)	883.4 (623.6, 1,016.4)	0.982
Cytokeratin 18 ^b	933.3 (433.9, 1,920.7)	838.1 (302.5, 1,916.6)	0.517
Elastin ^a	761.9 (643.2, 880.7)	684.3 (617.9, 750.7)	0.106
HSP47 ^a	407.5 (253.1, 562.0)	359.0 (241.7, 476.3)	0.914
HSP70 ^b	5,369.5 (4,440.2, 6,252.3)	5,515.1 (4,684.1, 6,567.3)	0.214
SS-B b ^a	4,033.8 (3,545.6, 4,790.4)	4,201.5 (3,866.6, 5,156.0)	0.387
Vimentin ^a	365.1 (282.1, 448.1)	432.6 (306.8, 558.3)	0.178
IgM autoantibodies			
Aggrecan ^b	84.0 (0.0, 465.1)	111.3 (0.0, 709.4)	0.117
Amyloid ^b	23.6 (1.0, 101.1)	18.9 (1.5, 85.9)	0.600
CENP-B ^b	3,885.5 (2,596.2, 8,974.4)	6,131.3 (2,557.9, 11,631.8)	0.014
Collagen I ^a	284.1 (183.4, 384.8)	257.0 (151.6, 362.5)	0.301
Collagen II ^b	463.6 (12.4, 1,147.2)	338.3 (8.6, 1,614.4)	0.954
Collagen IV ^b	9.2 (0.3, 278.7)	1.4 (0.0, 433.2)	0.520
Cytochrome C ^b	8,572.7 (7,913.6, 11,828.1)	9,157.1 (6,940.2, 10,326.5)	0.631
Cytokeratin 18 ^a	6,480.5 (3,560.2, 9,400.8)	5,846.3 (3,456.4, 8,236.1)	0.939
Elastin ^a	10,074.5 (9,025.2, 11,123.7)	10,311.0 (8,986.3, 11,635.7)	0.754
HSP47 ^b	374.2 (132.6, 801.7)	508.9 (177.9, 1,043.8)	0.078
HSP70 ^b	6,132.8 (4,592.6, 7,867.5)	6,077.0 (4,628.5, 9,365.8)	0.238
SS-B ^b	5,240.3 (4,630.9, 7,250.0)	5,471.9 (4,618.4, 6,972.6)	0.532
Vimentin ^a	815.8 (510.0, 1,121.5)	753.3 (477.8, 1,028.9)	0.829
Sputum cell counts			
Neutrophils (%) ^b	75.0 (53.3, 88.6)	70.5 (53.2, 85.9)	0.235
Macrophages (%) ^b	19.0 (6.7, 42.5)	23.3 (9.1, 44.2)	0.400
Eosinophils (%) ^b	2.5 (1.3, 4.0)	2.2 (1.2, 4.5)	0.722
Lymphocytes (%) ^b	1.2 (0.5, 1.7)	1.0 (0.4, 1.4)	0.345

Notes: ^aDay 1 – day 2 data with normal distribution or logarithmic normal distribution were analyzed by paired t-test. ^bNon-normally distributed day 1 – day 2 data were analyzed by Wilcoxon test.

Abbreviations: CENP-B, centromere protein B; HSP70, heat shock protein 70; IL, interleukin; INF- γ , interferon- γ ; sRAGE, the soluble receptor for advanced glycation end-products; SS-B, La/Sjögren syndrome type B antigen; TNF α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor.




Figure S1 Scatter plots of soluble protein markers with pair-wise comparisons of the two measurements. Scatter plots of (A) IL1 β ; (B) IL5; (C) IL6; (D) IL8; (E) IL10; (F) IL13; (G) IL33; (H) sRAGE; (I) TNF α ; and (J) VEGF.

Abbreviations: IL, interleukin; sRAGE, soluble receptor for advanced glycation end-products; TNF α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor.




Figure S2 Scatter plots of IgG autoantibodies with pair-wise comparisons of the two measurements. Scatter plots of (A) anti-aggrecaan IgG; (B) anti-amyloid IgG; (C) anti-CENP-B IgG; (D) anti-collagen I IgG; (E) anti-collagen II IgG; (F) anti-collagen IV IgG; (G) anti-cytochrome C IgG; (H) anti-cytokeratin 18 IgG; (I) anti-elasticin IgG; (J) anti-HSP47 IgG; (K) anti-HSP70 IgG; (L) anti-SS-B IgG; and (M) anti-vimentin IgG.

Abbreviations: CENP-B, centromere protein B; HSP47, heat shock protein 47; HSP70, heat shock protein 70; SS-B, La/Sjögren syndrome type B antigen.




Figure S3 Scatter plots of IgM autoantibodies with pair-wise comparisons of the two measurements. Scatter plots of (A) anti-aggrecan IgM; (B) anti-amyloid IgM; (C) anti-CENP-B IgM; (D) anti-collagen I IgM; (E) anti-collagen II IgM; (F) anti-collagen IV IgM; (G) anti-cytochrome C IgM; (H) anti-cytokeratin 18 IgM; (I) anti-elasticin IgM; (J) anti-HSP47 IgM; (K) anti-HSP70 IgM; (L) anti-SS-B IgM; (M) anti-vimentin IgM.

Abbreviations: CENP-B, centromere protein B; HSP47, heat shock protein 47; HSP70, heat shock protein 70; SS-B, La/Sjögren syndrome type B antigen.




Figure S4 Scatter plots of sputum cell counts with pair-wise comparisons of the two measurements. Scatter plots of (A) sputum neutrophils; (B) sputum macrophages; (C) sputum eosinophils; and (D) scatter plots of sputum lymphocytes.

Table S2 Comparison of measurements between groups

Soluble protein markers	Healthy	Stable COPD	P-value
IL1 β ^b	22.1 (13.3, 27.7)	25.8 (4.8, 101.9)	0.690
IL5 ^b	1.3 (0.2, 4.7)	6.1 (0.8, 10.1)	0.153
IL6 ^b	102.1 (21.9, 168.6)	381.8 (57.1, 797.5)	0.075
IL8 ^b	1,171.2 (863.2, 1,445.0)	1,302.6 (906.3, 1,721.0)	0.463
IL10 ^b	0.2 (0.0, 0.7)	0.6 (0.1, 2.2)	0.094
IL13 ^b	0.6 (0.3, 1.0)	0.4 (0.1, 1.0)	0.850
IL33 ^b	0.6 (0.0, 5.3)	2.9 (0.0, 14.4)	0.351
sRAGE ^b	339.6 (195.1, 3,188.6)	473.0 (16.4, 1,043.1)	0.201
TNF α ^b	28.9 (0.0, 44.4)	26.7 (0.0, 104.0)	0.680
VEGF ^b	2,549.7 (2,326.0, 3,093.1)	2,371.8 (1,574.1, 3,308.6)	0.438
IgG autoantibodies			
Aggrecan ^b	531.1 (474.3, 812.0)	569.6 (503.4, 622.5)	0.863
Amyloid ^b	86.7 (56.0, 95.2)	81.8 (38.4, 136.0)	0.825
CENP-B ^b	3,994.0 (2,907.8, 4,956.0)	2,847.6 (2,430.3, 3,878.2)	0.099
Collagen I ^a	356.8 (200.4, 513.2)	288.1 (243.9, 332.4)	0.357
Collagen II ^b	408.4 (293.6, 1,000.9)	263.6 (211.7, 536.9)	0.073
Collagen IV ^b	149.9 (41.2, 658.3)	104.8 (48.1, 172.6)	0.363
Cytochrome C ^b	665.9 (566.8, 763.6)	902.0 (697.4, 1,028.0)	0.008
Cytokeratin I8 ^b	843.0 (278.0, 2,817.3)	945.3 (508.9, 1,932.7)	0.712
Elastin ^b	706.2 (661.3, 809.3)	742.3 (546.5, 869.0)	0.863
HSP47 ^b	274.3 (166.0, 585.9)	206.0 (108.2, 645.3)	0.712
HSP70 ^b	5,640.2 (4,735.5, 7,712.8)	5,016.0 (4,255.8, 5,498.5)	0.176
SS-B ^b	3,771.4 (3,562.4, 4,298.6)	4,411.1 (3,419.8, 5,113.1)	0.228
Vimentin ^b	295.8 (284.3, 574.1)	301.6 (221.5, 474.8)	0.313
IgM autoantibodies			
Aggrecan ^b	0.0 (0.0, 30.8)	148.5 (1.2, 519.3)	0.006
Amyloid ^b	45.8 (0.0, 106.9)	13.6 (1.1, 106.9)	0.843
CENP-B ^b	3,659.3 (2,152.0, 6,249.1)	4,134.9 (2,908.3, 12,490.8)	0.325
Collagen I ^b	283.6 (95.6, 365.7)	199.1 (93.6, 383.1)	0.941
Collagen II ^b	463.9 (0.0, 1,252.4)	463.3 (109.0, 1,221.0)	0.336
Collagen IV ^b	1.2 (0.0, 321.8)	20.8 (1.1, 307.6)	0.359
Cytochrome C ^a	11,815.6 (9,060.0, 14,571.2)	8,710.0 (265.4, 791.2)	0.036
Cytokeratin I8 ^b	3,926.2 (1,233.7, 6,578.3)	4,046.7 (845.2, 12,122.9)	0.863
Elastin ^a	10,719.3 (8,817.8, 12,620.8)	9,769.0 (8,409.5, 11,128.6)	0.396
HSP47 ^b	232.2 (117.3, 930.8)	416.4 (127.0, 782.3)	0.806
HSP70 ^b	5,364.2 (3,779.5, 7,710.8)	6,418.4 (4,536.1, 7,907.4)	0.572
SS-B ^b	5,831.1 (4,664.5, 10,809.2)	5,000.8 (4,604.1, 6,589.2)	0.446
Vimentin ^b	677.7 (504.5, 1,471.3)	512.3 (278.7, 835.2)	0.268
Sputum cell counts			
Neutrophils (%) ^a	53.3 (39.3, 67.3)	77.4 (70.7, 84.2)	0.001
Macrophages (%) ^b	51.2 (30.2, 58.0)	15.1 (4.6, 23.9)	0.001
Eosinophils (%) ^a	1.6 (0.7, 2.5)	3.6 (2.5, 4.7)	0.004
Lymphocytes (%) ^b	1.4 (1.1, 2.0)	0.9 (0.4, 1.7)	0.249

Notes: ^aData with normal distribution were expressed as mean and 95% CI and analyzed by *t*-test. ^bData with non-normal distribution were expressed as median and interquartile range (IQR) and analyzed by Mann–Whitney *U* test.

Abbreviations: CENP-B, centromere protein B; HSP70, heat shock protein 70; IL, interleukin; INF- γ , interferon- γ ; sRAGE, the soluble receptor for advanced glycation end-products; SS-B, La/Sjögren syndrome type B antigen; TNF α , tumor necrosis factor- α ; VEGF, vascular endothelial growth factor.

International Journal of COPD

Dovepress

Publish your work in this journal

The International Journal of COPD is an international, peer-reviewed journal of therapeutics and pharmacology focusing on concise rapid reporting of clinical studies and reviews in COPD. Special focus is given to the pathophysiological processes underlying the disease, intervention programs, patient focused education, and self management protocols.

This journal is indexed on PubMed Central, MedLine and CAS. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/international-journal-of-chronic-obstructive-pulmonary-disease-journal>