



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



## Short communication

# PCR-based retrospective evaluation of diagnostic samples for emergence of porcine deltacoronavirus in US swine



Avanti Sinha\*, Phillip Gauger, Jianqiang Zhang, Kyoung-Jin Yoon, Karen Harmon

Department of Veterinary Diagnostic and Production Animal Medicine, College of Veterinary Medicine, Iowa State University, Ames, IA, USA

## ARTICLE INFO

## Article history:

Received 29 December 2014

Received in revised form 22 May 2015

Accepted 4 June 2015

## Keywords:

Porcine deltacoronavirus

PDCoV

Retrospective

RT-PCR

## ABSTRACT

Porcine deltacoronavirus (PDCoV) was first identified in Hong Kong in a regional surveillance study for Coronaviruses in 2012 and was detected for the first time in United States (US) swine in February 2014. However, it remains unknown if PDCoV had been introduced into the US prior to that time period. In the present study, 1734 clinical samples (903 cases) submitted to the Iowa State University Veterinary Diagnostic Laboratory (ISU VDL) for enteric disease diagnosis between October 2012 and December 2013 were tested retrospectively for PDCoV using a virus-specific real-time reverse transcription (RT) PCR targeting conserved region of the membrane gene. PDCoV genome was first detected in a fecal sample collected on August 19th 2013 from Minnesota. Subsequently, PDCoV was observed in samples collected on August 20th and August 27th from Iowa and on August 29th from Illinois. Therefore, with available samples submitted to the ISU VDL, it can be inferred that PDCoV has been present in US swine at least since August 2013.

© 2015 Elsevier B.V. All rights reserved.

## 1. Introduction

Coronaviruses belong to the order *Nidovirales*, family *Coronaviridae* and subfamily *Coronavirinae* and until recently, were subdivided into 3 genera including *Alphacoronavirus*, *Betacoronavirus*, and *Gammacoronavirus* (Carstens, 2010). In 2012, researchers in Hong Kong conducted a regional surveillance study on a variety of domestic and wild animal species for coronaviruses from 2007 through 2011 and identified coronaviruses in birds and mammals that formed a new genus subsequently named *Deltacoronavirus* (Woo et al., 2012; King et al., 2012). According to that study, porcine deltacoronavirus (PDCoV) has been present in pigs in Hong Kong since 2009. PDCoV is a single stranded, positive sense, enveloped RNA virus with a genome size varying between 25.4 and 26.6 Kb (Woo et al., 2012; Li et al., 2014).

The presence of PDCoV in the US was first announced by the Ohio Department of Agriculture in early February 2014 after which its genetic characterization was reported by several laboratories (Wang et al., 2014; Li et al., 2014; Marthaler et al., 2014). As of April 23, 2015, PDCoV has been detected in 19 US states according to the weekly report on PDCoV in the USDA/APHIS website ([www.aphis.usda.gov/animal-health/sectd](http://www.aphis.usda.gov/animal-health/sectd)). Nonetheless, it remains unknown if PDCoV had

been introduced into the US earlier than 2014 or if it was introduced into the US at the same time as porcine epidemic diarrhea virus (PEDV) as early as April, 2013 (Stevenson et al., 2013). In the present study, we performed retrospective PCR testing on samples which were submitted to the Iowa State University Veterinary Diagnostic Laboratory (ISU VDL) between October 2012 and December 2013 for investigation of enteric diseases, to determine the earliest date PDCoV was detected in US swine.

## 2. Materials and methods

### 2.1. Clinical samples

A total of 1734 clinical samples, including feces, fecal swabs, oral fluids, and environmental and intestinal specimens from cases of porcine diarrhea submitted to the ISU VDL from October 2012 to December 2013, were used in the study. Intestinal samples were fixed in formalin and then embedded in paraffin blocks (FFPE samples) at the time of receipt. FFPE blocks were stored at room temperature and all other specimens were preserved at  $-80^{\circ}\text{C}$  from the time of original receipt and testing until retrospective testing for PDCoV.

### 2.2. Sample processing and extraction

A small amount of feces, fecal swabs and environmental specimens (swabs/swiffers) were placed in 1 ml PBS and agitated

\* Corresponding author at: 1600 S. 16th Street, College of Veterinary Medicine, Iowa State University, Ames, IA 50011, USA.

E-mail address: [asinha@iastate.edu](mailto:asinha@iastate.edu) (A. Sinha).

manually for three seconds prior to the extraction step. Viral RNA extraction was conducted with 100  $\mu$ l of processed samples using the MagMAX™ viral RNA isolation kit (Life Technologies, Carlsbad, CA) and a Kingfisher 96 instrument (Thermo Scientific, Waltham, MA) following manufacturer's instructions. The viral RNA was eluted into 90  $\mu$ l of elution buffer. Viral RNA was extracted from the FFPE intestinal samples using the MagMAX™ FFPE Total nucleic acid isolation kit (Life Technologies, Carlsbad, CA) and a Kingfisher 96 instrument (Thermo Scientific, Waltham, MA) using manufacturer's recommendations. Viral RNA was eluted into 70  $\mu$ l of elution buffer. FFPE intestinal tissues from PDCoV positive cases submitted to the ISU VDL and PDCoV positive amplification control (PAC) were included as positive controls.

### 2.3. PDCoV real-time RT-PCR

The primers and probe were designed to target the conserved regions of the PDCoV membrane protein gene based on the nucleotide sequences of strains HKU15-44 and HKU15-155 deposited in GenBank (JQ065042.1 and JQ065043.1, respectively) (Woo et al., 2012). Primer and probe sequences were later verified as appropriate for the US virus after sequence information was available. The forward primer sequence was 5'-CGA CCA CAT GGC CCC AAT TC-3', the reverse primer sequence was 5'-CAG CTC TTG CCC ATG TAG CTT-3', and the probe sequence was 5'-CAC ACC AGT CGT TAA GCA TGG CAA GC-3'. The probe was labeled using the FAM/ZEN/3' Iowa Black detector (Integrated DNA Technologies, Coralville, IA, USA). Real-time RT-PCR was conducted on nucleic acid extracts using a Path-ID Multiplex One-Step RT-PCR kit (Life Technologies, Carlsbad, CA). Real-time RT-PCR was performed on an ABI 7500 Fast instrument (Life Technologies, Carlsbad, CA) with the following conditions: 1 cycle of 45 °C for 10 min, 1 cycle of 95 °C for 10 min, and 40 cycles of 95 °C for 15 s and 60 °C for 45 s.

## 3. Results

A retrospective study, described here, was conducted to determine if PDCoV was introduced at approximately the same time as PEDV or if this may have occurred independently. The 1734 clinical samples tested for PDCoV represents 903 ISU VDL cases from eighteen states submitted from early October 2012 to December 2013 to detect various enteric pathogens. A total of five samples tested positive for the presence of PDCoV RNA with  $C_T$  values of 14.6, 15.1, 28.1, 36.6 and 36.7, respectively. The states, pathogens and specimens tested from the 903 cases are given as a total and percentage in Table 1. The numbers of PDCoV positive cases are also included in Table 1.

The PDCoV genome was first detected in a fecal sample collected on August 19th 2013 from a pig in Minnesota with a history of diarrhea and positive for *Salmonella krefeld* but negative for PEDV by PCR followed by detection of PDCoV in another fecal sample collected on August 20th from a pig in Iowa with lesions of ileitis and positive for *Lawsonia intracellularis* but negative for PEDV by PCR. RNA extracted from a fixed intestine sample collected on August 27th 2013 from Iowa that was negative for PEDV and positive for Rotavirus group A, group B and group C also tested positive for PDCoV. Thereafter, PDCoV was detected in fecal samples collected on August 29th 2013 in two additional pigs from Illinois which were negative by PCR for PEDV, *transmissible gastroenteritis virus* (TGEV) and *L. intracellularis* and also negative by immunohistochemistry for PEDV, group A rotavirus and TGEV.

## 4. Discussion

A study conducted to evaluate the presence of Deltacoronavirus identified a novel porcine deltacoronavirus genome in

**Table 1**

Categorization of specimens submitted to ISU VDL between 2012 and 2013 for diagnosis of porcine enteric disease and tested in this study for PDCoV RNA.

Classifications	Cases, No. (%)	Positive cases
<b>State</b>		
Iowa	488 (54.0)	2
North Carolina	110 (12.2)	0
Illinois	57 (6.3)	2
Oklahoma	46 (5.1)	0
Ohio	42 (4.6)	0
Indiana	38 (4.2)	0
Minnesota	36 (3.9)	1
North Dakota	27 (2.9)	0
Nebraska	14 (1.5)	0
Pennsylvania	12 (1.3)	0
Missouri	8 (0.9)	0
Utah	6 (0.6)	0
Wisconsin	6 (0.6)	0
Texas	5 (0.5)	0
Arizona	4 (0.4)	0
South Dakota	2 (0.2)	0
Kansas	1 (0.1)	0
Virginia	1 (0.1)	0
<b>Sample types</b>		
Feces	784 (86.8)	
Fecal swab	248 (27.5)	
Intestine	49 (5.4)	
Oral fluid	26 (2.9)	
Environmental	19 (2.1)	
<b>Pathogens tested</b>		
TGEV	634 (70.2)	
Rotavirus	423 (46.8)	
<i>E. Coli</i>	209 (23.1)	
<i>Lawsonia</i>	137 (15.2)	
PEDV	125 (13.8)	
PRCV	30 (3.3)	

Note: TGEV, *transmissible gastroenteritis virus*; PEDV, *porcine epidemic diarrhea virus*; PRCV, *porcine respiratory coronavirus*.

swine reported as the HKU15 (strains HKU15-44 and HKU15-155) in Hong Kong, China (Woo et al., 2012). In early February 2014, PDCoV was detected and announced in the US by the Ohio Department of Agriculture. Soon thereafter, genome sequencing and analysis of the PDCoV strains present in the US revealed approximately 99% nucleotide similarity of these newly emerged strains to the porcine deltacoronavirus HKU15-44 and HKU15-155 from China (Wang et al., 2014; Li et al., 2014; Marthaler et al., 2014). It was unknown when PDCoV might have been introduced into the US considering an additional coronavirus (PEDV) had recently been detected in US swine in less than one year of detecting PDCoV (Stevenson et al., 2013). However, from the present study and with available samples submitted to the ISU VDL at the present time, it can be inferred that PDCoV has been present in US swine at least since August 2013, suggesting that the PDCoV introduction may have been independent of PEDV introduction into the US. Although the origin of PDCoV remains unknown at this time, detection of the deltacoronavirus after PEDV may suggest the potential of additional trans-boundary or foreign animal diseases to enter the US in the future.

## Acknowledgements

This work was supported by National Animal Health Laboratory Network (NAHLN) and the Iowa State University Veterinary Diagnostic Laboratory. The sponsors had no role in designing, data collection, interpretation of the results, or writing of the manuscript. The authors have no potential conflicts of interest with respect to the research, authorship and/or publication of this

manuscript. The authors thank Wendy Stensland, Laura Bradner, Joe Thompson and Joseph Brodie for assistance with testing and providing the samples for this study. The authors also thank Brett Crim for his technical assistance.

## References

- Carstens, E.B., 2010. Ratification vote on taxonomic proposals to the International Committee on Taxonomy of Viruses (2009). *Arch. Virol.* 155, 133–146.
- King, A.M.Q., Adams M.J., Carstens E.B., Lefkowitz E.J., 2012. *Virus Taxonomy: Classification and Nomenclature of Viruses: Ninth Report of the International Committee on Taxonomy of Viruses.*
- Li, G., Chen, Q., Harmon, K.M., Yoon, K.J., Schwartz, K.J., Hoogland, M.J., Gauger, P.C., Main, R.G., Zhang, J., 2014. Full-length genome sequence of porcine deltacoronavirus strain USA/IA/2014/8734. *Genome Announc.* 2.
- Marthaler, D., Jiang, Y., Collins, J., Rossow, K., 2014. Complete genome sequence of strain SDCV/USA/Illinois121/2014, a porcine deltacoronavirus from the United States. *Genome Announc.* 2.
- Stevenson, G.W., Hoang, H., Schwartz, K.J., Burrough, E.R., Sun, D., Madson, D., Cooper, V.L., Pillatzki, A., Gauger, P., Schmitt, B.J., Koster, L.G., Killian, M.L., Yoon, K.J., 2013. Emergence of Porcine epidemic diarrhea virus in the United States: clinical signs, lesions, and viral genomic sequences. *J. Vet. Diagn. Invest.* 25, 649–654.
- Wang, L., Byrum, B., Zhang, Y., 2014. Detection and genetic characterization of deltacoronavirus in pigs, Ohio, USA, 2014. *Emerging Infect. Dis.* 20, 1227–1230.
- Woo, P.C., Lau, S.K., Lam, C.S., Lau, C.C., Tsang, A.K., Lau, J.H., Bai, R., Teng, J.L., Tsang, C. C., Wang, M., Zheng, B.J., Chan, K.H., Yuen, K.Y., 2012. Discovery of seven novel Mammalian and Avian coronaviruses in the genus deltacoronavirus supports bat coronaviruses as the gene source of alphacoronavirus and betacoronavirus and avian coronaviruses as the gene source of gammacoronavirus and deltacoronavirus. *J. Virol.* 86, 3995–4008.