

## Regulation of Global Transcription in *Escherichia coli* by Rsd and 6S RNA

Avantika Lal,<sup>\*1</sup> Sandeep Krishna,<sup>†</sup> and Aswin Sai Narain Seshasayee<sup>\*1</sup>

<sup>\*</sup>National Centre for Biological Sciences, and <sup>†</sup>Simons Centre for the Study of Living Machines, National Centre for Biological Sciences, Tata Institute of Fundamental Research, Bangalore 560065, Karnataka, India

ORCID ID: 0000-0002-5827-0826 (A.L.)

**ABSTRACT** In *Escherichia coli*, the sigma factor  $\sigma^{70}$  directs RNA polymerase to transcribe growth-related genes, while  $\sigma^{38}$  directs transcription of stress response genes during stationary phase. Two molecules hypothesized to regulate RNA polymerase are the protein Rsd, which binds to  $\sigma^{70}$ , and the non-coding 6S RNA which binds to the RNA polymerase- $\sigma^{70}$  holoenzyme. Despite multiple studies, the functions of Rsd and 6S RNA remain controversial. Here we use RNA-Seq in five phases of growth to elucidate their function on a genome-wide scale. We show that Rsd and 6S RNA facilitate  $\sigma^{38}$  activity throughout bacterial growth, while 6S RNA also regulates widely different genes depending upon growth phase. We discover novel interactions between 6S RNA and Rsd and show widespread expression changes in a strain lacking both regulators. Finally, we present a mathematical model of transcription which highlights the crosstalk between Rsd and 6S RNA as a crucial factor in controlling sigma factor competition and global gene expression.

### KEYWORDS

gene regulation  
sigma factors  
RNA polymerase  
6S RNA  
Rsd

In bacteria, the core RNA polymerase ( $\alpha_2\beta\beta'\omega$ , referred to as E) transcribes all cellular genes. Proteins called sigma ( $\sigma$ ) factors bind to E forming  $E\sigma$  holoenzymes, and direct it to transcribe from specific promoters. *Escherichia coli* has seven sigma factors (Ishihama 2000) –  $\sigma^{70}$  (RpoD),  $\sigma^{38}$  (RpoS),  $\sigma^{32}$ ,  $\sigma^{54}$ ,  $\sigma^{28}$ ,  $\sigma^{24}$  and  $\sigma^{19}$ . Though it was previously reported that the cellular concentration of E exceeded that of sigma factors (Jishage and Ishihama 1995; Maeda *et al.* 2000), recent quantitation has shown that sigma factors exceed E under common culture conditions (Grigorova *et al.* 2006; Piper *et al.* 2009), and may compete to bind to E. This is supported both by *in vitro* assays demonstrating competition between sigma factors under limiting E (Maeda *et al.* 2000; Jishage *et al.* 2002; Laurie *et al.* 2003; Bernardo *et al.* 2006) and *in vivo* studies where increasing activity of one sigma factor decreased the activity of others and vice-versa (Farewell *et al.* 1998; Jishage *et al.* 2002; Laurie *et al.* 2003).

$\sigma^{70}$  (RpoD) is the housekeeping sigma factor, directing transcription of genes essential for growth. The major alternative sigma factor,  $\sigma^{38}$  (RpoS), is present in low concentration during exponential growth, but regulates several hundred genes (Rahman *et al.* 2006; Dong *et al.* 2008). Upon entry into stationary phase, while the concentrations of E and  $\sigma^{70}$  show little change,  $\sigma^{38}$  accumulates (Gentry *et al.* 1993; Tanaka *et al.* 1993; Piper *et al.* 2009; Table 1) and directs transcription of genes for multiple stress tolerance (Lange and Hengge-Aronis 1991; Hengge-Aronis *et al.* 1991; Hengge-Aronis and Fischer 1992; Weber *et al.* 2005). However,  $\sigma^{70}$  remains the most abundant sigma factor, and has the highest affinity for E (Maeda *et al.* 2000; Colland *et al.* 2002; Piper *et al.* 2009; Ganguly and Chatterji 2012), implying that additional regulators are needed for  $\sigma^{38}$  to compete effectively with  $\sigma^{70}$ .

An example of such a regulator is the Crl protein, which binds to  $\sigma^{38}$  and increases its affinity for E, and promotes transcription by the  $E\sigma^{38}$  holoenzyme at some promoters (Bougourd *et al.* 2004; Typas *et al.* 2007; England *et al.* 2008). The nucleotide ppGpp also increases the ability of alternative sigma factors to compete with  $\sigma^{70}$  (Jishage *et al.* 2002). However, ppGpp is produced transiently on entry into stationary phase (Cavanagh *et al.* 2010), and Crl has been shown to decrease in extended stationary phase (Bougourd *et al.* 2004).

On the other hand, two regulators - the protein Rsd and the non-coding 6S RNA - act on  $\sigma^{70}$ . Rsd binds to  $\sigma^{70}$ , sequestering it from E, and inhibits  $E\sigma^{70}$ -dependent transcription at several promoters

Copyright © 2018 Lal *et al.*

doi: <https://doi.org/10.1534/g3.118.200265>

Manuscript received March 23, 2018; accepted for publication April 18, 2018; published Early Online April 26, 2018.

This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Supplemental material available at Figshare: <https://doi.org/10.25387/g3.6167501>.

<sup>1</sup>Corresponding authors: Department of Pathology, Stanford University, 300 Pasteur Drive, Stanford, CA 94305; E-mail: [avilal@stanford.edu](mailto:avilal@stanford.edu); [aswin@ncbs.res.in](mailto:aswin@ncbs.res.in)

■ Table 1 Cellular concentrations of RNA polymerase, sigma factors, Rsd and 6S RNA

Molecule	Exponential	Stationary	Reference
Core RNA Polymerase	~4.3 $\mu$ M	~4.3 $\mu$ M	Piper <i>et al.</i> 2009
$\sigma^{70}$	~12.1 $\mu$ M	~12.0 $\mu$ M	
$\sigma^{38}$	Not Detected	~2.7 $\mu$ M	
Rsd	~5.5 $\mu$ M	~10.4 $\mu$ M	
6S RNA	~1000 molecules/cell	~10,000 molecules/cell	Wassarman and Storz 2000

*in vitro* (Jishage and Ishihama 1998). An Rsd null strain showed increased transcription from a  $\sigma^{70}$  dependent promoter, and reduced transcription from a  $\sigma^{38}$  dependent promoter, in stationary phase, while Rsd overexpression had the opposite effect (Jishage and Ishihama 1999). It was hypothesized that in stationary phase, Rsd reduces E $\sigma^{70}$  formation, and, by freeing E to bind  $\sigma^{38}$ , increases E $\sigma^{38}$  formation. However, a microarray experiment found no significant change in gene expression on knocking out Rsd, and even comparison of the knockout with an Rsd overexpressing strain found changed expression of only a few  $\sigma^{38}$ -dependent genes (Mitchell *et al.* 2007). Also, effects of Rsd were observed only during stationary phase, supposedly due to low Rsd levels in exponential phase. However, Rsd was later observed to be present at ~50% of the level of  $\sigma^{70}$  in exponential phase and ~90% in stationary phase (Piper *et al.* 2009) – raising the question of why no change in expression was seen in its knockout.

6S RNA is a non-coding RNA expressed from the *ssrS* gene, which binds to the E $\sigma^{70}$  holoenzyme (Wassarman and Storz 2000). It has been shown to block E $\sigma^{70}$  binding to a target promoter (Wassarman and Saecker 2006) and inhibit transcription from several promoters *in vitro* and *in vivo* (Wassarman and Storz 2000; Trotochaud and Wassarman 2004; Gildehaus *et al.* 2007; Cavanagh *et al.* 2008; Cavanagh *et al.* 2010). A 6S RNA knockout showed increased expression from some E $\sigma^{70}$  promoters containing extended -10 elements, and reduced expression from a few E $\sigma^{38}$  promoters, in stationary phase (Trotochaud and Wassarman 2004). It was suggested that sequestration of E $\sigma^{70}$  by 6S RNA allows  $\sigma^{38}$  to compete more effectively for E, increasing transcription by E $\sigma^{38}$ . An alternative hypothesis was that 6S RNA regulates a *trans*-acting factor important for E $\sigma^{38}$  activity. There have been two microarray-based genome-wide studies on 6S RNA knockouts. While the first (Cavanagh *et al.* 2008) showed that 6S RNA regulated hundreds of genes in stationary phase, and an extended -10 element and weak -35 element could make a promoter sensitive to 6S RNA, the second (Neusser *et al.* 2010) found no correlation of 6S RNA sensitivity with promoter sequence or sigma factor preference. There was also little overlap between the regulated genes found in the two experiments.

Rsd and 6S RNA are present at high concentrations during growth, increase in stationary phase, and are thereafter at high levels (Jishage and Ishihama 1998; Wassarman and Storz 2000; Piper *et al.* 2009). Yet their regulatory function remains controversial and studies have been contradictory. Both are hypothesized to reduce E $\sigma^{70}$  and increase E $\sigma^{38}$  formation in stationary phase. However, it is not clear whether Rsd affects the expression of  $\sigma^{70}$  target genes, nor whether it has any effect during growth – and if not, why not? Similarly, with 6S RNA, while studies concur that it reduces transcription from several  $\sigma^{70}$  promoters, one study found that it regulated  $\sigma^{38}$  target genes during stationary phase whereas another did not. Does it have such an effect? If so, is it a direct effect of E $\sigma^{70}$  sequestration or mediated by a *trans*-acting factor? What is that *trans*-acting factor? Finally, although both Rsd and 6S RNA have

been proposed to help  $\sigma^{38}$  compete with  $\sigma^{70}$ , one sequesters  $\sigma^{70}$  while the other sequesters the E $\sigma^{70}$  holoenzyme. What impact does this have on their regulatory effects? Why does the cell require two molecules to sequester  $\sigma^{70}$ ?

Here, we present a genome-wide investigation of the functions of Rsd and 6S RNA in *E. coli*. We used RNA-seq to identify genes regulated by Rsd and 6S RNA in five phases of growth and demonstrated that both regulate global transcription during exponential as well as stationary phase. We showed that both increase expression of  $\sigma^{38}$  target genes, with 6S RNA also regulating hundreds of  $\sigma^{70}$  targets, varying greatly with growth phase. We found evidence of substantial crosstalk between Rsd and 6S RNA, with each regulating the other's expression and non-additive effects on over a thousand genes. Finally, we developed a mathematical model of sigma factor competition in *E. coli*, which explains our experimental results and highlights crosstalk between Rsd and 6S RNA as a vital factor controlling sigma factor competition.

## MATERIALS AND METHODS

### Growth conditions

Luria-Bertani broth and agar (20 g/L) were used for routine growth. M9 defined medium (0.6% Na<sub>2</sub>HPO<sub>4</sub>, 0.3% KH<sub>2</sub>PO<sub>4</sub>, 0.05% NaCl, 0.01% NH<sub>4</sub>Cl, 0.1 mM CaCl<sub>2</sub>, 1 mM MgSO<sub>4</sub>, 5 × 10<sup>-4</sup>% Thiamin) supplemented with 0.5% glucose and 0.1% casamino acids was used for RNA-seq and validation. During strain construction, ampicillin or kanamycin were used at final concentrations of 100  $\mu$ g/ml and 50  $\mu$ g/ml respectively.

### Strain construction

Single gene deletions were achieved by  $\lambda$  Red recombination (Datsenko and Wanner 2000), using plasmids pKD46 and pKD13 and specific primers. This introduced a kanamycin resistance cassette into the chromosome. Knockout strains were selected on LB Kanamycin plates. In the *rsd* knockout, the resistance cassette was removed by FLP-mediated site-specific recombination using plasmid pCP20. The  $\Delta$ *rsd* $\Delta$ *ssrS* double knockout was generated by P1 transduction from single knockouts (Thomason *et al.* 2007). The 3x-FLAG epitope was added to the C-terminus of Rsd by PCR using plasmid pSUB11 as template (Uzzau *et al.* 2001) and introduced onto the MG1655 chromosome by  $\lambda$  Red recombination using specific primers. The *ssrS* knockout was moved into this strain using P1 transduction. Strain constructions were verified by PCR using specific primers and Sanger sequencing. Primer sequences are given in Table S4.

### RNA extraction and mRNA enrichment

Overnight cultures in M9 glucose were inoculated in 100 mL fresh M9 glucose to a final OD<sub>600</sub> of 0.02 and incubated at 37° with shaking. Two biological replicates were performed for each strain. Cells were collected by centrifugation at the early exponential (OD<sub>600</sub> ~0.3), mid-exponential (OD<sub>600</sub> ~0.8), transition to stationary (OD<sub>600</sub> ~1.6), stationary

(16 hr, OD<sub>600</sub> ~2), and late stationary (48 hr, OD<sub>600</sub> ~1.6) phases of growth. RNA was extracted using TRIzol (Invitrogen), following the manufacturer's protocol. Total RNA was treated with DNase I (Invitrogen, 18068-015) according to the manufacturer's protocol. Further precipitation of RNA and ribosomal RNA cleanup was achieved using the MICROBExpress bacterial mRNA purification Kit (Ambion, AM1905) according to the manufacturer's protocol. RNA concentration was determined using a Nanodrop 2000 (Thermo Scientific) and quality was checked by visualization on agarose gels.

### RNA-Seq

Sequencing libraries were prepared using TruSeq RNA sample preparation kit v2 (Illumina, RS-122-2001) according to the manufacturer's guidelines, checked for quality on an Agilent 2100 Bioanalyzer, and sequenced for 50 cycles from one end on an Illumina HiSeq1000 platform at the Centre for Cellular and Molecular Platforms, Bangalore. RNA-Seq data are summarized in Table S2. RNA-Seq reads were mapped to the *E. coli* K-12 MG1655 genome using BWA (Li and Durbin 2009).

### qRT-PCR for RNA-Seq validation

qRT-PCR was carried out using specific primers to selected mRNA targets (primers in Table S4, results in Table S3). 5 ng RNA was used for each RT-PCR reaction. TAKARA One-step SYBR PrimeScript RT-PCR kit II (RR086A) was used according to the manufacturer's protocol, on an Applied Biosystems ViiA 7 Real-Time PCR system.

### Western Blotting

Cells were grown as for RNA-seq. For stationary phase samples, 5 ml culture was harvested by centrifugation. For mid-exponential phase, 10 ml was harvested. Lysates were prepared and protein concentration was estimated using BCA assay (Thermo Fisher Scientific, 23227). Lysates containing equal amounts of protein were loaded onto an SDS-PAGE gel. Proteins were electroblotted onto a nitrocellulose membrane and probed with mouse primary antibody against the protein of interest followed by horseradish peroxidase-conjugated anti-mouse secondary antibody. The primary antibodies were: Mouse monoclonal antibody to RpoB (Neoclone, WP023), Mouse monoclonal antibody to  $\sigma^{70}$  (Neoclone, WP004), Mouse monoclonal antibody to  $\sigma^{38}$  (Neoclone, WP009), Mouse monoclonal anti-FLAG antibody (Sigma-Aldrich, F3165), Mouse monoclonal antibody to GroEL (Abcam, ab82592). Bands were visualized using SuperSignal West Dura Chemiluminescent Substrate (Thermo Fisher Scientific, 34076) and imaged using an ImageQuant LAS 4000 system (GE Healthcare Life Sciences). Band intensities were quantified using ImageJ (<http://imagej.nih.gov/ij>).

### Data sources

The *E. coli* K-12 MG1655 genome was downloaded from NCBI (NC\_000913.2). Gene coordinates were taken from RegulonDB v8.0 (Salgado *et al.* 2013). Lists of  $\sigma^{38}$  target genes and ppGpp regulated genes were obtained from (Weber *et al.* 2005) and (Traxler *et al.* 2008) respectively. Lists of genes regulated by 6S RNA were obtained from (Cavanagh *et al.* 2008) and (Neusser *et al.* 2010). Coordinates of RNA polymerase binding regions and their occupancy were obtained from (Cho *et al.* 2014). A list of 501 genes under constitutive  $\sigma^{70}$  promoters was obtained from (Shimada *et al.* 2014). Of these, we selected 270 genes which were not regulated by  $\sigma^{38}$  or other sigma factors according to RegulonDB (Salgado *et al.* 2013), (Weber *et al.* 2005), or our RNA-Seq from the  $\sigma^{38}$  knockout.




### Data Analysis

RNA-Seq reads were mapped to the *E. coli* K-12 MG1655 genome using BWA<sup>52</sup>, and reads mapping uniquely were used for further analysis. DESeq (Anders and Huber 2010) was used for differential expression analysis. Fold changes and p-values for all genes are given in the Supplementary Dataset. Genes with  $\leq 10$  reads mapping to them under all conditions were excluded from all analyses and plots. All statistical analyses were performed in R version 3.0.1.

### Mathematical model

The reactions in Figure 6A are represented by a set of differential equations that determine how the dynamical variables (levels of sigma factors, holoenzymes, etc.) change with time. These equations are given in File S2. Since the timescales on which these reactions occur is much faster than typical timescales of cell division, or processes such as stress responses in stationary phase, we assume that the levels of sigma factors, holoenzymes, etc., are in quasi steady-state in the cell. Parameters that determine the specific steady-state include the dissociation constants of the various complexes, total levels of Rsd, 6S RNA, E, sigma factors, and some others, listed in Table 2 with their default values. These values are altered in specific simulations as described in the results.

The model of holoenzyme formation (shaded in Figure 6A) describes the following reactions:



In steady state, the following equations must be fulfilled:

$$\frac{[E][\sigma^{70}]}{[E\sigma^{70}]} = K_{E\sigma^{70}} \quad (1)$$

$$\frac{[E][\sigma^{38}]}{[E\sigma^{38}]} = K_{E\sigma^{38}} \quad (2)$$

$$\frac{[Rsd][\sigma^{70}]}{[Rsd - \sigma^{70}]} = K_{Rsd} \quad (3)$$

$$\frac{[6SRNA][E\sigma^{70}]}{[6SRNA - E\sigma^{70}]} = K_{6S} \quad (4)$$

$$E_{total} = [E] + [E\sigma^{70}] + [E\sigma^{38}] + [6SRNA - E\sigma^{70}] \quad (5)$$

$$\sigma^{70}_{total} = [\sigma^{70}] + [E\sigma^{70}] + [Rsd - \sigma^{70}] + [6SRNA - E\sigma^{70}] \quad (6)$$

$$\sigma^{38}_{total} = [\sigma^{38}] + [E\sigma^{38}] \quad (7)$$

$$Rsd_{total} = [Rsd] + [Rsd - \sigma^{70}] \quad (8)$$








$$6SRNA_{total} = [6SRNA] + [6SRNA - E\sigma^{70}] \quad (9)$$

Steady-state levels of the dynamical variables were obtained from the above equations both by solving them numerically and by integrating the corresponding differential equations until they reached steady-state.

■ **Table 2** Default values of parameters used for simulations. Wherever possible, values are specific to stationary phase. These values are discussed in File S1

Parameter	Meaning	Value	Reference
Volume	Average cell volume	10 <sup>-15</sup> L	(Ali Azam <i>et al.</i> 1999)
E <sub>total</sub>	Total cellular concentration of RNA Polymerase	4.3 μM	(Piper <i>et al.</i> 2009)
σ <sup>70</sup> <sub>total</sub>	Total cellular concentration of σ <sup>70</sup>	12.0 μM	
σ <sup>38</sup> <sub>total</sub>	Total cellular concentration of σ <sup>38</sup>	2.7 μM	
Rsd <sub>total</sub>	Total cellular concentration of Rsd	10.4 μM	
6S RNA <sub>total</sub>	Total cellular concentration of 6S RNA	13 μM	(Wassarman and Storz 2000)
DNA <sub>total</sub>	Nonspecific binding sites per cell	4.6 × 10 <sup>6</sup>	MG1655 genome size
P <sub>70</sub> , P <sub>38</sub>	σ-cognate promoters per cell	200	(Mauri and Klumpp 2014)
K <sub>Eσ<sup>70</sup></sub>	Dissociation constant for E - σ <sup>70</sup> binding	3.3 nM	(Colland <i>et al.</i> 2002)
K <sub>Eσ<sup>38</sup></sub>	Dissociation constant for E - σ <sup>38</sup> binding	15.2 nM	
K <sub>NS</sub>	Dissociation constant for nonspecific binding of RNA Polymerase and DNA	10 <sup>-4</sup> M	(deHaseth <i>et al.</i> 1978)
K <sub>Rsd</sub>	Dissociation constant for Rsd - σ <sup>70</sup> binding	32 nM	(Sharma and Chatterji 2008)
K <sub>6S</sub>	Dissociation constant for 6S RNA - Eσ <sup>70</sup> binding	131 nM	(Beckmann <i>et al.</i> 2011)
K <sub>EσP</sub>	Dissociation constant for holoenzyme promoter binding	10 <sup>-7</sup> M	(deHaseth <i>et al.</i> 1998); (Bintu <i>et al.</i> 2005)
Operon length	Average operon length	1000 nt	RegulonDB (Salgado <i>et al.</i> 2013)
c	Rate of promoter clearance	0.005 s <sup>-1</sup>	(Smith and Sauer 1996); (Mauri and Klumpp 2014)
e	Rate of escape from elongation	0.021 s <sup>-1</sup>	(Proshkin <i>et al.</i> 2010)

The model with DNA (represented by the full schematic in Figure 6A) includes, in addition, the following reactions:



*c* represents the rate of promoter clearance and *e* represents the rate of transcript elongation. At steady state these fulfill the following equations, in addition to equations (1) - (4) and (8) - (9). *K<sub>NS</sub>* represents the dissociation constant for non-specific binding of RNA polymerase to DNA, which we assume is equal for E, Eσ<sup>70</sup> and Eσ<sup>38</sup>. We also assume the specific binding constants of Eσ<sup>70</sup> and Eσ<sup>38</sup> to their target promoters to be equal in stationary phase.

$$\frac{[E][DNA]}{[EDNA]} = \frac{[E\sigma^{70}][DNA]}{[E\sigma^{70}DNA]} = \frac{[E\sigma^{38}][DNA]}{[E\sigma^{38}DNA]} = K_{NS} \quad (10)$$

$$\frac{[E\sigma^{70}][P_{70}]}{[E\sigma^{70}P_{70}]} = \frac{[E\sigma^{38}][P_{38}]}{[E\sigma^{38}P_{38}]} = K_{E\sigma P} \quad (11)$$

$$E_{total} = [E] + [E\sigma^{70}] + [E\sigma^{38}] + [6SRNA - E\sigma^{70}] + [EDNA] + [E\sigma^{70}DNA] + [E\sigma^{38}DNA] + [E\sigma^{70}P_{70}] + [E\sigma^{38}P_{38}] + [E_{e70}] + [E_{e38}] \quad (12)$$

$$\sigma^{70}_{total} = [\sigma^{70}] + [E\sigma^{70}] + [Rsd - \sigma^{70}] + [6SRNA - E\sigma^{70}] + [E\sigma^{70}DNA] + [E\sigma^{70}P_{70}] \quad (13)$$

$$\sigma^{38}_{total} = [\sigma^{38}] + [E\sigma^{38}] + [E\sigma^{38}DNA] + [E\sigma^{38}P_{38}] \quad (14)$$

$$DNA_{total} = [DNA] + [EDNA] + [E\sigma^{70}DNA] + [E\sigma^{38}DNA] \quad (15)$$

$$P_{70total} = [P_{70}] + [E\sigma^{70}P_{70}] \quad (16)$$

$$P_{38total} = [P_{38}] + [E\sigma^{38}P_{38}] \quad (17)$$

Here, steady-state levels were obtained by integrating the corresponding differential equations until they reached steady-state.

### Data availability

RNA-Seq data are deposited with NCBI GEO under the accession number GSE74809. Supplementary material includes Figures S1-S12, Tables S1-S4, and Files S1-S3. Supplemental material available at Figshare: <https://doi.org/10.25387/g3.6167501>.


## RESULTS

### RNA-seq to identify genes regulated by Rsd and 6S RNA

Figure 1A shows a schematic of the binding activity of 6S RNA and Rsd. Rsd sequesters σ<sup>70</sup> and prevents it from binding to core RNA polymerase (E), while 6S RNA binds to the Eσ<sup>70</sup> holoenzyme and prevents it from binding promoters. To identify their effects on gene expression, we carried out RNA-seq to identify genes regulated by Rsd and / or 6S RNA in five growth phases.

Table S1 lists strains and plasmids used. Five strains: *E. coli* K-12 MG1655 (Wild-type), Rsd knockout (*Δrsd*), 6S RNA knockout (*ΔssrS*), 6S RNA-Rsd double knockout (*ΔrsdΔssrS*), and σ<sup>38</sup>/RpoS knockout (*ΔrpoS*) were used for RNA-seq. These had similar growth rates in M9 glucose (Figure S1). RNA-seq was performed in five growth phases:





**Figure 1** Rsd regulates RpoS target genes throughout growth. (A) Schematic showing the binding activity of Rsd and 6S RNA. (B) Boxplot of log<sub>2</sub> fold change in gene expression ( $\Delta rsd/wt$ ) for 313 genes whose expression is reduced at least twofold in  $\Delta rpoS/wild$ -type in EE phase, compared to all other genes. (C) Boxplot of log<sub>2</sub> fold change in gene expression ( $\Delta rsd/wild$ -type, stationary phase) for 634 genes whose expression is reduced at least twofold in  $\Delta rpoS/wild$ -type in stationary phase, compared to all other genes. p-values are for Wilcoxon Test.

early exponential (EE), mid-exponential (ME), transition to stationary (TS), stationary (S), and late stationary (LS; time points in Methods).

### Rsd increases $\sigma^{38}$ /RpoS activity throughout growth

We defined differentially expressed genes as genes whose expression changed  $\geq$ twofold in a mutant strain relative to wild-type, with FDR-adjusted p-value  $< 0.05$ . Using these criteria, the  $\Delta rsd$  strain showed only 16 differentially expressed genes. These included several non-coding RNAs (*ryfD*, *sokA*, *oxyS*, *sroH*, *sibD*) which were increased two- to sixfold in  $\Delta rsd$  during stationary phase. Notably, 6S RNA expression was altered in  $\Delta rsd$ ; 6S RNA was 2.3 times its wild-type level in stationary phase, but in ME phase was reduced to about half its wild-type level.

As very few genes were differentially expressed in  $\Delta rsd$ , we looked for smaller changes. We found that in all growth phases, genes whose expression was significantly reduced ( $\geq$ twofold,  $P < 0.05$ ) in  $\Delta rpoS$  also displayed reduced expression in  $\Delta rsd$ . The box plots in Figure 1B-C show the distribution of log<sub>2</sub> fold change in gene expression in  $\Delta rsd$  relative to wild-type.  $\sim 75\%$  of genes whose expression was significantly reduced in  $\Delta rpoS$  showed reduced expression (log<sub>2</sub> fold change  $< 0$ ) in  $\Delta rsd$ . Similar plots for other growth phases are in Figure S2A. This reduced expression was less than twofold in magnitude and so was not seen when searching for differentially expressed genes.

Though these fold changes are small (the median decrease in expression is 10%), we consider them important for several reasons. First, they represent a consistent and highly significant (Wilcoxon test  $P < 10^{-15}$  in all growth phases except LS) decrease in expression across hundreds of *rpoS* regulated genes, in all five growth phases. Second, the average overlap between the set of *rpoS* regulated genes in successive growth phases is only 54% - so it is not a single set of genes whose expression is reduced in  $\Delta rsd$ , but a different set in each growth phase. Third, the  $\Delta rsd$  samples showed high inter-replicate correlation in all growth phases (Table S2). Fourth, this trend of decreased expression held true when only previously reported  $\sigma^{38}$  targets (Weber *et al.* 2005) were considered (Figure S2B). Fifth, genes whose expression was increased  $\geq$ twofold in  $\Delta rpoS$  also showed increased expression in  $\Delta rsd$ , during EE, ME and stationary phases (Figure S2C). On the other hand, expression of genes controlled by constitutive  $\sigma^{70}$  target promoters (Shimada *et al.* 2014) was not substantially altered in any phase (Figure S3).

Thus, the Rsd knockout behaved like a  $\sigma^{38}$  knockout, only with smaller changes in gene expression. Western blots in stationary phase

showed that  $\sigma^{38}$  protein levels were similar in  $\Delta rsd$  and wild-type (Figure S4;  $\sigma^{38}$  level in  $\Delta rsd$  is 94% of wild-type); however, it is extremely difficult to obtain reliable measures of small differences in protein level using western blots. We cannot be certain based on this experiment whether the reduced expression of  $\sigma^{38}$  targets in  $\Delta rsd$  is due to reduced  $\sigma^{38}$  protein, reduced E- $\sigma^{38}$  association, or both. We attempt to address this question theoretically later.

### 6S RNA regulates distinct sets of genes in all phases of growth


What role does the E $\sigma^{70}$  - sequestering 6S RNA play in gene regulation? Is its function similar to that of the  $\sigma^{70}$ -sequestering Rsd? Our RNA-seq showed that the 6S RNA knockout ( $\Delta ssrS$ ) was very different from the Rsd knockout. To begin with, it showed  $\geq$ twofold differential expression of 447 genes in total.

Figures 2A and B show that there was very little overlap between genes differentially expressed in  $\Delta ssrS$  in successive growth phases, even in the two stationary phase time points. Only a few genes were differentially expressed in  $\Delta ssrS$  throughout growth, particularly *fau* (*ygfA*), which is downstream of 6S RNA in the same operon. *fau* expression was increased when 6S RNA was overexpressed in a wild-type background (Figure S5), indicating that this is at least partly a regulatory effect of 6S RNA and not merely a polar effect.

During exponential phase, expression of several genes for amino acid transporters (*artM*, *artI*, *hisP*, *hisQ*, *hisJ*, *hisM*, *tdcC*) and amino acid biosynthesis (*argH*, *argB*, *thrA*, *thrB*, *thrC*, *asnB*, *glyA*) was increased in  $\Delta ssrS$ , while expression of genes involved in stress responses (*rmf*, *appY*, *yadC*, *ybcM*, *yciF*, *gadW*, *ydeI*, *yodD*, *dps*, *hdeA*, *hslV*, *oppA*, *osmE*, *dosC*) was reduced. On the other hand, in LS phase, genes involved in global transcriptional regulation (*crp*, *hha*) and the TCA cycle (*sdhD*, *sdhC*, *gltA*, *aceB*, *ppc*, *sucA*) were upregulated. Thus, the physiological function of 6S RNA appears largely growth phase-dependent; nevertheless, there are certain patterns throughout growth.

### 6S RNA increases $\sigma^{38}$ /RpoS activity throughout growth

Like  $\Delta rsd$ , the  $\Delta ssrS$  strain showed reduced expression of  $\sigma^{38}$  target genes (Figures 3A and B, Figure S6, Wilcoxon test  $P < 10^{-15}$  in all growth phases except LS). This was not due to reduced  $\sigma^{38}$  protein, as western blots in stationary phase showed that the  $\Delta ssrS$  strain had higher  $\sigma^{38}$  protein than the wild-type (Figure S4). On the other hand, expression of genes under constitutive  $\sigma^{70}$  promoters was slightly increased in the ME, TS and LS phases (Figure S7).



**Figure 2** The 6S RNA regulon is highly growth-phase dependent. (A) Venn diagram showing genes whose expression is increased at least twofold in  $\Delta ssrS$ , in different phases of growth. (B) Venn diagram showing genes whose expression is reduced at least twofold in  $\Delta ssrS$ , in different phases of growth.

### 6S RNA regulates RpoB, Crl and Rsd

Since the wild-type expression of 6S RNA increased in successive growth phases (Figure 4A), its effect should increase with growth phase. Indeed, with the exception of the TS phase, the number of 6S RNA regulated genes increased with growth phase. A previous study (Cavanagh *et al.* 2010) showed increased ppGpp in a 6S RNA knockout during the TS phase, and we observed consistent changes in gene expression (Figure S8). Since ppGpp also favors competition of alternative sigma factors with  $\sigma^{70}$  (Jishage *et al.* 2002), increased ppGpp may reduce the effects of  $\Delta ssrS$  deletion during this phase.

We found 36 genes where 6S RNA had a dose-dependent effect throughout growth, increasing in each successive growth phase (Figure S9). For instance, expression of *rpoB* (encoding the  $\beta$  subunit of core RNA polymerase) was reduced in  $\Delta ssrS$ , and the magnitude of reduction increased with growth phase (Figure 4B). Reduced RpoB was validated in stationary phase by qRT-PCR (Table S3) and western blotting (Figure S10). Since excess  $E\sigma^{70}$  inhibits *rpoB* transcription (Fukuda *et al.* 1978), we suggest that deleting 6S RNA leads to higher free  $E\sigma^{70}$ , which proportionally represses *rpoB*. Since RpoB is limiting for the formation of core RNA polymerase (Piper *et al.* 2009), this implies that the cell compensates for 6S RNA loss by reducing RNA polymerase synthesis. Similarly, we observed decreased expression of genes encoding ribosomal proteins (Figure 4C), consistent with previous reports in stationary phase (Neusser *et al.* 2010).

6S RNA also represses some genes in a dose-dependent manner. Figure 4D shows that *crl* expression is increased in  $\Delta ssrS$ , and the magnitude of this effect increases with growth phase. As with the effect of reducing RNA polymerase synthesis to compensate for the loss of 6S RNA, increasing  $\sigma^{38}$  protein and Crl may be a means to compensate for reduced  $E\sigma^{38}$  activity in  $\Delta ssrS$  bacteria.

Since we observed that Rsd regulated 6S RNA expression, we checked whether 6S RNA in turn regulated Rsd. Our RNA-Seq showed that *rsd* was not differentially expressed in  $\Delta ssrS$ . To check if 6S RNA regulated Rsd post-transcriptionally, we added a 3xFLAG tag to the C-terminal of the Rsd protein, as in previous studies (Piper *et al.* 2009). Indeed, western blots showed that 3xFLAG-tagged Rsd was reduced in the  $\Delta ssrS$  background, in both ME and stationary phases (Figure 4E).


Finally, though 6S RNA binds to  $E\sigma^{70}$ , its effects are highly promoter-specific (Trotochaud and Wassarman 2004; Cavanagh *et al.* 2008; Cavanagh *et al.* 2010; Neusser *et al.* 2010). We did not observe a link between 6S RNA sensitivity and an extended -10 or weak -35 promoter sequence as reported in (Cavanagh *et al.* 2008) (Figure S11), nor with any other sequence features. Instead, during stationary phase, 6S RNA sensitivity was associated with low wild-type expression (Figure 4F) and low promoter occupancy by RNA polymerase (Figure 4G, based on ChIP-chip data from (Cho *et al.* 2014)). Thus, our data

support a model in which sequestration of RNA polymerase by 6S RNA primarily represses promoters that bind weakly to RNA polymerase.


### The Rsd/6S RNA double knockout shows differential expression of a distinct set of genes

We have discovered several instances of crosstalk between 6S RNA and Rsd; apart from the fact that both sequester  $\sigma^{70}$  in different forms, each regulates the other's expression, and both favor the activity of  $\sigma^{38}$ . We therefore asked whether the double knockout of Rsd and 6S RNA showed effects distinct from the single knockouts.

We observed that in every growth phase, hundreds of genes were differentially expressed in the double knockout relative to the wild-type. This far exceeded the number of differentially expressed genes in both single knockouts, demonstrating significant crosstalk between the two regulators. Figures 5A and B highlight genes that showed differential



**Figure 3** 6S RNA regulates RpoS target genes throughout growth. (A) Boxplot of log<sub>2</sub> fold change in gene expression ( $\Delta ssrS$ /wild-type, EE phase) for 313 genes whose expression is reduced at least twofold in  $\Delta rpoS$ /wild-type in EE phase, compared to all other genes. (B) Boxplot of log<sub>2</sub> fold change in gene expression ( $\Delta ssrS$ /wild-type, stationary phase) for 634 genes whose expression is reduced at least twofold in  $\Delta rpoS$ /wild-type in stationary phase, compared to all other genes. p-values are for Wilcoxon Test.



**Figure 4** 6S RNA controls RpoB, Crl and Rsd. (A) log<sub>2</sub> fold increase in 6S RNA levels in wild-type *E. coli* over successive growth phases (relative to its expression in EE phase), based on RNA-Seq. (B) log<sub>2</sub> fold change in *rpoB* expression in  $\Delta ssrS$ /wild-type, in successive growth phases. (C) log<sub>2</sub> fold change in the expression of 46 ribosomal protein genes in  $\Delta ssrS$ /wild-type, in successive growth phases. (D) log<sub>2</sub> fold change in *crl* expression in  $\Delta ssrS$ /wild-type, in successive growth phases. (E) Western blot for 3xFLAG-tagged Rsd in wild-type and  $\Delta ssrS$  backgrounds, in mid-exponential and stationary phases. (F) Boxplots showing the stationary phase wild-type expression level of genes that are reduced  $\geq$ twofold in  $\Delta ssrS$ /wild-type, genes that are not differentially expressed in  $\Delta ssrS$ , and genes that are increased  $\geq$ twofold in  $\Delta ssrS$ /wild-type. (G) Boxplots showing the stationary phase RNA polymerase occupancy (by ChIP-chip) of promoters belonging to genes that are reduced  $\geq$ twofold in  $\Delta ssrS$ /wild-type, genes that are not differentially expressed in  $\Delta ssrS$ , and genes that are increased  $\geq$ twofold in  $\Delta ssrS$ /wild-type. Only genes that were first in their transcription unit and were associated with a single RNA polymerase binding site, were included in (G). All p-values are for Wilcoxon Test.

expression in the double knockout, but less than twofold change in both single knockouts added together; there were 1780 such genes in total. These included genes encoding DNA Gyrase and Topoisomerase I, which maintain DNA supercoiling and regulate gene expression (Peter *et al.* 2004), the nucleoid-associated proteins HU, H-NS and StpA, the global transcriptional regulators ArcA, LRP and IHF, and the small RNA chaperone Hfq. This suggests that Rsd and 6S RNA act together, in a partially redundant manner, to control global gene expression and nucleoid structure.


### A theoretical model suggests an explanation for the behavior of Rsd and 6S RNA

Our RNA-Seq demonstrated that Rsd and 6S RNA regulate sigma factor competition and gene expression at a global scale; however, there are two major results that appear counter-intuitive. First, how is it that Rsd, which sequesters  $\sigma^{70}$ , increases transcription of  $\sigma^{38}$  target genes but has very little effect on  $\sigma^{70}$  target genes? Second, how does 6S RNA, which sequesters not only  $\sigma^{70}$ , but also core RNA polymerase which is required for transcription by all sigma factors, nevertheless increase transcription by  $\sigma^{38}$ ?

To answer these questions, we constructed a mathematical model of transcription during stationary phase, using parameters from literature (Table 2). We focused on stationary phase as that is when  $\sigma^{38}$ , 6S RNA and Rsd are at high concentrations. Our model is similar in structure to previous studies (Grigorova *et al.* 2006; Mauri and Klumpp 2014); however, it is the first to model stationary phase and to include both 6S RNA and Rsd. Consequently, our results differ from previous models.


Figure 6A shows a schematic of the model. Core RNA polymerase (E) binds to sigma factors ( $\sigma^{70}$  and  $\sigma^{38}$ ) forming holoenzymes ( $E\sigma^{70}$  and  $E\sigma^{38}$ ). Holoenzymes initiate transcription from target promoters ( $P_{70}$  and  $P_{38}$ ), releasing the sigma factor. The elongating RNA polymerase ( $E_{e70}$  and  $E_{e38}$ ) transcribes until released. Holoenzymes and E also bind DNA non-specifically. We focus on the steady state of this model, determined by equations (1) - (4) and (8) - (17) in Methods.

To understand how Rsd and 6S RNA regulate competition between sigma factors, we initially modeled the formation of  $E\sigma^{70}$  and  $E\sigma^{38}$  in the absence of DNA. This is represented by the shaded area in Figure 6A and steady-state equations (1) - (9) in Methods. Figure 6B(i) depicts what happens when Rsd is added to a system containing only E,  $\sigma^{70}$  and



**Figure 5** Rsd and 6S RNA have widespread non-additive effects on gene expression. Scatterplots of log<sub>2</sub> fold change in gene expression in the  $\Delta rsd\Delta ssrS$  double knockout vs. the sum of log<sub>2</sub> fold changes in the  $\Delta rsd$  and  $\Delta ssrS$  single knockouts, for each gene, in (A) Early exponential and (B) Stationary phase. Blue points represent genes that show differential expression ( $\geq$ twofold increase or decrease, FDR-adjusted p-value  $<$  0.05) in the double knockout, but less than twofold change in expression in both single knockouts added together.





**Figure 6** A mathematical model of Rsd and 6S RNA activity. (A) Schematic of the model. Shaded area represents reactions involved in holoenzyme formation (without DNA). (B) (i) Steady-state levels of  $E\sigma^{70}$  (black) and  $E\sigma^{38}$  (gray), computed from equations (1) - (9), as a function of total Rsd, when 6S RNA = 0. (ii) Same when total 6S RNA = 13  $\mu\text{M}$ . (iii) Steady-state levels of the 6S RNA- $E\sigma^{70}$  complex, as a function of total Rsd. (iv) Steady-state rate of transcription from  $E\sigma^{70}$  dependent promoters (black) and  $E\sigma^{38}$  dependent promoters (gray), as a function of total Rsd. Vertical dashed lines represent wild-type cellular concentrations in stationary phase. (v) Steady-state rate of transcription from  $E\sigma^{70}$  dependent promoters (black bars) and  $E\sigma^{38}$  dependent promoters (gray bars) in the wild-type, absence of Rsd, and simulated Rsd knockout. (C) (i) Steady-state rate of transcription from  $E\sigma^{70}$  dependent promoters (black) and  $E\sigma^{38}$  dependent promoters (gray), as a function of total 6S RNA. Vertical dashed lines represent wild-type cellular concentrations in stationary phase. (ii) Steady-state rate of transcription from  $E\sigma^{70}$  dependent promoters (black bars) and  $E\sigma^{38}$  dependent promoters (gray bars) in the wild-type, absence of 6S RNA, and simulated 6S RNA knockout.

$\sigma^{38}$ . Each value of Rsd on the x-axis corresponds to a separate 'run' of the model where we compute the steady-state for those parameter values, before moving on to the next Rsd value. We predict that as Rsd is increased, it sequesters  $\sigma^{70}$ , reducing  $E\sigma^{70}$  formation. This allows more E to bind  $\sigma^{38}$ , increasing the formation of  $E\sigma^{38}$ , consistent with previous predictions of Rsd function (Jishage and Ishihama 1999).

However, Figure 6B(ii) predicts that when 6S RNA is also present, Rsd increases  $E\sigma^{38}$  with little effect on  $E\sigma^{70}$ . How? This paradoxical result can be understood with Figure 6B(iii), which shows that the concentration of the 6S RNA- $E\sigma^{70}$  complex decreases as Rsd is increased. When 6S RNA is present, increasing Rsd still reduces E- $\sigma^{70}$  association; however, the reduction in  $E\sigma^{70}$  is compensated for by the release of  $E\sigma^{70}$  from its complex with 6S RNA, and so there is little change in the overall  $E\sigma^{70}$  level. This prediction remains when we include DNA in the model (represented by the complete schematic in Figure 6A). Figure 6B(iv) shows that simulating increased Rsd increases transcription by  $E\sigma^{38}$  with less effect on  $E\sigma^{70}$  transcription.

Our  $\Delta rsd$  strain also displayed a  $\sim 2.3$ -fold increase in 6S RNA during stationary phase. Figure 6B(v) shows the predicted rate of transcription from  $E\sigma^{70}$  and  $E\sigma^{38}$  target promoters in the wild-type and when Rsd = 0. The third pair of bars is an approximation of conditions in the  $\Delta rsd$  strain, where 6S RNA is increased 2.3-fold. We see that

increased expression of 6S RNA could reduce  $E\sigma^{70}$  dependent transcription in the  $\Delta rsd$  strain to almost the wild-type level, so that the main observable effect of knocking out Rsd would be reduced transcription by  $E\sigma^{38}$ .

This explains why, in both our RNA-seq and a previous study (Mitchell *et al.* 2007), Rsd increases transcription of  $E\sigma^{38}$  target genes, with little to no effect on  $E\sigma^{70}$ . We hypothesize that high levels of 6S RNA cause Rsd to act as a  $\sigma^{38}$  regulator and not a  $\sigma^{70}$  regulator, and, moreover, that Rsd controls the level of 6S RNA to modulate its own function.

We also observed a 2.4-fold increase in *oxyS* (a repressor of *rpoS* translation) in the Rsd knockout. We did not include this in the model due to lack of quantitative data on *oxyS*; however, it is possible that a small reduction in  $\sigma^{38}$  protein levels, perhaps due to increased *oxyS*, may contribute to reduced  $\sigma^{38}$  activity in the  $\Delta rsd$  strain. However, it is difficult to confirm this.

What is the effect of 6S RNA? Our model predicts that as 6S RNA levels increase, it sequesters  $E\sigma^{70}$ , reducing the amount of E available to bind to both sigma factors and thus inhibiting formation of both holoenzymes. This decreases the rate of transcription by both holoenzymes (Figure 6C(i)). However, our RNA-seq showed that deleting 6S RNA actually results in reduced transcription of  $E\sigma^{38}$



target genes, *i.e.*, 6S RNA increases transcription by  $E\sigma^{38}$ . This is also supported by single-promoter studies in (Trotochaud and Wassarman 2004). How is this possible? A 10-fold increase or decrease in any of the parameters was not sufficient to reproduce this observation.

Apart from losing 6S RNA, our  $\Delta ssrS$  strain also displayed reduced Rsd and RpoB, and increased  $\sigma^{38}$ . Figure 6C(ii) shows the predicted rate of transcription from  $E\sigma^{70}$  and  $E\sigma^{38}$  target promoters in the wild-type and when 6S RNA = 0. The third pair of bars is an approximation of conditions in the  $\Delta ssrS$  strain; Rsd and E are reduced by 50% and  $\sigma^{38}$  is increased by 50%.  $E\sigma^{38}$  transcription is now reduced to less than the wild-type level; in fact, within the default parameters of our model, reducing Rsd alone from 10.4  $\mu\text{M}$  to 8  $\mu\text{M}$  is sufficient to lower  $E\sigma^{38}$  transcription in the 6S RNA knockout below its wild-type level.

We also observed increased *crl* mRNA in  $\Delta ssrS$ . We have not modeled this due to lack of quantitative data on Crl. However, if the change in mRNA corresponds to increased Crl protein, it could partially mitigate the effect of reduced Rsd in the  $\Delta ssrS$  strain.

Thus, from our model we can make the strong claim that the binding reactions of Rsd and 6S RNA shown in Figure 6A cannot explain the reduced transcription of  $\sigma^{38}$  target genes in the  $\Delta ssrS$  strain. However, adding reduced Rsd levels to the model is sufficient. We therefore hypothesize that the reduced  $E\sigma^{38}$  transcription in the 6S RNA knockout is due to its indirect effects primarily via Rsd.

The quantitative predictions of this model would ideally be validated in detail by *in vitro* transcription experiments, measuring the change in transcription of  $\sigma^{38}$  and  $\sigma^{70}$  dependent promoters with Rsd and 6S RNA in varying concentrations. We predict that the effect of Rsd on  $\sigma^{70}$  dependent transcription should decrease as 6S RNA is increased. We also predict that increasing 6S RNA alone should reduce both  $\sigma^{38}$  and  $\sigma^{70}$  dependent transcription, but  $\sigma^{38}$  dependent transcription can be increased on increasing Rsd. To our knowledge, such studies have not been carried out. However, such experiments are necessarily limited to a few promoters. Some predictions may also be testable by looking at distributions of gene expression *in vivo*; for instance, increasing Rsd expression in  $\Delta ssrS$  back to the wild-type level should largely mitigate the reduced expression of  $\sigma^{38}$  target genes.

## DISCUSSION

Rsd and 6S RNA have long been known to regulate RNA polymerase in *E. coli*. Here we report for the first time that Rsd regulates gene expression from early exponential to stationary phase. Though it sequesters  $\sigma^{70}$ , its primary function is reducing transcription by the alternative sigma factor  $\sigma^{38}$ . Based on theoretical modeling, we suggest that this is due to 6S RNA, which minimizes the effect of Rsd on  $E\sigma^{70}$  levels. We show that Rsd regulates 6S RNA expression, thereby minimizing its own effect on  $E\sigma^{70}$ . Since Rsd overexpression increases transcription by  $\sigma^{24}$  and  $\sigma^{54}$  in a ppGpp<sup>0</sup> background (Laurie *et al.* 2003; Costanzo *et al.* 2008), Rsd may generally assist alternative sigma factors to associate with RNA polymerase, under suitable conditions.

6S RNA is more complex, regulating expression of >400 genes; however, we did not observe associations between promoter sequence and 6S RNA susceptibility as previously reported (Cavanagh *et al.* 2008). Our data are more similar to that of (Neusser *et al.* 2010), who reported reduced expression of *rpoB* and ribosomal genes in  $\Delta ssrS$  in stationary phase, and increased ppGpp without increase in *relA* (*spoT* levels were found unchanged in all studies). However, there is still relatively low overlap in the list of 6S RNA regulated genes (46 genes). This is likely because of differences in time points and media. Even within our dataset, there is little overlap between genes

regulated by 6S RNA at different time points. It seems that the 6S RNA regulon varies greatly under different conditions. This supports a model where 6S RNA acts as a background-level regulator operating on RNA polymerase, with gene-level outcomes depending strongly upon cellular environment; this could involve DNA topology, transcription factors, and other RNA polymerase-binding factors, all of which vary with growth phase (Dorman 2013). Regulators that interact with 6S RNA can be experimentally identified, for example by transposon insertion sequencing to identify genes whose disruption leads to a larger survival defect in a 6S RNA knockout than in a wild-type background, or by *in vitro* transcription experiments testing whether the ability of 6S RNA to regulate known targets is affected by the presence of other regulators in the medium.

Previous work using individual promoters (Trotochaud and Wassarman 2004) had suggested that 6S RNA might increase transcription by  $E\sigma^{38}$  during stationary phase. It was unknown whether this was a direct effect of 6S RNA, allowing  $\sigma^{38}$  to compete more effectively for E, or indirect, via a *trans*-acting factor important for  $\sigma^{38}$  activity. However, another study (Neusser *et al.* 2010) found no evidence of this link. We show that 6S RNA increases transcription by  $E\sigma^{38}$  globally, from early exponential to stationary phase, and show mathematically that this is extremely unlikely to be a direct effect of sequestration. We show that 6S RNA increases Rsd protein level and suggest that Rsd is primarily responsible for 6S RNA's effect on  $E\sigma^{38}$  activity.

It has been asked, given 6S RNA's function as a global regulator, why its deletion does not cause a growth defect. Our data, along with others (Cavanagh *et al.* 2010; Neusser *et al.* 2010), shows multiple feedback effects in the  $\Delta ssrS$  strain, where the cell reduces RNA polymerase expression to compensate for 6S RNA loss, and increases  $\sigma^{38}$  and possibly Crl to compensate for reduced  $E\sigma^{38}$  activity. Apart from these two, increased ppGpp may also have such a compensatory role during TS phase, while in LS phase, the 6S RNA knockout has higher expression of *ppk*, responsible for synthesis of polyphosphate - which activates transcription by  $E\sigma^{38}$  (Kusano and Ishihama 1997). In stationary phase, the Rsd/6S RNA double knockout shows reduced expression of DNA supercoiling enzymes; supercoiling regulates promoter binding by  $E\sigma^{70}$  and  $E\sigma^{38}$  (Kusano *et al.* 1996; Bordes *et al.* 2003); in ME phase, it shows reduced expression of Hfq, which increases  $\sigma^{38}$  expression (Zhang *et al.* 1998). These relationships, illustrated in Figure S12, demonstrate that competition between sigma factors is very tightly controlled, and 6S RNA is connected to multiple pathways involved in this process.

We see that 6S RNA and Rsd regulate each other's expression, and control each other's effect on gene expression; finally, we report that 1780 genes across 5 growth phases—almost 40% of the genes in the cell—are differentially expressed in the 6S RNA-Rsd double knockout but not in the single knockouts added together. This provides direct evidence that 6S RNA and Rsd have non-additive effects on gene expression, since the single knockouts cannot explain the widespread gene expression changes observed in the double knockout. Previous studies have only examined single knockouts; this underscores the importance of studying 6S RNA and Rsd as a unit that controls global RNA polymerase activity.

Given that 6S RNA homologs are widespread in bacteria, co-occurring with Rsd and with other RNA polymerase regulators such as the actinobacterial RbpA, and many species have two or three 6S RNA homologs with different expression patterns and potentially different binding partners (Barrick *et al.* 2005; Wehner *et al.* 2014), we suggest that studying the relationships between RNA polymerase regulators will give greater insights into transcriptional control across the bacterial kingdom.

## ACKNOWLEDGMENTS

Illumina sequencing was performed at the Centre for Cellular and Molecular Platforms. Strains and plasmids were obtained from CGSC. We thank Revathy Krishnamurthy and Harshvardhan for assistance with experiments. We also thank Prof. Dipankar Chatterji, Prof. Akira Ishihama, Prof. Karen Wassarman, Prof. Steve Busby, Dr. Dasaradhi Palakodeti, and Prof. M.M. Panicker for discussions.

## FUNDING STATEMENT

This work was supported by core NCBS funds, DBT grant BT/PR3695/BRB/10/979/2011 from the Department of Biotechnology, Government of India, and by grant IA/I/16/2/502711 from the Wellcome Trust-DBT India Alliance. S.K. is funded by the Simons Foundation.

## LITERATURE CITED

- Ali Azam, T., A. Iwata, A. Nishimura, S. Ueda, and A. Ishihama, 1999 Growth phase-dependent variation in protein composition of the *Escherichia coli* nucleoid. *J. Bacteriol.* 181: 6361–6370.
- Anders, S., and W. Huber, 2010 Differential expression analysis for sequence count data. *Genome Biol.* 11: R106. <https://doi.org/10.1186/gb-2010-11-10-r106>
- Barrick, J. E., N. Sudarsan, Z. Weinberg, W. L. Ruzzo, and R. R. Breaker, 2005 6S RNA is a widespread regulator of eubacterial RNA polymerase that resembles an open promoter. *RNA N Y N* 11: 774–784. <https://doi.org/10.1261/rna.7286705>
- Beckmann, B. M., O. Y. Burenina, P. G. Hoch, E. A. Kubareva, C. M. Sharma *et al.*, 2011 In vivo and in vitro analysis of 6S RNA-templated short transcripts in *Bacillus subtilis*. *RNA Biol.* 8: 839–849. <https://doi.org/10.4161/rna.8.5.16151>
- Bernardo, L. M. D., L. U. M. Johansson, D. Solera, E. Skärfstad, and V. Shingler, 2006 The guanosine tetraphosphate (ppGpp) alarmone, DksA and promoter affinity for RNA polymerase in regulation of sigma-dependent transcription. *Mol. Microbiol.* 60: 749–764. <https://doi.org/10.1111/j.1365-2958.2006.05129.x>
- Bintu, L., N. E. Buchler, H. G. Garcia, U. Gerland, T. Hwa *et al.*, 2005 Transcriptional regulation by the numbers: models. *Curr. Opin. Genet. Dev.* 15: 116–124. <https://doi.org/10.1016/j.gde.2005.02.007>
- Bordes, P., A. Conter, V. Morales, J. Bouvier, A. Kolb *et al.*, 2003 DNA supercoiling contributes to disconnect  $\sigma$ S accumulation from  $\sigma$ S-dependent transcription in *Escherichia coli*. *Mol. Microbiol.* 48: 561–571. <https://doi.org/10.1046/j.1365-2958.2003.03461.x>
- Bougdour, A., C. Lelong, and J. Geiselman, 2004 Crl, a low temperature-induced protein in *Escherichia coli* that binds directly to the stationary phase sigma subunit of RNA polymerase. *J. Biol. Chem.* 279: 19540–19550. <https://doi.org/10.1074/jbc.M314145200>
- Cavanagh, A. T., A. D. Klocko, X. Liu, and K. M. Wassarman, 2008 Promoter specificity for 6S RNA regulation of transcription is determined by core promoter sequences and competition for region 4.2 of sigma70. *Mol. Microbiol.* 67: 1242–1256. <https://doi.org/10.1111/j.1365-2958.2008.06117.x>
- Cavanagh, A. T., P. Chandrangsu, and K. M. Wassarman, 2010 6S RNA regulation of relA alters ppGpp levels in early stationary phase. *Microbiology* 156: 3791–3800. <https://doi.org/10.1099/mic.0.043992-0>
- Cho, B.-K., D. Kim, E. M. Knight, K. Zengler, and B. O. Palsson, 2014 Genome-scale reconstruction of the sigma factor network in *Escherichia coli*: topology and functional states. *BMC Biol.* 12: 4. <https://doi.org/10.1186/1741-7007-12-4>
- Colland, F., N. Fujita, A. Ishihama, and A. Kolb, 2002 The interaction between sigmaS, the stationary phase sigma factor, and the core enzyme of *Escherichia coli* RNA polymerase. *Genes Cells Devoted Mol Cell Mech* 7: 233–247. <https://doi.org/10.1046/j.1365-2443.2002.00517.x>
- Costanzo, A., H. Nicoloff, S. E. Barchinger, A. B. Banta, R. L. Gourse *et al.*, 2008 ppGpp and DksA likely regulate the activity of the extracytoplasmic stress factor sigmaE in *Escherichia coli* by both direct and indirect mechanisms. *Mol. Microbiol.* 67: 619–632. <https://doi.org/10.1111/j.1365-2958.2007.06072.x>
- Datsenko, K. A., and B. L. Wanner, 2000 One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* 97: 6640–6645. <https://doi.org/10.1073/pnas.120163297>
- deHaseth, P. L., M. L. Zupancic, and T. M. Record, 1998 RNA polymerase-promoter interactions: the comings and goings of RNA polymerase. *J. Bacteriol.* 180: 3019–3025. <https://doi.org/10.1021/bi00602a006>
- deHaseth, P. L., T. M. Lohman, R. R. Burgess, and M. T. Record, 1978 Nonspecific interactions of *Escherichia coli* RNA polymerase with native and denatured DNA: differences in the binding behavior of core and holoenzyme. *Biochemistry (Mosc.)* 17: 1612–1622. <https://doi.org/10.1021/bi00602a006>
- Dong, T., M. G. Kirchhof, and H. E. Schellhorn, 2008 RpoS regulation of gene expression during exponential growth of *Escherichia coli* K12. *Mol Genet Genomics* MGG 279: 267–277. <https://doi.org/10.1007/s00438-007-0311-4>
- Dorman, C. J., 2013 Genome architecture and global gene regulation in bacteria: making progress towards a unified model? *Nat. Rev. Microbiol.* 11: 349–355. <https://doi.org/10.1038/nrmicro3007>
- England, P. L., F. Westblade, G. Karimova, V. Robbe-Saule, F. Norel *et al.*, 2008 Binding of the unorthodox transcription activator, Crl, to the components of the transcription machinery. *J. Biol. Chem.* 283: 33455–33464. <https://doi.org/10.1074/jbc.M807380200>
- Farewell, A., K. Kvint, and T. Nyström, 1998 Negative regulation by RpoS: a case of sigma factor competition. *Mol. Microbiol.* 29: 1039–1051. <https://doi.org/10.1046/j.1365-2958.1998.00990.x>
- Fukuda, R., M. Taketo, and A. Ishihama, 1978 Autogenous regulation of RNA polymerase beta subunit synthesis in vitro. *J. Biol. Chem.* 253: 4501–4504.
- Ganguly, A., and D. Chatterji, 2012 A comparative kinetic and thermodynamic perspective of the  $\sigma$ -competition model in *Escherichia coli*. *Biophys. J.* 103: 1325–1333. <https://doi.org/10.1016/j.bpj.2012.08.013>
- Gentry, D. R., V. J. Hernandez, L. H. Nguyen, D. B. Jensen, and M. Cashel, 1993 Synthesis of the stationary-phase sigma factor sigma s is positively regulated by ppGpp. *J. Bacteriol.* 175: 7982–7989. <https://doi.org/10.1128/jb.175.24.7982-7989.1993>
- Gildehaus, N., T. Neußer, R. Wurm, and R. Wagner, 2007 Studies on the function of the riboregulator 6S RNA from *E. coli*: RNA polymerase binding, inhibition of in vitro transcription and synthesis of RNA-directed de novo transcripts. *Nucleic Acids Res.* 35: 1885–1896. <https://doi.org/10.1093/nar/gkm085>
- Grigорова, I. L., N. J. Phleger, V. K. Mutalik, and C. A. Gross, 2006 Insights into transcriptional regulation and sigma competition from an equilibrium model of RNA polymerase binding to DNA. *Proc. Natl. Acad. Sci. USA* 103: 5332–5337. <https://doi.org/10.1073/pnas.0600828103>
- Hengge-Aronis, R., and D. Fischer, 1992 Identification and molecular analysis of glgS, a novel growth-phase-regulated and rpoS-dependent gene involved in glycogen synthesis in *Escherichia coli*. *Mol. Microbiol.* 6: 1877–1886. <https://doi.org/10.1111/j.1365-2958.1992.tb01360.x>
- Hengge-Aronis, R., W. Klein, R. Lange, M. Rimmele, and W. Boos, 1991 Trehalose synthesis genes are controlled by the putative sigma factor encoded by rpoS and are involved in stationary-phase thermotolerance in *Escherichia coli*. *J. Bacteriol.* 173: 7918–7924. <https://doi.org/10.1128/jb.173.24.7918-7924.1991>
- Ishihama, A., 2000 Functional Modulation of *Escherichia coli* RNA Polymerase. *Annu. Rev. Microbiol.* 54: 499–518. <https://doi.org/10.1146/annurev.micro.54.1.499>
- Jishage, M., and A. Ishihama, 1995 Regulation of RNA polymerase sigma subunit synthesis in *Escherichia coli*: intracellular levels of sigma 70 and sigma 38. *J. Bacteriol.* 177: 6832–6835. <https://doi.org/10.1128/jb.177.23.6832-6835.1995>
- Jishage, M., and A. Ishihama, 1998 A stationary phase protein in *Escherichia coli* with binding activity to the major sigma subunit of RNA polymerase. *Proc. Natl. Acad. Sci. USA* 95: 4953–4958. <https://doi.org/10.1073/pnas.95.9.4953>

- Jishage, M., and A. Ishihama, 1999 Transcriptional organization and in vivo role of the *Escherichia coli* *rsd* gene, encoding the regulator of RNA polymerase sigma D. *J. Bacteriol.* 181: 3768–3776.
- Jishage, M., K. Kvint, V. Shingler, and T. Nyström, 2002 Regulation of  $\sigma$  factor competition by the alarmone ppGpp. *Genes Dev.* 16: 1260–1270. <https://doi.org/10.1101/gad.227902>
- Kusano, S., and A. Ishihama, 1997 Functional interaction of *Escherichia coli* RNA polymerase with inorganic polyphosphate. *Genes Cells Devoted Mol Cell Mech* 2: 433–441. <https://doi.org/10.1046/j.1365-2443.1997.13203301320330.x>
- Kusano, S., Q. Ding, N. Fujita, and A. Ishihama, 1996 Promoter selectivity of *Escherichia coli* RNA polymerase E sigma 70 and E sigma 38 holoenzymes. Effect of DNA supercoiling. *J. Biol. Chem.* 271: 1998–2004. <https://doi.org/10.1074/jbc.271.4.1998>
- Lange, R., and R. Hengge-Aronis, 1991 Identification of a central regulator of stationary-phase gene expression in *Escherichia coli*. *Mol. Microbiol.* 5: 49–59. <https://doi.org/10.1111/j.1365-2958.1991.tb01825.x>
- Laurie, A. D., L. M. D. Bernardo, C. C. Sze, E. Skärfstad, A. Szalewska-Palasz *et al.*, 2003 The Role of the Alarmone (p)ppGpp in sigma N Competition for Core RNA Polymerase. *J. Biol. Chem.* 278: 1494–1503. <https://doi.org/10.1074/jbc.M209268200>
- Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>
- Maeda, H., N. Fujita, and A. Ishihama, 2000 Competition among seven *Escherichia coli*  $\sigma$  subunits: relative binding affinities to the core RNA polymerase. *Nucleic Acids Res.* 28: 3497–3503. <https://doi.org/10.1093/nar/28.18.3497>
- Mauri, M., and S. Klumpp, 2014 A model for sigma factor competition in bacterial cells. *PLOS Comput. Biol.* 10: e1003845. <https://doi.org/10.1371/journal.pcbi.1003845>
- Mitchell, J. E., T. Oshima, S. E. Piper, C. L. Webster, L. F. Westblade *et al.*, 2007 The *Escherichia coli* Regulator of Sigma 70 Protein, Rsd, Can Up-Regulate Some Stress-Dependent Promoters by Sequestering Sigma 70. *J. Bacteriol.* 189: 3489–3495. <https://doi.org/10.1128/JB.00019-07>
- Neusser, T., T. Polen, R. Geissen, and R. Wagner, 2010 Depletion of the non-coding regulatory 6S RNA in *E. coli* causes a surprising reduction in the expression of the translation machinery. *BMC Genomics* 11: 165. <https://doi.org/10.1186/1471-2164-11-165>
- Peter, B. J., J. Arsuaga, A. M. Breier, A. B. Khodursky, P. O. Brown *et al.*, 2004 Genomic transcriptional response to loss of chromosomal supercoiling in *Escherichia coli*. *Genome Biol.* 5: R87. <https://doi.org/10.1186/gb-2004-5-11-r87>
- Piper, S. E., J. E. Mitchell, D. J. Lee, and S. J. W. Busby, 2009 A global view of *Escherichia coli* Rsd protein and its interactions. *Mol. Biosyst.* 5: 1943–1947. <https://doi.org/10.1039/b904955j>
- Proshkin, S., A. R. Rahmouni, A. Mironov, and E. Nudler, 2010 Cooperation between translating ribosomes and RNA polymerase in transcription elongation. *Science* 328: 504–508. <https://doi.org/10.1126/science.1184939>
- Rahman, M., M. R. Hasan, T. Oba, and K. Shimizu, 2006 Effect of *rpoS* gene knockout on the metabolism of *Escherichia coli* during exponential growth phase and early stationary phase based on gene expressions, enzyme activities and intracellular metabolite concentrations. *Biotechnol. Bioeng.* 94: 585–595. <https://doi.org/10.1002/bit.20858>
- Salgado, H., M. Peralta-Gil, S. Gama-Castro, A. Santos-Zavaleta, L. Muñiz-Rascado *et al.*, 2013 RegulonDB v8.0: omics data sets, evolutionary conservation, regulatory phrases, cross-validated gold standards and more. *Nucleic Acids Res.* 41: D203–D213. <https://doi.org/10.1093/nar/gks1201>
- Sharma, U. K., and D. Chatterji, 2008 Differential mechanisms of binding of anti-sigma factors *Escherichia coli* Rsd and bacteriophage T4 AsiA to *E. coli* RNA polymerase lead to diverse physiological consequences. *J. Bacteriol.* 190: 3434–3443. <https://doi.org/10.1128/JB.01792-07>
- Shimada, T., Y. Yamazaki, K. Tanaka, and A. Ishihama, 2014 The whole set of constitutive promoters recognized by RNA polymerase RpoD holoenzyme of *Escherichia coli*. *PLoS One* 9: e90447 (erratum: *PLoS One* 9: e100908). <https://doi.org/10.1371/journal.pone.0090447>
- Smith, T. L., and R. T. Sauer, 1996 Dual regulation of open-complex formation and promoter clearance by Arc explains a novel repressor to activator switch. *Proc. Natl. Acad. Sci. USA* 93: 8868–8872. <https://doi.org/10.1073/pnas.93.17.8868>
- Tanaka, K., Y. Takayanagi, N. Fujita, A. Ishihama, and H. Takahashi, 1993 Heterogeneity of the principal sigma factor in *Escherichia coli*: the *rpoS* gene product, sigma 38, is a second principal sigma factor of RNA polymerase in stationary-phase *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 90: 3511–3515. <https://doi.org/10.1073/pnas.90.8.3511>
- Thomason, L. C., N. Costantino, and D. L. Court, 2007 *E. coli* genome manipulation by P1 transduction. *Curr. Protoc. Mol. Biol.* Chapter 1: Unit 1.17. <https://doi.org/10.1002/0471142727.mb0117s79>
- Traxler, M. F., S. M. Summers, H.-T. Nguyen, V. M. Zacharia, J. T. Smith *et al.*, 2008 The global, ppGpp-mediated stringent response to amino acid starvation in *Escherichia coli*. *Mol. Microbiol.* 68: 1128–1148. <https://doi.org/10.1111/j.1365-2958.2008.06229.x>
- Trotochaud, A. E., and K. M. Wassarman, 2004 6S RNA Function Enhances Long-Term Cell Survival. *J. Bacteriol.* 186: 4978–4985. <https://doi.org/10.1128/JB.186.15.4978-4985.2004>
- Typas, A., C. Barembruch, A. Possling, and R. Hengge, 2007 Stationary phase reorganisation of the *Escherichia coli* transcription machinery by Crl protein, a fine-tuner of sigmas activity and levels. *EMBO J.* 26: 1569–1578. <https://doi.org/10.1038/sj.emboj.7601629>
- Uzzau, S., N. Figueroa-Bossi, S. Rubino, and L. Bossi, 2001 Epitope tagging of chromosomal genes in *Salmonella*. *Proc. Natl. Acad. Sci. USA* 98: 15264–15269. <https://doi.org/10.1073/pnas.261348198>
- Wassarman, K. M., and G. Storz, 2000 6S RNA regulates *E. coli* RNA polymerase activity. *Cell* 101: 613–623. [https://doi.org/10.1016/S0092-8674\(00\)80873-9](https://doi.org/10.1016/S0092-8674(00)80873-9)
- Wassarman, K. M., and R. M. Saecker, 2006 Synthesis-Mediated Release of a Small RNA Inhibitor of RNA Polymerase. *Science* 314: 1601–1603. <https://doi.org/10.1126/science.1134830>
- Weber, H., T. Polen, J. Heuveling, V. F. Wendisch, and R. Hengge, 2005 Genome-Wide Analysis of the General Stress Response Network in *Escherichia coli*: sigmaS-Dependent Genes, Promoters, and Sigma Factor Selectivity. *J. Bacteriol.* 187: 1591–1603. <https://doi.org/10.1128/JB.187.5.1591-1603.2005>
- Wehner, S., K. Damm, R. K. Hartmann, and M. Marz, 2014 Dissemination of 6S RNA among Bacteria. *RNA Biol.* 11: 1467–1478. <https://doi.org/10.4161/rna.29894>
- Zhang, A., S. Altuvia, A. Tiwari, L. Argaman, R. Hengge-Aronis *et al.*, 1998 The OxyS regulatory RNA represses *rpoS* translation and binds the Hfq (HF-I) protein. *EMBO J.* 17: 6061–6068. <https://doi.org/10.1093/emboj/17.20.6061>

Communicating editor: B. Andrews