




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

## Progress in Cardiovascular Diseases

journal homepage: [www.onlinepcd.com](http://www.onlinepcd.com)

## Why are COVID-19 effects less severe in Sub-Saharan Africa? Moving more and sitting less may be a primary reason



Lucy-Joy Wachira<sup>a,\*\*</sup>, Ross Arena<sup>b,c,\*</sup>, James F. Sallis<sup>d,h</sup>, Estelle V. Lambert<sup>e</sup>, Otieno Martin Ong'wen<sup>f</sup>, Deepika R. Laddu<sup>b,c</sup>, Vincent Onywera<sup>a</sup>, Adewale L. Oyeyemi<sup>g</sup>, On behalf of the HL-PIVOT Network

<sup>a</sup> Dept. of Physical Education and Exercise Science, School of Public Health and Applied Human Sciences, Kenyatta University, Nairobi, Kenya

<sup>b</sup> Department of Physical Therapy, College of Applied Science, University of Illinois, Chicago, IL, United States of America

<sup>c</sup> Healthy Living for Pandemic Event Protection (HL – PIVOT) Network, Chicago, IL, United States of America

<sup>d</sup> Herbert Wertheim School of Public Health and Human Longevity Science, University of California, San Diego, La Jolla, CA, United States of America

<sup>e</sup> Research Centre for Health through Physical Activity, Lifestyle and Sport (HPALS), Department of Human Biology, Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa

<sup>f</sup> AfyAfrica Orthopedic services, Nairobi, Kenya

<sup>g</sup> Department of Physiotherapy, University of Maiduguri, P.M.B 1069, Maiduguri, Borno State, Nigeria

<sup>h</sup> Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, Victoria, Australia

The world is entering a new phase of the coronavirus disease 2019 (COVID-19) health crisis with the lifting of social and physical distancing as well as lockdown restrictions to control the pandemic. Scientific evidence obtained during the COVID-19 pandemic to this point have brought clear themes to the forefront. One important theme pertains to who is at a higher risk for poorer outcomes if infected with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). Clearly, individuals with risk factors for chronic disease and one or more chronic disease diagnoses are at significantly higher risk for poor outcomes with SARS-CoV-2 infection.<sup>1,2</sup> Moreover, unhealthy lifestyle behaviors (i.e., physical inactivity, poor nutrition, smoking and excess body mass) are the leading cause for the high incidence and prevalence of chronic disease the world was facing well before the COVID-19 pandemic.<sup>3</sup> In fact, physical inactivity and chronic diseases were both characterized as pandemics prior to COVID-19.<sup>1,4,5</sup> In this context, decades of unhealthy lifestyle behaviors, both independently and through the association with chronic disease risk factors, *primed the pump* for poor COVID-19 outcomes.<sup>1,6–8</sup> Another emerging theme is the disproportionate burden of poor outcomes in underserved communities and underrepresented individuals.<sup>9–11</sup> A higher prevalence of unhealthy lifestyle characteristics and chronic disease certainly contributes to this disparity.<sup>12–14</sup> From an international perspective related to health inequalities, a common conclusion is that the burden of disease is usually highest in the lowest-income countries, especially countries in sub-Saharan Africa. This pattern is most starkly seen for a wide range of infectious diseases,<sup>15</sup> though rates of non-communicable diseases

are already substantial and are rising in the region.<sup>16,17</sup> As such, since the early days of the pandemic, there has been grave concern for the disastrous impact the COVID-19 pandemic would have in sub-Saharan Africa.

A recent New York Times article highlighted a surprising exception to the pattern of health inequities disfavoring Africa – COVID-19 pandemic outcomes.<sup>18</sup> The reporter described both anecdotal and quantitative evidence for why the anticipated devastating toll of COVID-19 in Africa, due to poverty, crowding, limited medical care, and inadequate distributions and uptake of vaccinations, had not occurred. A compilation of “excess deaths” in virtually every country throughout the pandemic by The Economist, among others, seems to generally support the low mortality of COVID-19 in sub-Saharan African countries.<sup>19,20</sup>

If confirmed by further analysis of available data, this pattern raises obvious questions about what might account for low COVID-19 mortality in Africa. The generally younger age distribution throughout Africa has been offered as one explanation,<sup>20</sup> and it is a credible one. But there are other explanations that should be explored, and physical activity (PA) is a promising candidate. Africa is currently the most physically active region of the world.<sup>21</sup> There is a sizeable scientific literature documenting numerous biological effects of PA relevant to infectious diseases, including widespread improvements in immune functioning, lower infection rates, reduced inflammation during infections, less severe outcomes of infections, and enhanced effectiveness of vaccines, especially among immunocompromised groups such as older adults.<sup>22,23</sup> If these benefits are meaningful, we would expect to see benefits of cardiorespiratory fitness and PA during the COVID-19 pandemic, which has in fact been the case.<sup>24–28</sup> A recent systematic review by the United States (US) Centers for Disease Control (CDC) included 25 studies specific to COVID-19 outcomes from multiple continents using a variety of measures of PA and fitness, as well as multiple COVID-19 outcomes. They found the evidence was consistent and conclusive that adults who were physically active or fit prior to a COVID-19 diagnosis were less likely to be hospitalized and die, with notable, but weaker

\* Correspondence to: R. Arena, Department of Physical Therapy, College of Applied Health Sciences, University of Illinois Chicago, 1919 W. Taylor Street (MC 898), Chicago, IL 60612.

\*\* Correspondence to: L.-J. Wachira, Dept. of Physical Education and Exercise Science, School of Public Health and Applied Human Sciences, Kenyatta University, P.O Box 438433 00100, Nairobi, Kenya.

E-mail address: [wachira.lucy@ku.ac.ke](mailto:wachira.lucy@ku.ac.ke) (L.-J. Wachira), [raarena@uic.edu](mailto:raarena@uic.edu) (R. Arena).

evidence related to ICU admission and being placed on a ventilator.<sup>29</sup> This analysis provided convincing evidence that low PA increases the risk for poor COVID-19 outcomes resulting in the CDC listing low PA as an established risk factor on their COVID-19 “people with certain medical conditions” webpage.<sup>30</sup> There is an additional recent large ( $n = 65,361$ ) study from South Africa confirming the main results of the review using device-based measures of PA in a large sample, with apparently larger effect sizes than have been reported in prior studies of similar design.<sup>31</sup> Thus, the benefits of physical activity for COVID-19 also apply in Africa, though this appeared to be a mainly higher-income sample.

To explain lower COVID-19 death rates throughout sub-Saharan Africa, it would have to be shown, first, that PA is generally higher in Africa than in higher-income regions. There is substantial evidence this is the case. Self-report data from the World Health Organization's (WHO) STEPs surveillance program show rates of insufficient PA are well below the global average for women and men.<sup>21</sup> There are several studies among smaller samples in sub-Saharan Africa using device-based PA measures providing further evidence that African adults are substantially more physically active than samples in the US<sup>32</sup> and Europe.<sup>33</sup> Numerous health benefits directly relevant to infectious diseases are associated with being physically active, one of which is lower levels of systemic inflammation.<sup>34,35</sup> Poor outcomes with SARS-CoV-2 have been associated with a *cytokine storm*,<sup>6</sup> and elevated systemic inflammation prior to infection may foretell an increased risk for this storm and an untoward clinical course.<sup>36</sup> There is some evidence to indicate systemic inflammation levels are lower in the sub-Saharan African population.<sup>37</sup> Thus, higher PA contributing to lower systemic inflammation may be one mechanism for improved COVID-19 outcomes in this region of the world. Obesity has been strongly linked to poor COVID-19 outcomes.<sup>38–40</sup> While obesity is on the rise in sub-Saharan African population,<sup>41,42</sup> the overall prevalence still lags behind other countries such as the US.<sup>43</sup> Perhaps interactions between higher PA levels and a lower body mass in sub-Saharan Africa have also contributed to improved COVID-19 outcomes. Recent data suggest physical inactivity was a stronger predictor than obesity of COVID-19 deaths in 53 Sub-Saharan Africa countries.<sup>44</sup> However, it should be noted again that obesity is on the rise in this region, serving as an ominous sign for the future.

A related potential explanation for the lower COVID-19 death rates in Africa is that many Africans spend more time outdoors, especially those who live outside of cities and in the rural areas.<sup>45–47</sup> More outdoor living could be protective from infectious diseases through at least two mechanisms. First, a recent systematic review summarized evidence from 33 studies that being in nature has beneficial effects on the immune and inflammation systems.<sup>48</sup> Second, though the distribution of physical activity outdoors versus indoors has not been well-studied among adults,<sup>49</sup> at least some subgroups of adults prefer to be active outdoors.<sup>50</sup> Many common leisure activities are performed outdoors. The commonly reported outdoor activity types among adults and adolescents in Sub-Saharan Africa are walking, running, cycling, digging, swimming, dancing, and playing soccer.<sup>47,51</sup> However, it is unclear whether being active and being outdoors have synergistic benefits relevant to decreased risk of chronic disease as well as decreased illness severity in the event of a viral infection.

In addition to many Africans spending more time outdoors, it is probable the majority of individuals did not significantly reduce their PA levels during the pandemic. Not because they did not care about the public health directives on movement restrictions, but because PA (i.e., *moving more and sitting less*) is embedded in their activities of daily living and economic survival. For instance, active transportation is part of normal life in sub-Saharan Africa, with a majority relying heavily on public transport, and a relatively low proportion of individuals having access to or owning a private vehicle.<sup>52</sup> The population in sub-Saharan Africa, the majority of whom are from resource deprived low-income families, depend less on motorized, digitized, labor saving

gadgets and equipment. Rather, this population relies on low technology activities (e.g., small informal business enterprises, farming) that involve extensive movement of light to moderate intensity to earn a living, including frequently interrupted sitting. Perhaps the benefits that come with energy expenditure and increased metabolism throughout the waking hours of a day in relation to COVID-19 infection rate and severity is worth exploring.

At the population level, there is evidence of a PA transition in sub-Saharan Africa that suggests disparity in physical activity by levels of urbanization and socioeconomic status,<sup>47,53</sup> with physical inactivity increasing with increasing urbanization and income.<sup>54</sup> Although data on disparities in COVID-19 infection rate and severity between Africa's urban and rural areas is scarce, anecdotal evidence suggests a sharp divide in urban-rural COVID-19 outcomes.<sup>55,56</sup> While fewer COVID-19 infections and less severity are often recorded in rural areas in Africa, the economic impact and social devastations, such as disruption in livelihoods and the agriculture and food system, are often more pronounced in rural than urban areas of Africa.<sup>56,57</sup> However, studies are needed to explore how each country's levels of urbanization, income and PA may interact to influence COVID-19 infection rate and severity in sub-Saharan Africa. Such studies could offer definitive answers to the hypothesis that PA in Africa may be, at least partly responsible for the lower incidence and prevalence of COVID-19 and its severe outcomes, when compared to those reported in the high-income Western countries.

In conclusion, the profound importance of being physical active to one's health is again being illustrated during the COVID-19 pandemic. While underserved communities and underrepresented individuals in many parts of the world are being disproportionately impacted by adverse COVID-19 outcomes, the sub-Saharan Africa region seems to be faring far better than would be expected. This is welcome news and the potential explanations for this favorable result should be examined. In this context, more studies identifying the benefits of PA and being outdoors prior to and during the COVID-19 pandemic are needed in sub-Saharan Africa. To inform evidence-based interventions in Africa, we suggest investigators should conduct PA and COVID-19 studies across the several phases of the behavioral epidemiological framework, including studies on establishing the link between physical activity and COVID-19, identifying social and structural factors that influence PA in the context of the COVID-19 pandemic, developing and evaluating interventions to increase PA during COVID-19, and translating research into practice and policy for current and future pandemics.<sup>58</sup> While the reasons for better outcomes are yet to be elucidated, higher levels of PA may be a primary protective contributor. If this proves to be the case, sub-Saharan Africa will serve as a shining example of how underserved communities can employ healthy living behaviors to prepare for and combat health crises, yet again illustrating *moving more and sitting less* is medicine that should be equitably administered around the world.

## References

1. Arena R, Lavie CJ. The global path forward - healthy living for pandemic event protection (HL - PIVOT). *Prog Cardiovasc Dis* 2021;64:96-101.
2. Liu B, Spokes P, He W, Kaldor J. High risk groups for severe COVID-19 in a whole of population cohort in Australia. *BMC Infect Dis* 2021;21:1685.
3. Sagner M, McNeil A, Puska P, et al. The P4 health spectrum - a predictive, preventive, personalized and participatory continuum for promoting Healthspan. *Prog Cardiovasc Dis* 2017;59:506-521.
4. Hall G, Laddu DR, Phillips SA, Lavie CJ, Arena R. A tale of two pandemics: how will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another? *Prog Cardiovasc Dis* 2021;64:108-110.
5. Kohl 3rd HW, Craig CL, Lambert EV, et al. The pandemic of physical inactivity: global action for public health. *Lancet* 2012;380:294-305.
6. Arena R, Bond S, Calvo IR, et al. Shelter from the cytokine storm: healthy living is a vital preventative strategy in the COVID-19 era. *Prog Cardiovasc Dis* 2021. <https://doi.org/10.1016/j.pcad.2021.06.008>. S0033-0620(21)00066-9, Online ahead of print.
7. Arena R, Hall G, Laddu DR, Phillips SA, Lavie CJ. A tale of two pandemics revisited: physical inactivity, sedentary behavior and poor COVID-19 outcomes reside in the

- same Syndemic City. *Prog Cardiovasc Dis* 2021. [https://doi.org/10.1016/j.pcad.2021.11.012.S0033-0620\(21\)00131-6](https://doi.org/10.1016/j.pcad.2021.11.012.S0033-0620(21)00131-6), Online ahead of print.
8. Arena R, Myers J, Kaminsky LA, et al. Current activities centered on healthy living and recommendations for the future: a position statement from the HL-PIVOT network. *Curr Probl Cardiol* 2021;46, 100823.
  9. Lee IJ, Ahmed NU. The devastating cost of racial and ethnic health inequity in the COVID-19 pandemic. *J Natl Med Assoc* 2021;113:114-117.
  10. Rentsch CT, Kidwai-Khan F, Tate JP, et al. Covid-19 by race and ethnicity: A national cohort study of 6 million United States veterans. medRxiv : the preprint server for health sciences; 2020.
  11. Fowler Davis S, Choppin S, Kelly S. Towards an understanding of population health data in a single NHS trust during COVID-19. *Healthcare (Basel, Switzerland)* 2022;10.
  12. Krishnaswami J, Sardana J, Daxini A. Community-engaged lifestyle medicine as a framework for health equity: principles for lifestyle medicine in low-resource settings. *Am J Lifestyle Med* 2019;13:443-450.
  13. Bantham A, Taverno Ross SE, Sebastião E, Hall G. Overcoming barriers to physical activity in underserved populations. *Prog Cardiovasc Dis* 2020;64:64-71. <https://doi.org/10.1016/j.pcad.2020.11.002>.
  14. Akam EY, Nuako AA, Daniel AK, Stanford FC. Racial disparities and cardiometabolic risk: new horizons of intervention and prevention. *Curr Diab Rep* 2022;22:129-136.
  15. Global, regional, and national age-sex-specific mortality for 282 causes of death in 195 countries and territories, 1980-2017: a systematic analysis for the global burden of disease study 2017. *Lancet* 2018;392:1736-1788.
  16. Bigna JJ, Noubiap JJ. The rising burden of non-communicable diseases in sub-Saharan Africa. *Lancet Glob Health* 2019;7:e1295-e1296.
  17. Gouda HN, Charlson F, Sorsdahl K, et al. Burden of non-communicable diseases in sub-Saharan Africa, 1990-2017: results from the global burden of disease study 2017. *Lancet Glob Health* 2019;7:e1375-e1387.
  18. Nolen S. *Trying to solve a covid mystery: Africa's low death rates*. New York Times. 2022.
  19. Economist T. Tracking covid-19 excess deaths across countries. <https://www.economist.com/graphic-detail/coronavirus-excess-deaths-tracker>. Accessed: 3/31/2022.
  20. Adams J, MacKenzie MJ, Amegah AK, et al. The conundrum of low COVID-19 mortality burden in sub-Saharan Africa: myth or reality? *Global Health Sci Pract* 2021;9: 433-443.
  21. Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *Lancet Glob Health* 2018;6:e1077-e1086.
  22. Chastin SFM, Abaraogu U, Bourgois JG, et al. Effects of regular physical activity on the immune system, vaccination and risk of community-acquired infectious disease in the general population: systematic review and Meta-analysis. *Sports Med* 2021;51: 1673-1686.
  23. Simpson RJ, Katsanis E. The immunological case for staying active during the COVID-19 pandemic. *Brain Behav Immun* 2020;87:6-7.
  24. Sallis R, Young DR, Tartof SY, et al. Physical inactivity is associated with a higher risk for severe COVID-19 outcomes: a study in 48 440 adult patients. *Br J Sports Med* 2021;55(19):1099-1105. <https://doi.org/10.1136/bjsports-2021-104080>. Epub 2021 Apr 13.
  25. Brawner CA, Ehrman JK, Bole S, et al. Inverse relationship of maximal exercise capacity to hospitalization secondary to coronavirus disease 2019. *Mayo Clin Proc* 2021;96:32-39.
  26. Kerrigan DJ, Brawner CA, Ehrman JK, Keteyian S. Cardiorespiratory fitness attenuates the impact of risk factors associated with COVID-19 hospitalization. *Mayo Clin Proc* 2021;96:822-823.
  27. Lavie CJ, Sanchis-Gomar F, Arena R. Fit is it in COVID-19, future pandemics, and overall healthy living. *Mayo Clin Proc* 2021;96:7-9.
  28. Lavie CJ, Sanchis-Gomar F, Arena R. In reply - cardiorespiratory fitness attenuates the impact of risk factors associated with COVID-19 hospitalization. *Mayo Clin Proc* 2021;96:823-824.
  29. CDC. Brief summary of findings on the association between physical inactivity and severe COVID-19 outcomes. <https://www.cdc.gov/coronavirus/2019-ncov/downloads/clinical-care/E-Physical-Inactivity-Review.pdf>.
  30. CDC. COVID-19: People with certain medical conditions. <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>. Accessed: 3/31/2022.
  31. Steenkamp L, Saggars RT, Bandini R, et al. Small steps, strong shield: directly measured, moderate physical activity in 65 361 adults is associated with significant protective effects from severe COVID-19 outcomes. *Br J Sports Med* 2022. <https://doi.org/10.1136/bjsports-2021-105159>. bjsports-2021-105159, Online ahead of print.
  32. Pratt M, Sallis JF, Cain KL, et al. Physical activity and sedentary time in a rural adult population in Malawi compared with an age-matched US urban population. *BMJ Open Sport Exerc Med* 2020;6, e000812.
  33. Brage S, Assah F, Msyamboza KP. Quantifying population levels of physical activity in Africa using wearable sensors: implications for global physical activity surveillance. *BMJ Open Sport Exerc Med* 2020;6, e000941.
  34. Hamer M, Sabia S, Batty GD, et al. Physical activity and inflammatory markers over 10 years: follow-up in men and women from the Whitehall II cohort study. *Circulation* 2012;126:928-933.
  35. Hamer M, Stamatakis E. Physical activity and risk of cardiovascular disease events: inflammatory and metabolic mechanisms. *Med Sci Sports Exerc* 2009;41:1206-1211.
  36. Ali N. Elevated level of C-reactive protein may be an early marker to predict risk for severity of COVID-19. *J Med Virol* 2020;92:2409-2411.
  37. Kohler IV, Anglewicz P, Kohler H-P, McCabe JF, Chilima B, Soldo BJ. Evaluating health and disease in sub-Saharan Africa: minimally invasive collection of plasma in the Malawi longitudinal study of families and health (MLSFH). *Genus* 2012;68:1-27.
  38. Alkhatib AL, Kreniske J, Zifodya JS, et al. BMI is associated with coronavirus disease 2019 intensive care unit admission in African Americans. *Obesity (Silver Spring)* 2020;28:1798-1801.
  39. Czernichow S, Bain SC, Capehorn M, et al. Costs of the COVID-19 pandemic associated with obesity in Europe: a health-care cost model. *Clin Obesity* 2021, e12442.
  40. Finer N, Garnett SP, Bruun JM. *COVID-19 and obesity*, 10. 2020:e12365.
  41. Price AJ, Crampton AC, Amberbir A, et al. Prevalence of obesity, hypertension, and diabetes, and cascade of care in sub-Saharan Africa: a cross-sectional, population-based study in rural and urban Malawi. *Lancet Diabet Endocrinol* 2018;6:208-222.
  42. Ramsay M, Crowther NJ, Angongo G, et al. As members of AWIG and the HAC. Regional and sex-specific variation in BMI distribution in four sub-Saharan African countries: the H3Africa AWI-gen study. *Glob Health Action* 2018;11:1556561.
  43. Tsao CW, Aday AW, Almarzooq ZI, et al. Heart disease and stroke statistics-2022 update: a report from the American Heart Association. *Circulation* 2022;145:e153-e639.
  44. Okeahalam C, Williams V, Otumbe K. Factors associated with COVID-19 infections and mortality in Africa: a cross-sectional study using publicly available data. *BMJ Open* 2020;10, e042750.
  45. Wachira L-JM. Lifestyle transition towards sedentary behavior among children and youth in sub-Saharan Africa: A narrative review. In: *Maerques A, Gouveia ER, eds. Sedentary behaviour - A contemporary view* London: Intech; 2021.
  46. Muthuri SK, Wachira L-JM, Leblanc AG, et al. Temporal trends and correlates of physical activity, sedentary behaviour, and physical fitness among school-aged children in sub-Saharan Africa: a systematic review. *Int J Environ Res Public Health* 2014;11: 3327-3359.
  47. Padrão P, Damasceno A, Silva-Matos C, Prista A, Lunet N. Physical activity patterns in Mozambique: urban/rural differences during epidemiological transition. *Prev Med* 2012;55:444-449.
  48. Andersen L, SSS Corazon, UKK Stigsdotter. Nature exposure and its effects on immune system functioning: a systematic review. *Int J Environ Res Public Health* 2021:18.
  49. Trost SG, Owen N, Bauman AE, Sallis JF, Brown W. Correlates of adults' participation in physical activity: review and update. *Med Sci Sports Exerc* 2002;34:1996-2001.
  50. Burton NW, Khan A, Brown WJ. How, where and with whom? Physical activity context preferences of three adult groups at risk of inactivity. *Br J Sports Med* 2012;46: 1125-1131.
  51. De Vos JCW, Du Toit D, Coetzee D. The types and levels of physical activity and sedentary behaviour of senior phase learners in Potchefstroom. *Health SA Gesondheid* 2016;21:372-380.
  52. Oyeyemi AL, Oyeyemi AY, Jidda ZA, Babagana F. Prevalence of physical activity among adults in a metropolitan Nigerian city: a cross-sectional study. *J Epidemiol* 2013;23:169-177.
  53. Assah F, Mbanya JC, Ekelund U, Wareham N, Brage S. Patterns and correlates of objectively measured free-living physical activity in adults in rural and urban Cameroon. *J Epidemiol Community Health* 2015;69:700-707.
  54. Lambert CT, Siefken K, Varela AR, Waqanivalu T, Schilkenporf N. Case study from Africa: Physical activity and safety from crime and traffic in Africa. In: *Oyeyemi AL, Kolbe-Alexander T, eds. Physical activity in low- and middle-income Countries*. Routledge; 2021. p. 171-186.
  55. Finn BM, Kobayashi LC. Structural inequality in the time of COVID-19: urbanization, segregation, and pandemic control in sub-Saharan Africa. *Dial Human Geogr* 2020;10:217-220.
  56. Ogunkola IO, Adebisi YA, Imo UF, Odey GO, Esu E, Lucero-Priso 3rd DE. Rural communities in Africa should not be forgotten in responses to COVID-19. *Int J Health Plann Manag* 2020;35:1302-1305.
  57. FAO, ECA and AUC. *Africa Regional Overview of Food Security and Nutrition 2020: Transforming Food Systems for Affordable Healthy Diets*. Accra: FAO. 2021.
  58. Sallis JF, Adlakh D, Oyeyemi A, Salvo D. An international physical activity and public health research agenda to inform coronavirus disease-2019 policies and practices. *J Sport Health Sci* 2020;9:328-334.