

Fourth, it is hypothesised by the authors that the pharmacokinetics and pharmacodynamics of propofol and desflurane are the same. They had discontinued desflurane 3 min before the end of the surgery, whereas propofol was stopped 10 min before. Moreover, nitrous oxide was used in the desflurane group, but not in the total intravenous anaesthesia (TIVA) group. It is established that the use of nitrous oxide impacts postoperative lung collapse and atelectasis process (7).

Although we applaud the authors for their commendable work, and also assume that they indeed observed a significant change in the postoperative pulmonary function in the balanced anaesthesia group compared with TIVA, it is difficult to be convinced of the attribution of such difference to desflurane, which hardly stays for few minutes after discontinuation. This needs introspection and further discussion. Although the clinical significance of any minor reduction in the postoperative pulmonary function in a healthy individual is minimal, it can be significant in obese patients and those having pulmonary comorbidities. Thus, we propose that further study will be required to clarify whether there is any impact of desflurane-based balanced anaesthesia on postoperative pulmonary function, by addressing the issues that we have raised, before we can abandon the use of this novel agent having many clinical advantages, especially the excellent quality of rapid recovery even after prolonged surgery.

#### Author's Reply

### Re: Attribution of Delayed Poor Lung Function to Desflurane-Based Balanced Anaesthesia Might be Inappropriate: Our Reply to Sharma et al.'s article

Sir, We read the comments by Karim et al. (1) on our published study (2) with great interest. Karim et al correctly suggest that the ventilator technique used, operative insult, the patient's ongoing pathophysiological insults, amount and type of intravenous fluid used and overall fluid balance affect postoperative pulmonary function. For this reason we conducted this study in patients of ASA physical status I/II undergoing mastoid surgery only. All received the same intraoperative ventilator technique and there was no significant blood loss or major fluid shifts.

Secondly, a standard analgesia regimen was used in all patients consisting of fentanyl  $2 \mu\text{g kg}^{-1}$  5 minutes before induction of anaesthesia followed by  $1 \mu\text{g kg}^{-1}$  repeated every hour. All patients received diclofenac  $1.5 \text{ mg kg}^{-1}$  and paracetamol  $1 \text{ gm}$  for postoperative analgesia. In the PACU, pain was to be

## References

1. Sharma S, Bhalotra AR, Awal S. Changes in Lung Function Parameters after Total Intravenous Anaesthesia and Balanced Anaesthesia with Desflurane: A Prospective Randomised Study. *Turk J Anaesthesiol Reanim* 2020; 48: 17-23. [\[CrossRef\]](#)
2. Saraswat V. Effects of anaesthesia techniques and drugs on pulmonary function. *Indian J Anaesth* 2015; 59: 557-64.
3. Güldner A, Kiss T, Serpa Neto A, Hemmes SNT, Canet J, Spieth PM, et al. Intraoperative protective mechanical ventilation for prevention of postoperative pulmonary complications: a comprehensive review of the role of tidal volume, positive end-expiratory pressure, and lung recruitment maneuvers. *Anesthesiology* 2015; 123: 692-713. [\[CrossRef\]](#)
4. Holte K, Jensen P, Kehlet H. Physiologic effects of intravenous fluid administration in healthy volunteers. *Anesth Analg* 2003; 96: 1504-9. [\[CrossRef\]](#)
5. Mills GH. Respiratory complications of anaesthesia. *Anaesthesia* 2018; 73(Suppl 1): 25-33. [\[CrossRef\]](#)
6. Lee LA, Caplan RA, Stephens LS, Posner KL, Terman GW, Voepel-Lewis T, et al. Postoperative opioid-induced respiratory depression: a closed claims analysis. *Anesthesiology* 2015; 122: 659-65. [\[CrossRef\]](#)
7. Yoshimura T, Ueda K, Kakinuma A, Sawai J, Nakata Y. Bronchial blocker lung collapse technique: nitrous oxide for facilitating lung collapse during one-lung ventilation with a bronchial blocker. *Anesth Analg* 2014; 118: 666-70. [\[CrossRef\]](#)

treated with morphine boluses  $1.5 \text{ mg}$  given at 5-min intervals to achieve adequate analgesia but this was not required in any patient.

Third, for accurate spirometry results, a well-cooperative patient is required. Before each assessment, it was ensured that the patient was warm and pain free so that there was no pain, shivering etc., which could interfere with the patients ability to breathe deeply and the patient could perform spirometry in a reliable manner. All patients underwent mastoid surgery and there was no requirement of postoperative sedative and analgesic drugs.

Fourthly, it was not hypothesised that the pharmacokinetics and pharmacodynamics of propofol and desflurane are the same. Desflurane was discontinued at the end of surgery as is recommended because of its low blood gas solubility and quick recovery whereas propofol was stopped 10 min before, keeping in mind its context sensitive half time. Nitrous oxide was used as part of a standard balanced anaesthesia technique but is not a part of a standard total intravenous anaesthesia (TIVA) technique and the purpose of this study was to compare these two techniques.

The aim of our study was to assess whether the lesser coughing and smoother emergence after use of TIVA with propofol leads to greater decreases in lung function postoperatively as compared to balanced anaesthesia with desflurane. We found that both anaesthetic techniques caused postoperative impairment in the lung function, but while TIVA caused a greater reduction in PFT in the early postoperative period, balanced anaesthesia with desflurane was associated with a greater reduction in forced vital capacity at 24 hours. We concluded that TIVA with propofol was not found to have detrimental effects on the lung function postoperatively as compared to a balanced anaesthesia technique with desflurane and suggested that further studies are required in patients who are at risk of developing postoperative pulmonary complications and have at no point suggested we abandon the use of balanced anaesthesia with desflurane.

## References

1. Karim HMR, Bhakta P. Attribution of Delayed Poor Lung Function to Desflurane-Based Balanced Anaesthe-

sia Might be Inappropriate: Our Reply to Sharma et al.'s article. Turk J Anaesthesiol Reanim 2020; DOI: 10.5152/TJAR.2020.97

2. Sharma S, Bhalotra AR, Awal S. Changes in Lung Function Parameters after Total Intravenous Anaesthesia and Balanced Anaesthesia with Desflurane: A Prospective Randomised Study. Turk J Anaesthesiol Reanim 2020; 48: 17-23. [\[CrossRef\]](#)

**Saurabh Sharma<sup>1</sup>** , **Anju Romina Bhalotra<sup>2</sup>** , **Shikha Awal<sup>3</sup>** 

<sup>1</sup>Department of Neuroanaesthesia, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, Kerala, India

<sup>2</sup>Department of Anaesthesiology, Maulana Azad Medical College and associated Lok Nayak Hospital, New Delhi, India

<sup>3</sup>Department of Anaesthesiology, Kalinga Hospital, Bhubaneswar, Orissa, India

Corresponding Author: Anju Romina Bhalotra E-mail: drakgk@yahoo.co.in  
DOI: 10.5152/TJAR.2020.030520

©Copyright 2020 by Turkish Society of Anaesthesiology and Reanimation - Available online at [www.turkjanaesthesiolreanim.org](http://www.turkjanaesthesiolreanim.org)