

## RESEARCH ARTICLE

# Effect of benzo(a)pyrene on oxidative stress and inflammatory mediators in astrocytes and HIV-infected macrophages

Asit Kumar, Namita Sinha, Sunitha Kodidela, Lina Zhou , Udai P. Singh , Santosh Kumar \*

Department of Pharmaceutical Sciences, College of Pharmacy, The University of Tennessee Health Science Center, Memphis, TN, United States of America

\* [ksantosh@uthsc.edu](mailto:ksantosh@uthsc.edu)



## OPEN ACCESS

**Citation:** Kumar A, Sinha N, Kodidela S, Zhou L, Singh UP, Kumar S (2022) Effect of benzo(a)pyrene on oxidative stress and inflammatory mediators in astrocytes and HIV-infected macrophages. *PLoS ONE* 17(10): e0275874. <https://doi.org/10.1371/journal.pone.0275874>

**Editor:** Michal Toborek, University of Miami, UNITED STATES

**Received:** January 20, 2022

**Accepted:** September 23, 2022

**Published:** October 14, 2022

**Copyright:** © 2022 Kumar et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its [Supporting Information](#) files.

**Funding:** This study was funded by the NIH grants DA047178 and MH125670 to Dr. Santosh Kumar. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Abstract

### Background

Benzo(a)pyrene (BaP), an important polycyclic aromatic hydrocarbons (PAH) component of cigarette/tobacco smoking, is known to cause adverse health effects and is responsible for various life-threatening conditions including cancer. However, it is not yet clear whether BaP contributes to the macrophage- and astrocyte-mediated inflammatory response.

### Methods

We examined the acute (up to 72 h) effects of BaP on the expression of antioxidant enzymes (AOEs), cytokines/chemokines, and cytochromes P450 (CYP) enzymes in astrocytic cell lines, SVGA, and chronically HIV-infected U1 macrophage. The treated cells were examined for mRNA, protein levels of CYPs, AOEs superoxide dismutase-1 (SOD1) and catalase (CAT), cytokines/chemokines, using Western blot, multiplex ELISA, and reactive oxygen species (ROS) by flow cytometry analysis.

### Results

Upon acute exposure, BaP (1  $\mu$ M) showed a significant increase in the mRNA levels of CYPs (CYP1A1 and CYP1B1), and pro-inflammatory cytokine IL-1 $\beta$  in SVGA cells following BaP for 24, 48, and 72h. In addition, we observed a significant increase in the mRNA levels of SOD1 and CAT at 24h of BaP treatment. In contrast, BaP did not exert any change in the protein expression of AOEs and CYP enzymes. In U1 cells, however, we noticed an interesting increase in the levels of MCP-1 as well as a modest increase in TNF $\alpha$ , IL-8 and IL-1 $\beta$  levels observed at 72 h of BaP treatment but could not reach to statistically significant level.

### Conclusions

Overall, these results suggest that BaP contributes in part to macrophage and astrocyte-mediated neuroinflammation by mainly inducing IL-1 $\beta$  and MCP-1 production, which is likely to occur with the involvement of CYP and/or oxidative stress pathways.

## Introduction

Cigarette smoking is known to cause the accumulation of toxic metabolites and induction of oxidative stress and can contribute to the onset or progression of several diseases including chronic obstructive pulmonary disease (COPD) [1–4], and coronary heart disease (CHD) [5–7], and cancer [8–10]. Smoking can initiate a series of events that include induction of inflammatory cytokines/chemokines, altering antioxidant enzymes (AOEs), and metabolism of smoking constituents through dysregulation of cytochromes P450 (CYP) enzymes [11–13]. Nicotine is the most studied constituent of tobacco and one of the extensively used substances among substance or drug users [14]. In our previous studies, we have shown that nicotine induces oxidative stress through the CYP2A6-mediated pathway in U937 monocytic cells and SVGA cells, an immortalized astrocytic cell line [11, 15]. Moreover, BaP, a known carcinogenic polycyclic aromatic hydrocarbon (PAHs) present in cigarette smoke, has been shown to increase the expression of CYP1A and AOEs superoxide dismutase-1, (SOD1) and catalase (CAT), resulting in reactive oxygen species (ROS) generation and cytotoxicity in U937 cells [16].

Glial cells including astrocytes are important to brain cells that mediate effects in the brain upon tobacco smoking, perhaps via CYP and oxidative stress pathways, not only in HIV-infected but also in general populations [17, 18]. Astrocytes are not primarily infected with HIV but are possible through cell-cell contact and receptor-mediated endocytosis, however, they may become activated as a result of proximity to HIV-infected macrophages and microglia [19, 20]. SVGA cells have been extensively studied to show the effect of astrocyte activation and subsequent neuroinflammation [15, 21, 22]

Similar to inflammation, the expression of AOEs and CYPs might also be altered by smoking and tobacco-related products. Thus, there is a need to identify the constituent(s) of tobacco and the underlying mechanism that is responsible for smoking-associated inflammation and oxidative stress in astrocytes. We hypothesize that BaP, a major carcinogen in a cigarette can influence the expression of specific cytokines/chemokines and AOEs (SOD1 and CAT) in astrocytes, that may be associated with smoking-associated toxicity and progression of inflammatory conditions in smokers. Further, we have shown that BaP enhances HIV replication in differentiated U1 macrophages by a CYP-mediated oxidative stress pathway [23]. But to date, it is not yet clear whether BaP contributes to the macrophage and brain resident astrocyte-mediated inflammatory response. Therefore, the objective of this study was to determine the effect of BaP on the acute inflammatory and oxidative stress mediators in the astrocytic and HIV-infected differentiated U1 macrophages.

## Materials and methods

### Cell culture and treatment

The human SVGA astrocytic cell line was maintained as previously discussed [15]. In brief, SVGA cells were cultured in DMEM supplemented with 10% (v/v) Fetal bovine serum (FBS), 1% penicillin, and streptomycin (40 U/ml and 40 mg/ml, respectively) at 37°C with 5% CO<sub>2</sub>. Astrocytes were treated with BaP (0.01–1 μM, Millipore Sigma, Cat No. B1760) every 24 h for up to 72 h for acute treatment. A stock solution of BaP was prepared in 100% dimethyl sulfoxide (DMSO; Millipore Sigma). The final concentration of DMSO in the cell culture media was not more than 0.3%. BaP concentration was maintained constant at every treatment.

Chronically infected HIV-promonocytic (U1) cell lines were obtained from the NIH AIDS Reagent Program (Germantown, MD). In this study, we used U1, instead of uninfected U937 cells, because U937 cells are non-HIV and have already been characterized for oxidative stress

and inflammatory regulation in the presence of tobacco constituents [11, 16]. It's now important to study HIV-infected macrophages (U1), which are juxtaposed to astrocytic cells and could activate astrocytic cells during HIV infection in the brain. The cells were cultured in RPMI 1640 media containing 10% (v/v) FBS, 1% penicillin, and streptomycin (40 U/ml and 40 mg/ml, respectively) at 37°C with 5% CO<sub>2</sub> [23]. To differentiate the cells into macrophages, 0.8 million cells were seeded in 1.5 ml of media containing 100 nM phorbol 12-myristate 13-acetate (PMA) in each well of a 6-well plate. After 3 days, the media containing PMA and non-adherent cells were removed and the differentiated cells were washed with phosphate buffer saline. The cells were topped with fresh 1 ml media and treated with different concentrations of BaP (0.01–1 μM) every 24 h for up to 72 h for acute treatment. BaP concentration was maintained constant at every treatment.

### Western blotting

For western blot, cells were lysed with RIPA buffer and cellular protein was quantified using Pierce BCA Protein Assay Kit (ThermoFisher Scientific). Each sample was loaded onto 10% SDS-PAGE gel and the rest of the procedures were performed as described previously [13, 24]. We evaluated the expression of proteins associated with AOE (SOD1 and CAT), pro-inflammatory cytokine (IL-1β), and CYP enzymes (CYP1A1, CYP1B1, CYP2A6) and quantified by Western blot. Following SDS-PAGE, proteins were transferred onto PVDF membrane and blocked in Li-Cor blocking buffer (LI-COR Biosciences, Lincoln, NE) for 1 h. After blocking, membranes were incubated overnight at 4°C with primary antibodies, including SOD1 mouse mAb (1:500 dilution, catalog no. sc-101523, Santa Cruz Biotechnology), and CAT mouse mAb (1:500 dilution, catalog no. sc-365738, Santa Cruz Biotechnology), IL-1β rabbit pAb (1:1000 dilution, catalog no. 16806-1AP, Proteintech), CYP2A6 rabbit mAb (1:1000 dilution, catalog no. ab3570, Abcam), CYP1A1 mouse mAb (1:500 dilution, catalog no. sc-48432, Santa Cruz Biotechnology), CYP1B1 mouse mAb (1:500 dilution, catalog no. sc-374228, Santa Cruz Biotechnology), and β-Actin mouse mAb (1:1000 dilution, catalog no. 3700, Cell Signaling). The next day, after washing (three times for 5 min each in PBS containing 0.2% Tween 20), membranes were incubated with either goat anti-mouse or goat anti-rabbit secondary antibody (1:10000 dilution, LI-COR Biosciences) for 1 h at room temperature in the dark. After 1 h membranes were washed and scanned using the Li-Cor Scanner (LI-COR Biosciences). Densitometry analyses of the proteins were performed using LI-COR Image Studio Software (v.5.2, Nebraska, USA).

### LDH activity

The cytotoxic effect of BaP in U1 and SVGA cells was evaluated by LDH release using the CyQUANT™ LDH Cytotoxicity Assay Kit (Catalog no. C20300, ThermoFisher Scientific) according to the manufacturer's instruction. Briefly, U1 and SVGA cells were exposed to BaP (0.01–1 μM) for 24, 48, and 72 h. Cell culture supernatants were collected at indicated time points and transferred to a 96-well plate. The reaction mixture was added to the cell culture supernatant for 30 min at room temperature. Background absorbance 680 nm was subtracted from 490 nm absorbance to determine the LDH activity.

### Measurement of reactive oxygen species (ROS)

ROS was quantified by flow cytometry analysis using the fluorescence dye 5-(and-6)-chloromethyl 2',7'-dichlorodihydrofluorescein diacetate (CM-H<sub>2</sub>DCFDA) (ThermoFisher Scientific). The treated cells were thoroughly washed with PBS and resuspended in 5 μM of CM-H<sub>2</sub>DCFDA in PBS. The cells were then incubated at room temperature in the dark for 45

minutes in CM-H<sub>2</sub>DCFDA and subsequently washed and resuspended in 300  $\mu$ L of PBS. ROS generated in the cells was detected and data was analyzed using flow cytometry built-in software (Agilent NovoCyte).

### mRNA Real-Time PCR

RNA was isolated using an RNeasy mini kit (Qiagen). For cDNA synthesis, total RNA was reverse transcribed using the High-Capacity cDNA Reverse Transcription Kit (ThermoFisher Scientific) as previously described [25]. In brief, every 20 ml of reverse transcription (RT) reaction consisted of 2 ml of 10 RT buffer, 0.8 ml of 25 dNTP mix (100 mM), 2.0 ml of 10 RT random primers, 1.0 ml of MultiScribe reverse transcriptase, 4.2 ml of nuclease-free water, and 10 ml of the RNA sample. Thermal cycling conditions used for reverse transcription are 10 minutes at 25°C, 120 minutes at 37°C, and finally 5 minutes at 85°C followed by a stop at 4°C. The synthesized cDNA was the template for the real-time PCR amplification that was carried out by Quant Studio 6 Pro real-time PCR System (ThermoFisher Scientific), using TaqMan Universal Master Mix II, no UNG (ThermoFisher Scientific) as per the company's protocol. Reaction conditions were 10 minutes at 95°C, followed by 40 cycles of 15 seconds at 95°C, 60 seconds at 60°C, followed by the final plate read. The following primers were used in this study: Hs00533490\_m1 SOD1 (human), Hs00156308\_m1 CAT (human), Hs00868409\_s1 CYP2A6 (human), Hs01054794\_m1 CYP1A1 (human), Hs00164383\_m1 CYP1B1 (human), Hs01555410\_m1 IL-1 $\beta$  (human), and Hs99999903\_m1 Actin (human). Actin served as an internal control for sample normalization and the comparative cycle threshold method ( $2^{-\Delta\Delta C_t}$ ) was used for data quantification as described previously [26, 27].

### Multiplex ELISA

The levels of cytokines (pro-inflammatory: TNF- $\alpha$ , IL-1 $\beta$ , IL-6, IL-8, IL-18; anti-inflammatory: IL-1RA, IL-10) and chemokines (MCP-1 and RANTES) were measured in the cell culture supernatant using customized human 9-Plex ProcartaPlex<sup>TM</sup> multiplex immunoassay (ThermoFisher Scientific) as per previously described method [13]. Briefly, SVGA cells were treated with BaP (1  $\mu$ M) for 24, 48, and 72 h. Cell culture supernatants were collected at indicated time points and transferred to a 96-well ELISA plate. The samples and standards were incubated with a magnetic 96-well ELISA plate for 1 h at room temperature. Afterward, wells were washed, and subsequently, detection antibody, streptavidin-PE, and reading buffer were added followed by a plate read. Quantification was achieved as per the manufacturer's kit operating manual. The concentration of cytokines and chemokines was expressed as pg/ml.


### Statistical analysis

The data were analyzed using Graph-Pad version 5.0 (GraphPad Software Inc., San Diego, CA, USA). Values of all experiments are represented as mean  $\pm$  SEM of at least three independent experiments. Values were compared using Student's t-test (two groups) or one-way ANOVA with Tukey's test (multiple comparisons), when appropriate. The level of significance was set at \* $p \leq 0.05$ , \*\* $p \leq 0.01$ , \*\*\* $p \leq 0.001$ .

## Results

### Effect of BaP on AOE and IL-1 $\beta$ expression in U1 macrophages

Differentiated U1 cells were treated with different concentrations of BaP (0.01, 0.1, 1  $\mu$ M) every 24 h up to 72 h to determine the effect of BaP on AOE (SOD1 and CAT) and pro-inflammatory cytokine (IL-1 $\beta$ ) (Fig 1). Before analyzing AOE and IL-1 $\beta$ , we performed an



**Fig 1.** LDH activity and protein expression of AOE and cytokine IL-1 $\beta$  (A, B). (A) LDH activity was performed in U1 cells treated with different concentrations of BaP (daily treatment up to 72 h). (B) Representative Western blots of AOE (SOD1 and CAT) and IL-1 $\beta$  in U1 cells (Western blot at 72

h) treated with different concentrations of BaP (daily treatment up to day 3). Untreated controls were considered as 100% in the densitometry analysis. (C) Densitometry analysis. (D) mRNA expression was analyzed in U1 cells (at 24, 48, and 72 h) treated with 1  $\mu\text{M}$  concentration of BaP (daily treatment). Statistical analyses were carried out by using one-way ANOVA (multiple comparison) and Student's t-test (two groups). Results are expressed as means  $\pm$  S.E.M of  $n = 3-4$  experiments.

<https://doi.org/10.1371/journal.pone.0275874.g001>

LDH assay to analyze the cytotoxicity of BaP on U1 cells. We observed that treatment of U1 cells with BaP for 24, 48, and 72 h did not show a statistically significant increase in LDH activity (Fig 1A), indicating no detectable cytotoxicity with the selected doses of BaP. Next, we performed a Western blot and observed an upward trend in the protein expression of SOD1, CAT, and IL-1 $\beta$  with increasing concentrations of BaP at 72 h (Fig 1B and 1C, S1 Fig). However, none of these could reach a statistically significant level. To evaluate whether BaP exerts effects at the transcriptional level on SOD1, CAT, and IL-1 $\beta$ , U1 cells were treated with 1  $\mu\text{M}$  BaP for 24, 48, and 72 h (Fig 1D). Similar to protein expression, BaP did not significantly affect the SOD1, CAT, and IL-1 $\beta$  gene expression in U1 cells even at different time points.

### Effect of BaP on Reactive Oxygen Species (ROS) in U1 macrophages


To study the effect of BaP on ROS, U1 cells were treated with 1  $\mu\text{M}$  BaP for 24, 48, and 72 h. We noticed that 1  $\mu\text{M}$  does show maximum effect without any toxicity is the reason to use this dose. In the flow cytometry analysis, we observed a significant increase in the ROS with BaP treatment at 24 and 48 h (Fig 2A and 2B). However, there was not much effect on ROS with BaP concentration at 72 h time point. This might be due to the threshold level of ROS with time or other factors that required further investigations.

### Effect of BaP on cytokines/chemokines in U1 macrophages

To determine whether BaP alters the cytokines and chemokines, we measured the extracellular release of proinflammatory cytokines (IL-1 $\beta$ , TNF $\alpha$ , IL-6, IL-18), anti-inflammatory cytokines (IL-10, IL-1RA), and chemokines (IL-8, RANTES, MCP-1) in BaP (1  $\mu\text{M}$ ) treated U1 cells supernatant. Our results demonstrated not much change in the levels of cytokines and chemokines in U1 cells supernatants treated with BaP at 24, 48, and 72 h (Fig 3). However, we noticed an interesting increase in the levels of MCP-1 as well as a modest increase in TNF $\alpha$ , IL-8 and IL-1 $\beta$  levels observed at 72 h of BaP treatment but could not reach to statistically significant level.

### BaP induces IL-1 $\beta$ expression in SVGA cells

Having observed an upward trend in the expression of SOD1, CAT, and IL-1 $\beta$  upon treatment with BaP in the U1 macrophages, we set out to investigate changes of SOD1 and CAT involved in the clearance of oxidative stress and IL-1 $\beta$  involved in pro-inflammatory immune response induced by BaP in SVGA cells. We treated the SVGA cells with different concentrations of BaP (0.01, 0.1, 1  $\mu\text{M}$ ) every 24 h up to 72 h to determine the effect of BaP on AOE (SOD1 and CAT), and pro-inflammatory cytokine IL-1 $\beta$ . Before analyzing AOE and IL-1 $\beta$  in SVGA cells upon BaP treatment, we performed an LDH assay to analyze whether BaP causes cell cytotoxicity. We observed that treatment of SVGA cells with BaP for 24, 48, and 72 h did not show a significant increase in LDH activity when compared to untreated cells (Fig 4A), indicating no detectable cytotoxicity with the BaP in SVGA cells. Though not statistically significant, we observed an increase in the expression of SOD1 and CAT at a 1  $\mu\text{M}$  concentration of BaP (Fig 4B and 4C, S2 Fig). Notably, cellular expression of IL-1 $\beta$  was dose-dependently elevated by BaP treatment in SVGA cells. In addition, at 1  $\mu\text{M}$  concentration, BaP significantly increased




**Fig 2.** Effect of BaP on ROS in U1 cells (A, B). (A) The U1 cells were treated with 1  $\mu$ M BaP for 24, 48, and 72 h. ROS level using flow cytometry. (B) Data is quantified using fluorescent cell count that were measured in %. Statistical analyses were carried out by using Student's t-test (two groups each time point). Results are expressed as means  $\pm$  S.E.M of  $n = 3$  experiments. \*\* $P \leq 0.01$  in comparison with untreated control.

<https://doi.org/10.1371/journal.pone.0275874.g002>

IL-1 $\beta$  protein ( $P \leq 0.05$ ; ANOVA; Fig 4C) expression in SVGA cells when compared to untreated cells.

### Time-dependent effect of BaP on IL-1 $\beta$ , AOE, and CYP enzymes in SVGA cells

Having demonstrated that BaP dose-dependently induces intracellular levels of IL-1 $\beta$ , a cytokine associated with a pro-inflammatory immune response, we then set out to evaluate the




**Fig 3. Levels of cytokines and chemokines in BaP-treated U1 cells.** Multiplex ELISA analyses of pro-inflammatory cytokine (IL-1 $\beta$ , TNF $\alpha$ , IL-6, IL-18), anti-inflammatory cytokine (IL-10, IL-1RA), and chemokines (IL-8, RANTES, MCP-1) were performed in U1 cells (at 24, 48, and 72 h) treated with 1  $\mu$ M concentration of BaP (daily treatment). Statistical analyses were carried out by using Student's t-test. Results are expressed as means  $\pm$  S.E.M of  $n = 3-4$  experiments.

<https://doi.org/10.1371/journal.pone.0275874.g003>

changes in the IL-1 $\beta$  expression at mRNA and protein levels in SVGA cells at different time points. Moreover, we analyzed the mRNA and protein expression levels of AOE (SOD1 and CAT) and CYP enzymes (CYP2A6, CYP1A1, and CYP1B1), which are known to increase oxidative stress upon tobacco exposure, in SVGA cells at different time points (Fig 5A and 5B) after BaP (1  $\mu$ M) exposure (Fig 5A–5C). IL-1 $\beta$  mRNA expression was significantly upregulated in SVGA cells at 24 h ( $P \leq 0.001$ ; Student's t-test; Fig 5A), 48 h ( $P \leq 0.001$ ; Student's t-test; Fig 5A) and 72 h ( $P \leq 0.001$ ; Student's t-test; Fig 5A) post-BaP treatment when compared to untreated cells. Similarly, CYP1A1 and CYP1B1 mRNA expression was also increased after the






**Fig 4.** LDH activity and protein expression of AOE and cytokine IL-1 $\beta$  (A, B). (A) LDH activity was performed in SVGA cells treated with different concentrations of BaP (daily treatment up to 72 h). (B) Representative Western blots of AOE (SOD1 and CAT) and IL-1 $\beta$  in SVGA cells (Western blot at 72 h) treated with different concentrations of BaP (daily treatment up to day 3). Untreated controls were considered as 100% in the densitometry analysis. (C) Densitometry analysis. Statistical analyses were carried out by using one-way ANOVA. Results are expressed as means  $\pm$  S.E.M of  $n = 4$  experiments. \* $P \leq 0.05$ , in comparison with untreated control.

<https://doi.org/10.1371/journal.pone.0275874.g004>

BaP treatment at 24 h ( $P \leq 0.001$ ; Student's t-test; Fig 5A), 48 h ( $P \leq 0.001$ ; Student's t-test; Fig 5A) and 72 h ( $P \leq 0.001$ ; Student's t-test; Fig 5A). Whereas SOD1 and CAT mRNA expression was found to have notably increased at 24 h ( $P \leq 0.01$ ; Student's t-test; Fig 5A) only. However, CYP2A6 mRNA expression was not altered at any time point after the BaP treatment. Next, the protein expression of IL-1 $\beta$ , AOE, and CYP enzymes was measured in SVGA cells treated with BaP (1  $\mu$ M) at 24, 48, and 72 h (Fig 5B and 5C). IL-1 $\beta$  protein expression was significantly increased in SVGA cells at 48 h ( $P \leq 0.01$ ; Student's t-test; Fig 5B and 5C, S3 Fig) and 72 h ( $P \leq 0.01$ ; Student's t-test; Fig 5B and 5C, S3 Fig) post-BaP treatment when compared to untreated




**Fig 5.** mRNA and protein expression of CYP enzymes, AOE and cytokine IL-1 $\beta$  (A-C). (A) mRNA expression was analyzed in SVG cells (at 24, 48, and 72 h) treated with 1  $\mu$ M concentration of BaP (daily treatment). (B) Representative western blots of IL-1 $\beta$ , AOE (SOD1 and CAT), and CYP enzymes (CYP2A6, CYP1B1, CYP1A1) in SVG cells (Western blot at 24, 48, and 72 h) treated with 1  $\mu$ M concentration of BaP (daily treatment). Untreated controls were considered as 100% in the densitometry analysis. (C) Densitometry analysis. Statistical analyses were

carried out by using Student's t-test. Results are expressed as means ± S.E.M of n = 4 experiments. \*\*P ≤0.01, \*\*\*P ≤0.001 in comparison with untreated control.

<https://doi.org/10.1371/journal.pone.0275874.g005>

cells. Together, these data indicate that BaP exposure results in a significant increase in IL-1β expression in SVGA cells during the acute cellular phase. Moreover, we performed a western blot of AOE (SOD1 and CAT) and CYP enzymes (CYP2A6, CYP1A1, and CYP1B1) in SVGA cells after BaP treatment for different time points (Fig 5B and 5C, S3 Fig). However, none of these were altered after the BaP treatment observed up to 72 h.



**Fig 6.** Effect of BaP on ROS in SVG cells (A, B). (A) The SVG cells were treated with 1 μM BaP for 24, 48, and 72 h. ROS level using flow cytometry. (B) Data is quantified using fluorescent cell count that were measured in %. Statistical analyses were carried out by using Student's t-test (two groups each time point). Results are expressed as means ± S.E.M of n = 3 experiments.


<https://doi.org/10.1371/journal.pone.0275874.g006>

## Effect of BaP on Reactive Oxygen Species (ROS) in SVGA cells

Next, to examine the effect of BaP on ROS, SVGA cells were treated with 1  $\mu$ M BaP for 24, 48, and 72 h. We analyzed the level of ROS by flow cytometry analysis. In the flow cytometry analysis, we did not see any significant changes in the ROS with BaP treatment up to 72 h (Fig 6A and 6B).

## BaP induces differential effects on cytokines/chemokines in SVGA cells

Since IL-1 $\beta$  expression was enhanced in SVGA cells upon BaP treatment compared to control, we next investigated whether the release of other cytokines and chemokines also altered after BaP treatment. Several cytokines and chemokines have been reported to be altered in smokers with and without systemic and brain diseases [28, 29]. Therefore, we assessed nine cytokines/chemokines (IL-1 $\beta$ , IL-6, TNF $\alpha$ , IL-18, IL-8, IL-1RA, IL-10, RANTES, MCP-1) expression profiles in BaP (1  $\mu$ M) treated SVGA cells supernatant at 24, 48, and 72 h. We observed that BaP significantly elevated IL-1 $\beta$  levels at 48 h ( $P \leq 0.05$ ; Student's t-test; Fig 7) and 72 h ( $P \leq 0.01$ ; Student's t-test; Fig 7) when compared to respective control cells. Notably, IL-6 cytokine, which is known to play a dual role during infections, was robustly increased in SVGA cells upon BaP treatment at 24 h ( $P \leq 0.01$ ; Student's t-test; Fig 7) and 48 h ( $P \leq 0.05$ ; Student's t-test; Fig 7). However, at 72 h, the IL-6 level was not found statistically significant, although an



**Fig 7. Levels of cytokines and chemokines in BaP-treated SVGA cells.** Multiplex ELISA analyses of pro-inflammatory cytokine (IL-1 $\beta$ , IL-6, TNF $\alpha$ , IL-18), anti-inflammatory cytokine (IL-10), and chemokines (RANTES) were performed in SVG cells (at 24, 48, and 72 h) treated with 1  $\mu$ M concentration of BaP (daily treatment). Statistical analyses were carried out by using Student's t-test (two groups each time point). Results are expressed as means  $\pm$  S.E.M of n = 3–4 experiments. \* $P \leq 0.05$  and \*\* $P \leq 0.01$  in comparison with untreated control.

<https://doi.org/10.1371/journal.pone.0275874.g007>

upward trend was observed. Furthermore, exposure of BaP to SVGA cells did not significantly alter the levels of TNF $\alpha$ , IL-18, IL-10, and RANTES at either 24, 48, or 72 h when compared to respective controls. IL-8, IL-1RA, and MCP-1 levels were not found in the detectable range.

## Discussion

Our prior findings have shown that cigarette smoke condensate (CSC) exposure enhanced HIV viral load and oxidative stress in HIV-infected monocytes/macrophages [30, 31]. Further, we noticed an increase in HIV viral load, oxidative stress, and nicotine metabolism with possible involvement of CYP enzymes in *ex vivo* samples obtained from HIV-infected smokers [30]. Taken together, these studies suggest an association of CYP-mediated nicotine metabolism and subsequent oxidative stress with HIV replication in tobacco smokers. A comparison of oxidative stress levels after nicotine and CSC exposure revealed that the induction of ROS by CSC is much higher than ROS induction by nicotine [11, 15, 31], suggesting the involvement of other PAH components of CSC such as BaP in the induction of oxidative stress. We recently studied the chronic effect of BaP, an important PAH component of CSC, in HIV-infected monocytes/macrophages cells and observed that BaP increases CYP1A1 expression, ROS levels, and cytotoxicity [16, 23]. The excessive ROS production caused by BaP likely disturbed the redox homeostasis, causing oxidative stress, which resulted in cytotoxicity in U937 cells [16]. Furthermore, we demonstrated that oxidative stress generated by the CYP1A1-mediated metabolism of BaP, triggers the redox-sensitive transcription factor, NF- $\kappa$ B that leads to HIV replication [16]. In the present study, we have demonstrated the effect of BaP on the AOE and cytokines/chemokines as inflammatory mediators in U1 and macrophages neighboring SVGA cells. Overall, our findings suggest that BaP can enhance ROS in U1 macrophages and inflammatory modulators, in particular, IL-1 $\beta$  in SVGA cells, which may potentially contribute to the tobacco/smoking-mediated neuroinflammation.

Upon exposure to smoking/tobacco products, cells activate their antioxidant mechanism to counteract oxidative stress [32, 33]. Oxidative stress induced due to an imbalance of free radicals/ROS and antioxidants has been implicated in the pathogenesis and progression of inflammatory diseases and HIV [34, 35]. Oxidative stress is alleviated by activation of an antioxidant defense mechanism that involves activation of antioxidant-related pathways, production of non-enzymatic antioxidants, and induction of antioxidant enzymes [36]. Activation of the transcription factor Nrf2 via the Keap1-Nrf2-ARE signaling pathway regulates the basal and inducible expression of AOE (SOD, CAT, GPx) and counteract ROS levels to maintain physiological homeostasis [36]. Although under normal physiological conditions, these AOE exert protective function against the adverse effects of xenobiotics, any perturbations in AOE function or ROS generation could lead to oxidative stress [37]. In many pathophysiological conditions, including cigarette smoking, excessive production of reactive species may accelerate the inactivation of nitric oxide and cause endothelial dysfunction that leads to inflammation, loss of vasodilation, platelet aggregation, and impaired modulation of vascular growth, and dysregulation of vascular remodeling [38]. While at the physiological level, ROS participates in various cellular processes including adaptation to hypoxia and regulation of autophagy, immunity, differentiation, and longevity [39]. Antioxidant functions are mainly dominated by three key enzymes SOD1, CAT, and glutathione peroxidase [40, 41]. In our previous studies [16, 23], we used low (100 nM for 7 days) and high (1  $\mu$ M for 3 days) concentrations of BaP to study its effects on HIV replication and oxidative stress pathway in U1 and U937 cells. We showed that while chronic exposure to BaP at 100nM induced HIV replication in U1 cells, it failed to induce the oxidative stress mediators SOD1 and CAT at mRNA and protein levels.

In the present study, we investigated the acute inflammatory effects of up to 1  $\mu\text{M}$  BaP in U1 macrophages and SVGA cells. Several prior *in vitro* and *in vivo* studies have used BaP concentration, which is comparable to or higher than 1  $\mu\text{M}$  [23, 42–47]. Although 1  $\mu\text{M}$  is higher than the physiological BaP brain concentration in humans, relatively high concentrations are routinely used to determine acute effects of BaP. A relatively high BaP concentration in acute treatment may correlate to near physiological BaP concentration in a particular region of the brain in heavy and chronic cigarette smokers. Therefore, the concentration of BaP 0.01–1  $\mu\text{M}$  up to 3 days (acute treatment) was chosen in our study to investigate the effects of BaP on oxidative stress and inflammatory mediators in U1 and SVGA cells. Moreover, BaP 1  $\mu\text{M}$  concentration is toxic to the cells when treated for 7 days. Consistent with the previous findings, even at 1  $\mu\text{M}$  BaP for 3 days, we observed no significant alteration in the mRNA expression of SOD1 and CAT in U1 macrophages. However, an upward trend was observed in the protein expression of SOD1 and CAT, suggesting mRNA decay or post-transcriptional modification. Possibly due to this increase in protein levels, we observed BaP-mediated production of ROS at early time points in U1 macrophages. In SVGA cells, we observed an increase in the mRNA expression of SOD1 and CAT at an early time point after BaP exposure, no significant changes were observed in the protein expression of SOD1 and CAT, which may be the reason associated with no effect on the production of ROS in SVGA. More, importantly, unlike in U1 cells, a significant increase in the IL-1 $\beta$  at both mRNA and protein levels in the SVGA astrocyte was observed. Our results suggest that BaP exposure in SVGA cells induces notable oxidative stress, and produces a strong pro-inflammatory activation state, similar to the immune response after a xenobiotic encounter in the brain. The results also suggest that astrocytic cells are relatively more susceptible to tobacco constituent, BaP, compared to macrophages.

Proinflammatory cytokines are believed to drive the neuroinflammatory events in many neurodegenerative diseases including HIV-associated neurocognitive disorder [48–50]. Cytokines including IL-1 levels have been found to be elevated in sera and monocytes/macrophages in HIV patients, and in the cerebrospinal fluid (CSF) of HIV patients suffering from HAND [51, 52]. During HIV infection, IL-1 $\beta$  and other cytokines has been induced by HIV accessory protein TAT, in T cells and monocytes/macrophages, major target cells for HIV in the blood [53, 54]. Exposure of BaP induces ROS and IL-8 production in normal human epidermal keratinocytes through the activation of the aryl hydrocarbon receptor signaling pathway [55], whereas the production of IL-1 $\alpha$ , IL-6, TNF- $\alpha$ , and GM-CSF was not affected by the BaP. Another study also suggests that BaP exacerbates colonic damage via inducing TNF- $\alpha$  and IL-1 $\beta$  expression in the mucosa of mice challenged with BaP (62.5mg/kg-250mg/kg) when compared with control [56]. In our study, IL-1 $\beta$  produced from the BaP (1  $\mu\text{M}$ ) treated SVGA cells was found to be increased substantially than untreated cells. The mRNA expression was found significantly increased at 24, 48, and 72 h, whereas the magnitude of IL-1 $\beta$  protein expression at intracellular levels was found significantly increased at 48 h and 72 h post-treatment with BaP, suggesting that BaP treatment causes upregulation of the expression of IL-1 $\beta$ . These data clearly show that following BaP treatment, IL-1 $\beta$  is induced within SVGA cells which results in an increased inflammatory milieu inside the cells. Though, the expression of SOD1 and CAT were not found significantly altered at any time point of investigation.

The current study also investigated whether BaP-induced pro-inflammatory cytokine IL-1 $\beta$  in SVGA cells is consistent with the induction of CYP enzymes *in vitro*. In a prior study, we have shown that CSC exposure increases oxidative stress and induction of CYP enzymes in U937 and/or U1 macrophages [31]. CYP enzyme, in particular, CYP2A6 was shown to be involved in nicotine metabolism and oxidative stress in U937 cells [11]. These studies suggested the involvement of CYP enzymes in the production of oxidation stress and increased HIV viral load in HIV-infected smokers. A prior study also showed that BaP (5  $\mu\text{M}$ ) increases

the expression of CYP enzymes (CYP1A1 and CYP1B1), production of ROS, and cytotoxicity in lung epithelial cells (BEAS-2B cells) after 24 h treatment [57]. It has been earlier shown that CYP enzymes regulate the production of ROS via several mechanisms that include regulation of gene transcription and modulation of interactions between protein components of the p450 monooxygenase resulting in its poor activity, coupling, and stability [58]. We earlier studied the effect of BaP in U937 and/or primary monocytic cells and observed that BaP (100nM) increases cytotoxicity, which is likely through the CYP1A1 and oxidative stress mediators [16]. Recently, we investigated whether BaP affects HIV viral load and induced the oxidative stress pathway via the CYP pathway *in vitro* [23] and observed that BaP enhances the HIV replication in macrophages by a CYP-mediated oxidative stress pathway followed by the NF- $\kappa$ B pathway. Furthermore, a significant increase was observed in the gene expression of CYP1A1 in addition to enhanced ROS production and cytotoxicity in U1 cells. These findings suggest that BaP-induced ROS occurs via CYP1A1-mediated metabolic activation of BaP [23]. A study performed in a human lung cell line also indicated a significant role of CYP1A1 in the formation of BaP to its hydrolyzed carcinogenic form BaP-diol-epoxides, while both CYP1A1 and CYP1B1 were found to be involved in the metabolism of BaP [59]. Therefore, in the present study, we chose to investigate the BaP effect on the induction of CYP enzymes. The expression of CYP2A6, CYP1A1, and CYP1B1, which are the major tobacco metabolizing CYPs, were examined in SVGA cells after BaP (1  $\mu$ M) exposure daily for 72 h. The result did not show significant changes in the mRNA and protein expression of CYP2A6. At the mRNA level, we did see significant changes in the expression of CYP enzymes (CYP1A1 and CYP1B1) in SVGA cells upon BaP treatment, which are potentially associated with the BaP effects on IL-1 $\beta$ . However, we did not observe the significant expression of CYP1A1 and CYP1B1 at protein levels, perhaps due to post translational modifications or instability of the protein after extraction. The discrepancy in these results with our earlier investigation could be due to the use of different, monocytic (U937 and U1) and astrocytic (SVGA) cell lines. Chi et al, in their study, showed similar CYP1A1/CYP1B1 induction or preferential CYP1A1 induction in tissue-specific tobacco-related squamous cell carcinoma cell lines [60]. In contrast to their investigation of CYP enzymes in cell lines, their investigation of gingival tissue revealed preferential induction of CYP1B1 among smokers [60]. Thus, monocyte and SVGA astrocytes may show differential inflammatory responses and expression of CYP enzymes in response to BaP exposure. In the present study though not at the protein level, we did see an increase in mRNA levels of CYP1A1 and CYP1B1 possibly suggesting involvement of CYPs in the BaP-induced IL-1 $\beta$  production in SVGA cells. There are no changes observed on ROS upon BaP treatment in SVGA cells possibly associated with no change in the levels of AOE at protein levels. Other than the expected release of IL-1 $\beta$  upon BaP treatment in SVGA cells, the release of other cytokines mainly IL-6 was also found to be increased. A prior study showed that CYP1A1 overexpression augmented TNF- $\alpha$  and IL-6 production in RAW264.7 cells by enhancing JNK/AP-1 signaling [61]. Other studies also suggested the involvement of JNK/AP-1/ATF2 and NF- $\kappa$ B/IKK- $\beta$  signaling in the production of IL-6 in microglia and monocytes/macrophages, respectively [62–64]. However, further study is needed to examine the mechanism of IL-1 $\beta$  and IL-6 production in the context of BaP and SVGA cells. Future studies could be conducted on other glial cells to determine the comprehensive role of BaP on neuroinflammation. Overall, in our study we showed the effect of BaP on the astrocytes and macrophages that BaP predominantly induce IL-1 $\beta$  production and presents clinical application of cytokine therapy in combination with cART that may play an important role in offering new strategies for modulation and treatment of smoking-mediated HIV pathogenesis and HIV-associated neuropathogenesis.

## Supporting information

**S1 Fig. Original western blot images for Fig 1B.**  
(PDF)

**S2 Fig. Original western blot images for Fig 4B.**  
(PDF)

**S3 Fig. Original western blot images for Fig 5B.**  
(PDF)

## Acknowledgments

The authors are grateful to Krina Vipul Patel (University of Tennessee Health Science center) for critical reading and editing of the manuscript.

## Author Contributions

**Conceptualization:** Santosh Kumar.

**Data curation:** Asit Kumar, Namita Sinha, Sunitha Kodidela, Lina Zhou, Udai P. Singh.

**Formal analysis:** Asit Kumar, Udai P. Singh, Santosh Kumar.

**Funding acquisition:** Santosh Kumar.

**Supervision:** Santosh Kumar.

**Writing – original draft:** Asit Kumar.

**Writing – review & editing:** Namita Sinha, Sunitha Kodidela, Lina Zhou, Udai P. Singh, Santosh Kumar.

## References

1. King PT. Inflammation in chronic obstructive pulmonary disease and its role in cardiovascular disease and lung cancer. *Clin Transl Med.* 2015; 4(1):68. Epub 2015/07/30. <https://doi.org/10.1186/s40169-015-0068-z> PMID: 26220864; PubMed Central PMCID: PMC4518022.
2. Tetley TD. Inflammatory cells and chronic obstructive pulmonary disease. *Curr Drug Targets Inflamm Allergy.* 2005; 4(6):607–18. Epub 2007/02/20. <https://doi.org/10.2174/156801005774912824> PMID: 17305517.
3. Willemse BW, ten Hacken NH, Rutgers B, Lesman-Leegte IG, Postma DS, Timens W. Effect of 1-year smoking cessation on airway inflammation in COPD and asymptomatic smokers. *Eur Respir J.* 2005; 26(5):835–45. Epub 2005/11/03. <https://doi.org/10.1183/09031936.05.00108904> PMID: 16264044.
4. Ind PW. COPD disease progression and airway inflammation: uncoupled by smoking cessation. *Eur Respir J.* 2005; 26(5):764–6. Epub 2005/11/03. <https://doi.org/10.1183/09031936.05.00102805> PMID: 16264034.
5. Stallones RA. The association between tobacco smoking and coronary heart disease. *Int J Epidemiol.* 2015; 44(3):735–43. Epub 2015/07/16. <https://doi.org/10.1093/ije/dyv124> PMID: 26174518.
6. Oshunbade AA, Kassahun-Yimer W, Valle KA, Hamid A, Kipchumba RK, Kamimura D, et al. Cigarette Smoking, Incident Coronary Heart Disease, and Coronary Artery Calcification in Black Adults: The Jackson Heart Study. *J Am Heart Assoc.* 2021; 10(7):e017320. Epub 2021/03/24. <https://doi.org/10.1161/JAHA.120.017320> PMID: 33754833; PubMed Central PMCID: PMC8174312.
7. Salehi N, Janjani P, Tadbiri H, Rozbahani M, Jalilian M. Effect of cigarette smoking on coronary arteries and pattern and severity of coronary artery disease: a review. *J Int Med Res.* 2021; 49(12):3000605211059893. Epub 2021/12/03. <https://doi.org/10.1177/03000605211059893> PMID: 34855538; PubMed Central PMCID: PMC8647272.
8. Blot WJ, McLaughlin JK, Winn DM, Austin DF, Greenberg RS, Preston-Martin S, et al. Smoking and drinking in relation to oral and pharyngeal cancer. *Cancer Res.* 1988; 48(11):3282–7. Epub 1988/06/01. PMID: 3365707.



9. Loeb LA, Ernster VL, Warner KE, Abbotts J, Laszlo J. Smoking and lung cancer: an overview. *Cancer Res.* 1984; 44(12 Pt 1):5940–58. Epub 1984/12/01. PMID: [6388830](#).
10. Begum S. Molecular changes in smoking-related lung cancer. *Expert Rev Mol Diagn.* 2012; 12(1):93–106. Epub 2011/12/03. <https://doi.org/10.1586/erm.11.84> PMID: [22133122](#).
11. Jin M, Earla R, Shah A, Earla RL, Gupte R, Mitra AK, et al. A LC-MS/MS method for concurrent determination of nicotine metabolites and role of CYP2A6 in nicotine metabolism in U937 macrophages: implications in oxidative stress in HIV + smokers. *J Neuroimmune Pharmacol.* 2012; 7(1):289–99. Epub 2011/06/10. <https://doi.org/10.1007/s11481-011-9283-6> PMID: [21655912](#); PubMed Central PMCID: [PMC4118724](#).
12. Haque S, Kodidela S, Sinha N, Kumar P, Cory TJ, Kumar S. Differential packaging of inflammatory cytokines/ chemokines and oxidative stress modulators in U937 and U1 macrophages-derived extracellular vesicles upon exposure to tobacco constituents. *PLoS One.* 2020; 15(5):e0233054. Epub 2020/05/21. <https://doi.org/10.1371/journal.pone.0233054> PMID: [32433651](#); PubMed Central PMCID: [PMC7239484](#).
13. Kumar A, Sinha N, Haque S, Kodidela S, Wang T, Martinez AG, et al. Nicotine self-administration with menthol and audiovisual cue facilitates differential packaging of CYP2A6 and cytokines/chemokines in rat plasma extracellular vesicles. *Sci Rep.* 2021; 11(1):17393. Epub 2021/09/01. <https://doi.org/10.1038/s41598-021-96807-5> PMID: [34462474](#); PubMed Central PMCID: [PMC8405708](#).
14. How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General. Publications and Reports of the Surgeon General. Atlanta (GA) 2010.
15. Ande A, Earla R, Jin M, Silverstein PS, Mitra AK, Kumar A, et al. An LC-MS/MS method for concurrent determination of nicotine metabolites and the role of CYP2A6 in nicotine metabolite-mediated oxidative stress in SVGA astrocytes. *Drug Alcohol Depend.* 2012; 125(1–2):49–59. Epub 2012/04/14. <https://doi.org/10.1016/j.drugalcdep.2012.03.015> PMID: [22498344](#); PubMed Central PMCID: [PMC3413753](#).
16. Ranjit S, Midde NM, Sinha N, Patters BJ, Rahman MA, Cory TJ, et al. Effect of Polyaryl Hydrocarbons on Cytotoxicity in Monocytic Cells: Potential Role of Cytochromes P450 and Oxidative Stress Pathways. *PLoS One.* 2016; 11(9):e0163827. Epub 2016/09/30. <https://doi.org/10.1371/journal.pone.0163827> PMID: [27684561](#); PubMed Central PMCID: [PMC5042547](#).
17. Shah A, Kumar S, Simon SD, Singh DP, Kumar A. HIV gp120- and methamphetamine-mediated oxidative stress induces astrocyte apoptosis via cytochrome P450 2E1. *Cell Death Dis.* 2013; 4:e850. Epub 2013/10/12. <https://doi.org/10.1038/cddis.2013.374> PMID: [24113184](#); PubMed Central PMCID: [PMC3824683](#).
18. Ande A, McArthur C, Kumar A, Kumar S. Tobacco smoking effect on HIV-1 pathogenesis: role of cytochrome P450 isozymes. *Expert Opin Drug Metab Toxicol.* 2013; 9(11):1453–64. Epub 2013/07/05. <https://doi.org/10.1517/17425255.2013.816285> PMID: [23822755](#); PubMed Central PMCID: [PMC4007120](#).
19. Borrajo A, Spuch C, Penedo MA, Olivares JM, Agis-Balboa RC. Important role of microglia in HIV-1 associated neurocognitive disorders and the molecular pathways implicated in its pathogenesis. *Ann Med.* 2021; 53(1):43–69. Epub 2020/08/26. <https://doi.org/10.1080/07853890.2020.1814962> PMID: [32841065](#); PubMed Central PMCID: [PMC7877929](#).
20. Lutgen V, Narasipura SD, Barbian HJ, Richards M, Wallace J, Razmpour R, et al. HIV infects astrocytes in vivo and egresses from the brain to the periphery. *PLoS Pathog.* 2020; 16(6):e1008381. Epub 2020/06/12. <https://doi.org/10.1371/journal.ppat.1008381> PMID: [32525948](#); PubMed Central PMCID: [PMC7289344](#).
21. Cao L, Fu M, Kumar S, Kumar A. Methamphetamine potentiates HIV-1 gp120-mediated autophagy via Beclin-1 and Atg5/7 as a pro-survival response in astrocytes. *Cell Death Dis.* 2016; 7(10):e2425. Epub 2016/10/21. <https://doi.org/10.1038/cddis.2016.317> PMID: [27763640](#); PubMed Central PMCID: [PMC5133984](#).
22. Cao L, Walker MP, Vaidya NK, Fu M, Kumar S, Kumar A. Cocaine-Mediated Autophagy in Astrocytes Involves Sigma 1 Receptor, PI3K, mTOR, Atg5/7, Beclin-1 and Induces Type II Programmed Cell Death. *Mol Neurobiol.* 2016; 53(7):4417–30. Epub 2015/08/06. <https://doi.org/10.1007/s12035-015-9377-x> PMID: [26243186](#); PubMed Central PMCID: [PMC4744147](#).
23. Ranjit S, Sinha N, Kodidela S, Kumar S. Benzo(a)pyrene in Cigarette Smoke Enhances HIV-1 Replication through NF-kappaB Activation via CYP-Mediated Oxidative Stress Pathway. *Sci Rep.* 2018; 8(1):10394. Epub 2018/07/12. <https://doi.org/10.1038/s41598-018-28500-z> PMID: [29991690](#); PubMed Central PMCID: [PMC6039513](#).
24. Kodidela S, Gerth K, Sinha N, Kumar A, Kumar P, Kumar S. Circulatory Astrocyte and Neuronal EVs as Potential Biomarkers of Neurological Dysfunction in HIV-Infected Subjects and Alcohol/Tobacco Users. *Diagnostics (Basel).* 2020; 10(6). Epub 2020/06/03. <https://doi.org/10.3390/diagnostics10060349> PMID: [32481515](#); PubMed Central PMCID: [PMC7345258](#).

25. Kumar A, Henry RJ, Stoica BA, Loane DJ, Abulwerdi G, Bhat SA, et al. Neutral Sphingomyelinase Inhibition Alleviates LPS-Induced Microglia Activation and Neuroinflammation after Experimental Traumatic Brain Injury. *J Pharmacol Exp Ther*. 2019; 368(3):338–52. Epub 2018/12/20. <https://doi.org/10.1124/jpet.118.253955> PMID: 30563941; PubMed Central PMCID: PMC6367691.
26. Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method. *Nat Protoc*. 2008; 3(6):1101–8. Epub 2008/06/13. <https://doi.org/10.1038/nprot.2008.73> PMID: 18546601.
27. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-</sup>(Delta Delta C(T)) Method. *Methods*. 2001; 25(4):402–8. Epub 2002/02/16. <https://doi.org/10.1006/meth.2001.1262> PMID: 11846609.
28. Strzelak A, Ratajczak A, Adamiec A, Feleszko W. Tobacco Smoke Induces and Alters Immune Responses in the Lung Triggering Inflammation, Allergy, Asthma and Other Lung Diseases: A Mechanistic Review. *Int J Environ Res Public Health*. 2018; 15(5). Epub 2018/06/09. <https://doi.org/10.3390/ijerph15051033> PMID: 29883409; PubMed Central PMCID: PMC5982072.
29. Sgambato JA, Jones BA, Caraway JW, Prasad GL. Inflammatory profile analysis reveals differences in cytokine expression between smokers, moist snuff users, and dual users compared to non-tobacco consumers. *Cytokine*. 2018; 107:43–51. Epub 2017/12/09. <https://doi.org/10.1016/j.cyto.2017.11.013> PMID: 29217402.
30. Ande A, McArthur C, Ayuk L, Awasom C, Achu PN, Njinda A, et al. Effect of mild-to-moderate smoking on viral load, cytokines, oxidative stress, and cytochrome P450 enzymes in HIV-infected individuals. *PLoS One*. 2015; 10(4):e0122402. Epub 2015/04/17. <https://doi.org/10.1371/journal.pone.0122402> PMID: 25879453; PubMed Central PMCID: PMC4399877.
31. Rao P, Ande A, Sinha N, Kumar A, Kumar S. Effects of Cigarette Smoke Condensate on Oxidative Stress, Apoptotic Cell Death, and HIV Replication in Human Monocytic Cells. *PLoS One*. 2016; 11(5): e0155791. Epub 2016/05/21. <https://doi.org/10.1371/journal.pone.0155791> PMID: 27203850; PubMed Central PMCID: PMC4874604.
32. Kamceva G, Arsova-Sarafinovska Z, Ruskovska T, Zdravkovska M, Kamceva-Panova L, Stikova E. Cigarette Smoking and Oxidative Stress in Patients with Coronary Artery Disease. *Open Access Maced J Med Sci*. 2016; 4(4):636–40. Epub 2016/12/29. <https://doi.org/10.3889/oamjms.2016.117> PMID: 28028404; PubMed Central PMCID: PMC5175512.
33. Foronjy R D'Armiento J. The Effect of Cigarette Smoke-derived Oxidants on the Inflammatory Response of the Lung. *Clin Appl Immunol Rev*. 2006; 6(1):53–72. Epub 2006/01/01. <https://doi.org/10.1016/j.cair.2006.04.002> PMID: 23997664; PubMed Central PMCID: PMC3755630.
34. Ivanov AV, Valuev-Elliston VT, Ivanova ON, Kochetkov SN, Starodubova ES, Bartosch B, et al. Oxidative Stress during HIV Infection: Mechanisms and Consequences. *Oxid Med Cell Longev*. 2016; 2016:8910396. Epub 2016/11/11. <https://doi.org/10.1155/2016/8910396> PMID: 27829986; PubMed Central PMCID: PMC5088339.
35. Uttara B, Singh AV, Zamboni P, Mahajan RT. Oxidative stress and neurodegenerative diseases: a review of upstream and downstream antioxidant therapeutic options. *Curr Neuropharmacol*. 2009; 7(1):65–74. Epub 2009/09/02. <https://doi.org/10.2174/157015909787602823> PMID: 19721819; PubMed Central PMCID: PMC2724665.
36. Lee S, Hu L. Nrf2 activation through the inhibition of Keap1-Nrf2 protein-protein interaction. *Med Chem Res*. 2020; 29(5):846–67. Epub 2020/05/12. <https://doi.org/10.1007/s00044-020-02539-y> PMID: 32390710; PubMed Central PMCID: PMC7207041.
37. Mates JM. Effects of antioxidant enzymes in the molecular control of reactive oxygen species toxicology. *Toxicology*. 2000; 153(1–3):83–104. Epub 2000/11/25. [https://doi.org/10.1016/s0300-483x\(00\)00306-1](https://doi.org/10.1016/s0300-483x(00)00306-1) PMID: 11090949.
38. Cai H, Harrison DG. Endothelial dysfunction in cardiovascular diseases: the role of oxidant stress. *Circ Res*. 2000; 87(10):840–4. Epub 2000/11/14. <https://doi.org/10.1161/01.res.87.10.840> PMID: 11073878.
39. Sena LA, Chandel NS. Physiological roles of mitochondrial reactive oxygen species. *Mol Cell*. 2012; 48(2):158–67. Epub 2012/10/30. <https://doi.org/10.1016/j.molcel.2012.09.025> PMID: 23102266; PubMed Central PMCID: PMC3484374.
40. Ighodaro OM, Akinloye OA. First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria Journal of Medicine*. 2019; 54(4):287–93. <https://doi.org/10.1016/j.ajme.2017.09.001>
41. Lei XG, Zhu JH, Cheng WH, Bao Y, Ho YS, Reddi AR, et al. Paradoxical Roles of Antioxidant Enzymes: Basic Mechanisms and Health Implications. *Physiol Rev*. 2016; 96(1):307–64. Epub 2015/12/19. <https://doi.org/10.1152/physrev.00010.2014> PMID: 26681794; PubMed Central PMCID: PMC4839492.

42. Stephanou P, Konstandi M, Pappas P, Marselos M. Alterations in central monoaminergic neurotransmission induced by polycyclic aromatic hydrocarbons in rats. *Eur J Drug Metab Pharmacokinet.* 1998; 23(4):475–81. Epub 1999/05/14. <https://doi.org/10.1007/BF03189998> PMID: 10323330.
43. Myers JN, Harris KL, Rekhadevi PV, Pratap S, Ramesh A. Benzo(a)pyrene-induced cytotoxicity, cell proliferation, DNA damage, and altered gene expression profiles in HT-29 human colon cancer cells. *Cell Biol Toxicol.* 2021; 37(6):891–913. Epub 2021/01/08. <https://doi.org/10.1007/s10565-020-09579-5> PMID: 33411230.
44. Moir D, Viau A, Chu I, Withey J, McMullen E. Pharmacokinetics of benzo[a]pyrene in the rat. *J Toxicol Environ Health A.* 1998; 53(7):507–30. Epub 1998/04/30. <https://doi.org/10.1080/009841098159114> PMID: 9561966.
45. Das M, Seth PK, Mukhtar H. Distribution of benzo(a)pyrene in discrete regions of rat brain. *Bull Environ Contam Toxicol.* 1985; 35(4):500–4. Epub 1985/10/01. <https://doi.org/10.1007/BF01636545> PMID: 4052652.
46. Das L, Patel B, Patri M. Adolescence benzo[a]pyrene treatment induces learning and memory impairment and anxiolytic like behavioral response altering neuronal morphology of hippocampus in adult male Wistar rats. *Toxicol Rep.* 2019; 6:1104–13. Epub 2019/11/14. <https://doi.org/10.1016/j.toxrep.2019.10.014> PMID: 31720231; PubMed Central PMCID: PMC6838974.
47. Appel BR, Guirguis G, Kim IS, Garbin O, Fracchia M, Flessel CP, et al. Benzene, benzo(a)pyrene, and lead in smoke from tobacco products other than cigarettes. *Am J Public Health.* 1990; 80(5):560–4. Epub 1990/05/01. <https://doi.org/10.2105/ajph.80.5.560> PMID: 2327532; PubMed Central PMCID: PMC1404640.
48. Wang XM, Zhang YG, Li AL, Long ZH, Wang D, Li XX, et al. Relationship between levels of inflammatory cytokines in the peripheral blood and the severity of depression and anxiety in patients with Parkinson's disease. *Eur Rev Med Pharmacol Sci.* 2016; 20(18):3853–6. Epub 2016/10/14. PMID: 27735031.
49. Sciacca FL, Ferri C, Licastro F, Veglia F, Biunno I, Gavazzi A, et al. Interleukin-1B polymorphism is associated with age at onset of Alzheimer's disease. *Neurobiol Aging.* 2003; 24(7):927–31. Epub 2003/08/21. [https://doi.org/10.1016/s0197-4580\(03\)00011-3](https://doi.org/10.1016/s0197-4580(03)00011-3) PMID: 12928052.
50. Griffin WS, Mrak RE. Interleukin-1 in the genesis and progression of and risk for development of neuronal degeneration in Alzheimer's disease. *J Leukoc Biol.* 2002; 72(2):233–8. Epub 2002/08/01. PMID: 12149413; PubMed Central PMCID: PMC3835694.
51. Scott-Algara D, Vuillier F, Marasescu M, de Saint Martin J, Dighiero G. Serum levels of IL-2, IL-1 alpha, TNF-alpha, and soluble receptor of IL-2 in HIV-1-infected patients. *AIDS Res Hum Retroviruses.* 1991; 7(4):381–6. Epub 1991/04/01. <https://doi.org/10.1089/aid.1991.7.381> PMID: 1676897.
52. Kreuzer KA, Dayer JM, Rockstroh JK, Sauerbruch T, Spengler U. The IL-1 system in HIV infection: peripheral concentrations of IL-1beta, IL-1 receptor antagonist and soluble IL-1 receptor type II. *Clin Exp Immunol.* 1997; 109(1):54–8. Epub 1997/07/01. <https://doi.org/10.1046/j.1365-2249.1997.4181315.x> PMID: 9218824; PubMed Central PMCID: PMC1904717.
53. Yang Y, Wu J, Lu Y. Mechanism of HIV-1-TAT induction of interleukin-1beta from human monocytes: Involvement of the phospholipase C/protein kinase C signaling cascade. *J Med Virol.* 2010; 82(5):735–46. Epub 2010/03/26. <https://doi.org/10.1002/jmv.21720> PMID: 20336759; PubMed Central PMCID: PMC2849807.
54. Ben Haij N, Planes R, Leghmari K, Serrero M, Delobel P, Izopet J, et al. HIV-1 Tat Protein Induces Production of Proinflammatory Cytokines by Human Dendritic Cells and Monocytes/Macrophages through Engagement of TLR4-MD2-CD14 Complex and Activation of NF-kappaB Pathway. *PLoS One.* 2015; 10(6):e0129425. Epub 2015/06/20. <https://doi.org/10.1371/journal.pone.0129425> PMID: 26090662; PubMed Central PMCID: PMC4474861.
55. Tsuji G, Takahara M, Uchi H, Takeuchi S, Mitoma C, Moroi Y, et al. An environmental contaminant, benzo(a)pyrene, induces oxidative stress-mediated interleukin-8 production in human keratinocytes via the aryl hydrocarbon receptor signaling pathway. *J Dermatol Sci.* 2011; 62(1):42–9. Epub 2011/02/15. <https://doi.org/10.1016/j.jdermsci.2010.10.017> PMID: 21316925.
56. Ajayi BO, Adedara IA, Farombi EO. Benzo(a)pyrene induces oxidative stress, pro-inflammatory cytokines, expression of nuclear factor-kappa B and deregulation of wnt/beta-catenin signaling in colons of BALB/c mice. *Food Chem Toxicol.* 2016; 95:42–51. Epub 2016/06/25. <https://doi.org/10.1016/j.fct.2016.06.019> PMID: 27338711.
57. Zhu W, Cromie MM, Cai Q, Lv T, Singh K, Gao W. Curcumin and vitamin E protect against adverse effects of benzo[a]pyrene in lung epithelial cells. *PLoS One.* 2014; 9(3):e92992. Epub 2014/03/26. <https://doi.org/10.1371/journal.pone.0092992> PMID: 24664296; PubMed Central PMCID: PMC3963982.

58. Zangar RC, Davydov DR, Verma S. Mechanisms that regulate production of reactive oxygen species by cytochrome P450. *Toxicol Appl Pharmacol*. 2004; 199(3):316–31. Epub 2004/09/15. <https://doi.org/10.1016/j.taap.2004.01.018> PMID: 15364547.
59. Uppstad H, Ovrebo S, Haugen A, Mollerup S. Importance of CYP1A1 and CYP1B1 in bioactivation of benzo[a]pyrene in human lung cell lines. *Toxicol Lett*. 2010; 192(2):221–8. Epub 2009/11/03. <https://doi.org/10.1016/j.toxlet.2009.10.025> PMID: 19879933.
60. Chi AC, Appleton K, Henriod JB, Krayner JW, Marlow NM, Bandyopadhyay D, et al. Differential induction of CYP1A1 and CYP1B1 by benzo[a]pyrene in oral squamous cell carcinoma cell lines and by tobacco smoking in oral mucosa. *Oral Oncol*. 2009; 45(11):980–5. Epub 2009/07/07. <https://doi.org/10.1016/j.oraloncology.2009.05.562> PMID: 19576839; PubMed Central PMCID: PMC2764788.
61. Tian LX, Tang X, Zhu JY, Luo L, Ma XY, Cheng SW, et al. Cytochrome P450 1A1 enhances inflammatory responses and impedes phagocytosis of bacteria in macrophages during sepsis. *Cell Commun Signal*. 2020; 18(1):70. Epub 2020/05/06. <https://doi.org/10.1186/s12964-020-0523-3> PMID: 32366266; PubMed Central PMCID: PMC7199371.
62. Jain SK, Singh S, Khajuria A, Guru SK, Joshi P, Meena S, et al. Pyrano-isochromanones as IL-6 inhibitors: synthesis, in vitro and in vivo antiarthritic activity. *J Med Chem*. 2014; 57(16):7085–97. Epub 2014/08/12. <https://doi.org/10.1021/jm500901e> PMID: 25111439.
63. Kumar A, Bhatia HS, de Oliveira AC, Fiebich BL. microRNA-26a modulates inflammatory response induced by toll-like receptor 4 stimulation in microglia. *J Neurochem*. 2015; 135(6):1189–202. Epub 2015/09/17. <https://doi.org/10.1111/jnc.13364> PMID: 26376347.
64. Singh A, Chibber P, Kolimi P, Malik TA, Kapoor N, Kumar A, et al. Rohitukine inhibits NF-kappaB activation induced by LPS and other inflammatory agents. *Int Immunopharmacol*. 2019; 69:34–49. Epub 2019/01/22. <https://doi.org/10.1016/j.intimp.2019.01.015> PMID: 30665042.