

RESEARCH ARTICLE

Open Access

Genome-wide DNA methylation changes in skeletal muscle between young and middle-aged pigs

Long Jin^{1†}, Zhi Jiang^{2†}, Yudong Xia^{3†}, Ping'er Lou¹, Lei Chen⁴, Hongmei Wang¹, Lu Bai¹, Yanmei Xie¹, Yihui Liu¹, Wei Li¹, Bangsheng Zhong¹, Junfang Shen⁵, An'an Jiang¹, Li Zhu¹, Jinyong Wang⁴, Xuewei Li^{1*} and Mingzhou Li^{1*}

Abstract

Background: Age-related physiological, biochemical and functional changes in mammalian skeletal muscle have been shown to begin at the mid-point of the lifespan. However, the underlying changes in DNA methylation that occur during this turning point of the muscle aging process have not been clarified. To explore age-related genomic methylation changes in skeletal muscle, we employed young (0.5 years old) and middle-aged (7 years old) pigs as models to survey genome-wide DNA methylation in the *longissimus dorsi* muscle using a methylated DNA immunoprecipitation sequencing approach.

Results: We observed a tendency toward a global loss of DNA methylation in the gene-body region of the skeletal muscle of the middle-aged pigs compared with the young group. We determined the genome-wide gene expression pattern in the *longissimus dorsi* muscle using microarray analysis and performed a correlation analysis using DMR (differentially methylated region)-mRNA pairs, and we found a significant negative correlation between the changes in methylation levels within gene bodies and gene expression. Furthermore, we identified numerous genes that show age-related methylation changes that are potentially involved in the aging process. The methylation status of these genes was confirmed using bisulfite sequencing PCR. The genes that exhibited a hypomethylated gene body in middle-aged pigs were over-represented in various proteolysis and protein catabolic processes, suggesting an important role for these genes in age-related muscle atrophy. In addition, genes associated with tumorigenesis exhibited aged-related differences in methylation and expression levels, suggesting an increased risk of disease associated with increased age.

Conclusions: This study provides a comprehensive analysis of genome-wide DNA methylation patterns in aging pig skeletal muscle. Our findings will serve as a valuable resource in aging studies, promoting the pig as a model organism for human aging research and accelerating the development of comparative animal models in aging research.

Keywords: DNA methylation, Skeletal muscle, Pig, Aging, MeDIP-seq, DNMTs

Background

Aging is a nearly universal, chronic process that is shared by all organisms. The most prominent feature of the aging process in mammals is a gradual loss of function at the cellular, tissue and organismal levels. 'Aging epigenetics' is an emerging field that has generated exciting revelations. A global loss of DNA methylation has been identified as

an age-related epigenetic alteration [1]. Many studies have revealed that DNA methylation plays an important role in aging and in the development of various diseases [2,3]. Therefore, a survey of epigenetic signatures that change with age might be useful to identify biomarkers of aging and age-associated diseases, which could potentially be used to make clinical diagnoses and prolong the lifespan.

The aging process and its underlying mechanisms have been studied extensively in rodent models [4,5]. The sequencing and analysis of the pig genome [6] will greatly accelerate the development of the pig as a biomedical

* Correspondence: xuewei.li@sicau.edu.cn; mingzhou.li@163.com

†Equal contributors

¹Institute of Animal Genetics and Breeding, College of Animal Science and Technology, Sichuan Agricultural University, Ya'an, Sichuan 625014, China
Full list of author information is available at the end of the article

model for many diseases in humans, such as obesity and diabetes [7-9]. However, few aging studies have been carried out using pigs as models. Pigs age at a rate of approximately 5 years to every 1 year of human life, resulting in an average life expectancy of 15–20 years. Therefore, pigs could serve as an appealing model for studying aging because of their relatively longer lifespan than rodents and similar metabolic features, cardiovascular systems and proportional organ sizes relative to those of humans [10-12].

Notably, an increase in the incidence of age-related pathologies mostly begins at approximately the mid-point of a species' life span [13-15]. It is well documented that the remarkable structural and functional changes that occur in skeletal muscle during aging, including a reduction of muscle mass and increased apoptosis [14,16-19], are initiated at the mid-point of the lifespan [14]. Consequently, studies focusing on DNA methylation changes in skeletal muscle during the mid-life period compared with the young stage are long overdue.

In this study, we used the pig as a model to perform a genome-wide survey of differences in DNA methylation and gene expression in a representative skeletal muscle (*longissimus dorsi* muscle, LDM) between two age stages in female pigs: young (0.5 years old) and middle-aged (MA, 7 years old) [10,11]. We identified the patterns of methylation in the pig genome and the age-related differentially methylated regions (DMRs), then performed functional enrichment analysis for genes exhibiting DMRs. We found more genes showing a hypomethylated gene body in the middle-aged pigs than in the young pigs; these genes were potentially involved in aging processes, such as the development of muscular atrophy. We believe that this study will serve as a valuable resource for aging studies while also promoting the pig as a model organism for human aging research and accelerating the considerable development of comparative animal models for aging research.

Results

Global DNA methylation analysis

Epigenetic alterations, such as global DNA hypomethylation, have been shown to progressively accumulate during aging [1]. We therefore first investigated the global DNA methylation status of six types of tissue by performing direct colorimetric quantification of methylated DNA. The global methylation level in the heart was significantly decreased in the MA pigs compared with the younger pigs (Figure 1), whereas the methylation levels in five other tissues showed a slight (but not significant) decrease in the MA pigs. This global loss of DNA methylation in older pigs was consistent with studies in humans, supporting a potential role of this type of epigenetic alteration in age-related gene regulation.




Figure 1 Global methylation status of six tissues in young and middle-aged pigs. "5-mC%" denotes the ratio of methylated cytosines to all nucleotides in the genome. (Student's t-test, ** $P < 0.01$).

DNA methyltransferases (*DNMTs*) are crucial for the establishment and maintenance of genomic methylation patterns [20]. To determine whether the global loss of DNA methylation observed in the MA pigs was associated with alterations in *DNMTs*, we evaluated the mRNA transcript levels of *DNMT1* (the major maintenance methyltransferase), *DNMT3a* and *3b* (two *de novo* methylation methyltransferase) in these six tissues. For *DNMT1* and *DNMT3a*, no significant differences in gene expression were observed in most of the tissues between the two age groups (Additional file 1), whereas *DNMT3b* showed increased mRNA levels in most of the tissues (except the ovary) in the MA pigs. Previous studies have revealed increased expression of both *DNMT3a* and *DNMT3b* in aging fibroblasts and in the aging human liver [21,22]. However, the expression level of *DNMT3a* showed no difference between young and MA pigs. Our result suggests that *DNMT3b*, rather than *DNMT3a*, may regulate the genomic methylation pattern in a tissue-specific manner in pigs during aging.

Summary of methylated DNA immunoprecipitation sequencing (MeDIP-seq) data

Approximately 46 gigabases (Gb) of MeDIP-seq data were generated from six skeletal muscle samples (approximately 7.64 Gb per sample), among which 81% (approximately 35.31 Gb) of the clean reads were aligned to the porcine reference genome build 9.2. After removing ambiguous reads from the clean reads, 75% of the reads (approximately 36.56 Gb) were uniquely aligned across all of the aligned reads. The reads that showed the same mapping locations in each sample were regarded as potentially duplicated clones generated via PCR amplification and treated as the

same read. Consequently, we acquired 24.46 Gb (approximately 92% of the unique mappable reads) of uniquely aligned non-duplicated reads (Additional file 2). CpG sites covered by a read depth of more than 10 were scored as high-confidence CpG sites. On average, 34.32% of the CpG sites met this threshold (Additional file 3).

To study methylation changes on a genome-wide scale, we measured methylation levels along the chromosomes in the samples using a 1 Mb sliding window to smooth the distribution (Additional file 4). Correlations between methylation levels and genomic features were assessed. We found that the methylation levels across the chromosomes were negatively correlated with the chromosomal length (Pearson's $r = 0.633$, $P = 0.0036$) and positively correlated with the GC content ($r = 0.787$, $P = 6.49 \times 10^{-5}$), single-nucleotide polymorphism (SNP) density ($r = 0.549$, $P = 0.0149$) and gene density ($r = 0.516$, $P = 0.0236$). In addition, a strong positive correlation was observed with the ratio between the observed and expected numbers of CpG sites ($\text{CpG}_{o/e}$) ($r = 0.931$, $P = 7.41 \times 10^{-9}$), which agreed with the results of a previous study of porcine DNA methylomes (Additional file 5) [8]. However, methylation levels were not correlated with the density of repeat regions ($r = 0.348$, $P = 0.145$). Among these genomic features, the $\text{CpG}_{o/e}$ ratio showed the highest correlation with the methylation level. The gene density also exhibited a significant correlation with the methylation level, which may be due to the relatively higher GC content in the gene regions being examined (Additional file 6), suggesting the potential role of methylation changes in the regulation of gene transcription [23]. Previous studies have demonstrated a strong genetic component of the variation in DNA methylation profiles [24], and a potential role has been suggested for CpG-SNPs in genetic variation of the epigenome [25]. Previous studies demonstrated that the level of methylation contributes to variation in the substitution rates at CpG sites [26,27]. We observed a positive correlation between the SNP density and methylation level, suggesting that the genetic variation reflected by SNPs may have a substantial impact on local methylation patterns and gene expression.

We defined 24 categories of functional genomic elements and further classified the promoters into three types based on their CpG sites. Each type of promoter was then classified according to its distance from the transcription start site (Additional file 7) [8]. We also classified CpG islands and CGI shores into four categories according to their genomic locations, as described in previous studies [24,28]. We found that intermediate CpG promoters (ICPs) exhibited a relatively higher methylation status than did high CpG promoters (HCPs) and low CpG promoters (LCPs) (two-way ANOVA, $P = 2.87 \times 10^{-37}$). The methylation levels within the distal (D), proximal (P) and intermediate (I) regions of promoters also showed significant differences (two-way


ANOVA, $P = 1.44 \times 10^{-7}$) (Additional file 7). This result agreed with a previous finding that methylation occurs more frequently at ICPs [29]. Our data also suggested that a relatively higher methylation level within gene bodies is a general phenomenon in mammals [30], and it has been correlated with gene expression levels [28]. We observed that the methylation level of exons was higher than that of introns ($P = 9.8 \times 10^{-9}$) (Additional file 7), reflecting the higher GC content of exons compared with their surrounding introns and further indicating the possible different roles of exons and introns in the regulation of gene transcription [31]. Recently, DNA methylation at CpG island (CGI) shores has been demonstrated to play a more important role in gene regulation than that of the CGIs themselves [32]. We observed distinct methylation levels for CGI shores in various genomic locations (Additional file 7), which may suggest the distinctive roles of these CGI shores in regulating gene expression.

Differential DNA methylation in the subtelomeric regions of young and MA pigs

We surveyed the chromosomal profiles of DNA methylation and found that the subtelomeric regions showed significant hypermethylation compared with non-subtelomeric regions for all chromosomes (Figure 2A), which was consistent with the heavily methylated status of the subtelomeric region previously reported in mice [33]. Epigenetic modifications of the subtelomeric region correlate with telomere elongation, which is closely related to aging. Interestingly, we found that the methylation levels of the subtelomeric regions of each chromosome exhibited distinct patterns between the young and MA pigs (Figure 2B and 2C; Additional file 4). The average methylation status of the subtelomeric regions of all of the chromosomes of the MA pigs was significantly lower than that of the young pigs (Figure 2D). Previous studies of humans revealed that low levels of subtelomeric methylation may contribute toward increasing the levels of telomeric repeat-containing RNA, whose transcription originates in the subtelomere and is accompanied by a reduction in telomerase activity [34], thereby controlling the telomere length [35]. We therefore also measured telomere length using a qPCR assay (Figure 2D). The ratio of the telomere signal to the signal of the single-copy gene (T/S) (see "Methods") was relatively lower in the MA pigs than in the young pigs, suggesting that the average telomere length was shorter in the MA pigs [36]. This result indicated that hypomethylation in the subtelomeric regions of the MA pigs may be accompanied by telomere attrition and involved in the aging process.

Differentially methylated regions (DMRs) associated with aging

We identified 9,234 DMRs (with a Benjamini-Hochberg corrected $P < 0.01$, FDR < 0.01) in skeletal muscle between



the young and MA pigs, representing approximately 0.064% of the length of the genome and approximately 0.28% of the total number of CpGs in the genome (Table 1). We performed hierarchical clustering of the samples using the DMRs of each genomic element and differentially expressed genes. Among all of the clustering plots, six samples could be clustered into two groups: young pigs and MA pigs, although the topological structure within the groups showed subtle differences (Additional file 8). This result revealed a high correlation between the biological replicates and indicated experimental reliability and relative epigenetic concordance within each group.

To further explore potential distribution biases in the age-related DMRs, we analyzed the percentage of CpGs


within the DMRs in each of the 24 genomic elements. Among promoters, more DMRs were enriched in the ICPs compared with HCPs and LCPs. ICPs generally contain weak CpG islands and are prone to regulation by DNA methylation [29]. Our results indicated that not only did ICPs exhibit the highest methylation levels compared with the other two promoter classes (Additional file 7), but DMRs were also observed more frequently in ICPs (Figure 3A). These observations demonstrated that ICPs are more susceptible to methylation and suggested that this type of alteration of the methylation status of ICPs may play an important role in modulating gene expression relevant to several biological processes, such as aging. Meanwhile, although a previous study found that the

Table 1 Summary of differentially methylated regions (DMRs)

DMR type	Number of DMRs	Percentage of genomic length*	Percentage of genomic CpGst
Age-related DMRs (n = 6)	9,234	0.064	0.280

*Total length of all DMRs relative to the length (approximately 2.60 billion bp) of the pig genome (Sscrofa 9.2).

†The number of CpGs among the total DMRs relative to the total number of CpGs (approximately 26.91 M) in the pig genome (Sscrofa 9.2).



distal (D) regions of promoters contained more DMRs [8], no significant enrichment of DMRs was observed in the present study except in the distal regions of HCPs, suggesting that various types of DMRs may participate in distinct biological processes (Figure 3A). There is increasing evidence that methylation in the gene body affects gene expression in plants [37,38] and mammals [30]. Our analysis indicated that more DMRs are located in gene bodies (1,720) than in promoters (185) (Additional file 9), which may suggest that gene bodies are more susceptible to changes compared to promoters during the aging process. The first exon contained relatively few DMRs within the gene body (Figure 3B), which may be the result of certain motifs overlapping between the promoter and


the first exon. We also found that the majority of DMRs occurred at CGI shores (two-way ANOVA, $P = 0.001$), whereas the numbers of DMRs did not significantly vary across gene features ($P = 0.348$), consistent with previous reports on human cancer (Figure 3C) [32,39-41].

Gene-body DNA methylation and gene expression

The influence of the methylation status of promoter regions is considered to be an important mechanism regulating gene transcription. Nevertheless, investigation of the specific roles of DNA methylation in gene bodies is long overdue. To explore whether intragenic methylation affects gene expression, we performed a correlation analysis of DMR-mRNA pairs and observed a significant negative correlation ($r = -0.206$, $P = 3.186 \times 10^{-7}$) between changes in the methylation levels in the gene body and gene expression (Figure 4). Whether methylation in the gene body inhibits or induces transcription remains unclear [30,31,37,42]. Nevertheless, we conclude here that gene-body methylation reduced gene expression, possibly via an intragenic DNA methylation-induced decrease in the Pol II elongation efficiency [42].

Functional enrichment analysis for genes with DMRs

To examine the potential function of the genes that showed a differential methylation status, we performed an enrichment analysis for genes with DMRs in their promoters and genes that were hyper- and hypomethylated in the gene body associated with age. Because DNA methylation concentrates at gene bodies on the X chromosome



due to X chromosome inactivation (Xi), we excluded the DMRs on chromosome X from this analysis. A total of 185 genes that exhibited DMRs in their promoters were mapped to their human orthologs, and 657 and 1,063 genes with DMRs in their gene bodies that mapped to human genes showed hyper- and hypomethylation in the MA pigs, respectively (Figure 5A). More DMRs were observed in gene bodies rather than promoters, and the DMRs in gene bodies were more prone to be hypomethylated with age (Figure 5B). The finding that more genes were hypomethylated in the MA group also supported the theory of progressive global methylation loss during aging. The genes displaying DMRs in their promoters were not significantly enriched in any biological process or molecular function, which may due to the relatively low abundance of genes with DMRs in their promoters. The gene-body hypermethylated genes were significantly enriched for the processes 'GTPase regulator activity' (30 genes, $P = 0.03$), 'ATP binding' (79 genes, $P = 0.03$) and 'protein kinase activity' (39 genes, $P = 0.04$) (Additional file 10). Notably, the gene-body hypomethylated genes showed significant enrichment for various processes related to proteolysis ($P = 4.55 \times 10^{-4}$), protein catabolism ($P = 5.84 \times 10^{-8}$), and energy metabolism (GO: energy derivation by oxidation of organic compounds, $P = 0.04$) (Additional file 10). Previous studies have indicated that catabolic processes increase in aged muscles [43]. Our findings suggested that the hypomethylated genes observed in the *longissimus dorsi* muscle of the MA pigs were involved in protein degradation and may be responsible for muscular atrophy, which is one of the general structural and phenotypic changes observed in aging muscle [14].

Genes involved in the aging process

To further highlight the potential roles of genes involved in aging, we considered the intersection of genes that presented DMRs in their promoters and gene bodies with the known age-related genes deposited in the Human Ageing Genomic

Resources (HAGR) database [44]. Among the 288 genes potentially involved in the human aging process according to the HAGR-GenAge database, we did not identify any genes with DMRs in their promoters, whereas there were 12 known age-related genes included in the list of genes with DMRs in their gene bodies (Fisher's exact test, $P = 0.024$) (Figure 5C).

A potential role of DNA methylation of the gene body in regulating gene expression has been proposed [28,31,37,45]. We therefore investigated genes with DMRs in the gene body. Several genes included in the list of well-characterized human age-related genes exhibited differentially methylated gene bodies, which was confirmed using the bisulfite sequencing PCR (BSP) approach, and were shown to cause changes in mRNA expression (Figure 6). For example, *FoxO3* (forkhead box subfamily O member 3) has been well documented to be an essential transcription factor involved in the atrophy of muscles and myotubes [46]. *FoxO3* stimulates lysosomal proteolysis in muscle by activating autophagy and proteasomal pathways, and inducing the expression of autophagy-related genes [47]. We observed up-regulated expression and a lower gene-body methylation status of this gene in MA pigs (Figure 6A), in accordance with previous findings in aging muscle. Another gene, *FGFR1* (fibroblast growth factor receptor 1), which shows an opposite function to that of *FoxO3*, could inhibit the atrophy of skeletal muscle [48]. Down-regulated expression of *FGFR1*, together with hypermethylation of its gene body, was observed in MA pig skeletal muscle (Figure 6B). *GRB2* (growth factor receptor-bound protein 2) is critical for cell proliferation, and emerging evidence indicates that *GRB2* plays a role in tumorigenesis and is over-expressed in tumors [49]. In addition, a reduced expression level of *GRB2* in skeletal muscle contributes to increased insulin sensitivity [50,51]. Microarray mRNA expression and genome-wide methylation data on *GRB2* revealed an increased expression level and hypomethylation in the gene body of this gene, respectively (Figure 6C).





Figure 5 Differentially methylated (DM) genes in gene bodies and promoters. (A) Venn diagram of the numbers of genes showing promoter and gene-body DMRs. **(B)** Comparison of the genomic CpG percentage (number of CpGs in DMRs versus the total number of CpGs) and the genomic length percentage (DMR length versus the total length of DMRs) between gene-body and promoter regions. **(C)** Gene-body-DM genes that overlap with known age-related genes in the HAGR database.



Discussion

This study provides a comprehensive analysis of genome-wide DNA methylation patterns in the skeletal muscle of aging pigs. Similar to previous reports in humans and mice, a global loss of methylation induced by transcriptional changes in *DNMT3b* was observed in various tissues of the MA pigs, suggesting that this type of epigenetic alteration is common in aging mammals. Although global DNA hypomethylation and promoter CpG island hypermethylation have been observed to progressively accumulate during aging [52], the present study identified more DMRs in gene bodies than in promoters, and gene-body hypomethylation was observed in more genes in the MA pigs (Figure 5 and Additional file 9). It is therefore reasonable to assume that during the loss of global methylation during aging, there is a greater tendency toward hypomethylation in the gene body rather than the promoter. It should be noted that based on our current data, it is not feasible to identify methylation changes on the X chromosome because of Xi [53]. Further research using SNP data could allow allele-specific analysis of DNA methylation to identify the specific methylation changes on chromosome X [54].

It is believed that increased gene-body methylation correlates with increased transcription [30,31,37]; although some researchers have proposed that intragenic

methylation might reduce gene expression [42]. Our data suggested that methylation in the gene body reduces gene expression (Figure 4). However, gene-body methylation is only one of the many factors that influence gene expression. Further studies focusing on the methylation of specific regions that exhibit distinct gene regulation contexts are needed to elucidate the complicated epigenetic mechanism underlying aging and its associations with disease.

Previous reports have indicated that increased protein catabolism occurs in aging skeletal muscles [55,56]. Structural and functional changes associated with aging, such as reductions in the muscle mass and muscle fibers, have been observed across a wide range of species, from worms to mammals [14]. This type of epigenetic alteration of skeletal muscle with aging (Figure 6A and 6B; Additional file 10), consistent with findings in other species, was observed in pigs for the first time in the present work. Interestingly, genes related to tumorigenesis, as well as insulin sensitivity, exhibited a relatively higher expression level in the skeletal muscle of the MA pigs compared with younger pigs (Figure 6C), suggesting a higher risk of developing diseases with increased age.

Our results will promote further development of the pig as a model organism for human aging research.

Most of the studies carried out in pigs to date have been conducted in neonatal or very young animals (generally when the pigs reach peak commercial value at approximately 6 months of age), before they reach the age of 1 year [57,58]. Limited studies have been carried out on relatively older pigs (aged 2 years or more). Here, younger (0.5 years old) and middle-aged (7 years old) pigs were examined to investigate DNA methylation changes during the aging process. Although the aging process differs across species, with human aging showing major differences from the aging of most other species because of the relatively longer lifespan of humans, many species, including humans, pigs and mice, exhibit similarities with respect to aging muscle, muscular protein catabolism and muscle atrophy [59]. However, the time course of the muscle function changes occurring in pigs remains to be determined in further studies. In addition, pigs have a longer lifespan than that of rodents and present similar metabolic features, cardiovascular systems and proportional organ sizes to those humans [7]. Consequently, pigs can serve as a good biomedical model for human studies on the chronic aging process and its associated diseases [6,8,9]. However, only two age groups: young and middle-aged pigs were used in our study, and examining pigs of additional consecutive ages is necessary to further elucidate the changes in epigenetic modifications associated with age, as well as the ultimately complicated mechanisms underlying the aging process.

Conclusions

In summary, the present study provides a comprehensive analysis of genome-wide DNA methylation patterns in the skeletal muscle of aging pigs. We identified remarkable DNA methylation changes, such as a tendency toward hypomethylation in gene bodies in the *longissimus dorsi* muscle of MA pigs. Furthermore, we identified numerous genes that exhibited age-related methylation changes and are potentially involved in the aging process. These genes are mainly related to protein catabolism, suggesting that predisposition to amyotrophy emerges during middle age. This study will serve as a valuable resource for aging studies, promoting the pig as a model organism for human aging research and accelerating the development of comparative animal models in aging research.

Methods

Animals

A total of six healthy female pigs (Chinese Jinhua breed) were used in this study from two age groups: 0.5 and 7 years old, representing young and middle-aged pigs, respectively. Each age group included three individuals, which were regarded as biological replicates. The animals were reared in the same environment and fed the same diet *ad libitum* during the experimental period. Food was

withheld from the animals on the night before they were slaughtered. All experimental procedures and sample collection were approved by the Institutional Animal Care and Use Committee of the College of Animal Science and Technology of Sichuan Agricultural University, Sichuan, China, under permit No. DKY-B20121403.

Tissue collection

Six types of tissues (brain, liver, ovary, spleen, heart and *longissimus dorsi* muscle) were rapidly sampled from each carcass and immediately frozen in liquid nitrogen. All tissue samples were stored at -80°C until DNA and total RNA extraction.

Measurement of the global DNA methylation status

DNA from each collected tissue was extracted using the DNeasy Blood & Tissue Kit (Qiagen). Global DNA methylation was evaluated using the MethylFlash™ Methylated DNA Quantification Kit (Epigentek). The amount of input DNA for each assay was 100 ng to ensure optimal quantification. The experiments were performed according to the manufacturer's instructions.

Quantitative PCR analysis of the DNMT genes

Total RNA (10 μg) was extracted from the six muscle samples using TRIzol (Invitrogen). RNase-free DNase I (TaKaRa) was used to remove genomic DNA from the RNA samples. cDNA was synthesized using PrimeScript RT Master Mix (TaKaRa). Quantitative real-time PCR (q-PCR) was performed using SYBR Premix Ex Taq (TaKaRa) in the CFX96 Real-Time PCR Detection System (Bio-Rad). The primers used for q-PCR are listed in Additional file 11. All measurements were performed in parallel with a negative control (no cDNA template), and each RNA sample was analyzed in triplicate. Porcine *ACTB*, *TBP* and *TOP2B* were used as endogenous control genes [60,61]. The gradient dilution PCR assays for these three reference genes showed stable, high amplification efficiencies (Additional file 12). Relative expression levels were then calculated using the $\Delta\Delta\text{Ct}$ method [62].

Methylated DNA immunoprecipitation sequencing

DNA was extracted using the DNeasy Blood & Tissue Kit (Qiagen) and then eluted using 10 mM Tris·Cl, pH 7.5. The quality of the isolated DNA was then measured using a NanoDrop spectrophotometer. The ratio of the absorbance at 260 nm versus 280 nm (A_{260}/A_{280}) provides an estimate of the purity of the DNA. The A_{260}/A_{280} value should be 1.8 to 2.0 for each DNA sample to guarantee quality. The initial volume of DNA for each sample should be at least 5 μg to ensure the success of subsequent MeDIP-seq experiments. The protocol for MeDIP-seq and detailed information on the construction of a MeDIP DNA library were provided in a previous report

Functional enrichment analysis for genes with DMRs

The DAVID (Database for Annotation, Visualization and Integrated Discovery) web server (<http://david.abcc.ncifcrf.gov/>) was used to perform functional enrichment analysis of Gene Ontology (GO) and KEGG pathway categories [66]. Genes with DMRs in their promoters and gene bodies were mapped to their respective human orthologs and then submitted to DAVID for enrichment analysis, which included GO biological processes (GO-BP), molecular function (GO-MF) terminologies and KEGG pathway categories. Only GO-BP, GO-MF or KEGG-pathway terms with a Benjamini-corrected *P* value less than 0.05 were considered to be significant and therefore included in the list.

Gene expression microarray

Total RNA (10 µg) was extracted from the six samples using TRIzol (Invitrogen) and further purified using an RNeasy column (Qiagen). The integrity of the total RNA was confirmed using a Bioanalyzer 2100 and the RNA 6000 Nano LabChip Kit (Agilent Technologies). Detailed information on the workflow of the microarray experiment is provided in a previous report from our group [8]. First, we mapped 43,603 probes (60 mer in length) to the pig reference genome while allowing up to one mismatch, which resulted in 27,955 probes (64.11%) that were uniquely mapped. Among these uniquely mapped probes, 4,983 (11.43%) were uniquely mapped to exons in Ensembl genes (more than 60% sequence overlap). Multiple probes that mapped to the same or different exons of a specific gene were excluded. Therefore, only 3,074 probes (7.05%), which uniquely represented 3,074 genes and were considered to represent high-confidence gene expression data, were used in the subsequent analysis.

Differentially expressed genes were identified using the MultiExperiment Viewer (MeV) [67], and this software was also employed to perform subsequent hierarchical clustering of samples. The gene expression microarray data have been submitted to the GEO database under accession number GSE49791.

Bisulfite sequencing PCR

Methylation Primer Express Software V1.0 was used to design bisulfite sequencing PCR (BSP) primers, which are provided in Additional file 13. The bisulfite conversion of genomic DNA was performed using the EZ DNA Methylation-Gold™ Kit (Zymo Research, D5006). PCR was carried out using ZymoTaq™ PreMix (Zymo Research, E2004). The PCR product was then purified using the DNA Clean & Concentrator - 25™ Kit (Zymo Research, D4005), and the PCR product was cloned into the TA vector pCR2.1 (Invitrogen, K2000-01). Ten subclones were selected for each gene and subsequently sequenced

using an ABI 3730 DNA sequencer (Applied Biosystems). All of the sequences were analyzed using BiQ Analyzer V2.0 software [68].

Additional files

Additional file 1: Relative mRNA expression levels of DNMTs in six tissues between young and middle-age pigs. The expression levels were normalized to the maximum value obtained in the two groups. (Student's *t*-test, ***P* < 0.01, **P* < 0.05).

Additional file 2: Summary of MeDIP-seq data production.

Low-quality reads were filtered out of the raw reads, and the clean reads were then used in further analyses. "% aligned" is the percentage of clean reads aligned to the pig reference genome (version 9.2). "% unique" is the percentage of reads uniquely aligned across all of the aligned reads. The reads showing the same mapping locations in each sample were considered to be potentially duplicated clones generated via PCR amplification during sequencing library construction and were therefore removed from the analysis. "% non-duplicate alignment" is the percentage of uniquely aligned non-duplicated reads over all of the uniquely aligned reads.

Additional file 3: Percentage of CpGs showing an average coverage that meets the read depth threshold over all samples. Values are the means ± s.d. (*n* = 6).

Additional file 4: Genome-wide distribution of the DNA methylation levels. To compare the DNA methylation state between samples, the read depth was normalized to the overall average number of reads in each group. The CpG_{o/e} ratio, SNPs density, numbers of genes, repeats and CGIs were all calculated over 1 M Mb sliding windows.

Additional file 5: Pearson's correlation between DNA methylation levels and chromosomal features.

Additional file 6: Box plots of the percentages of GC content in the promoter, gene-body, and intergenic regions. Box-plot edges indicate the 25th and 75th percentiles; central bars indicate the medians; and whiskers indicate the non-outlier extremes. The significance was evaluated using *t*-tests.

Additional file 7: Comparison of methylation levels in various categories of genomic elements. The average methylation level for each group was plotted. Then DNA methylation levels (average read depth of two age groups) of various categories were compared. (A) Methylation levels across various promoter categories were compared using two-way ANOVA. Intermediate CpG promoters (ICPs) exhibited a relatively higher methylation status relative to those of high CpG promoters (HCPs) and low CpG promoters (LCPs) (*P* = 2.87 × 10⁻³⁷). The methylation levels within the distal (D), proximal (P) and intermediate (I) regions also showed significant differences (*P* = 1.44 × 10⁻⁷). (B) Comparison of methylation levels between exons and introns was performed using Student's *t*-test. The exon methylation was higher than the intron methylation (*P* = 9.8 × 10⁻⁹). (C) Methylation levels between CGIs and CGI shores were compared using two-way ANOVA. The average methylation in CGI shores was significantly higher than that in CGIs (*P* = 2.17 × 10⁻³).

Additional file 8: Hierarchical clustering of samples using DMRs in various genomic elements and differentially expressed (DE) mRNAs. Clustering was performed using MultiExperiment Viewer software. The distance metric applied for clustering was Pearson correlation across samples.

Additional file 9: List of DMRs.

Additional file 10: Over-represented functional gene categories for DMRs.

Additional file 11: Information on primers used to perform q-PCR.

Additional file 12: Amplification efficiencies of gradient dilution PCR assays for the *ACTB*, *TBP* and *TOP2B* genes.

Additional file 13: Information on primers used to perform BSP.

Abbreviations

LDM: *Longissimus dorsi* muscle; MA: Middle-aged; DMRs: Differentially methylated regions; DNMTs: DNA methyltransferases; MeDIP-seq: Methylated DNA immunoprecipitation sequencing; HCP: High CpG promoter;

ICP: Intermediate CpG promoter; LCP: Low CpG promoter; CGI: CpG island; D: Distal; P: Proximal; I: Intermediate; HAGR: Human ageing genomic resources; BSP: Bisulfite sequencing PCR.

Competing interests

The authors declared that they have no competing interests.

Authors' contributions

LJ, ZJ, YX, ML and XL designed the experiment. LJ, PL, LZ, JW, ML and XL participated in sample collection. LJ, HW, LB and YMX made substantial contributions to the laboratory work. LJ, ZJ, YX, LC, WL, YL, JS, ML and XL carried out the data analysis. LJ, YD, PL, ML and XL drafted the manuscript. LJ, LC, BZ, AJ, ML and XL participated in revising the manuscript for important intellectual content. All authors read and approved the final manuscript.

Acknowledgments

This work was supported by grants from the National Special Foundation for Transgenic Species of China (2014ZX0800950B), the National Natural Science Foundation of China (31301942, 31101701 and 31372284), the National High Technology Research and Development Program of China (863 Program) (2013AA102502), the Fund of Fok Ying-Tung Education Foundation (141117), the Fund for Distinguished Young Scientists of Sichuan Province (2013JQ0013), the Specialized Research Fund of Ministry of Agriculture of China (NYCYTX-009), the Program for Changjiang Scholars and Innovative Research Team in University (IRT13083), International Science & Technology Cooperation Program of China (2014DFA31260).

Author details

¹Institute of Animal Genetics and Breeding, College of Animal Science and Technology, Sichuan Agricultural University, Ya'an, Sichuan 625014, China. ²Novogene Bioinformatics Institute, Beijing 100083, China. ³E-GENE, Shenzhen, Guangdong 518173, China. ⁴Chongqing Academy of Animal Science, Chongqing 402460, China. ⁵BGI-Tech, BGI-Shenzhen, Shenzhen 518083, China.

Received: 1 May 2014 Accepted: 31 July 2014

Published: 5 August 2014

References

- Wilson VL, Smith RA, Ma S, Cutler RG: **Genomic 5-methyldeoxycytidine decreases with age.** *J Biol Chem* 1987, **262**(21):9948–9951.
- Movassagh M, Choy MK, Goddard M, Bennett MR, Down TA, Foo RS: **Differential DNA methylation correlates with differential expression of angiogenic factors in human heart failure.** *PLoS One* 2010, **5**(1):e8564.
- Mastroeni D, Grover A, Delvaux E, Whiteside C, Coleman PD, Rogers J: **Epigenetic changes in Alzheimer's disease: decrements in DNA methylation.** *Neurobiol Aging* 2010, **31**(12):2025–2037.
- Maegawa S, Hinkal G, Kim HS, Shen L, Zhang L, Zhang J, Zhang N, Liang S, Donehower LA, Issa JP: **Widespread and tissue specific age-related DNA methylation changes in mice.** *Genome Res* 2010, **20**(3):332–340.
- Serra C, Tangherlini F, Rudy S, Lee D, Toraldo G, Sandor NL, Zhang A, Jasuja R, Bhasin S: **Testosterone improves the regeneration of old and young mouse skeletal muscle.** *J Gerontol A Biol* 2013, **68**(1):17–26.
- Groenen MA, Archibald AL, Uenishi H, Tuggle CK, Takeuchi Y, Rothschild MF, Rogel-Gaillard C, Park C, Milan D, Megens HJ, Li S, Larkin DM, Kim H, Frantz LA, Caccamo M, Ahn H, Aken BL, Anselmo A, Anthon C, Auvin L, Badaoui B, Beattie CW, Bendixen C, Berman D, Blecha F, Blomberg J, Bolund L, Bosse M, Botti S, Buije Z, et al: **Analyses of pig genomes provide insight into porcine demography and evolution.** *Nature* 2012, **491**(7424):393–398.
- Spurlock ME, Gabler NK: **The development of porcine models of obesity and the metabolic syndrome.** *J Nutr* 2008, **138**(2):397–402.
- Li M, Wu H, Luo Z, Xia Y, Guan J, Wang T, Gu Y, Chen L, Zhang K, Ma J, Liu Y, Zhong Z, Nie J, Zhou S, Mu Z, Wang X, Qu J, Jing L, Wang H, Huang S, Yi N, Wang Z, Xi D, Wang J, Yin G, Wang L, Li N, Jiang Z, Lang Q, Xiao H, et al: **An atlas of DNA methylomes in porcine adipose and muscle tissues.** *Nat Commun* 2012, **3**:850.
- Fan N, Lai L: **Genetically Modified Pig Models for Human Diseases.** *J Genet Genomics* 2013, **40**(2):67–73.
- Peters RC, Johnson WA: **Ageing and Miniature Swine Heart and Liver Plasma Membranes.** In *Advances in Swine in Biomedical Research*, Volume 2. Edited by Tumbleson ME. New York: Springer; 1997:581–593.
- Bhathena S, Berlin E, Johnson W: **The minipig as a model for the study of aging in humans: selective responses of hormones involved in carbohydrate and lipid metabolism in different sexes.** In *Advances in Swine in Biomedical Research*, Volume 2. Edited by Bhathena SJ. New York: Plenum; 1996:571–580.
- Larsen MO, Rolin B, Wilken M, Carr RD, Svendsen O, Bollen P: **Parameters of glucose and lipid metabolism in the male Gottingen minipig: influence of age, body weight, and breeding family.** *Comp Med* 2001, **51**(5):436–442.
- Jemal A, Siegel R, Xu J, Ward E: **Cancer statistics, 2010.** *Ca Cancer J Clin* 2010, **60**(5):277–300.
- Nair KS: **Ageing muscle.** *Am J Clin Nutr* 2005, **81**(5):953–963.
- Heyn H, Li N, Ferreira HJ, Moran S, Pisanò DG, Gomez A, Diez J, Sanchez-Mut JV, Setien F, Carmona FJ, Puca AA, Sayols S, Pujana MA, Serra-Musach J, Iglesias-Platas I, Formiga F, Fernandez AF, Fraga MF, Heath SC, Valencia A, Gut IG, Wang J, Esteller M: **Distinct DNA methylomes of newborns and centenarians.** *Proc Natl Acad Sci U S A* 2012, **109**(26):10522–10527.
- Dirks A, Leeuwenburgh C: **Apoptosis in skeletal muscle with aging.** *Am J Physiol Reg I* 2002, **282**(2):R519–R527.
- Navarro A, Lopez-Cepero JM, Sanchez del Pino MJ: **Skeletal muscle and aging.** *Front Biosci* 2001, **6**:D26–D44.
- Janssen I, Heymsfield SB, Wang ZM, Ross R: **Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr.** *J Appl Physiol* 2000, **89**(1):81–88.
- Pollack M, Phaneuf S, Dirks A, Leeuwenburgh C: **The role of apoptosis in the normal aging brain, skeletal muscle, and heart.** *Ann NY Acad Sci* 2002, **959**:93–107.
- Kim GD, Ni J, Kelesoglu N, Roberts RJ, Pradhan S: **Co-operation and communication between the human maintenance and de novo DNA (cytosine-5) methyltransferases.** *Embo J* 2002, **21**(15):4183–4195.
- Lopatina N, Haskell JF, Andrews LG, Poole JC, Saldanha S, Tollefsbol T: **Differential maintenance and de novo methylating activity by three DNA methyltransferases in aging and immortalized fibroblasts.** *J Cell Biochem* 2002, **84**(2):324–334.
- Xiao Y, Word B, Starlard-Davenport A, Haefele A, Lyn-Cook BD, Hammons G: **Age and gender affect DNMT3a and DNMT3b expression in human liver.** *Cell Biol Toxicol* 2008, **24**(3):265–272.
- Kim JK, Samaranyake M, Pradhan S: **Epigenetic mechanisms in mammals.** *Cell Mol Life Sci* 2009, **66**(4):596–612.
- Bell JT, Pai AA, Pickrell JK, Gaffney DJ, Pique-Regi R, Degner JF, Gilad Y, Pritchard JK: **DNA methylation patterns associate with genetic and gene expression variation in HapMap cell lines.** *Genome Biol* 2011, **12**(1):R10.
- Shoemaker R, Deng J, Wang W, Zhang K: **Allele-specific methylation is prevalent and is contributed by CpG-SNPs in the human genome.** *Genome Res* 2010, **20**(7):883–889.
- Xia J, Han L, Zhao Z: **Investigating the relationship of DNA methylation with mutation rate and allele frequency in the human genome.** *BMC Genomics* 2012, **13**(Suppl 8):S7.
- Mugal CF, Ellegren H: **Substitution rate variation at human CpG sites correlates with non-CpG divergence, methylation level and GC content.** *Genome Biol* 2011, **12**(6):R58.
- Maunakea AK, Nagarajan RP, Bilenyk M, Ballinger TJ, D'Souza C, Fouse SD, Johnson BE, Hong C, Nielsen C, Zhao Y, Turecki G, Delaney A, Varhol R, Thiessen N, Shchors K, Heine VM, Rowitch DH, Xing X, Fiore C, Schillebeeckx M, Jones SJ, Haussler D, Marra MA, Hirst M, Wang T, Costello JF: **Conserved role of intragenic DNA methylation in regulating alternative promoters.** *Nature* 2010, **466**(7303):253–257.
- Weber M, Hellmann I, Stadler MB, Ramos L, Paabo S, Rebhan M, Schubeler D: **Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome.** *Nat Genet* 2007, **39**(4):457–466.
- Ball MP, Li JB, Gao Y, Lee JH, LeProust EM, Park IH, Xie B, Daley GQ, Church GM: **Targeted and genome-scale strategies reveal gene-body methylation signatures in human cells.** *Nat Biotechnol* 2009, **27**(4):361–368.
- Laurent L, Wong E, Li G, Huynh T, Tsigos A, Ong CT, Low HM, Kin Sung KW, Rigoutsos I, Loring J, Wei CL: **Dynamic changes in the human methylome during differentiation.** *Genome Res* 2010, **20**(3):320–331.
- Irizarry RA, Ladd-Acosta C, Wen B, Wu Z, Montano C, Onyango P, Cui H, Gabo K, Rongione M, Webster M, Ji H, Potash JB, Sabuncyan S, Feinberg AP: **The human colon cancer methylome shows similar hypo- and hypermethylation at conserved tissue-specific CpG island shores.** *Nat Genet* 2009, **41**(2):178–186.

33. Gonzalo S, Jaco I, Fraga MF, Chen T, Li E, Esteller M, Blasco MA: **DNA methyltransferases control telomere length and telomere recombination in mammalian cells.** *Nat Cell Biol* 2006, **8**(4):416–424.
34. Ng LJ, Croypley JE, Pickett HA, Reddel RR, Suter CM: **Telomerase activity is associated with an increase in DNA methylation at the proximal subtelomere and a reduction in telomeric transcription.** *Nucleic Acids Res* 2009, **37**(4):1152–1159.
35. Blasco MA: **Telomeres and human disease: ageing, cancer and beyond.** *Nat Rev Genet* 2005, **6**(8):611–622.
36. Ji G, Liu K, Okuka M, Liu N, Liu L: **Association of telomere instability with senescence of porcine cells.** *BMC Cell Biol* 2012, **13**:36.
37. Cokus SJ, Feng S, Zhang X, Chen Z, Merriman B, Haudenschild CD, Pradhan S, Nelson SF, Pellegrini M, Jacobsen SE: **Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning.** *Nature* 2008, **452**(7184):215–219.
38. Lister R, O'Malley RC, Tonti-Filippini J, Gregory BD, Berry CC, Millar AH, Ecker JR: **Highly integrated single-base resolution maps of the epigenome in Arabidopsis.** *Cell* 2008, **133**(3):523–536.
39. Davies MN, Volta M, Pidsley R, Lunnon K, Dixit A, Lovestone S, Coarfa C, Harris RA, Milosavljevic A, Troakes C, Al-Sarraj S, Dobson R, Schalkwyk LC, Mill J: **Functional annotation of the human brain methylome identifies tissue-specific epigenetic variation across brain and blood.** *Genome Biol* 2012, **13**(6):R43.
40. Feber A, Wilson GA, Zhang L, Presneau N, Idowu B, Down TA, Rakyanc VK, Noon LA, Lloyd AC, Stupka E, Schiza V, Teschendorff AE, Schroth GP, Flanagan A, Beck S: **Comparative methylome analysis of benign and malignant peripheral nerve sheath tumors.** *Genome Res* 2011, **21**(4):515–524.
41. Rao X, Evans J, Chae H, Pirose J, Kim S, Yan P, Huang RL, Lai HC, Lin H, Liu Y, Miller D, Rhee JK, Huang YW, Gu F, Gray JW, Huang TM, Nephew KP: **CpG island shore methylation regulates caveolin-1 expression in breast cancer.** *Oncogene* 2013, **32**(38):4519–4528.
42. Lorincz MC, Dickerson DR, Schmitt M, Groudine M: **Intragenic DNA methylation alters chromatin structure and elongation efficiency in mammalian cells.** *Nat Struct Mol Biol* 2004, **11**(11):1068–1075.
43. Carmeli E, Coleman R, Reznick AZ: **The biochemistry of aging muscle.** *Exp Gerontol* 2002, **37**(4):477–489.
44. de Magalhaes JP, Budovsky A, Lehmann G, Costa J, Li Y, Fraifeld V, Church GM: **The Human Ageing Genomic Resources: online databases and tools for biogerontologists.** *Aging Cell* 2009, **8**(1):65–72.
45. Zilberman D, Gehring M, Tran RK, Ballinger T, Henikoff S: **Genome-wide analysis of Arabidopsis thaliana DNA methylation uncovers an interdependence between methylation and transcription.** *Nat Genet* 2007, **39**(1):61–69.
46. Mammucari C, Milan G, Romanello V, Masiero E, Rudolf R, Del Piccolo P, Burden SJ, Di Lisi R, Sandri C, Zhao J, Goldberg AL, Schiaffino S, Sandri M: **FoxO3 controls autophagy in skeletal muscle in vivo.** *Cell Metab* 2007, **6**(6):458–471.
47. Zhao J, Brault JJ, Schild A, Cao P, Sandri M, Schiaffino S, Lecker SH, Goldberg AL: **FoxO3 coordinately activates protein degradation by the autophagic/lysosomal and proteasomal pathways in atrophying muscle cells.** *Cell Metab* 2007, **6**(6):472–483.
48. Eash J, Olsen A, Breur G, Gerrard D, Hannon K: **FGFR1 inhibits skeletal muscle atrophy associated with hindlimb suspension.** *BMC Musculoskel Dis* 2007, **8**:32.
49. Giubellino A, Burke TR Jr, Bottaro DP: **Grb2 signaling in cell motility and cancer.** *Expert Opin Ther Tar* 2008, **12**(8):1021–1033.
50. Liu X, Liu M, Zhang J, Bai X, Ramos F, Van Remmen H, Richardson A, Liu FY, Dong LQ, Liu F: **Downregulation of Grb2 contributes to the insulin-sensitizing effect of calorie restriction.** *Am J Physiol Endoc M* 2009, **296**(5):E1067–E1075.
51. Morley JE: **Diabetes and aging: epidemiologic overview.** *Clin Geriatr Med* 2008, **24**(3):395–405. v.
52. Fraga MF, Agrelo R, Esteller M: **Cross-talk between aging and cancer: the epigenetic language.** *Ann Ny Acad Sci* 2007, **1100**:60–74.
53. Hellman A, Chess A: **Gene body-specific methylation on the active X chromosome.** *Science* 2007, **315**(5815):1141–1143.
54. Harris RA, Wang T, Coarfa C, Nagarajan RP, Hong C, Downey SL, Johnson BE, Fouse SD, Delaney A, Zhao Y, Olshen A, Ballinger T, Zhou X, Forsberg KJ, Gu J, Echipare L, O'Geen H, Lister R, Pelizzola M, Xi Y, Epstein CB, Bernstein BE, Hawkins RD, Ren B, Chung WY, Gu H, Bock C, Gnirke A, Zhang MQ, Haussler D, et al: **Comparison of sequencing-based methods to profile DNA methylation and identification of monoallelic epigenetic modifications.** *Nat Biotechnol* 2010, **28**(10):1097–1105.
55. Carmeli E, Reznick AZ, Coleman R, Carmeli V: **Muscle strength and mass of lower extremities in relation to functional abilities in elderly adults.** *Gerontology* 2000, **46**(5):249–257.
56. Zahn JM, Sonu R, Vogel H, Crane E, Mazan-Mamczarz K, Rabkin R, Davis RW, Becker KG, Owen AB, Kim SK: **Transcriptional profiling of aging in human muscle reveals a common aging signature.** *PLoS Genet* 2006, **2**(7):e115.
57. Takahashi J, Misawa M, Iwahashi H: **Oligonucleotide microarray analysis of age-related gene expression profiles in miniature pigs.** *PLoS One* 2011, **6**(5):e19761.
58. Zhao X, Mo D, Li A, Gong W, Xiao S, Zhang Y, Qin L, Niu Y, Guo Y, Liu X, Cong P, He Z, Wang C, Li J, Chen Y: **Comparative analyses by sequencing of transcriptomes during skeletal muscle development between pig breeds differing in muscle growth rate and fatness.** *PLoS One* 2011, **6**(5):e19774.
59. Fraga MF, Esteller M: **Epigenetics and aging: the targets and the marks.** *Trends Genet* 2007, **23**(8):413–418.
60. Erkens T, Van Poucke M, Vandesompele J, Goossens K, Van Zeveren A, Peelman LJ: **Development of a new set of reference genes for normalization of real-time RT-PCR data of porcine backfat and longissimus dorsi muscle, and evaluation with PPARC1A.** *BMC Biotechnol* 2006, **6**:41.
61. Nygard AB, Jorgensen CB, Cirera S, Fredholm M: **Selection of reference genes for gene expression studies in pig tissues using SYBR green qPCR.** *BMC Mol Biol* 2007, **8**:67.
62. Livak KJ, Schmittgen TD: **Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-Delta Delta C(T)} Method.** *Methods* 2001, **25**(4):402–408.
63. Li R, Yu C, Li Y, Lam T-W, Yiu S-M, Kristiansen K, Wang J: **SOAP2: an improved ultrafast tool for short read alignment.** *Bioinformatics* 2009, **25**(15):1966–1967.
64. Archibald AL, Bolund L, Churcher C, Fredholm M, Groenen MA, Harlizius B, Lee KT, Milan D, Rogers J, Rothschild MF, Uenishi H, Wang J, Schook LB: **Pig genome sequence-analysis and publication strategy.** *BMC Genomics* 2010, **11**:438.
65. Cawthon RM: **Telomere measurement by quantitative PCR.** *Nucleic Acids Res* 2002, **30**(10):e47.
66. Da Wei Huang BTS, Lempicki RA: **Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources.** *Nat Protoc* 2008, **4**(1):44–57.
67. Saeed A, Sharov V, White J, Li J, Liang W, Bhagabati N, Braisted J, Klapa M, Currier T, Thiagarajan M: **TM4: a free, open-source system for microarray data management and analysis.** *Biotechniques* 2003, **34**(2):374.
68. Bock C, Reither S, Mikeska T, Paulsen M, Walter J, Lengauer T: **BiQ Analyzer: visualization and quality control for DNA methylation data from bisulfite sequencing.** *Bioinformatics* 2005, **21**(21):4067–4068.

doi:10.1186/1471-2164-15-653

Cite this article as: Jin et al.: Genome-wide DNA methylation changes in skeletal muscle between young and middle-aged pigs. *BMC Genomics* 2014 **15**:653.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

