

REVIEW

Open Access

# Reflections on the immunology of tuberculosis: will we ever unravel the skein?

Maurizio de Martino\*, Luisa Galli, Elena Chiappini

## Abstract

Many and large dumps exist in our knowledge about *Mycobacterium tuberculosis* infection and disease in infants and children. We still do not understand why some individuals do acquire and others do not acquire the infection in the presence of the same risk factors. We do not understand why some individuals convert from latent to active tuberculosis and why other individuals convert from active to inactive tuberculosis even without treatment. As a matter of fact the immune system mounts a bouncing, robust and polyedral defence against *Mycobacterium tuberculosis*, but the bacillus is so much artful and dextrous that it has ahead from this immunological fierce accoutrements. *Mycobacterium tuberculosis* survival, multiplication, and transmission are largely favoured by the immune mechanisms. The granuloma itself is more *bacillus*- than host-protective. These abilities make *Mycobacterium tuberculosis* one of more successful human pathogens, but dumps in our knowledge and the counterproductive immunity hinder development of new diagnostics, therapies and vaccines. This occurs in front of an infection which engages one third of the world population and a disease which kills in a year about 1.5 million individuals worldwide. Understanding mechanisms and meaning of immune response in tuberculosis marks out the foundations of strategies with a view to prepare effective vaccines and reliable diagnostic tools as well as to build up therapeutic weapons. To gain these objectives is vital and urgent considering that tuberculosis is a common cause of morbidity and is a leading cause of death.

Tuberculosis is a paradox disconcerting for immunologists since a good immune response is developed in most individuals, but this response does not just eliminate the infection. On the contrary it aids the survival of *Mycobacterium tuberculosis*, assists its replication and transmission and induces the bacterium to adopt a silent underhand state from which it can reactivate as it pleases [1-4].

In addition, we would like to understand why some individuals do acquire whilst others do not acquire the infection (exposure conditions being equal), why some rare individuals get rid of infection, and how some individuals (without any treatment) are able to convert an active into an inactive infection [5].

## In spite of innate immunity, mycobacteria organize their bridgehead

Paraphrasing a famous aphorism of general Erwin Rommel, with regard to the amphibious warfare, but

well appropriate in the contest of tuberculosis, *all is decided the first day, which gets the longest day*.

Sometimes, soon after inhalation of *Mycobacterium tuberculosis*, bacilli are phagocitized by alveolar macrophages which can kill them and settle the case. This probably depends the coincidence of lucky factors including an intrinsic capacity of macrophages to generate reactive oxygen intermediates (ROI) and consequent microbicidal activity, a favorable inflammatory environment at the infection site, and not outstanding pathogenetic abilities of the inhaled strain [4].

If bacilli survive during the *longest day* (this is the commonest occurrence) they have won the fight. They proliferate at a logarithmic rate within dendritic cells and alveolar macrophages inducing the production of proinflammatory cytokines including interleukin (IL)-1- $\alpha$ , IL-1 $\beta$ , tumor necrosis factor *Ianus Bifrons*, IL-6, IL-12, and interferon (IFN)- $\gamma$  [2-4].

The role of these cytokines calls to mind the myth of *Ianus bifrons*: IFN- $\gamma$  regulates T-cell response to mycobacterial infection, promotes antigen presentation, activates

\* Correspondence: maurizio.demartino@unifi.it  
Department of Health Sciences, Meyer Children University Hospital,  
University of Florence, Florence, Italy

phagocytosis, production of ROI, TNF- $\alpha$  mediates early inflammatory responses against infectious agents also stimulating IL-1 and IL-6 production. The susceptibility to mycobacterial infection of TNF- $\alpha$  deficient mice (which is unable to undertake the formation of granuloma: but, it desirable or not? See afterwards), the increased susceptibility to *bacillus* of Calmette and Guérin (BCG; formerly *vaccin Bilié de Calmette et Guérin*) infection in mice who received anti-TNF- $\alpha$  antibodies and the mycobacterial susceptibility of human treated with TNF- $\alpha$  blockers should suggest its protective role. Also IL-6 is fundamental in developing the early inflammatory response. Its absence in mice may inhibit the maintenance of the protective (may be) IL-17 and causes a delayed IFN- $\gamma$  production and consequently determines an increased mycobacterial burden. It is well known that IL-12 promotes the TH-1 response and that its deficiency is associated in humans with an increased susceptibility to mycobacterial infection [3,4,6,7]

This first immune mechanism is largely mediated by molecules of the innate immunity. A large number of them are involved, including multiple pattern recognition receptors (that is toll-like receptors), C-type lectin receptor family, cytosolic pattern-recognition receptors, pyrin domain containing 3, and nucleotide-binding oligomerization domain protein 2. These molecules recognize as many again a number of mycobacterial products such as lipoproteins, lipomannans, and DNA [8-12].

The main consequence of recognition of mycobacterial molecules by innate immunity molecules is not only the expression of pro-inflammatory cytokines, but also the expression of cellular adhesion molecules and chemokines which mobilize and activate macrophages, dendritic cells, and neutrophils [11].

In many other infectious diseases this activity should be considered commendable and useful [1]. Instead, in tuberculosis, the inflow of these phagocytic cells supplies a lot of cellular niches that allow the *mycobacteria* to grow without hindrance [1,13,14]. The mycobacterial population grows and spreads to freshly recruited adjacent uninfected cells. This is preceded (admirable strategy) by a prolonged survival of the infected cell caused by the *Mycobacterium tuberculosis* itself. In fact, *Mycobacterium tuberculosis* inhibits the cell apoptosis also allowing a larger number of bacteria to be accumulated before bacteria go out of the dead cell [15]. By means of this mechanism, the induction of the adaptive immunity is delayed [15]. Also cell death is regulated by the *Mycobacterium of tuberculosis*. In fact, the virulence factor ESX1 type VII secretion system allows the bacteria to induce cell death, allowing them a well-timed exit [16]. It is well known that *Mycobacterium bovis* (used in the BCG vaccine) is devoid of the ESX1 type VII secretion system [17].

In few words: during this first stage of the immune response managed by the innate immunity, the *Mycobacterium tuberculosis* overcomes in big styles.

### The armed equilibrium during the adaptive immunity phase

Whereas the adaptive immunity to influenza virus is recruited after 20 hours, innate immunity against tubercular infection is activated after more than 40 days. Likely, this delay is due to the delayed transport by dendritic cells of live *mycobacteria* from the lungs to the draining lymph nodes [1,18]. This delayed transport is caused by the inhibition of transport (which should be stimulated by ligands for the CC-chemokine receptor 7) mediated by *Mycobacterium tuberculosis* [18]. The activation of the adaptive immunity, with an accumulation of effector CD4<sup>+</sup> and CD8<sup>+</sup> T-cells in the lungs, opens a period of trench warfare during which the bacterial load remains stable. Thus the adaptive immunity is not able to eliminate the infection but not even to reduce its entity. Since during this period of apparent immunological equilibrium *mycobacteria* accumulate mutations and a subset of bacteria insist to replicate [1].

Some of the mechanisms which cause this defeat are proper to immune cells and include expansion of Foxp3-expressing regulatory T-cells, defective antigen recognition, and reduced macrophage major histocompatibility complex expression and antigen processing [19]. Other mechanisms are proper to the *Mycobacterium tuberculosis* and include the resistance to the activation of macrophage function mediated by IFN- $\gamma$  and blinding of specific CD4<sup>+</sup> T-cell receptors caused by the downregulation of some genes which determine the vanishing of sensible microbial antigens. During this period, nitric oxide, carbon monoxide, and hypoxia predominate in the infection microenvironment causing the expression of a regulon (controlled by the signal transduction system DosR-DosS) which allows *Mycobacteria* use lipids as an alternative energy source [20].

During the adaptive immunity phase, a balance between T<sub>H</sub>1 (and consequent IFN- $\gamma$  production) and T<sub>H</sub>17 (and consequent IL-17 production which causes neutrophil recruitment and tissue damage) cells can control bacterial growth and limit immunopathology [21].

During this phase  $\gamma\delta$  T-cells play a significant defensive role [22]. *Mycobacterium tuberculosis* activates  $\gamma\delta$  T-cells but also infected monocytes are efficient accessory cells for  $\gamma\delta$  T-cells in a non-major histocompatibility complex-restricted also by means of IL-15. Analogously to  $\alpha\beta$  T-cell receptor<sup>+</sup>,  $\gamma\delta$  T-cells optimize their function interacting with adhesion molecules and other molecules including CD40-CD40L, CD28-B7.1/7.2.  $\gamma\delta$  T-cells produce both TNF- $\alpha$  and IFN- $\gamma$ . Release of IFN- $\gamma$  is more efficient than that from CD4<sup>+</sup> T-cells.

During this second phase, *Mycobacteria* do not gain ground, but gain a crucial strategic victory: they preserve and reinforce their bridgehead. Thus far the infected subject has the latent tuberculosis. One third of the world's population have been infected with *Mycobacterium tuberculosis* and have this form of tuberculosis [23].

### The attack: from latent to active tuberculosis

At the right moment (right by the point of view of *Mycobacterium tuberculosis*) the resuscitation-promoting factor may be activated, even years and years after initial infection, resuscitating bacteria from the nutrient-starved state and determining the establishment of an active symptomatic disease. Sometimes (especially in areas with a high prevalence of infection) active disease is determined by re-infection with a second strain. The latent infection turn into an active infection. This change may be due to *mycobacteria* as well as to the host [1].

Differently from viruses (which have fixed programs of expression of their genes), *mycobacteria* are able to change expression of their genes according to necessities and stimuli which are derived from the environment. Thus *mycobacteria* can change the expression of their surface antigens and evade T-cell recognition. In mice *mycobacteria* persist in the lungs also because, as soon as CD4<sup>+</sup> and CD8<sup>+</sup> T-cells enter the lungs, ESAT6 and Ag85B antigen expression is downregulated [24].

The host counterpart of re-activation may be an immunological low on guard [25]. Mechanisms are very poorly understood. Some model exist, including anti-TNF monoclonal antibodies used in a number of autoimmune disease, treatments with steroids, and the CD4<sup>+</sup> loss in patients with human immunodeficiency virus-type 1 (HIV-1) infection. Interestingly, re-activation in HIV-1 patients occurs largely before the nadir of CD4<sup>+</sup> cells, because HIV-1 selectively eliminates *Mycobacterium tuberculosis* antigen-specific CD4<sup>+</sup> cells at a higher rate than other antigen-specific CD4<sup>+</sup> cells.

possible that this is a more general mechanism of T-cell exhaustion in tuberculosis since the mycobacterial load in the lungs is inversely related to the proportion of blood antigen-specific T-cells. In addition, unfavourable polymorphisms of chemokines may alter the cell trafficking in the lungs a reduce the recruitment of cells which should contain the infection. It has been reported that some patients with latent TB have signatures similar to those in patients with active TB. They manifest a specific 86-transcript active TB. This signature is dominated by a neutrophil-driven IFN-inducible gene profile, consisting of both IFN- $\gamma$  and type I IFN- $\alpha\beta$  signalling [26]. These findings underscore the role of IFNs and neutrophils in the development of active tuberculosis

More vague are other possible mechanisms [1]. One is the supposed longer delay in the development of adaptive

immunity (observed in the diabetic mice and possibly caused by difficulties in the movement of dendritic cells from the lungs to the lymph nodes) which could explain the increased frequency of re-activation in the diabetic patient. It is an old story that thinnish people (besides their nutritional status) are more prone to tubercular re-activation. This association may depend on leptin which regulates, besides appetite, also growth and activity of T<sub>H</sub>1 cells.

As a proof that this second stage of immune response managed by adaptive immunity is a dynamic process of shooting war, the patients may present fever and *erythema nodosum* [27].

### Only sometimes immunity holds on tightly

Even though the mechanisms are unexplored, it is known that active infection is not a single track. In fact, a substantial but anyway minority portion of patients with active tuberculosis can succeed in getting rid of infection and lead to an inactive tuberculosis. Whereas patients with latent tuberculosis do not manifest chest-X-rays findings, subjects with inactive tuberculosis have chest-X-rays changes [1].

Memory T cells are distinguished by function and migratory capacity. Central memory T-cells have high proliferative activity and produce IL-2, whereas effector memory T-cells produce both IL-2 and IFN- $\gamma$ . Circulating effector memory T-cells indicates that an antigen-specific response persists, whereas circulating central memory T-cells indicate that infection has been controlled [28].

Subjects with inactive tuberculosis evidence CD4<sup>+</sup> central memory T-cell response. By contrast, patients with active tuberculosis evidence CD4<sup>+</sup> effector memory T-cell responses [29].

### *Mycobacteria* flood

The ultimate objective of all living beings is to disseminate as much as possible their genetic stock. It could seem unbelievable, but the adaptive immunity does favour the transmission of *mycobacteria* [1].

For transmission *mycobacteria* need a large surface exposed to the exterior. This surface allows the *mycobacteria* are expelled by an infected (and coughing) patient with active tuberculosis, reach and infect a previously uninfected individual [30,31]

A large surface is obtained by means the development of cavitory tuberculosis which is the result of lung tissue destruction. There are several lines of evidence, deriving from the HIV-1 infection model, suggesting that T-cells contribute to the development of cavitory tuberculosis: 1) HIV-1 infected subjects less frequently have cavitory tuberculosis; 2) HIV-1 infected patients less frequently transmit *mycobacteria* as compared to HIV-1 uninfected people; 3) in HIV-1 infected patients more the CD4<sup>+</sup>

T-cells and more the frequency of cavitory tuberculosis. Mechanisms are unknown. In very general terms it can be supposed that T-cells promote a high level of inflammation which damage lung tissue [32].

As a matter of fact, strains of *Mycobacterium tuberculosis* diverged genetically many and many years ago (to give an idea: during the upper paleolithic, roughly at the end of the last ice age), but all strains have conserved epitopes which are just that recognized by T-cells [33]. When something is preserved by a species for a long time during evolution it means that it is fundamental for survival of the species. From this we may conclude that for *Mycobacterium tuberculosis* it is an evolutionary advantage to be recognized by T-cells [34,35].

### **Granulomata: friend or foe?**

Richard Morton discovered in 1679 granuloma (at the time termed tubercles) and described it in his book entitled *Phthisiologia* more than two centuries before the discovery of the agent causing tuberculosis [36].

Epithelial granulomata are the hallmark of tuberculosis. For a long time it has been believed that these evolutionarily primitive structures control the infection since they wall out bacilli. Unfortunately, it is not so [37-39].

An (apparently) hardened and dynamic arsenal of well organized immunity cells is concentrated in tubercular granulomata. T-cells, B-cells, neutrophils, fibroblasts, macrophages (infected or uninfected), epithelioid cells derived from macrophages, foamy macrophages, and multinucleated giant Langerhans cells make up the tubercular granulomata. Granulomata development starts shortly after infection. At the beginning an innate granuloma develops which is composed by neutrophils and alveolar macrophages. Unfortunately, *mycobacteria* within granulomata are able to disarm alveolar macrophages arresting the phagolysosome fusion thus blocking the killing of internalized *mycobacteria*. *Mycobacteria* also drive in infected and harmless (by the point of view of *mycobacteria*) macrophages the production of chemoattractant and pro-inflammatory cytokines. In addition, the release by *mycobacteria* of the 6 kDa early secretory antigenic target causes the activation of the epithelium which stimulates the recruitment of macrophages causing the expression of matrix metalloproteinase-9 (MMP-9). Also the expression of MMP-1 is driven by *mycobacteria* [40-42]

The result of this frantic fascination towards immune cells is the development of the immune granuloma and the multiplication of fresh *pabulum* for *mycobacteria*. Mechanisms involved in the promotion and development of granulomata are mostly driven by *mycobacteria* and this clearly suggests that all is done in the *mycobacteria*'s own interest [42].

TNF is believed to be fundamental for defence against *Mycobacteria* and for granuloma formation. However,

its iatrogenic inhibition disrupts granuloma allowing replication of *mycobacteria* and thus increases the efficacy of chemotherapies [43].

Thus, contrary to the traditional view of a protective role of granulomata, it is now convincing that granulomata favour the persistence and diffusion of the *mycobacterium* more than its segregation or even its elimination.

### **B-cells and antibodies: intelligence agents, ancillary services, or useless?**

The role of humoral immunity in tuberculosis is, at the very least, evanescent. On the one hand there are the inconsistent findings in studies of passive immunization, the irrelevance of humoral deficiencies and B-cell immune defects in *Mycobacterium tuberculosis* infection and disease and the ambiguous results in knockout mice besides the general rule that antibodies are nearly ineffective against intracellular agents [44-47].

On the other hand, there are well known exceptions to this rule (infections caused by species of *Leishmania*, *Chlamydia*, *Francisella*), the levels of specific antibodies correlate to the efficacy of BCG vaccine, mice knockout for polymeric immunoglobulin receptors have a higher susceptibility to *mycobacterial* infections (is there a role of mucosal dimeric IgA?), in animal models monoclonal antibodies against *mycobacterial* antigens have a beneficial effect in tuberculosis. In addition, there is evidence that B-cells are a fundamental component of granulomata and follicle-like B-cell aggregates (site of antigen presentation to T-cells?) are a characteristic feature of granulomatous progression in *mycobacterial* infection.

In a broader view it is well known that B-cells and humoral immunity collaborate with T-cells and cellular immunity in almost all infectious diseases and B-cells act as antigen-presenting cells to T-cells and/or polarize T-cell activity influencing the production of cytokines. B-cells are deeply involved in the production in the lungs of the anti-inflammatory IL-10.

It is a peculiar finding that sometimes these mechanisms work in the presence of B-cells, but in the absence of antibodies. Sometimes the opposite occurs. Even more singularly, in tuberculosis and particularly in active tuberculosis there is an significantly elevated prevalence of subjects with autoantibodies [48].

### **Conclusions**

Tuberculosis is a major international public health. Great efforts are being made by science to oppose this plague. But, candidly, translational results are still stalling.

Each human (and his/her immune mechanisms) is very similar but also largely differ from other humans (and their immune mechanisms). Maybe that personalized medicine is the way to unravel the skein. Results of how immune gene polymorphisms and, in particular,

polymorphisms concerning how toll-like receptor and autophagy-related genes influence tuberculosis susceptibility and clinical course are consistent with line of thinking [49-51]. Maybe that some immunological dogma should be revised. One of many is the ambiguous role of IFN- $\gamma$  [52]. Maybe that some branch of research should be implemented. One is that concerning the unaccountably underestimated role of  $\gamma\delta$  T-cells [22]. Maybe that, if we want to win the fight against *Mycobacterium tuberculosis*, we must renounce to conformism. It is time for a different way of thinking.

#### Abbreviations

BCG: bacillus Calmette Guerin; IFN: interferon; IL: interleukin; MMP: metalloproteinase; ROI: rective oxygen intermeffiate; TNF: tumor necrosis factor.

#### Competing interests

The authors declare that they have no competing interests.

#### Authors' contributions

MdM conceived the idea, collected data from the literature, and drafted the manuscript. EC and LG helped to draft and critically reviewed the manuscript.

#### Declarations

Funding for this supplement has come from a grant of the Italian Health Ministry /Young Research Project.

This article has been published as part of *BMC Infectious Diseases* Volume 14 Supplement 1, 2014: Highlights in Pediatric Tuberculosis. The full contents of the supplement are available online at <http://www.biomedcentral.com/bmcinfectdis/supplements/14/S1>

Published: 8 January 2014

#### References

- Ernst JD: The immunological life cycle of tuberculosis. *Nature Rev* 2012, **12**:581-591.
- Zumla A, Raviglione M, Hafner R, von Reyn F: Tuberculosis. *N Engl J Med* 2013, **368**:745-755.
- Cooper AM, Torrado E: Protection versus pathology in tuberculosis: recent insights. *Curr Opin Immunol* 2012, **24**:431-437.
- Zuñiga J, Torres-Garcia D, Santos Mendoza T, Rodriguez-Reyna TS, Granados J, Yunis EJ: Cellular and humoral mechanisms involved in the control of tuberculosis. *Clin Develop Immunol* 2012, ID 193923 doi: 10.1155/2012/193923.
- Roy RB, Whittaker E, Kampmann B: Current understanding of the immune response to tuberculosis in children. *Curr Opin Infect Dis* 2012, **25**:250-257.
- Cooper AM, Adams LB, Dalton DK, Appleberg R, Ehlers S: IFN- $\gamma$  and NO in mycobacterial diseases: new jobs for old hands. *Trends Microbiol* 2002, **10**:221-222.
- Kaufmann SHE: How can immunology contribute to the control of tuberculosis? *Nature Rev Immunol* 2001, **1**:20-30.
- Kim JS, Kim WS, Choi HG, Jang B, Lee K, Park JH, Kim HJ, Cho SN, Shin SJ: Mycobacterium tuberculosis RpfB drives Th1-type T cell immunity via a TLR4-dependent activation of dendritic cells. *J Leukoc Biol* 2013, [Epub ahead of print].
- Bafica A, Scanga CA, Feng CG, Leifer C, Cheever A, Sher A: TLR9 regulates Th1 responses and cooperates with TLR2 in mediating optimal resistance to Mycobacterium tuberculosis. *J Exp Med* 2005, **202**:1715-1724.
- Banaiee N, Kincaid EZ, Buchwald U, Jacobs WR Jr, Ernst JD: Potent inhibition of cytokines in the balance of protection and pathology during mycobacterial infections. *Adv Exp Med Biol* 2013, **783**:121-140.
- Thada S, Valluri VL, Gaddam SL: Influence of toll-like receptor gene polymorphisms to tuberculosis susceptibility in humans. *Scand J Immunol* 2013, **78**:221-229.
- Torrado E, Cooper AM: Macrophage responses to IFN- $\gamma$  by live virulent *Mycobacterium tuberculosis* is independent of mature mycobacterial lipoproteins but dependent on TLR2. *J Immunol* 2006, **176**:3019-3027.
- Chackerman AA, Alt JM, Perera TV, Dascher CC, Behar SM: Dissemination of *Mycobacterium tuberculosis* is influenced by host factors and precedes the initiation of T cell immunity. *Infect Immun* 2002, **70**:4501-4509.
- Huynh KK, Joshi SA, Brown EJ: A delicate dance: host response to mycobacteria. *Curr Opin Immunol* 2011, **23**:464-472.
- Blomgran R, Desvignes L, Briken V, Ernst JD: Mycobacterium tuberculosis inhibits neutrophil apoptosis, leading to delayed activation of naive CD4 T cells. *Cell Host Microbe* 2012, **11**:81-90.
- Abdallah AM, Bestebroer J, Savage ND, de Punder K, van Zon M, Wilson L, Korbee CJ, van der Sar AM, Ottenhoff TH, van der Wel NN, Bitter W, Peters PJ: Mycobacterial secretion systems ESX-1 and ESX-5 play distinct roles in host cell death and inflammasome activation. *J Immunol* 2011, **187**:4744-4753.
- Pym A, Brodin P, Brosch R, Huerre M, Cole S: Loss of RD1 contributed to the attenuation of the live tuberculosis vaccines *Mycobacterium bovis* BCG and *Mycobacterium microti*. *Mol Microbiol* 2002, **46**:709-717.
- Scott-Browne JP, Scott-Browne JP, Shafiani S, Tucker-Heard G, Ishida-Tsubota K, Fontenot JD, Rudensky AY, Bevan MJ, Urdahl KB: Expansion and function of Foxp3-expressing T regulatory cells during tuberculosis. *J Exp Med* 2007, **204**:2159-2169.
- Reiley WW, Calayag MD, Wittmer ST, Huntington JL, Pearl JE, Fountain JJ, Martino CA, Roberts AD, Cooper AM, Winslow GM, Woodland DL: ESAT-6-specific CD4 T cell responses to aerosol *Mycobacterium tuberculosis* infection are initiated in the mediastinal lymph nodes. *Proc Natl Acad Sci USA* 2008, **105**:10961-10966.
- Black GF, Thiel BA, Ota MO, Parida SK, Adegbola R, Boom WH, Dockrell HM, Franken KL, Friggen AH, Hill PC, Klein MR, Lalor MK, Mayanja H, Schoolnik G, Stanley K, Welding K, Kaufmann SH, Walz G, Ottenhoff TH: Immunogenicity of novel DosR regulon-encoded candidate antigens of *Mycobacterium tuberculosis* in three high-burden populations in Africa. *Clin Vaccine Immunol* 2009, **16**:1203-1212.
- Torrado E, Cooper AM: IL-17 and Th17 cells in tuberculosis. *Cytokine Growth Factor Rev* 2010, **21**:455-462.
- Meraviglia S, El Daker S, Dieli F, Martini F, Martino A:  $\gamma\delta$  T cells cross-link innate and adaptive immunity in *Mycobacterium tuberculosis* infection. *Clin Dev Immunol* 2011, ID 587315. doi: 10.1155/2011/587315.
- World Health Organization: Tuberculosis., Fact sheet no104, February 2013.
- Rogerson BJ, Jung YJ, LaCourse R, Ryan L, Enright N, North RJ: Expression levels of *Mycobacterium tuberculosis* antigen-encoding genes versus production levels of antigen-specific T cells during stationary level lung infection in mice. *Immunology* 2006, **118**:195-201.
- Geldmacher C, Zumla A, Hoelscher M: Interaction between HIV and *Mycobacterium tuberculosis*: HIV-1-induced CD4 T-cell depletion and the development of active tuberculosis. *Curr Opin HIV AIDS* 2012, **7**:268-275.
- Berry MP, Graham CM, McNab FW, Xu Z, Bloch SA, Oni T, Wilkinson KA, Banchereau R, Skinner J, Wilkinson RJ, Quinn C, Blankenship D, Dhawan R, Cush JJ, Mejias A, Ramilo O, Kon OM, Pascual V, Banchereau J, Chaussabel D, O'Garra A: An interferon inducible neutrophil-driven blood transcriptional signature in human tuberculosis. *Nature* 2010, **466**:973-977.
- Poulsen A: Some clinical features of tuberculosis. *Acta Tuberc. Scand* 1950, **24**:311-346, I. Incubation period.
- Harari A, Dutoit V, Cellerai C, Bart P-A, Du Pasquier RA, Pantaleo G: Functional signatures of protective antiviral T-cell immunity in human virus infections. *Immunol Rev* 2006, **211**:236-254.
- Millington KA, Gooding S, Hinks TS, Reynolds DJ, Lalvani A: Mycobacterium tuberculosis-specific cellular immune profiles suggest bacillary persistence decades after spontaneous cure in untreated tuberculosis. *J Infect Dis* 2010, **202**:1685-1689.
- Rodrigo T, Caylà JA, García de Olalla P, Galdós-Tangüis H, Jansà JM, Miranda P, Brugal T: Characteristics of tuberculosis patients who generate secondary cases. *Int J Tuberc Lung Dis* 1997, **1**:352-357.
- Kaplan G, Post FA, Moreira AL, Wainwright H, Kreiswirth BN, Tanverdi M, Mathema B, Ramaswamy SV, Walther G, Steyn LM, Barry CE 3rd, Bekker LG: Mycobacterium tuberculosis growth at the cavity surface: a microenvironment with failed immunity. *Infect Immun* 2003, **71**:7099-7108.
- Kwan CK, Ernst JD: HIV and tuberculosis: a deadly human syndemic. *Clin Microbiol Rev* 2011, **24**:351-376.

33. Kapur V, Whittam TS, Musser JM: **Is *Mycobacterium tuberculosis* 15,000 years old?** *J Infect Dis* 1994, **170**:1348-1349.
34. Comas I, Chakravarti J, Small PM, Galagan J, Niemann S, Kremer K, Ernst JD, Gagneux S: **Human T cell epitopes of *Mycobacterium tuberculosis* are evolutionarily hyperconserved.** *Nature Genet* 2010, **42**:498-503.
35. Blouin Y, Hauck Y, Soler C, Fabre M, Vong R, Dehan C, Cazajous G, Massoure PL, Kraemer P, Jenkins A, Garnotel E, Pourcel C, Vergnaud G: **Significance of the identification in the Horn of Africa of an exceptionally deep branching *Mycobacterium tuberculosis* clade.** *PLoS One* 2012, **7**:e52841, doi: 10.1371/journal.pone.0052841.
36. Morton R: **Phthisiologia: or, a treatise of consumptions: wherein the difference, nature, causes, signs, and cure of all sorts of consumptions are explained.** W and J Innys, London; 1720.
37. Shaler CR, Horvath CN, Jeyanathan M, Xing Z: **Within the enemy's camp: contribution of the granuloma to the dissemination, persistence and transmission of *Mycobacterium tuberculosis*.** *Front Immunol* 2013, **4**, doi: 10.3389/fimmu.2013.00030.
38. Guirado E, Schlesinger LS: **Modeling the *Mycobacterium tuberculosis* granuloma – the critical battlefield in host immunity and disease.** *Front Immunol* 2013, **4**, doi:10.3389/fimmu.2013.00098.
39. Ramakrishnan L: **Revisiting the role of the granuloma in tuberculosis.** *Nat Rev Immunol* 2012, **12**:352-366.
40. Volkman HE, Pozos TC, Zheng J, Davis JM, Rawls JF, Ramakrishnan L: **Tuberculous granuloma induction via interaction of a bacterial secreted protein with host epithelium.** *Science* 2010, **327**:466-469.
41. Ramakrishnan L: **Looking within the zebrafish to understand the tuberculous granuloma.** *Adv Exp Med Biol* 2013, **783**:251-266.
42. Davis JM, Ramakrishnan L: **The role of granuloma in expansion and dissemination of early tuberculous infection.** *Cell* 2009, **136**:37-49.
43. Bourigault ML, Vacher R, Rose S, Olleros ML, Janssens JP, Quesniaux VF, Garcia I: **Tumor necrosis factor neutralization combined with chemotherapy enhances *Mycobacterium tuberculosis* clearance and reduces lung pathology.** *Am J Clin Exp Immunol* 2013, **2**:124-134.
44. Phuah JY, Mattila JT, Lin PL, Flynn JL: **Activated B cells in the granulomas of nonhuman primates infected with *Mycobacterium tuberculosis*.** *Am J Pathol* 2012, **181**:508-514.
45. Achkar JM, Casadevall A: **Antibody-mediated immunity against tuberculosis: implications for vaccine development.** *Cell Host Microbe* 2013, **13**:250-262.
46. Kozakiewicz L, Phuah J, Flynn J, Chan J: **The role of B cells and humoral immunity in *Mycobacterium tuberculosis* infection.** *Adv Exp Med Biol* 2013, **783**:225-250.
47. Maglione PJ, Chan J: **How B cells shape the immune response against *Mycobacterium tuberculosis*.** *Eur J Immunol* 2009, **39**:676-686.
48. Shen CY, Hsieh SC, Yu CL, Wang JY, Lee LN, Yu CJ: **Autoantibody prevalence in active tuberculosis: reactive or pathognomonic?** *BMJ Open* 2013, **3**, doi:pii: e002665. 10.1136/bmjopen-2013-002665.
49. Thada S, Valluri VL, Gaddam SL: **Influence of toll-like receptor gene polymorphisms to tuberculosis susceptibility in humans.** *Scand J Immunol* 2013, **78**:221-229.
50. Azad AK, Sadee W, Schlesinger LS: **Innate gene polymorphisms in tuberculosis.** *Infect Immun* 2012, **80**:3343-3359.
51. Songane M, Kleinnijenhuis J, Netea MG, van Crevel R: **The role of autophagy in host defence against *Mycobacterium tuberculosis* infection.** *Tuberculosis* 2012, **92**:388-396.
52. Abebe F: **Is interferon-gamma the right marker for bacille Calmette-Guérin-induced immune protection? The missing link in our understanding of tuberculosis immunology.** *Clin Exp Immunol* 2012, **169**:213-219.

doi:10.1186/1471-2334-14-S1-S1

**Cite this article as:** de Martino *et al.*: Reflections on the immunology of tuberculosis: will we ever unravel the skein? *BMC Infectious Diseases* 2014 **14**(Suppl 1):S1.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

