









ORIGINAL RESEARCH

Hospital Variation in Time to Endovascular Treatment for Ischemic Stroke: What Is the Optimal Target for Improvement?

Sanne J. den Hartog , MD; Hester F. Lingsma , MD, PhD; Pieter-Jan van Doormaal , MD; Jeannette Hofmeijer , MD, PhD; Lonneke S. F. Yo, MD, PhD; Charles B. L. M. Majoie , MD, PhD; Diederik W. J. Dippel , MD, PhD; Aad van der Lugt , MD, PhD; Bob Roozenbeek , MD, PhD; on behalf of the MR CLEAN Registry investigators*

BACKGROUND: Time to reperfusion in patients with ischemic stroke is strongly associated with functional outcome and may differ between hospitals and between patients within hospitals. Improvement in time to reperfusion can be guided by between-hospital and within-hospital comparisons and requires insight in specific targets for improvement. We aimed to quantify the variation in door-to-reperfusion time between and within Dutch intervention hospitals and to assess the contribution of different time intervals to this variation.

METHODS AND RESULTS: We used data from the MR CLEAN (Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands) Registry. The door-to-reperfusion time was subdivided into time intervals, separately for direct patients (door-to-computed tomography, computed tomography-to-computed tomography angiography [CTA], CTA-to-groin, and groin-to-reperfusion times) and for transferred patients (door-to-groin and groin-to-reperfusion times). We used linear mixed models to distinguish the variation in door-to-reperfusion time between hospitals and between patients. The proportional change in variance was used to estimate the amount of variance explained by each time interval. We included 2855 patients of 17 hospitals providing endovascular treatment. Of these patients, 44% arrived directly at an endovascular treatment hospital. The between-hospital variation in door-to-reperfusion time was 9%, and the within-hospital variation was 91%. The contribution of case-mix variables on the variation in door-to-reperfusion time was marginal (2%–7%). Of the between-hospital variation, CTA-to-groin time explained 83%, whereas groin-to-reperfusion time explained 15%. Within-hospital variation was mostly explained by CTA-to-groin time (33%) and groin-to-reperfusion time (42%). Similar results were found for transferred patients.

CONCLUSIONS: Door-to-reperfusion time varies between, but even more within, hospitals providing endovascular treatment for ischemic stroke. Quality of stroke care improvements should not only be guided by between-hospital comparisons, but also aim to reduce variation between patients within a hospital, and should specifically focus on CTA-to-groin time and groin-to-reperfusion time.

Key Words: brain ischemia ■ quality improvement ■ reperfusion ■ stroke ■ thrombectomy

In patients with acute ischemic stroke, shorter times to endovascular treatment (EVT) are strongly associated with more favorable outcomes.^{1,2} This association is found on several outcomes, such as

mortality, reperfusion grade after EVT, and the functional outcome, measured with the modified Rankin Scale score at 3 months.^{1,2} Because of this established strong association, process measures, such

Correspondence to: Sanne Jacoba den Hartog, Erasmus MC, University Medical Center, Department of Neurology, Department of Radiology and Nuclear Medicine, Department of Public Health, Room Ee2240, PO Box 2040, 3000 CA Rotterdam, the Netherlands. E-mail: s.denhartog@erasmusmc.nl

*A complete list of the MR CLEAN Registry investigators can be found in the Appendix at the end of the article.

Supplemental Material for this article is available at <https://www.ahajournals.org/doi/suppl/10.1161/JAHA.121.022192>

For Sources of Funding and Disclosures, see page 11.

© 2021 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

JAHA is available at: www.ahajournals.org/journal/jaha

CLINICAL PERSPECTIVE

What Is New?

- In patients with acute ischemic stroke, variation in door-to-reperfusion time is mostly explained by computed tomography angiography-to-groin time and groin-to-reperfusion time, and less by patient characteristics.
- There is more variation in door-to-reperfusion time within hospitals than between hospitals.

What Are the Clinical Implications?

- Quality of stroke care improvements should not only be guided by between-hospital comparisons, but also aim to reduce variation between patients within a hospital, and should specifically focus on computed tomography angiography-to-groin time and groin-to-reperfusion time.

Nonstandard Abbreviations and Acronyms

CTA	computed tomography angiography
EVT	endovascular treatment
MR CLEAN	Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands

as door-to-reperfusion time, are suitable for measuring quality of stroke care. Quality improvement can be guided by the comparison of hospitals with each other and by the comparison of patients within a hospital. Insight in variation between and within hospitals, including the contribution of different time intervals (door-to-computed tomography [CT], CT-to-CT angiography [CTA], CTA-to-groin, and groin-to-reperfusion times) to this variation, may contribute to improvement of quality of ischemic stroke care.

We aimed to quantify the variation in door-to-reperfusion time between and within Dutch EVT hospitals and to assess the contribution of different time intervals to this variation.

METHODS

We used data from the MR CLEAN (Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands) Registry. This is a prospective observational study in all 17 centers performing EVT in the Netherlands. All patients who underwent EVT for acute ischemic stroke in the anterior or posterior circulation between March 16, 2014, and November 1, 2017, were

registered. EVT was defined as entry into the angiography suite and having arterial puncture. Details on the study design and methods were described previously.³ The central medical ethics committee of the Erasmus MC, University Medical Center Rotterdam, the Netherlands, evaluated the study protocol and granted permission to perform the study as a registry (MEC-2014-235). In compliance with the General Data Protection Regulation, source data are not available for other researchers. Information about analytic methods, study materials, and scripts of the statistical analyses are available from the corresponding author on reasonable request.

Patients

For the purpose of this study, we used the following inclusion criteria: age of ≥ 18 years; groin puncture within 6.5 hours after stroke onset; treatment in a MR CLEAN Registry trial center; intracranial proximal arterial occlusion in the anterior circulation (intracranial carotid artery, internal carotid artery terminus, middle [M1/M2] cerebral artery, or anterior [A1/A2] cerebral artery) demonstrated by CTA, magnetic resonance angiography, or digital subtraction angiography. The analyzed data comprised patients who were treated with EVT between March 16, 2014, and November 1, 2017. We focused this analysis on patients who presented at the emergency departments of the participating hospitals. Patients with in-hospital stroke were excluded from this analysis.

Outcomes

The primary outcome measure was door-to-reperfusion time, defined as the total time in minutes between arrival of the patient in an EVT hospital and the end of the EVT procedure. The latter was defined as time Expanded Thrombolysis in Cerebral Infarction 2B was achieved, or if no reperfusion was achieved, time of the last contrast bolus.³ Analyses were stratified for patients presented directly at an EVT hospital (direct patients) and patients transferred from a primary stroke center (transferred patients). The total door-to-reperfusion time was divided in intervals formed by the sequential performed imaging and treatments during the entire process from door to reperfusion. The time intervals for direct patients were the following: door-to-CT, CT-to-CTA, CTA-to-groin, and groin-to-reperfusion (or last contrast bolus) times. In transferred patients, diagnostic imaging (CT and CTA) was already performed in the primary stroke center. Therefore, we limited the time intervals for this group to door-to-groin time and groin-to-reperfusion time (or last contrast bolus). We focused on intervals that can directly be influenced by the stroke teams in the EVT hospitals.

Missing Data

All baseline data were reported as crude data. For each time interval, we investigated extreme minimum and maximum outliers. These outliers were considered as incorrect and recoded as missing (Table S1). All missing variables were imputed with multiple imputation with R (package, MICE) based on relevant covariates and outcomes.

Statistical Analysis

For descriptive purposes, we grouped hospitals into tertiles based on the mean door-to-reperfusion time per hospital for patients directly presented at an EVT hospital. We used an ordinal logistic regression model to analyze the association between door-to-reperfusion time (as a continuous variable) and the modified Rankin Scale score at 3 months and presented common odds ratio (OR) with 95% CI.

We used linear mixed models to examine the amount of variation in the door-to-reperfusion time between and within hospitals, explained by each time interval. All models included hospital-specific random intercepts to account for patient clustering within each hospital.^{4,5}

We started with model 1, which only contained a hospital-specific random intercept and no covariates. This model provides the intraclass correlation coefficient, which describes the proportion of the total variance that is attributable to clustering within hospitals, in our case the between-hospital variance in door-to-reperfusion time. The remaining total variance is attributable to within-hospital variation between patients. To investigate the contribution of case-mix on the between-hospital and within-hospital variation in door-to-reperfusion time, we added case-mix variables to the model. Model 2 contained the following variables: a hospital-specific random intercept, age, sex, history of atrial fibrillation, history of hypertension, history of diabetes, history of myocardial infarction, history of peripheral artery disease, history of ischemic stroke, history of hyperlipidemia, prestroke modified Rankin Scale score, baseline National Institutes of Health Stroke Scale score, onset-to-door time, admission during off hours, and the anatomical location of occluded artery. To investigate how much of the between-hospital and within-hospital variation in door-to-reperfusion time was attributable to each time interval, after adjustment for case-mix, we added each time interval to model 2 in a cumulative way. Model 3A contains a hospital-specific random intercept, case-mix, and door-to-CT time. Model 3B contains a hospital-specific random intercept, case-mix, door-to-CT time, and CT-to-CTA time. Model 3C contains a hospital-specific random intercept, case-mix, door-to-CT time, CT-to-CTA time, and CTA-to-groin time. Model 3D contains a

hospital-specific random intercept, case-mix, door-to-CT time, CT-to-CTA time, CTA-to-groin time, and groin-to-reperfusion time. The proportional change in variance was used to estimate the amount of variance explained by each model compared with model 1.^{6,7} To calculate the attribution of each variable individually to the variance, the proportional changes in variance of the models were subtracted.

In a separate analysis, we investigated the influence of potentially delaying factors on the door-to-reperfusion time of patients who presented directly at the EVT hospital, by adding systolic blood pressure (≥ 185 mm Hg) requiring blood pressure-lowering therapy, intravenous alteplase treatment, and general anesthesia to the case-mix adjusted model. For transferred patients, models 1 and 2 were the same as described above. Model 3A contains a hospital-specific random intercept, case-mix, and door-to-groin time. Model 3B contains a hospital-specific random intercept, case-mix, door-to-groin time, and groin-to-reperfusion time. We considered repetition of imaging (CT or CTA) in an EVT hospital to be a delaying factor for transferred patients. All statistical analyses were performed with R statistical software (version 3.6.1).

$$PCVh = \frac{Vh1 - Vh3}{Vh1}$$

h indicates hospital level (between-hospital variance) could be replaced by p: patient level (within-hospital variance); h1 or p1, model with only hospital as random intercept; h3 or p3, model with case-mix variables and time intervals; PCVh, proportional change in variance at hospital level; and V, variance.

RESULTS

In total, 3637 patients were registered in the MR CLEAN Registry between March 16, 2014, and November 1, 2017. First, we excluded 457 patients, mostly because of occlusion in the posterior circulation or treatment starting after 6.5 hours from the onset of symptoms (Figure 1). Then, we excluded 325 patients with in-hospital stroke. The remaining 2855 patients were used for analysis. From these patients, 1244 (44%) presented directly at an EVT hospital and 1611 (56%) were transferred from a primary stroke center to an EVT hospital.

Patient Characteristics

For patients who arrived directly at an EVT hospital, median door-to-reperfusion time was 147 minutes (interquartile range, 116–185 minutes), and the median varied between hospitals from 121 to 184 minutes. The median door-to-reperfusion time for transferred patients was 93 minutes (interquartile range, 70–283




Figure 1. Flowchart of MR CLEAN (Multicenter Randomized Clinical Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands) Registry patients selected for analysis.

EVT indicates endovascular treatment.

minutes) and varied between hospitals from 75 to 112 minutes. There were only minor differences in age and medical history between the tertiles of hospitals (Table 1). There was a difference in the percentage of transferred patients between the tertiles (62% versus 61% versus 37%). There was also a difference in the use of general anesthesia between the tertiles (7% versus 27% versus 54%). Of the case-mix variables, only location of occluded artery, onset-to-door time, and admission during off hours were associated with door-to-reperfusion time (Table S2). An increased door-to-reperfusion time was associated with worse functional outcome (modified Rankin Scale score at 3 months) for direct and transferred patients (per 10 minutes, common OR 0.92 [95% CI: 0.91–0.94] and common OR 0.93 [95% CI: 0.91–0.95], respectively).

Variation Between Hospitals for Patients Directly Presented at an EVT Hospital

Of the total variation in door-to-reperfusion time for patients directly presented at an EVT hospital, 9% was attributable to between-hospital variation (Table 2). Case-mix variables explained only 2% of the total between-hospital variation in door-to-reperfusion time (Figure 2A). CTA-to-groin puncture

time explained 83% of the between-hospital variation in case-mix adjusted door-to-reperfusion time, whereas groin-to-reperfusion time explained an additional 15%. The correlation between these time intervals and door-to-reperfusion time is shown in Figure S1A and S1B. In a separate analysis, delaying factors in door-to-reperfusion time, such as a high baseline systolic blood pressure (≥ 185 mm Hg), treatment with intravenous alteplase treatment, or general anesthesia, explained 8% of the between-hospital variation in door-to-reperfusion time.

Variation Within Hospitals for Patients Directly Presented at an EVT Hospital

Of the total variation in door-to-reperfusion time for patients directly presented at an EVT hospital, 91% was attributable to within-hospital variation (Table 2 and Figure 3). Case-mix variables explained 7% of the total within-hospital variation in door-to-reperfusion time (Figure 2A). CT-to-CTA time explained an additional 13%, CTA-to-groin time explained 33%, and groin-to-reperfusion time explained 42% of the within-hospital variation in door-to-reperfusion time. Delaying factors explained 3% of the within-hospital variation in door-to-reperfusion time.

Table 1. Baseline Characteristics per Hospital Tertiles of the Mean Door-to-Reperfusion Time of Direct Patients

Characteristic	Total	Tertile 1 (range, 128–146 min)	Tertile 2 (range, 146–158 min)	Tertile 3 (range, 163–188 min)	P value
No. of patients	2855	1006	1277	572	
No. of centers	17	5	6	6	
Age, y	72 (61–81) (2855)	73 (63–81)	71 (59–79)	72 (62–82)	0.072
Men, %	52 (1491/2855)	52	52	53	0.951
Atrial fibrillation, %	24 (662/2818)	27	21	24	0.002
Hypertension, %	52 (1455/2797)	59	49	46	<0.001
Diabetes, %	16 (440/2833)	15	16	14	0.619
Myocardial infarction, %	14 (381/2798)	15	14	9	0.003
Peripheral artery disease, %	9 (246/2798)	13	6	8	<0.001
Previous ischemic stroke, %	16 (463/2832)	17	16	15	0.650
Hyperlipidemia, %	30 (816/2728)	39	25	25	<0.001
Baseline NIHSS score	16 (11–19) (2816)	16 (11–20)	15 (11–19)	16 (11–19)	0.083
Prestroke modified Rankin scale score, %					0.105
0	70 (1943/2793)	68	68	75	
1	13 (363/2793)	15	13	10	
2	7 (193/2793)	7	8	5	
≥3	11 (294/2793)	10	11	10	
Transfer from primary stroke center, %	56 (1611/2855)	62	61	37	<0.001
Intravenous alteplase treatment, %	80 (2278/2847)	80	80	80	0.743
Level of occlusion, %*					0.010
ICA	5 (130/2740)	4	5	7	
ICA-T	22 (590/2740)	23	20	22	
M1	58 (1590/2740)	58	60	56	
M2	15 (409/2740)	15	15	16	
Other (M3/anterior)	1 (21/2740)	0.2	1	0.4	
Collaterals, %					0.054
Grade 0	6 (171/2673)	8	5	7	
Grade 1	37 (978/2673)	37	35	39	
Grade 2	39 (1028/2673)	37	40	39	
Grade 3	19 (496/2673)	18	20	16	
Onset-to-door time, min	135 (65–188) (2753)	140 (81–191)	137 (70–189)	103 (52–175)	<0.001
Off hours, %†	64 (1837/2855)	64	65	63	0.491
General anesthesia, %	26 (686/2677)	7	27	54	<0.001
Systolic blood pressure (≥185 mm Hg), %	9 (249/2784)	8	10	9	0.164

Categorical variables are presented as percentage (n/N). Continuous variables are presented as median (interquartile range) (N). ICA indicates intracranial carotid artery; ICA-T, internal carotid artery terminus; middle (M1/M2/M3) cerebral artery; and NIHSS, National Institutes of Health Stroke Scale.

*Based on computed tomography angiography.

†Admission between 5:00 PM and 8:00 AM, on weekends, or a national holiday.

Variation Between and Within Hospitals for Transferred Patients

Of the total variation in door-to-reperfusion time for transferred patients, 3% was attributable to between-hospital variation (Table 3). This between-hospital

variation was explained by door-to-groin time (56%) and groin-to-reperfusion time (44%) (Figure 2B). The delaying factor of repetition of imaging explained 31% of the between-hospital variation in door-to-reperfusion time. The within-hospital variation in door-to-reperfusion

Table 2. Multilevel Regression Analysis of Door-to-Reperfusion Time of Patients Directly Presented at an EVT Hospital

Variable	Model 1 Empty model	Model 2 Case-mix	Model 3A Door-CT time	Model 3B CT-CTA time	Model 3C CTA-groin time	Model 3D Groin-reperfusion time
Proportional change in variance*						
Between hospitals	Reference	0.02	-0.14	-0.32	0.85	1.00
Within hospitals	Reference	0.07	0.12	0.25	0.58	1.00
ICC	0.09	0.09				

Model 1: hospital. Model 2: hospital and case-mix. Model 3A: hospital, case-mix, and door-to-CT time. Model 3B: hospital, case-mix, door-to-CT time, and CT-CTA time. Model 3C: hospital, case-mix, door-to-CT time, CT-CTA time, and CTA-to-groin time. Model 3D: hospital, case-mix, door-to-CT time, CT-CTA time, CTA-to-groin time, and groin-to-reperfusion time. The ICC describes the proportion of the total variance that is attributable to clustering within hospitals, in our case the between-hospital variance in door-to-reperfusion time. The remaining total variance is attributable to within-hospital variation between patients. The proportional change in variance describes the change of the between-hospital and within-hospital variation in door-to-reperfusion time in each model compared with model 1. The individual attribution of each added variable on the variation in door-to-reperfusion time can be calculated by subtracting the proportional changes in variance of each model. These numbers are shown in Figure 2. CT indicates computed tomography; CTA, CT angiography; EVT, endovascular treatment; and ICC, intraclass correlation coefficient.

*A negative sign indicates that the time interval increased the variance.

time of transferred patients was explained by case-mix (3%), door-to-groin time (40%), and groin-to-reperfusion time (57%).

DISCUSSION

In this study, we quantified between-hospital and within-hospital variation in door-to-reperfusion time in patients with acute ischemic stroke treated with EVT in Dutch EVT hospitals. We found that door-to-reperfusion time varies between hospitals, but even more within hospitals. This variation in door-to-reperfusion time is mostly explained by CTA-to-groin time and groin-to-reperfusion time, and less by patient characteristics.

Our results imply that workflow improvement strategies should primarily target the reduction of variation within hospitals. This is in line with a study that showed that most variability in quality of care occurred at the patient level (82%) within hospitals.⁸ The ability to identify the cause of variation is fundamental for health care improvement.⁹ Variation is often seen as a source of errors or issues with the system.¹⁰ Investigation of causes and attributable factors on this variation may help in reducing variation and improving procedures. The within-hospital variation in door-to-reperfusion time is much larger than the between-hospital variation in door-to-reperfusion time. We do not have one clear explanation for the large within-hospital variation in door-to-reperfusion time. We tried to explain the within-hospital variation in door-to-reperfusion time by adding potentially delaying factors (eg, systolic blood pressure ≥ 185 mm Hg) requiring blood pressure-lowering therapy, intravenous alteplase treatment, and general anesthesia) to the model. However, these variables did not explain the variation. There are many other factors that could influence the time intervals and could be different

between patients. For example, there are structure measures, such as the availability of an interventionist or angiographic suite and hospital case volume. But there are also process measures, such as the persistence of the interventionist to achieve reperfusion and difficulties in procedures as tortuosity of the vascular tree. Most of these factors are difficult to measure. Almost no case-mix variable was associated with door-to-reperfusion time. However, admission during off hours showed a significant association with the door-to-reperfusion time of direct patients. Previous research showed that admission during off hours was associated with a slight delay in start of endovascular treatment in patients with acute ischemic stroke.¹¹ This may mean that there is potential for improvements of workflow times during off hours.

The between-hospital variation is low. This could be attributable to the standardization of processes in the treatment with EVT of patients with an ischemic stroke. Moreover, Dutch EVT hospitals need to meet various high-quality standards, which could reduce the between-hospital variation. The total variation in door-to-reperfusion time is therefore mainly explained by within-hospital variation.

CTA-to-groin time and groin-to-reperfusion time were important drivers of between-hospital and within-hospital variation in door-to-reperfusion time. Of all time intervals, CTA-to-groin time is the largest. This probably explains why CTA-to-groin time contributed most to the variation in door-to-reperfusion time for patients directly presented at an EVT hospital. Various improvement strategies focusing on reduction of between-hospital variation in door-to-reperfusion times have been suggested. A meta-analysis about different strategies for workflow improvement showed that the most promising workflow intervention types concerned anesthetic management, in-hospital patient transfer management, prehospital management, teamwork,




Figure 2. The contribution of each added variable to the variation in door-to-reperfusion time. A, Direct patients. B, Transferred patients. CT indicates computed tomography; and CTA, CT angiography.

and feedback on performance.¹² Performance feedback was the most effective intervention in this meta-analysis. To better estimate the (magnitude of the) effect of performance feedback, a stepped wedge cluster randomized controlled trial can be used. With this study design, hospitals instead of individual patients are randomized subsequently (in “steps”), which makes it possible to take time trends into account. Such a study has recently been started in the Netherlands: Performance feedback on quality of care in hospitals performing thrombectomy for ischemic stroke.¹³ In this study, the effectiveness of monitoring and performance feedback is being investigated. This feedback consists of process and outcome measures, shown in 3 monthly feedback reports. Local quality improvement teams of every hospital will make improvement plans and can evaluate their actions every 3 months. This study aims to reduce the between-hospital variation in process measures and hopefully will give insight in

delaying factors on time to treatment. However, quality improvement should not only focus on the comparison of hospitals and between-hospital variation, but also on within-hospital variation.

Our study has some limitations. We did not have information on whether CT-perfusion was performed, and we know that this varied between hospitals at the moment of our data registration. We expect the influence on variation in door-to-reperfusion time to be small, because we observed that the time interval of CT-to-CTA (in which CT-perfusion is performed) did not explain the between-hospital variation in door-to-reperfusion time.

We included all relevant case-mix variables available in our data, but we could have missed variables associated with door-to-reperfusion time. However, based on our analyses, we expect that case-mix variables are less important in the explanation of variation in door-to-reperfusion time.




Figure 3. Density plots per hospital of door-to-reperfusion time of patients directly presented at an endovascular treatment hospital (crude data).

CONCLUSIONS

Door-to-reperfusion time varies between, but even more within, hospitals providing endovascular treatment for ischemic stroke. Quality of stroke care

improvements should not only be guided by between-hospital comparisons, but also aim to reduce variation between patients within a hospital, and should specifically focus on CTA-to-groin time and groin-to-reperfusion time.

Table 3. Multilevel Regression Analysis of Door-to-Reperfusion Time of Transferred Patients

Variable	Model 1 Empty model	Model 2 Case-mix	Model 3A Door-groin time	Model 3B Groin-reperfusion time
Proportional change in variance*				
Between hospitals	Reference	-0.07	0.56	1.00
Within hospitals	Reference	0.03	0.43	1.00
ICC	0.03	0.04		

Model 1: hospital. Model 2: hospital and case-mix. Model 3A: hospital, case-mix, and door-to-groin time. Model 3B: hospital, case-mix, door-to-groin time, and groin-to-reperfusion time. The ICC describes the proportion of the total variance that is attributable to clustering within hospitals, in our case the between-hospital variance in door-to-reperfusion time. The remaining total variance is attributable to within-hospital variation between patients. The proportional change in variance describes the change of the between-hospital and within-hospital variation in door-to-reperfusion time in each model compared with model 1. The individual attribution of each added variable on the variation in door-to-reperfusion time can be calculated by subtracting the proportional changes in variance of each model. These numbers are shown in Figure 2. ICC indicates intraclass correlation coefficient.

*A negative sign indicates that the time interval increased the variance.

APPENDIX

MR CLEAN Registry investigators

Diederik W. J. Dippel (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Aad van der Lugt (Department of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center, Rotterdam), Charles B. L. M. Majoie (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Yvo B. W. E. M. Roos (Department of Neurology, Amsterdam UMC, University of Amsterdam, Amsterdam), Robert J. van Oostenbrugge (Department of Neurology, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Wim H. van Zwam (Department of Radiology and Nuclear Medicine, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Jelis Boiten (Department of Neurology, Haaglanden Medical Center, The Hague), Jan Albert Vos (Department of Radiology, Sint Antonius Hospital, Nieuwegein), Ivo G. H. Jansen (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Maxim J. H. L. Mulder (Department of Neurology, Radiology and Nuclear Medicine, Erasmus MC, University Medical Center, Rotterdam), Robert-Jan B. Goldhoorn (Department of Neurology, Radiology and Nuclear Medicine, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Kars C. J. Compagne (Department of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center, Rotterdam), Manon Kappelhof (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Josje Brouwer (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Sanne J. den Hartog (Department of Neurology, Radiology and Nuclear Medicine, Public Health, Erasmus MC, University Medical Center, Rotterdam), Wouter H. Hinsenveld (Department of Neurology, Radiology and Nuclear Medicine, Maastricht University Medical Center and

Cardiovascular Research Institute Maastricht), Bob Roozenbeek (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Adriaan C. G. M. van Es (Department of Radiology, Leiden University Medical Center, Leiden), Bart J. Emmer (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Jonathan M. Coutinho (Department of Neurology, Amsterdam UMC, University of Amsterdam, Amsterdam), Wouter J. Schonewille (Department of Neurology, Sint Antonius Hospital, Nieuwegein), Marieke J. H. Wermer (Department of Neurology, Leiden University Medical Center, Leiden), Marianne A. A. van Walderveen (Department of Radiology, Leiden University Medical Center, Leiden), Julie Staals (Department of Neurology, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Jeannette Hofmeijer (Department of Neurology, Rijnstate Hospital, Arnhem), Jasper M. Martens (Department of Radiology, Rijnstate Hospital, Arnhem), Geert J. Lycklama à Nijeholt (Department of Radiology, Haaglanden MC, The Hague), Sebastiaan F. de Bruijn (Department of Neurology, HAGA Hospital, The Hague), Lukas C. van Dijk (Department of Neurology, HAGA Hospital, The Hague), H. Bart van der Worp (Department of Neurology, University Medical Center Utrecht), Rob H. Lo (Department of Radiology, University Medical Center Utrecht), Ewoud J. van Dijk (Department of Neurology, Radboud University Medical Center, Nijmegen), Hieronymus D. Boogaarts (Department of Neurosurgery, Radboud University Medical Center, Nijmegen), J. de Vries (Department of Radiology, Radboud University Medical Center, Nijmegen), Paul L. M. de Kort (Department of Neurology, Elisabeth-TweeSteden ziekenhuis, Tilburg), Julia van Tuijl (Department of Neurology, Elisabeth-TweeSteden ziekenhuis, Tilburg), Jo P. Peluso (Department of Radiology, Elisabeth-TweeSteden ziekenhuis, Tilburg), Puck Fransen (Department of Neurology, Isala Klinieken, Zwolle), Jan S. P. van den Berg (Department of Neurology, Isala Klinieken, Zwolle), Boudewijn A. A.

M. van Hasselt (Department of Radiology, Isala Klinieken, Zwolle), Leo A. M. Aerden (Department of Neurology, Reinier de Graaf Gasthuis, Delft), René J. Dallinga (Department of Radiology, Reinier de Graaf Gasthuis, Delft), Maarten Uyttenboogaart (Department of Neurology, University Medical Center Groningen), Omid Eschgi (Department of Radiology, University Medical Center Groningen), Reinoud P. H. Bokkers (Department of Radiology, University Medical Center Groningen), Tobien H. C. M. L. Schreuder (Department of Neurology, Atrium Medical Center, Heerlen), Roel J. J. Heijboer (Department of Radiology, Atrium Medical Center, Heerlen), Koos Keizer (Department of Neurology, Catharina Hospital, Eindhoven), Lonneke S. F. Yo (Department of Radiology, Catharina Hospital, Eindhoven), Heleen M. den Hertog (Department of Neurology, Isala Klinieken, Zwolle), Emiel J. C. Sturm (Department of Radiology, Medisch Spectrum Twente, Enschede), Paul J. A. M. Brouwers (Department of Neurology, Medisch Spectrum Twente, Enschede), Marieke E. S. Sprengers (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Sjoerd F. M. Jenniskens (Department of Radiology, Radboud University Medical Center, Nijmegen), René van den Berg (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Albert J. Yoo (Department of Radiology, Texas Stroke Institute, Texas), Ludo F. M. Beenen (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Alida A. Postma (Department of Radiology and Nuclear Medicine, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Stefan D. Roosendaal (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Bas F. W. van der Kallen (Department of Radiology, Haaglanden MC, The Hague), Ido R. van den Wijngaard (Department of Radiology, Haaglanden MC, The Hague), Joost Bot (Department of Radiology, Amsterdam UMC, Vrije Universiteit van Amsterdam, Amsterdam), Pieter-Jan van Doormaal (Department of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center, Rotterdam), Anton Meijer (Department of Radiology, Radboud University Medical Center, Nijmegen), Elyas Ghariq (Department of Radiology, Haaglanden MC, The Hague), Marc P. van Proosdij (Department of Radiology, Noordwest Ziekenhuisgroep, Alkmaar), G. Menno Krietemeijer (Department of Radiology, Catharina Hospital, Eindhoven), Dick Gerrits (Department of Radiology, Medisch Spectrum Twente, Enschede), Wouter Dinkelaar (Department of Radiology and Nuclear Medicine, Erasmus MC, University Medical Center, Rotterdam), Auke P. A. Appelman (Department of Radiology, University Medical Center Groningen), Bas Hammer (Department of Radiology, HAGA

Hospital, The Hague), Sjoert Pegge (Department of Radiology, Radboud University Medical Center, Nijmegen), Anouk van der Hoorn (Department of Radiology, University Medical Center Groningen), Saman Vinke (Department of Neurosurgery, Radboud University Medical Center, Nijmegen), HZwenneke Flach (Department of Radiology, Isala Klinieken, Zwolle), Hester F. Lingsma (Department of Public Health, Erasmus MC, University Medical Center, Rotterdam), Naziha el Ghannouti (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Martin Sterrenberg (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Wilma Pellikaan (Department of Neurology, Sint Antonius Hospital, Nieuwegein), Rita Sprengers (Department of Neurology, Amsterdam UMC, University of Amsterdam, Amsterdam), Marjan Elfrink (Department of Neurology, Rijnstate Hospital, Arnhem), Michelle Simons (Department of Neurology, Rijnstate Hospital, Arnhem), Marjolein Vossers (Department of Radiology, Rijnstate Hospital, Arnhem), Joke de Meris (Department of Neurology, Haaglanden Medical Center, The Hague), Tamara Vermeulen (Department of Neurology, Haaglanden Medical Center, The Hague), Annet Geerlings (Department of Neurology, Radboud University Medical Center, Nijmegen), Gina van Vemde (Department of Neurology, Isala Klinieken, Zwolle), Tiny Simons (Department of Neurology, Atrium Medical Center, Heerlen), Gert Messchendorp (Department of Neurology, University Medical Center Groningen), Nynke Nicolaj (Department of Neurology, University Medical Center Groningen), Hester Bongenaar (Department of Neurology, Catharina Hospital, Eindhoven), Karin Bodde (Department of Neurology, Reinier de Graaf Gasthuis, Delft), Sandra Kleijn (Department of Neurology, Medisch Spectrum Twente, Enschede), Jasmijn Lodico (Department of Neurology, Medisch Spectrum Twente, Enschede), Hanneke Droste (Department of Neurology, Medisch Spectrum Twente, Enschede), Maureen Wollaert (Department of Neurology, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Sabrina Verheesen (Department of Neurology, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), D. Jeurissen (Department of Neurology, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Erna Bos (Department of Neurology, Leiden University Medical Center), Yvonne Drabbe (Department of Neurology, HAGA Hospital, The Hague), Michelle Sandiman (Department of Neurology, HAGA Hospital, The Hague), Nicoline Aldering (Department of Neurology, Rijnstate Hospital, Arnhem), Berber Zweedijk (Department of Neurology, University Medical Center Utrecht), Jocova Vervoort (Department of Neurology, Elisabeth-TweeSteden ziekenhuis, Tilburg),

Eva Ponjee (Department of Neurology, Isala Klinieken, Zwolle), Sharon Romviel (Department of Neurology, Radboud University Medical Center, Nijmegen), Karin Kanselaar (Department of Neurology, Radboud University Medical Center, Nijmegen), Denn Barning (Department of Radiology, Leiden University Medical Center), Esmee Venema (Department of Public Health, Erasmus MC, University Medical Center, Rotterdam), Vicky Chalos (Department of Neurology, Public Health, Erasmus MC, University Medical Center, Rotterdam), Ralph R. Geuskens (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Tim van Straaten (Department of Neurology, Radboud University Medical Center, Nijmegen), Saliha Ergezen (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Roger R. M. Harmsma (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Daan Muijres (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Anouk de Jong (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam), Olvert A. Berkhemer (Department of Neurology, Erasmus MC, University Medical Center, Rotterdam, Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam, Department of Radiology, Maastricht University Medical Center and Cardiovascular Research Institute Maastricht), Anna M. M. Boers (Department of Radiology and Nuclear Medicine, Biomedical Engineering and Physics, Amsterdam UMC, University of Amsterdam, Amsterdam), J. Huguët (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), P. F. C. Groot (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Marieke A. Mens (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Katinka R. van Kranendonk (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Kilian M. Treurniet (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Manon L. Tolhuisen (Department of Radiology and Nuclear Medicine, Biomedical Engineering and Physics, Amsterdam UMC, University of Amsterdam, Amsterdam), Heitor Alves (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Annick J. Weterings (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Eleonora L. F. Kirkels (Department of Radiology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Eva J. H. F. Voogd (Department of Neurology, Rijnstate Hospital, Arnhem), Lieve M. Schupp (Department of Radiology

and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Sabine L. Collette (Department of Neurology, Radiology, University Medical Center Groningen), Adrien E. D. Groot (Department of Neurology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Natalie E. LeCouffe (Department of Neurology and Nuclear Medicine, Amsterdam UMC, University of Amsterdam, Amsterdam), Praneeta R. Konduri (Department of Biomedical Engineering and Physics, Amsterdam UMC, University of Amsterdam, Amsterdam), Haryadi Prasetya (Department of Biomedical Engineering and Physics, Amsterdam UMC, University of Amsterdam, Amsterdam), Nerea Arrarte-Terreros (Department of Biomedical Engineering and Physics, Amsterdam UMC, University of Amsterdam, Amsterdam), Lucas A. Ramos (Department of Biomedical Engineering and Physics, Amsterdam UMC, University of Amsterdam, Amsterdam). All locations are in the Netherlands, unless otherwise indicated.

ARTICLE INFORMATION

Received May 31, 2021; accepted November 19, 2021.

Affiliations

Department of Neurology (S.J.d.H., D.W.J.D., B.R.); Department of Radiology and Nuclear Medicine (S.J.d.H., P.-J.v.D., A.v.d.L., B.R.); and Department of Public Health (S.J.d.H., H.F.L.), Erasmus MC, University Medical Center, Rotterdam, the Netherlands ; Department of Neurology, Rijnstate, Arnhem, the Netherlands (J.H.), Department of Radiology and Nuclear Medicine, Catharina Hospital, Eindhoven, the Netherlands (L.S.F.Y.), and Department of Radiology and Nuclear Medicine, Amsterdam University Medical Centers, Location AMC, Amsterdam, the Netherlands (C.B.L.M.M.).

Sources of Funding

The MR CLEAN (Multicenter Randomized Clinical Trial of Endovascular Treatment of Acute Ischemic Stroke) Registry was partly funded by Stichting Toegepast Wetenschappelijk Instituut voor Neuromodulatie, Erasmus MC, University Medical Center, Maastricht University Medical Center, and Amsterdam University Medical Center.

Disclosures

Drs Dippel and van der Lugt report funding from the Dutch Heart Foundation, Brain Foundation Netherlands, The Netherlands Organisation for Health Research and Development, Health Holland Top Sector Life Sciences and Health, and unrestricted grants from Penumbra Inc, Stryker European Operations BV, Medtronic, Thrombolytic Science, LLC, and Cerenovus for research, all paid to institution. Dr Majoie reports grants from CVON/ Dutch Heart Foundation, European Commission, TWIN Foundation, Dutch Health Evaluation Program, and Stryker, paid to institution; and is minority shareholder of Nico-lab. The remaining authors have no disclosures to report.

Supplemental Material

Tables S1–S2

Figure S1

REFERENCES

1. Mulder M, Jansen IGH, Goldhoorn RB, Venema E, Chalos V, Compagne KCJ, Roozenbeek B, Lingsma HF, Schonewille WJ, van den Wijngaard IR, et al. Time to endovascular treatment and outcome in acute ischemic

- stroke: MR CLEAN registry results. *Circulation*. 2018;138:232–240. doi: 10.1161/CIRCULATIONAHA.117.032600
2. Jahan R, Saver JL, Schwamm LH, Fonarow GC, Liang LI, Matsouaka RA, Xian Y, Holmes DN, Peterson ED, Yavagal D, et al. Association between time to treatment with endovascular reperfusion therapy and outcomes in patients with acute ischemic stroke treated in clinical practice. *JAMA*. 2019;322:252–263. doi: 10.1001/jama.2019.8286
 3. Jansen IGH, Mulder M, Goldhoorn RB, MR CLEAN Registry investigators. Endovascular treatment for acute ischaemic stroke in routine clinical practice: prospective, observational cohort study (MR CLEAN registry). *BMJ*. 2018;360:k949. doi: 10.1136/bmj.k949
 4. Abel G, Elliott MN. Identifying and quantifying variation between health-care organisations and geographical regions: using mixed-effects models. *BMJ Qual Saf*. 2019;28:1032–1038. doi: 10.1136/bmjqs-2018-009165
 5. Lingsma HF, Steyerberg EW, Eijkemans MJ, Dippel DW, Scholte Op Reimer WJ, Van Houwelingen HC, Netherlands Stroke Survey Investigators. Comparing and ranking hospitals based on outcome: results from the Netherlands stroke survey. *QJM*. 2010;103:99–108. doi: 10.1093/qjmed/hcp169
 6. Merlo J, Chaix B, Ohlsson H, Beckman A, Johnell K, Hjerpe P, Råstam L, Larsen K. A brief conceptual tutorial of multilevel analysis in social epidemiology: using measures of clustering in multilevel logistic regression to investigate contextual phenomena. *J Epidemiol Community Health*. 2006;60:290–297. doi: 10.1136/jech.2004.029454
 7. Merlo J, Yang M, Chaix B, Lynch J, Råstam L. A brief conceptual tutorial on multilevel analysis in social epidemiology: investigating contextual phenomena in different groups of people. *J Epidemiol Community Health*. 2005;59:729–736. doi: 10.1136/jech.2004.023929
 8. Reeves MJ, Gargano J, Maier KS, Broderick JP, Frankel M, LaBresh KA, Moomaw CJ, Schwamm L. Patient-level and hospital-level determinants of the quality of acute stroke care: a multilevel modeling approach. *Stroke*. 2010;41:2924–2931. doi: 10.1161/STROKEAHA.110.598664
 9. Ferguson J. Reducing unwanted variation in healthcare clears the way for outcomes improvement. Health Catal Prod Serv Outcomes Improve. 2017. Available at: <https://www.healthcatalyst.com/insights/reducing-variation-in-healthcare-to-boost-improvement/>. Accessed May 3, 2021.
 10. James C. Manufacturing's prescription for improving healthcare quality. *Hosp Top*. 2005;83:2–8. doi: 10.3200/HTPS.83.1.2-8
 11. Hinsenvelde WH, de Ridder IR, van Oostenbrugge RJ, Vos JA, Groot AE, Coutinho JM, Lycklama à Nijeholt GJ, Boiten J, Schonewille WJ, Dippel DWJ, et al. Workflow intervals of endovascular acute stroke therapy during on- versus off-hours: the MR CLEAN registry. *Stroke*. 2019;50:2842–2850. doi: 10.1161/STROKEAHA.119.025381
 12. Janssen PM, Venema E, Dippel DWJ. Effect of workflow improvements in endovascular stroke treatment. *Stroke*. 2019;50:665–674. doi: 10.1161/STROKEAHA.118.021633
 13. den Hartog SJ, Amini M, van Leeuwen N, Kuhrij LS, Nederkoorn PJ, Lingsma HF, van der Lugt A, Dippel DWJ & Roozenbeek B Performance feedback on quality of care in hospitals performing thrombectomy for ischemic stroke (perfeqtos trial). Poster presented at: ESOC May 22, 2019 Milan, Italy. URL [https://journals.sagepub.com/doi/full/10.1177/2396987319848124#_i528]

SUPPLEMENTAL MATERIAL

Table S1. Extreme minimum and maximum outliers of each time interval, for patients directly presented at an EVT hospital and for transferred patients.

Time interval	Extreme minimum outlier	Extreme maximum outlier	% missing values including outliers
Door-CT time	0 minutes	> 200 minutes	3%
CT-CTA time	none*	> 200 minutes	3%
CTA-groin time	≤ 5 minutes	> 300 minutes	3%
Groin-reperfusion time			
Direct	none*	> 300 minutes	10%
Transferred	none*	> 300 minutes	9%
Door-groin time	none*	none*	5%
Door-reperfusion time			
Direct	none*	> 500 minutes	11%
Transferred	none*	> 500 minutes	13%

* there were no outliers


Table S2. Multilevel model, association between case-mix variables and door-to-reperfusion time.

	Direct patients	Transferred patients
Case-mix variables	adjusted beta, 95% CI	adjusted beta, 95% CI
Age	0.03 (-0.20-0.26)	-0.04 (-0.21-0.12)
Men	-2.23 (-8.48-4.03)	2.81 (-1.33-6.96)
Atrial fibrillation	-1.58 (-8.93-5.76)	-2.21 (-7.03-2.62)
Hypertension	4.30 (-2.37-10.97)	1.84 (-3.02-6.71)
Myocardial infarction	-4.50 (-13.33-4.33)	2.10 (-4.02-8.21)
Previous stroke	0.10 (-8.13-8.34)	1.14 (-4.55-6.83)
Peripheral artery disease	9.37 (-1.32-20.06)	-3.34 (-11.16-4.47)
Diabetes Mellitus	5.16 (-3.24-13.56)	2.78 (-3.17-8.74)
Hyperlipidemia	-6.75 (-13.73-0.24)	0.12 (-6.57-6.80)
NIHSS at baseline, per point	-0.09 (-0.59-0.42)	0.37 (-0.01-0.75)
Pre-stroke modified Rankin Scale		
0	reference	reference
1	4.46 (-4.76-13.67)	-2.16 (-8.52-4.20)
2	6.24 (-5.35-17.84)	0.25 (-8.18-8.69)
≥ 3	5.70 (-4.08-15.48)	-0.99 (-9.26-7.27)
Location of occluded artery		
M1	reference	reference
M2	8.02 (0.21-15.82)	-0.72 (-7.52-6.09)
Intracranial ICA	20.83 (6.26-35.40)	8.42 (-1.85-18.69)
ICA-T	13.31 (5.66-20.97)	7.47 (1.81-13.13)
Other (Anterior/M3)	28.98 (-2.08-60.05)	18.52 (-9.44-46.49)
Onset to door time, minutes	-0.05 (-0.09--0.01)	-0.06 (-0.09--0.03)
Admission during off hours*	22.10 (16.11-28.09)	1.43 (-3.46-6.32)

* Admission between 5:00 PM and 8:00 AM, on weekends, or a national holiday
 CI, confidence interval, NIHSS, National Institutes of Health Stroke Scale, M1/M2/M3, 1st, 2nd and 3rd segment of the middle cerebral artery, ICA, internal carotid artery, ICA-T, internal carotid artery terminus.

Figure S1. A) Correlation between CTA-to-groin time and door-to-reperfusion time. B) Correlation between groin-to-reperfusion time and door-to-reperfusion time.

A



B

