

Research Article

Enhanced Production of Polymyxin E in *Paenibacillus polymyxa* by Replacement of Glucose by Starch

Zhiliang Yu , Zhongqi Sun, Jianhua Yin , and Juanping Qiu 

College of Biotechnology and Bioengineering, Zhejiang University of Technology, Hangzhou 310014, China

Correspondence should be addressed to Zhiliang Yu; zlyu@zjut.edu.cn and Juanping Qiu; qiujiang@zjut.edu.cn

Received 2 May 2018; Revised 12 September 2018; Accepted 30 September 2018; Published 14 October 2018

Guest Editor: Artur Alves

Copyright © 2018 Zhiliang Yu et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Polymyxin E or colistin, produced by *Paenibacillus polymyxa*, is an important antibiotic against Gram-negative pathogens. The objective of this study is to evaluate the effect of starch in fermentation medium on colistin biosynthesis in *P. polymyxa*. The results indicated that replacement of glucose by starch stimulated colistin production and biosynthesis rate. Overall, the stimulation extent was starch concentration-dependent. As expected, addition of starch induced the expression of *amyE* encoding amylase and increased amylase activity in fermentation solution. Additionally, replacement of glucose by starch resulted in residue reducing sugar and pH of fermentation mixture low relative to glucose as the sole sugar source. At the molecular level, it was found that replacement of glucose by starch has enhanced the relative expression level of *ccpA* encoding catabolite control protein A. Therefore, the repression of starch utilization by glucose could be probably relieved. In addition, use of starch stimulated the expression of regulatory gene *spo0A* but repressed the expression of another regulatory gene *abrB*. As a result, the expression of genes directly involved in colistin biosynthesis and secretion increased, indicating that at the transcriptional level *spo0A* and *abrB* played opposite roles in regulating colistin biosynthesis in *P. polymyxa*. Taken together, our data demonstrated that starch instead of glucose can promote colistin production probably by affecting the expression of colistin biosynthesis-related genes, as well as reducing the repression of glucose to a secondary metabolic product.

1. Introduction

Polymyxin E, also called colistin, is an important old antibiotic known for around six decades for treatment of infection caused by Gram-negative pathogens [1, 2]. Later studies showed that colistin can also kill Gram-positive bacteria [3, 4]. Currently, its clinical use is broadly restricted regarding its toxicity mainly to the kidney and nervous system [5]. In recent, the occurrence of Gram-negative multidrug-resistant pathogens which are resistant to many available antibiotics has revived its clinical application in healthcare centers, since colistin is broadly considered as one of the last-line options of antibiotic therapy for multidrug-resistant bacteria. Therefore, its market demand is increasing [6, 7].

Colistin is composed of two parts: a cyclic heptapeptide and a tripeptide side chain which is acylated by a fatty acid at the amino terminus [8, 9]. In total, colistin has ten orderly assembled amino acid residues [10]. Among them, six are L-2,4-diaminobutyric acids (L-Dabs). L-Dab is biosynthesized by 2,4-diaminobutyrate aminotransferase (EctB) which is

encoded by *ectB* [11]. Colistin can be biosynthesized by a multienzyme nonribosomal peptide synthetase system (NRPS) in *Paenibacillus polymyxa* [3, 4]. The phosphopantetheinyl transferase (Sfp) encoded by *sfp* is important for colistin biosynthesis due to its activation function on NRPS [12, 13]. A gene cluster including five open reading frames, *pmxA*, *pmxB*, *pmxE*, *pmxC*, and *pmxD*, encoding three synthetases PmxA, PmxB, and PmxE, and two membrane transporters PmxC and PmxD, respectively, has been characterized for biosynthesis and secretion of colistin in *P. polymyxa* [10, 14]. It has been determined that colistin biosynthesis is negatively regulated by *AbrB*, a DNA-binding protein, by directly binding to the upstream region of *pmxA* [11]. The expression of *abrB* itself is negatively controlled by *Spo0A*, another DNA-binding protein, encoded by *spo0A* [15]. These two genes play opposite roles in regulation of colistin production.

So far, colistin has been best characterized with respect to its structure and biosynthesis, antibacterial mechanism and bacterial resistance, and toxicity and derivatives. In contrast, extremely little is known about medium optimization for its

fermentation output. It has been reported that colistin production depends on the inorganic phosphate concentration [16]. Further optimization showed that addition of L-Dab as well as its precursor aspartic acid to fermentation medium containing appropriate PO_4^{3-} stimulates colistin production [16]. However, other studies showed that addition of either L-Dab or aspartic acid to medium after 35 h fermentation significantly inhibits colistin production by suppressing the expression of *pmxA* and *pmxE*, as well as *ectB* in another producer strain [17]. Although it has been found that corn meal in the medium is essential for the better production of colistin [16], glucose is the most widely used carbon source for colistin biosynthesis [18]. In our previous study, glucose was also used as a sugar source for colistin production in *P. polymyxa* C12 and its production reached around 6.2×10^4 U/mL (2600 $\mu\text{g}/\text{mL}$) in flask level [17]. As a concern, glucose has been widely found to repress the accumulation of secondary metabolic compounds in microbes through carbon catabolite repression (CCR) [19]. CCR in microbes is regarded as the mechanism in which bacteria preferentially utilize the rapidly metabolizable carbon source (normally glucose). As a result, the utilization of secondary carbon resource is repressed. CCR is considered to be a part of the global control system and therefore it affects many genes [20]. In Gram-positive bacteria, the catabolite control protein A (CcpA) is the master regulator of CCR. Various physiological processes in Gram-positive bacteria are regulated by CcpA [21–23].

In this study, we attempt to substitute starch for glucose and investigate the effect of the sugar source on colistin production in *P. polymyxa* C12. In addition, the effect of the sugar source on amylase activity and relative expression of genes associated with colistin biosynthesis was also evaluated.

2. Materials and Methods

2.1. Strain and Culture Conditions. Colistin-producer *P. polymyxa* C12 [17] used in this study was frozen at -80°C in our lab at Zhejiang University of Technology, China. Unless otherwise stated, *P. polymyxa* was firstly cultivated on a culture medium agar plate (10 g/L of beef extract, 15 g/L of peptone, 10 g/L of glucose, 2 g/L of yeast extract, 3 g/L of NaCl, 0.1 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, and 20 g/L of agar, pH 7.0) at 30°C for 2 d. Then, a ring of *P. polymyxa* was transferred to 50 mL of seed medium (30 g/L of soybean meal, 5 g/L of soybean oil, 0.1 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 15 g/L $(\text{NH}_4)_2\text{SO}_4$, 0.77 g/L of KH_2PO_4 , 0.7 g/L of CaCO_3 , and 10 g/L of glucose, pH 7.0) in a 250 mL flask for incubation at 30°C for 24 h with a shaking at 200 rpm. Next, 5 mL of cell culture was transferred to 50 mL of fermentation medium (23.9 g/L of soybean meal powder (Zhejiang Qianjiang Biochemical Co., Ltd., China), 21.1 g/L of soybean cake powder (Zhejiang Qianjiang Biochemical Co., Ltd., China), 10 g/L of soybean oil, 0.1 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 25 g/L $(\text{NH}_4)_2\text{SO}_4$, 0.77 g/L of KH_2PO_4 , 1 g/L of CaCO_3 , and 45 g/L of glucose, pH 7.0) in a 250 mL flask at 30°C with shaking at 200 rpm for fermentation. A certain amount of glucose from 20 g/L to 45 g/L in the fermentation medium was replaced by starch if necessary. Unless otherwise

specified, *P. polymyxa* was fermented for 96 h and 0.5 mL of samples was collected per 12 h. The colistin concentration and relevant gene expression were determined by HPLC and quantitative real-time PCR (qRT-PCR), respectively.

2.2. Measurement of Cell Growth. Unless otherwise stated, the bacterial biomass of the cultured cells was determined based on the value of colony forming unit (CFU) [24, 25]. First, the cells were collected after centrifugation at 5,000 g for 5 min. After washing twice with 0.5 mL of fresh broth culture medium (10 g/L of beef extract, 15 g/L of peptone, 10 g/L of glucose, 2 g/L of yeast extract, 3 g/L of NaCl, and 0.1 g/L of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, pH 7.0), the cells were then resuspended in 0.5 mL of fresh broth culture medium. Then, the cells were tenfold gradiently diluted. Finally, 100 μL of cells was spread to a culture medium agar plate for growth. After cultivation at 30°C for 2 d, CFU were counted.

2.3. HPLC Analysis of Colistin. One milliliter of fermentation liquor and 9 mL of ultrapure water were mixed. Then, 1 mL of diluted fermentation liquor was centrifuged at 10,000 g for 10 min and the supernatant was collected and filtered with 0.45 μm hydrophilic microporous membrane (Millipore). Analysis of colistin was performed using binary gradient model of an HPLC system (SHIMADZU, Japan). Twenty μL of supernatant sample was injected into a reverse-phase column, YMC Pack ODS-A (150 \times 4.6 mm I.D., 5 μm), eluted at 33°C , and analyzed in a mixed solvent of acetonitrile (22%) and water containing 0.223% Na_2SO_4 (78%), at a constant flow of 1 mL/min. The mixed solvent was prepared by mixing pure acetonitrile in A-pump with water containing 0.223% Na_2SO_4 in B-pump. Separation program was set as follows: 0~20 min, 22% A-pump and 78% B-pump; 21~30 min, 90% A-pump and 10% B-pump; 31~40 min, 22% A-pump and 78% B-pump. Colistin peak was determined at wavelength of 240 nm. Colistin concentration produced was calculated based on the extracted correlation between the concentration of standard colistin (Zhejiang Qianjiang Biochemical Co., Ltd., China) and the corresponding peak area in HPLC. One unit is equal to 0.0418 μg of colistin [17]. Colistin was dissolved in 1 mL of pure water to make 2×10^5 U/mL solution. Then, colistin was fivefold gradiently diluted with water to make serial colistin solutions. For HPLC analysis, 20 μL of standard colistin was injected.

2.4. Detection of Reducing Sugar and Assay of α -Amylase Activity. After fermentation, the cell mixture was centrifuged at 10,000 g for 10 min and the supernatant was collected. Unless otherwise specified, the amylase activity in the fermentation supernatant (crude enzyme solution) was determined by measuring the reducing sugar generated during the reaction [26]. In general, 0.2 mL of 0.1 M citrate-phosphate buffer with pH 7.0 containing 1% (w/v) soluble starch was preheated at 30°C for 5 min. Then, 0.05 mL of crude enzyme solution appropriately diluted with sterilized ultrapure water was added and mixed thoroughly. After incubation at 30°C for 30 min, the reaction was terminated by adding 1 mL of DNS reagent containing 182 g/L Rochelle salt,

TABLE 1: Sequences of primer pairs for PCR amplification of genes.

Genes	Nucleotide sequences (5'-3')		Product sizes (bp)
	Forward primers	Reverse primers	
<i>amyE</i>	ATGGTCCACAATCCTGTT	CCTCATGTTCTTCCCTCA	1319
<i>ccpA</i>	TGGTCAGCAAACGCATCG	AAACCTCAGACCCGCAAG	832

TABLE 2: Sequences of primer pairs for real-time PCR analysis of gene expression.

Genes	Nucleotide sequences (5'-3')		Product sizes (bp)
	Forward primers	Reverse primers	
<i>pmxA</i>	TCAACTCGCTCAGAAGCGTT	TTGTACGGAAACCGACGGAG	105
<i>pmxB</i>	ATGAAATCTTTGTTTGAAAA	CCAGGACGTACACCCTCAAC	111
<i>pmxC</i>	TATTCCCAGCTCATCACGC	TCGGAAGCGAACGACCATTT	107
<i>pmxD</i>	TGTTTCGTTCAACGCCTCGTA	GCTTGCAAACGCTCGGTAAA	118
<i>pmxE</i>	CACTTTGCCTGAAACGACCG	GCCAGAATGCGTTCATACCG	111
<i>spo0A</i>	TCGCAGAATCCCGCAACATA	CGGTTGTGGAGTCAGGTTCA	103
<i>abrB</i>	AAATACGGAACAGCCCGTCC	TCGCTCGCTGTCTTCAAAT	114
<i>ectB</i>	CAGTGGATACGGTCTGCCAA	CTCCGACAAACGCTAGCTGA	113
<i>sfp</i>	GTACCTCCTGCGCAAAGTGA	CACGACAGAGGGCTTTACGA	110
<i>amyE</i>	TCTGGGCGGAACGATTTTGA	CGAGTGCCGCCCTATTGTAT	110
<i>ccpA</i>	ATCAATTCCGGCTGCTTCCA	CACCGCCAAATCGCAATGAT	102

21 g/L NaOH, 6.3 g/L dinitrosalicylic acid, 5 g/L crystalline phenol, and 5 g/L Na₂SO₃. Next, the mixture was placed in boiling water for 5 min, followed by cooling down to room temperature. Finally, the absorbance value of mixture at 540 nm was measured [26]. The concentration of reducing sugar was determined based on the extracted correlation between standard glucose over a range of concentrations and the corresponding absorbance values at 540 nm. One unit of amylase activity was defined as the amount of enzyme required to release 1 μ g of glucose equivalent per minute under the assay condition using glucose as the standard.

2.5. PCR Amplification and Sequence Retrieval of Genes. The sequences of *pmxABCDE*, *spo0A*, *abrB*, *ectB*, and *sfp* have been amplified and collected in our previous study [17]. In this study, the primers (Table 1) for *amyE* encoding amylase and *ccpA* encoding CcpA in PCR reaction were designed based on the complete genome sequence of *Paenibacillus polymyxa* SC2 (GenBank access no. CP002213.2). Bacterial genomic DNA was extracted using a bacterial genomic DNA extraction kit (GE, USA). PCR reaction was performed as reported previously [27, 28]. In brief, the gene fragments were amplified in 50 μ L containing 37 μ L of ddH₂O, 5 μ L of 10X EasyTaq buffer, 4 μ L of 2.5 mM dNTPs, 100 nM forward primer, 100 nM reverse primer, 1 ng genomic DNA, and 1 U Taq DNA polymerase (TaKaRa, Dalian, China) with denaturation at 94°C for 5 min followed by 30 cycles of 1 min at 94°C, 50 s at 55°C, 90 s at 72°C, and a final 10 min extension at 72°C. At the end of reaction, PCR product was cooled to 4°C for further use. After size confirmation on 1.0% agarose gel, the desired amplicons were purified using a gel extraction kit (Qiagen, CA, USA) for TA cloning with pMD19-T simple vector (TaKaRa, Dalian, China). After sequencing by Sangon

Biotech (Shanghai, China), the gene sequences were collected and compared with the reference genes in GenBank for confirmation.

2.6. qPCR Analysis of Gene Expression. In brief, 0.5 mL of the bacterial cells was pelleted after centrifugation at 8,000 rpm for 10 min at 4°C and the total RNA was extracted using an RNAiso Plus kit (TaKaRa, Dalian, China) according to the manufacturer's instructions. RNA purity was spectrophotometrically evaluated based on OD_{260 nm}/OD_{280 nm} ratio. Then, 300 ng of DNA-free RNA was reversely transcribed to cDNA in a 10 μ L volume using PrimeScript™ RT Master Mix (Perfect Real Time) kit (Toyobo, Tokyo, Japan). After appropriate dilution, the obtained cDNA was used for amplification of target gene fragment with primer sets (Table 2) [17] by using the SYBR green Premix Ex Taq™ (Tli RNaseH Plus) kit. A master mixture was prepared and each well of reaction contained the following reagents: 5 μ L of SYBR Green Master Mix, 0.2 μ L forward primer and reverse primer, respectively, and 3.6 μ L of ddH₂O to a total of 9 μ L. After addition of 1 μ L of each diluted cDNA sample to each well, the PCR was run on CFX Connect Real-Time System (Bio-Rad, Hercules, CA) with an amplification protocol consisting of an initial denaturation at 95°C for 10 min, followed by 40 cycles of denaturation at 95°C for 15 s and annealing/elongation at 60°C for 30 s. Immediately after the final cycle of PCR, melting curve was analyzed to retrieve the specificity of the reaction based on the observation of melting temperature from the product [29].

The cycle threshold (C_T) for each PCR was determined using StatView software which automatically set the threshold signal at the log phase of amplification curve. The amplification efficiency of gene was retrieved from the slope of




FIGURE 1: Influence of replacement of glucose by starch on colistin production in *P. polymyxa*. (a) Colistin production along fermentation; (b) growth curve of *P. polymyxa* along fermentation; (c) rate of colistin biosynthesis along fermentation; (d) colistin production per cell biomass after 96 h fermentation. 45S: 45 g/L starch; 40S+5G: 40 g/L starch plus 5 g/L glucose; 30S+15G: 30 g/L starch plus 15 g/L glucose; 20S+25G: 20 g/L starch plus 25 g/L glucose; 45G: 45 g/L glucose.

that linear regression according to the formula $E=10^{(-1/\text{slope})}$. Several dilutions of each cDNA sample were assayed for the gene of interest in order to obtain a linear regression between the C_T values (ranging from 15 to 35 cycles) and the log of cDNA. The 116 bp of 16S rRNA gene fragment ranging from 16SF (5'-GAGAAGAAAGCCCCGGCTAA-3') to 16SR (5'-ACCAGACTTAAAGAGCCGCC-3') was used as the internal control to verify that there was an equal amount of target cDNA in all samples. The expression of target gene relative to 16S rRNA gene was calculated as described in report [30].

2.7. Data Analysis and Availability. Unless otherwise specified, triplicate reactions per experiment were performed. All data were presented as mean \pm standard error and tested for statistical significance based on analysis of variance

(ANOVA) followed by Dunnett's post hoc test using StatView 5.0 program. When the probability (p) was less than 0.05 and 0.01, the values were considered significantly (*) and very significantly (**) different, respectively.

3. Results

3.1. Dependence of Starch Concentration on Colistin Biosynthesis. To investigate the effect of starch on colistin accumulation in *P. polymyxa* C12, different amounts of glucose in fermentation medium were replaced by starch. As shown in Figure 1(a), colistin was undetectable within the first 24 h using 45 g/L glucose as the sole sugar source. Then, its production rapidly increased up to 48 h, followed by almost a constant in the remaining period. The highest yield of colistin was around 8.5×10^4 U/mL. Instead, the replacement of




FIGURE 2: Influence of replacement of glucose by starch on amylase and gene expression. (a) Amylase activity along fermentation; (b) amylase activity per cell biomass after 96 h fermentation; (c) relative expression level of *amyE*. 45S: 45 g/L starch; 40S+5G: 40 g/L starch plus 5 g/L glucose; 30S+15G: 30 g/L starch plus 15 g/L glucose; 20S+25G: 20 g/L starch plus 25 g/L glucose; 45G: 45 g/L glucose. The statistically significant results are related to the condition of 45 g/L glucose.

glucose by starch in fermentation medium gave detectable colistin at 12 h. Next, colistin production rapidly increased also up to 48 h, followed by a moderate increase in the remaining period. Overall, the more the glucose was replaced by starch, the higher the colistin was produced. The highest yield of colistin with 40 g/L starch plus 5 g/L glucose was around 1.66×10^5 U/mL, approximately one time higher than the one with 45 g/L glucose. The 45 g/L starch as the sole sugar source showed a similar result to 40 g/L starch plus 5 g/L glucose. Figure 1(b) indicated that the proportion of sugar source clearly affects the cell growth. Overall, the high proportion of glucose was beneficial to cell accumulation for early stage of fermentation. In contrast, the high proportion of starch was beneficial to cell accumulation for later stage of fermentation. Most probably, the use of glucose is faster than that of starch. Figure 1(c) showed that the rate of colistin biosynthesis with 45 g/L glucose as the sole sugar source rapidly increased and then decreased. The highest rate of colistin biosynthesis was 4.8×10^3 U/(mL·h) at 48 h. Similarly, the rate of colistin biosynthesis with all mixtures of glucose and starch rapidly increased and then decreased. Interestingly, the highest rate of colistin biosynthesis overall appeared earlier and higher with the increase of replacement of glucose with starch. The highest rate of colistin biosynthesis with 40 g/L starch plus 5 g/L glucose reached 6.7×10^3 U/(mL·h) at 36 h. Figure 1(d) further showed that the replacement of glucose with starch enhanced the colistin production per biomass. Overall, the more the glucose was replaced, the higher the colistin per biomass was produced. The highest yield of colistin per biomass with 40 g/L starch plus 5 g/L glucose was around 2.1×10^3 U/(mL·10⁶ CFU), approximately two times higher than the one with 45 g/L glucose. The 45 g/L starch as the sole sugar source displayed a similar result to 40 g/L starch plus 5 g/L glucose. All these data congruously indicated that

the replacement of glucose with starch stimulates the colistin accumulation in *P. polymyxa*.

3.2. Effect of Starch on Amylase Activity and Relative Expression of *amyE*. Starch should be decomposed by amylase before use in fermentation. Therefore, amylase activity was monitored. As shown in Figure 2(a), amylase activity with 45 g/L glucose as the sole sugar source can be detected at 48 h. Then, it increased to 11 U/mL at 60 h, followed by almost a constant in the remaining period. Amylase activities with both 40 g/L starch plus 5 g/L glucose and 45 g/L starch were around 100 U/mL at 48 h and 72 h, respectively, eight times higher than the one with 45 g/L glucose. Figure 2(b) further showed that amylase activity per 10⁶ CFU with 40 g/L starch plus 5 g/L glucose was around 1.48 U/mL, eight times higher than 0.17 U/mL of amylase activity per 10⁶ CFU with 45 g/L glucose. Figure 2(c) indicated that the relative expression level of *amyE* encoding amylase significantly increased with increase of starch replaced for glucose. All these results supported the reports that the transcription of *amyE* is strongly increased by starch [31] but repressed by glucose [32].

3.3. Effect of Starch on pH and Reducing Sugar Formation. Carbon source could affect reducing sugar and accordingly fermentation output [33, 34]. Figure 3(a) showed that the residue reducing sugars in fermentation with original glucose ranging from 15 g/L to 45 g/L decreased within the first 48 h and kept almost constant in the second 48 h. Most probably, the consumption of glucose in fermentation medium could result in the decrease of reducing sugar within the first 48 h. The higher the original concentration of glucose in fermentation medium was, the faster the reducing sugar decreased within the first 48 h. On the contrary, the residue reducing sugar in fermentation with original glucose ≤ 15 g/L




FIGURE 3: Influence of replacement of glucose by starch on residue reducing sugar (a) and pH (b) during fermentation. 45S: 45 g/L starch; 40S+5G: 40 g/L starch plus 5 g/L glucose; 30S+15G: 30 g/L starch plus 15 g/L glucose; 20S+25G: 20 g/L starch plus 25 g/L glucose; 45G: 45 g/L glucose. The statistically significant results are related to the condition of 45 g/L glucose.




FIGURE 4: Effect of replacement of glucose by starch on relative expression level of genes involved in regulation of colistin biosynthesis. (a) *ccpA*; (b) *abrB*; (c) *spo0A*. 45G: 45 g/L glucose; 20S+25G: 20 g/L starch plus 25 g/L glucose; 40S+5G: 40 g/L starch plus 5 g/L glucose. The statistically significant results are related to the condition of 45 g/L glucose.

increased within the first 48 h and kept almost constant in the second 48 h. Most probably, the rate of decomposition of starch in fermentation medium would surpass the rate of reducing sugar consumption within the first 48 h, thus resulting in the increase of reducing sugar. Figure 3(a) further indicated that the higher the original concentration of glucose was, the higher the concentration of residue reducing sugar was at the end of fermentation. The residue reducing sugars in fermentation medium with 45 g/L glucose and 45 g/L starch as original sugar were 13.6 g/L and 4.1 g/L, respectively, at 96 h. Figure 3(b) showed that the pH of fermentation solution with different sugar sources displayed a similar pattern, rapid decrease within the first 48 h and slight increase within the second 48 h. Overall, the higher the original glucose

concentration in fermentation medium was, the higher the pH of fermentation solution was within the second 48 h, which is negatively correlated with colistin production (Figure 1). The fact that the overall difference in pH of fermentation solution derived from proportion of sugar source is visible, but not remarkable, is worth noting.

3.4. Effect of Starch on Relative Expression of Genes for Regulation of Colistin Biosynthesis. CcpA encoded by *ccpA* is the master regulator of CCR in Gram-positive bacteria and it can affect the expression of the *abrB* [35]. Both *abrB* and *spo0A* are believed to be associated with colistin production [17]. Therefore, the relative expression of these three genes was investigated. Figure 4 showed that *ccpA* and *spo0A*




FIGURE 5: Continued.




FIGURE 5: Effect of replacement of glucose by starch on relative expression level of genes directly involved in colistin biosynthesis and secretion. (a) *pmxA*; (b) *pmxB*; (c) *pmxC*; (d) *pmxD*; (e) *pmxE*; (f) *ectB*; (g) *sfp*. 45G: 45 g/L glucose; 20S+25G: 20 g/L starch plus 25 g/L glucose; 40S+5G: 40 g/L starch plus 5 g/L glucose. The statistically significant results are related to the condition of 45 g/L glucose.

gave overall similar patterns in relative gene expression. The higher the original concentration of glucose in fermentation medium was, the lower the relative expression of either *ccpA* or *spo0A* was. Interestingly, *abrB* had the opposite pattern in relative gene expression. The higher the original concentration of glucose in fermentation medium was, the higher the relative expression of *abrB* was. All these results indicated that the replacement of glucose by starch can stimulate the expression of both *ccpA* and *spo0A* but repress the expression of *abrB*, which in turn stimulates colistin production (Figure 1).

3.5. Effect of Replacement of Glucose by Starch on the Relative Expression of Genes Directly Involved in Colistin Biosynthesis. As shown above, the replacement of glucose by starch can promote the relative expression of genes associated with regulation of colistin production. Therefore, the relative expression levels of genes directly involved in colistin biosynthesis and secretion were examined. Figure 5 showed that the replacement of glucose by starch stimulated the relative expressions of *pmxABCDE*, *ectB*, and *sfp*. The more the glucose was replaced, the higher the relative expressions of those genes were, indicating that the replacement of glucose by starch can promote the expression of those genes and in turn increased colistin production (Figure 1).

4. Discussion

Colistin is broadly used to treat the infection of Gram-negative pathogens, particularly prevalent multidrug-resistant bacteria. It is produced by *P. polymyxa*. To date, very few reports dealt with the medium optimization for improvement of colistin production. In the present study, the effect of replacement of glucose by starch in fermentation

medium on colistin production as well as transcription level of colistin biosynthesis-related genes was investigated. It was found that addition of starch could improve the production and biosynthesis rate of colistin (Figure 1). Moreover, the improvement extent was positively correlated with the amount of glucose replaced by starch (Figure 1). Our data further showed that the replacement of glucose by starch could refine two important fermentation factors, residue reducing sugar and pH (Figure 3). It seems that low concentration of residue reducing sugar and pH is better for colistin biosynthesis in *P. polymyxa* (Figure 1), but the detailed correlation mechanism needs to be further explored.

It has been found that the use of glucose represses the amylase activity and sporulation [36]. A report has shown that, relative to other carbon sources, glucose causes the strongest CCR, reducing the production of secondary metabolite [37]. It has been revealed that CCR is achieved by the global transcription regulator CcpA. The expression of *ccpA* results in the reduction of CCR [22]. Therefore, CcpA positively regulates secondary metabolism [23]. Our results showed that the replacement of glucose by starch could increase the relative expression of *ccpA* (Figure 4). As a result, the replacement of glucose by starch increased the transcription of *amyE* and amylase activity (Figure 2), which in turn is probably conducive to colistin (secondary metabolite) accumulation in *P. polymyxa* (Figure 1). Therefore, our findings are in line with the reports [22, 23, 36].

It has been demonstrated that Spo0A positively regulates secondary metabolism [38]. Our results indicated that the use of starch enhanced the relative expression level of *spo0A* (Figure 4) and subsequently increased colistin production (Figure 1), suggesting that *spo0A* also positively affects colistin biosynthesis in *P. polymyxa* at the transcriptional level. Therefore, our findings are in line with the report [38]. It has been found that the expression of *abrB* is negatively regulated

by Spo0A [15]. Our results also indicated that the relative expression level of *abrB* decreased with the increase of *spo0A* expression (Figure 4). There is evidence to show that AbrB negatively regulates colistin biosynthesis by directly binding to the upstream region of *pmxA* [11]. Thus, the decrease of *abrB* expression by adding starch (Figure 4) enhanced the relative expression of *pmxABCDE*, a gene cluster for colistin biosynthesis (Figure 5). As a result, colistin accumulation increased (Figure 1).

Data Availability

All the data used to support the findings of this study are available from the corresponding author upon request.

Conflicts of Interest

The authors have declared that no conflicts of interest exist.

Authors' Contributions

Zhiliang Yu and Juanping Qiu contributed equally to this work.

Acknowledgments

This study was supported by the National Natural Science Foundation of China (31670114) and Zhejiang Provincial Natural Science Foundation of China (LY16C010002) for which the authors are thankful.

References

- [1] D. R. Storm, K. S. Rosenthal, and P. E. Swanson, "Polymyxin and related peptide antibiotics," *Annual Review of Biochemistry*, vol. 46, pp. 723–763, 1977.
- [2] Z. Yu, W. Qin, J. Lin, S. Fang, and J. Qiu, "Antibacterial mechanisms of polymyxin and bacterial resistance," *BioMed Research International*, vol. 2015, Article ID 679109, 11 pages, 2015.
- [3] Z. Yu, Y. Cai, W. Qin, J. Lin, and J. Qiu, "Polymyxin E induces rapid *Paenibacillus polymyxa* death by damaging cell membrane while Ca^{2+} can protect cells from damage," *PLoS ONE*, vol. 10, no. 8, Article ID e0135198, 2015.
- [4] Z. Yu, L. Zhang, W. Qin, J. Yin, and J. Qiu, "Exogenous catalase stimulates the polymyxin E-induced rapid killing of *Paenibacillus polymyxa*," *International Journal of Peptide Research and Therapeutics*.
- [5] M. E. Falagas and A. Michalopoulos, "Polymyxins: old antibiotics are back," *The Lancet*, vol. 367, no. 9511, pp. 633–634, 2006.
- [6] J. Li, R. L. Nation, R. W. Milne, J. D. Turnidge, and K. Coulthard, "Evaluation of colistin as an agent against multi-resistant Gram-negative bacteria," *International Journal of Antimicrobial Agents*, vol. 25, no. 1, pp. 11–25, 2005.
- [7] A. S. Levin, A. A. Barone, J. Penço et al., "Intravenous colistin as therapy for nosocomial infections caused by multidrug-resistant *Pseudomonas aeruginosa* and *Acinetobacter baumannii*," *Clinical Infectious Diseases*, vol. 28, no. 5, pp. 1008–1011, 1999.
- [8] H. Tsubery, I. Ofek, S. Cohen, M. Eisenstein, and M. Fridkin, "Modulation of the hydrophobic domain of polymyxin B nonapeptide: effect on outer-membrane permeabilization and lipopolysaccharide neutralization," *Molecular Pharmacology*, vol. 62, no. 5, pp. 1036–1042, 2002.
- [9] N. I. Martin, H. Hu, and M. M. Moake, "Isolation, structural characterization, and properties of mattacin (polymyxin M), a cyclic peptide antibiotic produced by *Paenibacillus kobensis* M," *The Journal of Biological Chemistry*, vol. 278, no. 15, pp. 13124–13132, 2003.
- [10] S.-K. Choi, S.-Y. Park, and R. Kim, "Identification of a polymyxin synthetase gene cluster of *Paenibacillus polymyxa* and heterologous expression of the gene in *Bacillus subtilis*," *Journal of Bacteriology*, vol. 191, no. 10, pp. 3350–3358, 2009.
- [11] S.-Y. Park, S.-K. Choi, J. Kim, T.-K. Oh, and S.-H. Park, "Efficient production of polymyxin in the surrogate host *Bacillus subtilis* by introducing a foreign *ectB* gene and disrupting the *abrB* gene," *Applied and Environmental Microbiology*, vol. 78, no. 12, pp. 4194–4199, 2012.
- [12] L. E. N. Quadri, P. H. Weinreb, M. Lei, M. M. Nakano, P. Zuber, and C. T. Walsh, "Characterization of Sfp, a *Bacillus subtilis* phosphopantetheinyl transferase for peptidyl carrier protein domains in peptide synthetases," *Biochemistry*, vol. 37, no. 6, pp. 1585–1595, 1998.
- [13] M. Sunbul, N. J. Marshall, Y. Zou, K. Zhang, and J. Yin, "Catalytic turnover-based phage selection for engineering the substrate specificity of Sfp phosphopantetheinyl transferase," *Journal of Molecular Biology*, vol. 387, no. 4, pp. 883–898, 2009.
- [14] B. Niu, C. Rueckert, J. Blom, Q. Wang, and R. Borriss, "The genome of the plant growth-promoting rhizobacterium *Paenibacillus polymyxa* M-1 contains nine sites dedicated to nonribosomal synthesis of lipopeptides and polyketides," *Journal of Bacteriology*, vol. 193, no. 20, pp. 5862–5863, 2011.
- [15] M. Perego, G. B. Spiegelman, and J. A. Hoch, "Structure of the gene for the transition state regulator, *abrB*: regulator synthesis is controlled by the *spo0A* sporulation gene in *Bacillus subtilis*," *Molecular Microbiology*, vol. 2, no. 6, pp. 689–699, 1988.
- [16] Y. Kuratsu, Y. Arai, K. Inuzuka, and T. Suzuki, "Stimulatory effect of aspartic acid on colistin production by *Bacillus polymyxa*," *Agricultural and Biological Chemistry*, vol. 47, no. 11, pp. 2607–2612, 1983.
- [17] Z. Yu, C. Guo, and J. Qiu, "Precursor amino acids inhibit polymyxin E biosynthesis in *Paenibacillus polymyxa*, probably by affecting the expression of polymyxin E biosynthesis-associated genes," *BioMed Research International*, vol. 2015, Article ID 690830, 11 pages, 2015.
- [18] J. Wang, Y. Tao, Z. Xu, R. Xu, P. Cen, and X. Wang, "An integrated high throughput strategy to screen mutants of *Paenibacillus polymyxa* with high polymyxin E-productivity," *World Journal of Microbiology and Biotechnology*, vol. 24, no. 9, pp. 1885–1891, 2008.
- [19] J. S. Rokem, A. E. Lantz, and J. Nielsen, "Systems biology of antibiotic production by microorganisms," *Natural Product Reports*, vol. 24, no. 6, pp. 1262–1287, 2007.
- [20] J. Deutscher, "The mechanisms of carbon catabolite repression in bacteria," *Current Opinion in Microbiology*, vol. 11, no. 2, pp. 87–93, 2008.
- [21] K. Seidl, C. Goerke, C. Wolz, D. Mack, B. Berger-Bächi, and M. Bischoff, "Staphylococcus aureus CcpA affects biofilm formation," *Infection and Immunity*, vol. 76, no. 5, pp. 2044–2050, 2008.

- [22] Y. Tang, W. Wu, X. Zhang, Z. Lu, J. Chen, and W. Fang, "Catabolite control protein a of *Streptococcus suis* type 2 contributes to sugar metabolism and virulence," *Journal of Microbiology*, vol. 50, no. 6, pp. 994–1002, 2012.
- [23] J. Varga, V. L. Stirewalt, and S. B. Melville, "The CcpA protein is necessary for efficient sporulation and enterotoxin gene (cpe) regulation in *Clostridium perfringens*," *Journal of Bacteriology*, vol. 186, no. 16, pp. 5221–5229, 2004.
- [24] Z. Yu, Y. Zhu, W. Qin, J. Yin, and J. Qiu, "Oxidative stress induced by polymyxin E involves in rapid killing of *Paenibacillus polymyxa*," *BioMed Research International*, vol. 2017, Article ID 5437139, 12 pages, 2017.
- [25] Z. Yu, Y. Yang, Y. Wang, J. Yin, and J. Qiu, "Reactive oxygen species-scavenging system is involved in l-amino acid oxidase accumulation in *Pseudoalteromonas* sp. B3," *3 Biotech*, vol. 7, no. 5, 2017.
- [26] G. L. Miller, "Use of dinitrosalicylic acid reagent for determination of reducing sugar," *Analytical Chemistry*, vol. 31, no. 3, pp. 426–428, 1959.
- [27] Z. Yu, "Optimization of PCR amplification for sensitive capture of *Methanopyrus* isoleucyl-tRNA synthetase gene in environmental samples," *Annals of Microbiology*, vol. 60, no. 4, pp. 757–762, 2010.
- [28] Y. Huang, Y. Zeng, Z. Yu, J. Zhang, H. Feng, and X. Lin, "In silico and experimental methods revealed highly diverse bacteria with quorum sensing and aromatics biodegradation systems - A potential broad application on bioremediation," *Bioresource Technology*, vol. 148, pp. 311–316, 2013.
- [29] Z. Yu, J. Wang, J. Lin, M. Zhao, and J. Qiu, "Exploring regulation genes involved in the expression of L-amino acid oxidase in *Pseudoalteromonas* sp. Rf-1," *PLoS ONE*, vol. 10, no. 3, Article ID e0122741, 2015.
- [30] T. D. Schmittgen and K. J. Livak, "Analyzing real-time PCR data by the comparative CT method," *Nature Protocols*, vol. 3, no. 6, pp. 1101–1108, 2008.
- [31] A. Lachmund, U. Urmann, K. Minol, S. Wirsal, and E. Ruttkowski, "Regulation of α -amylase formation in *Aspergillus oryzae* and *Aspergillus nidulans* transformants," *Current Microbiology*, vol. 26, no. 1, pp. 47–51, 1993.
- [32] F. G. Priest, "Effect of glucose and cyclic nucleotides on the transcription of α -Amylase mRNA in *Bacillus subtilis*," *Biochemical and Biophysical Research Communications*, vol. 63, no. 3, pp. 606–610, 1975.
- [33] R. P. Desai, T. Leaf, E. Woo, and P. Licari, "Enhanced production of heterologous macrolide aglycones by fed-batch cultivation of *Streptomyces coelicolor*," *Journal of Industrial Microbiology and Biotechnology*, vol. 28, no. 5, pp. 297–301, 2002.
- [34] P. Poudel, Y. Tashiro, H. Miyamoto, H. Miyamoto, Y. Okugawa, and K. Sakai, "Direct starch fermentation to l-lactic acid by a newly isolated thermophilic strain, *Bacillus* sp. MC-07," *Journal of Industrial Microbiology and Biotechnology*, vol. 42, no. 1, pp. 143–149, 2015.
- [35] C. Ren, Y. Gu, Y. Wu et al., "Pleiotropic functions of catabolite control protein CcpA in Butanol-producing *Clostridium acetobutylicum*," *BMC Genomics*, vol. 13, no. 1, article no. 349, 2012.
- [36] N. Shih and R. G. Labbé, "Effect of glucose on sporulation and extracellular amylase production by *Clostridium perfringens* type A in a defined medium," *Current Microbiology*, vol. 29, no. 3, pp. 163–169, 1994.
- [37] K. D. Singh, M. H. Schmalisch, J. Stülke, and B. Görke, "Carbon catabolite repression in *Bacillus subtilis*: Quantitative analysis of repression exerted by different carbon sources," *Journal of Bacteriology*, vol. 190, no. 21, pp. 7275–7284, 2008.
- [38] I.-H. Huang, M. Waters, R. R. Grau, and M. R. Sarker, "Disruption of the gene (*spo0A*) encoding sporulation transcription factor blocks endospore formation and enterotoxin production in enterotoxigenic *Clostridium perfringens* type A," *FEMS Microbiology Letters*, vol. 233, no. 2, pp. 233–240, 2004.