

EPP0999**The effect of job stress appraisals on mental health among health professionals: The mediating role of work engagement**

A. Chaabouni*

Department Of Psychology And Pedagogic Science, St Mary's University, Twickenham, United Kingdom

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1244

Introduction: Health professionals face an increased risk of developing mental health difficulties due to work-related stress. It has been demonstrated that work engagement has a protective role on mental health from work-related stress. The majority of the research on the psychological impact of job stress among health professionals focused on the work-related stressors or the type of stressors as challenges or hindrances. However, the impact might depend on an individual's appraisal of challenges and hindrances.

Objectives: - Examine the effects of job appraisals on mental health.
- Establish the role of work engagement as a mediator between them.

Methods: An online survey was completed by 196 health professionals and included questionnaires about job appraisals, stressors (variety of tasks, responsibility and cooperation with colleagues), work engagement, anxiety and depression.

Results: Appraising stressors as challenges did not have any direct impact on mental health, whereas hindrance appraisals had a negative influence. Participants who appraised cooperation with colleagues as challenging reported lower levels of depression through higher work engagement ($B = -0.17$, 95% CI $[-0.354, -0.027]$). Appraising variety of tasks as a hindrance predicted higher levels of depression through lower work engagement ($B = 0.150$, 95% CI $[0.041, 0.289]$). Participants appraising the other two stressors as hindrances were more anxious and depressed through lower work engagement.

Conclusions: The negative psychological impact of hindrance appraisals was persistent, whereas the positive impact of challenge appraisals through work engagement depended on the stressor. Stress interventions may need to consider both the type of appraisal and the type of stressor.

Keywords: health professionals; work engagement; Depression; Anxiety

EPP0998**Social networks, depressive symptoms and quality of life in the elderly: Results of an intergenerational program**

A.P. Amaral*, M.J. Nascimento and C. Rocha

Coimbra Health School, Polytechnic of Coimbra, Coimbra, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1245

Introduction: Intergenerational programs involving children and the elderly promote intergenerational interactions and can positively affect physical and mental health, and the quality of life of the elderly.

Objectives: To test the effects of an intergenerational intervention in social isolation, depressive symptoms and quality of life of the elderly.

Methods: This study employed a pretest-posttest design. Measures: Portuguese versions of Geriatric Depression Scale, Lubben Social Networks Scale and WHOQOL-OLD. Participants: 12 elderly, 75% females, with mean age of 80.8 years ($sd=8.8$) and 20 kindergarten children (65% female) with mean age of 4,1 years ($sd=0.79$). The intervention ran for 6 weeks, with 11 intergenerational group sessions, each range between 30 and 120 minutes. A nonparametric paired samples tests was conducted to evaluate the impact of the intervention.

Results: After the intervention, when comparing elderly with and without depressive symptoms, results showed significant differences in the total value of quality of life ($p=.048$) and in the facets: 1) Sensory functioning, 2) Autonomy, 3) Past, present and future activities ($p=.003$; $p=.018$; $p=.030$, respectively). 12,5% of de elderly with depressive symptoms before the intervention no longer have depressive symptoms after the intervention. Regarding social networks, there were no significant differences ($p=.576$) between the mean values of the two assessments.

Conclusions: The implemented intergenerational program was effective in promoting quality of live and minimized depressive symptoms. After the intervention, the number of the elderly without depressive symptoms have increased and these presented a higher quality of life. Finally, longitudinal studies with a large sample are needed to consolidate results.

Keywords: quality of life; Elderly; Intergenerational; depressive symptoms

EPP0999**Does the crisis intervention team (CIT) training improve police officers' knowledge, attitude, and mental health stigma?**N. Veluri^{1*} and Z. Mansuri²

¹N/a, American University of Integrative Sciences, School of Medicine, St.Michale, Barbados and ²Department Of Psychiatry, Boston Children's Hospital/Harvard Medical School, Boston, United States of America

*Corresponding author.

doi: 10.1192/j.eurpsy.2021.1246

Introduction: The Crisis Intervention Team (CIT) training was developed to educate police officers regarding the complexity of mental health (MHI) issues, and better prepare them for crisis encounters with persons with mental illness (PwMI).

Objectives: To determine if CIT training improves police officers' knowledge, attitude, and stigma about mental health issues.

Methods: A systematic review followed the PRISMA protocol and was conducted on the PubMed database (Figure 1). Search strings were "crisis intervention team training," "crisis intervention team," "CIT," "effectiveness," and "police." Inclusion eligibility required primary studies using surveys that measured the CIT training outcomes (i.e., knowledge, attitude, and stigma). Literature/narrative reviews or opinions were excluded.