


Unifying Pathophysiological Explanations for Sports-Related Concussion and Concussion Protocol Management: Literature Review

Praveen Satarasinghe¹, D Kojo Hamilton²,
Robert J Buchanan^{1,3} and Michael T Koltz^{1,3} 

¹Department of Neurosurgery, Dell Medical School, The University of Texas at Austin, Austin, TX, USA. ²Department of Neurosurgery, University of Pittsburgh, Pittsburgh, PA, USA. ³Department of Neurosurgery, Seton Brain and Spine Institute, Austin, TX, USA.

Journal of Experimental Neuroscience
Volume 13: 1–10
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1179069518824125



ABSTRACT

OBJECTIVE: There is a plethora of theories about the pathophysiology behind a sport-related concussion. In this review of the literature, the authors evaluated studies on the pathophysiology of sport-related concussion and professional athlete return-to-play guidelines. The goal of this article is to summarize the most common hypotheses for sport-related concussion, evaluate if there are common underlying mechanisms, and determine if correlations are seen between published mechanisms and the most current return-to-play recommendations.

METHODS: Two authors selected papers from the past 5 years for literature review involving discussion of sport-related concussion and pathophysiology, pathology, or physiology of concussion using mutually agreed-upon search criteria. After the articles were filtered based on search criteria, pathophysiological explanations for concussion were organized into tables. Following analysis of pathophysiology, concussion protocols and return-to-play guidelines were obtained via a Google search for the major professional sports leagues and synthesized into a summary table.

RESULTS: Out of 1112 initially identified publications, 53 met our criteria for qualitative analysis. The 53 studies revealed 5 primary neuropathological explanations for sport-related concussion, regardless of the many theories talked about in the different papers. These 5 explanations, in order of predominance in the articles analyzed, were (1) tauopathy, (2) white matter changes, (3) neural connectivity alterations, (4) reduction in cerebral perfusion, and (5) gray matter atrophy. Pathology may be sport specific: white matter changes are seen in 47% of football reports, tauopathy is seen in 50% of hockey reports, and soccer reports 50% tauopathy as well as 50% neural connectivity alterations. Analysis of the return-to-play guidelines across professional sports indicated commonalities in concussion management despite individual policies.

CONCLUSIONS: Current evidence on pathophysiology for sport-related concussion does not yet support one unifying mechanism, but published hypotheses may potentially be simplified into 5 primary groups. The unification of the complex, likely multifactorial mechanisms for sport-related concussion to a few common explanations, combined with unique findings within individual sports presented in this report, may help filter and link concussion pathophysiology in sport. By doing so, the authors hope that this review will help guide future concussion research, treatment, and management.

KEYWORDS: Concussion, sports, return to play, protocol, pathophysiology, neuropathology

RECEIVED: October 2, 2018. **ACCEPTED:** December 19, 2018.

TYPE: Traumatic Brain Injury (TBI) and Chronic Traumatic Encephalopathy (CTE) - Review

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Michael T Koltz, Department of Neurosurgery, Dell Medical School, The University of Texas at Austin, Austin, TX 78712, USA.
Email: mtkoltz@gmail.com

Introduction

According to the Centers for Disease Control (CDC) and the Brain Injury Research Institute, approximately 1.6 to 3.8 million individuals face sport-related concussions in the United States every year and the CDC estimates that roughly 10% of all contact sport athletes suffer from at least 1 concussion yearly, with brain injuries taking the top spot in causes of sport-related injuries and the cause of death in children and young adults.¹ Contact sports can be broken down into either full or limited contact classifications.² Full contact describes sports such as football, hockey, soccer, basketball, and boxing, whereas limited contact describes sports such as baseball, softball, cheerleading, and volleyball. The sport and position played within the sport may be a

contributing risk factor for sport-related concussions which given the number of participants in such activities may help support the statistics of increasing frequency of sport-related concussion.

Neurodegenerative conditions may be linked to repetitive concussion, with 1 potential neuropathological sequela being chronic traumatic encephalopathy (CTE).^{2,3} Repetitive concussions may be a risk factor for cognitive decline and neurodegenerative disease such as CTE, and therefore return-to-play (RTP) guidelines play a pivotal role in player development. Rapid return to play, an underdiagnosis of the problem, may place the participant at a “health risk.” Delayed return to play, an overdiagnosis of the problem, may place the participant at an “opportunity for success risk.”



Currently, a plethora of concussion literature exists; however, theories are variable, information is not consolidated for easy future investigation, and no article has yet attempted to concisely summarize the breadth of knowledge on sport-related concussion. The variability in theories on concussion pathophysiology and mechanism is the result of the medical and scientific community's high output of research attempting to elucidate the complex, largely unknown pathophysiology. Given the unknown nature of sport-related concussion, in combination with American culture's love for sport, a firestorm of opinions have been expressed in America's media platform.⁴⁻⁶ No clear pathologic definition or description of concussion exists, leading to a lack of consensus in the use of the term, lack of agreement on the neuropathological progression of brain damage associated with a concussion, and, ultimately, difficulty deciding how to clinically distinguish and treat a concussion compared with other traumatic brain injuries (TBIs).⁷⁻⁸ As evidenced by the confusion that still exists among even experts in sport-related concussion, the understanding of concussion is still a problem in the modern day.

In this article, the authors systematically analyzed the current literature for theories on the pathophysiological process underlying a sport-related concussion, as well as the RTP guidelines instituted within different contact sports where players are at risk for concussion. Through review of the literature, the authors of this article sought to collect pathophysiological explanations and management guidelines for different sport-related concussions, concisely present the information, and identify commonalities and differences that may shed light and coherently synthesize opinions on sport-related concussion.

Methods

Overview

For this study, there were 2 sets of information that were collected: (1) pathophysiology of sport-related concussion and (2) RTP guidelines following concussion associated with different contact sports. Thus, the methods consisted of 2 separate protocols for information collection (see section "Search"). Our methods for this study of concussion pathophysiology in sport followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.

Search

For information set 1 (pathophysiology), a review of the literature was performed by the authors to identify studies that investigated pathophysiology of sport-related concussion. For information set 2 (RTP guidelines), websites for major league sports concussion protocols (eg, Major League Soccer [MLS], National Hockey League [NHL], National Basketball Association [NBA], National Football League [NFL], and




Figure 1. Article search criteria for review of the literature on sports-related concussion.

Major League Baseball [MLB]) and National Collegiate Athletic Association (NCAA) concussion protocols were referenced for RTP guidelines. As this study primarily looked into sport-related concussion in college and professional athletes, analysis of NCAA and professional sport guidelines was deemed appropriate.

The PubMed database was used to collect information set 1. In PubMed, MeSH term "concussion" was used to find all articles in records that were indexed under or related to concussion. To further narrow down the search, under the "concussion" MeSH search, "pathophysiology," "pathology," and "physiology" checkboxes were selected to identify articles that mentioned pathophysiology of concussion. Finally, the keyword "sports" was added to the search to only select published articles related to sport-related concussion (see Figure 1).

After clearing the previous search history on Google, an incognito window was used to collect information set 2. To obtain RTP concussion guidelines for major league sports, a Google search was completed using the phrase "2018 concussion return to play guidelines" followed by the name of the major league for each sport (see Figure 1). To obtain RTP concussion guidelines for NCAA sports, the same phrase "2018 concussion return to play guidelines" followed by "NCAA" was completed.

Inclusion criteria

For information set 1, because the authors' goal in this article was to conduct a recent review of the literature on the pathophysiology of sport-related concussion, specific criteria were incorporated into the search strategy through PubMed to take into account publication date, publication language, publication full-length text availability, and publication content (see Figure 2). Only articles published in the past 5 years were included in this review to summarize and offer an analysis of the most recently published and up-to-date literature.

For information set 2, the most recent RTP guidelines from the official websites of the major league sports and NCAA were collected. If the most recent RTP guidelines were not viewable through the official websites, publically available data from media articles reporting the concussion policies were used. Obtaining the most up-to-date concussion RTP guidelines was prioritized.

Full-length article criteria

For information set 1, after using the search strategy and filtering further articles out through the inclusion criteria, 208 full-length articles were identified and read by both the primary and senior authors. Studies that did not mention both




Figure 2. Article selection for review of the literature on sports-related concussion.

sports and concussion were excluded. Because the authors wanted to identify primary literature on concussion pathophysiology, systematic reviews, and other review articles that had not been eliminated via application of the inclusion criteria were excluded. Studies that discussed sports below the collegiate level were excluded, as the goal of this article was to review sport-related concussion at the collegiate and professional levels. Data were extracted from the articles independently based on an agreed-upon format for the tables in section “Results.” Pathophysiology of concussion was the primary finding sought, along with the mentioned sport in each article, lead author, and date of publication.

Results

For information set 1, comprehensive screening of 208 full-length articles resulted in 53 articles for analysis. The 53 studies were summarized for pathophysiology of concussion and categorized by sport (see Tables 1 to 7). The articles summarized in the tables cover concussion pathophysiology in individual sports, including football, hockey, basketball, soccer, baseball, boxing, fighting, and rugby, as well as multiple sports.

As illustrated in the final summary table of all of the articles analyzed in our study, the primary neuropathological explanations for sport-related concussion are (1) tauopathy, (2) white matter tract changes, (3) neural connectivity alterations, (4) reduction in cerebral perfusion, and (5) gray matter

Table 1. Summary of football articles on sports-related concussion.

FIRST AUTHOR	JOURNAL OF PUBLICATION	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Mez et al ⁹	JAMA	Tauopathy and neurofibrillary tangle (NFT) deposition
Slobounov et al ¹⁰	Neuroimage Clin	Reduced neural connections in isthmus cingulate cortex (ICC) and hippocampus and reduced cerebral blood flow
Coughlin et al ¹¹	JAMA Neurol	White matter changes and increase in translocator protein (TSPO)
Cherry et al ¹²	Acta Neuropathol Commun	Hyperphosphorylated p-tau, neuroinflammation, and chronic microglial activation
Bahrami et al ¹³	Radiology	White matter tract changes and decrease in fractional anisotropy (FA)
Strain et al ¹⁴	J Neurotrauma	White matter changes
Amen et al ¹⁵	J Alzheimers Dis	Reduced cerebral perfusion and blood flow
Meier et al ¹⁶	J Neurotrauma	Dentate gyrus atrophy, reduction in kynurenine, and reduction in hippocampal volume
Koerte et al ¹⁷	J Neurotrauma	Cavum septum pellucidi (CSP)
Wang et al ¹⁸	J Neurotrauma	Reduced cerebral perfusion and blood flow
Meier et al ¹⁹	J Neurotrauma	Reduction in cortical thickness
Gardner et al ²⁰	J Neurotrauma	Cavum septum pellucidi (CSP)
Poole et al ²¹	Dev Neuropsychol	Alteration in brain metabolism
Davenport et al ²²	J Neurotrauma	White matter changes
Strain et al ²³	Neurology	White matter changes

Table 2. Summary of hockey articles on sports-related concussion.

FIRST AUTHOR	JOURNAL OF PUBLICATION	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Shahim et al ²⁴	Neurology	Tauopathy and reduction in S100B
Manning et al ²⁵	Neurology	White matter tract changes and decreased mean diffusivity
Shahim et al ²⁶	Neurology	Increase in serum neurofilament (NfL)
Wright et al ²⁷	PLoS One	Myelin disruption

Table 3. Summary of basketball articles on sports-related concussion.

FIRST AUTHOR	JOURNAL OF PUBLICATION	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Yengo-Kahn et al ²⁸	Phys Sportsmed	White matter tract changes and angular acceleration alteration

Table 4. Summary of soccer articles on sports-related concussion.

FIRST AUTHOR	JOURNAL OF PUBLICATION	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Ling et al ²⁹	Acta Neuropathol	Amyloid angiopathy, TDP-43 proteinopathy, hippocampal sclerosis, and corticobasal degeneration
Di Virgillio et al ³⁰	EBioMedicine	Corticomotor system inhibition
Grinberg et al ³¹	J Alzheimers Dis	TDP-43 proteinopathy and hippocampal sclerosis
Hwang et al ³²	J Neurotrauma	Dysfunction in vestibular processing
Lipton et al ³³	Radiology	White matter microstructural changes

Table 5. Summary of baseball articles on sports-related concussion.

FIRST AUTHOR	JOURNAL OF PUBLICATION	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Zimmer et al ³⁴	J Athl Train	Structural atrophy and angular acceleration alteration

Table 6. Summary of other sports (boxing/fighting/rugby) articles on sports-related concussion.

FIRST AUTHOR	JOURNAL OF PUBLICATION	SPORT	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Bernick et al ³⁵	Br J Sports Med	Fighting	Reduction in thalamic volumes
Johnson et al ³⁶	J Neurotrauma	Rugby	Alteration in neural connectivity
Nesselius et al ³⁷	PLoS One	Boxing	NfL elevation and axonal injury
Nesselius et al ³⁸	PLoS One	Boxing	Increased NfL

atrophy (see Table 8). Several articles mentioned multiple of the 5 concussion mechanisms listed above. Of these articles, 32% highlighted tauopathy, 28% highlighted white matter tract changes, 25% highlighted neural connectivity alterations, 19% highlighted a reduction in cerebral perfusion, and 11% highlighted gray matter atrophy. Note that the percentages add up to over 100% because several articles mentioned

more than 1 neuropathological explanation for sport-related concussion.

There was also variability in pathophysiological explanation for concussion across different sports. Pathology may be sport specific: white matter changes are seen in 47% of football reports, tauopathy is seen in 50% of hockey reports, and soccer reports 50% tauopathy as well as 50% neural connectivity alterations.

Table 7. Summary of articles on sports-related concussion referencing all sports.

FIRST AUTHOR	JOURNAL OF PUBLICATION	NEUROPATHOLOGICAL EXPLANATION FOR CONCUSSION
Di Battista et al ³⁹	J Neuroimmunol	Tauopathy
Churchill et al ⁴⁰	Front Neurol	Elevated FA, reduction in white matter, reduction in functional connectivity, and alteration in brain metabolites
Gill et al ⁴¹	Neurology	Tauopathy
Reches et al ⁴²	Brain Inj	Changes in functional network connectivity
Safinia et al ⁴³	J Vasc Interv Neurol	Tauopathy
Papa ⁴⁴	Sports Med Arthrosc	S100 and NfL alterations. Tauopathy
Babcock et al ⁴⁵	Brain Inj	Glial fibrillary acidic protein (GFAP) alteration
Herrera et al ⁴⁶	J Neurotrauma	White matter tract changes
Merchant-Borna et al ⁴⁷	J Neurotrauma	Neuroinflammation and inflammatory gene expression changes
Kontos et al ⁴⁸	Brain Imaging Behav	Reduced brain network connectivity
Militana et al ⁴⁹	Brain Imaging Behav	Altered functional connectivity
Goswami et al ⁵⁰	Brain Struct Funct	Reduction in mean diffusivity and orbitofrontal, cortical, and temporal atrophy
Pham et al ⁵¹	PLoS One	Elevation in cellular prion protein (PrPC)
Schulte et al ⁵²	J Athl Train	Elevation in S100B
Tremblay et al ⁵³	Brain	White matter tract changes
Czerniak et al ⁵⁴	Brain Imaging Behav	Alterations in functional connectivity
Murugavel et al ⁵⁵	J Neurotrauma	White matter tract changes and reduction in diffusivity
Bazarian et al ⁵⁶	PLoS One	White matter tract changes
Ji et al ⁵⁷	J Neurotrauma	White matter tract changes
Kiechle et al ⁵⁸	PLoS One	Increase in S100B
De Beaumont et al ⁵⁹	BMC Neurol	Reduction in M1 glutamate concentration and metabolite alteration
Dettwiler et al ⁶⁰	J Neurotrauma	Activation of prefrontal cortex (PFC) and inferior parietal lobe

Table 8. Primary neuropathological explanations for sports-related concussion.

CONCUSSION PATHOPHYSIOLOGY MECHANISM	STRUCTURAL OR FUNCTIONAL DEFICIT	NUMBER OF ARTICLES MENTIONING MECHANISM
Tauopathy/proteinopathy	Structural	17
White matter tract changes	Both	15
Neural connectivity alterations	Functional	13
Reduction in cerebral perfusion	Functional	10
Gray matter atrophy	Structural	6

Total number of articles exceeds n=53 because several articles mentioned multiple concussion mechanisms.

For information set 2, concussion RTP guidelines from the NFL, NBA, MLS, MLB, and NHL were summarized, along with the NCAA concussion RTP guidelines (see Table 9). Across

all the major sports leagues, concussion RTP protocols, regardless of the number of specific steps and unique intricacies, include 3 common stages: (1) rest, (2) re-exertion, and (3) return to play.

Table 9. Summary of concussion RTP guidelines.

SPORT, LEAGUE (LEVEL)	YEAR OF MOST RECENT GUIDELINES	SOURCE OF INFORMATION	MAJOR POINTS FROM RTP CONCUSSION PROTOCOL
Football, NFL (pro)	2017/2018	NFL Head, Neck and Spine Committee, playsmartplaysafe.com ⁶¹	(1) Players first rest until signs and symptoms disappear. (2) Players then undergo light aerobic exercise under oversight of medical staff. (3) Players start strength activities and cardiovascular activities. (4) Players start non-contact football activities like throwing and catching. (5) Players can resume football activities and full contact after clearance by team physician and examination by independent neurological consultant.
Hockey, NHL (pro)	2016/2017	NHL Concussion Education Program ⁶²	(1) Player cannot return to practice or game on the same day event occurred, irrespective of symptom resolution. (2) Player may return to unrestricted play when there is complete recovery of concussion symptoms at rest, no emergence of concussion symptoms at competitive exertion levels, and player has been judged by club physician and consulting neuropsychologist. (3) There is no mandatory period of withdrawal time following concussion. (4) Club physician is solely responsible for making RTP decisions based on the above parameters.
Soccer, MLS (pro)	2017/2018	US Soccer National Team Concussion Evaluation and Management Program ⁶³	(1) Rest until asymptomatic. (2) Light aerobic exercise. (3) Moderate aerobic exercise. (4) Sport-specific training. (5) Non-contact, full exertion drills. (6) Begin contact heading training. (7) Full contact training. (8) Return to competition when instructed by team physician.
Basketball, NBA (pro)	2017/2018	NBA concussion protocol program, via NBC Sports ⁶⁴	(1) No evidence of concussion symptoms at rest. (2) Evaluation by physician. (3) Successful completion of NBA exertion protocol. (4) Team physician and director of NBA concussion program agreement to return player to play.
Baseball, MLB (pro)	2011/2012	MLB Players Association concussion policy, via ESPN ⁶⁵	(1) All symptoms at rest and with exertion cleared, including baseball activities. (2) ImPACT (immediate post-concussion assessment and cognitive testing) results have returned to baseline. (3) Club physician feels it is safe for player to return to competition.
All sports, NCAA (collegiate)	2017/2018	NCAA Sport Science Institute ⁶⁶	(1) Light aerobic exercise, no resistance training until asymptomatic. (2) Sport-specific activities with no head impact until asymptomatic. (3) Non-contract sports drills with resistance training until asymptomatic. (4) Unrestricted training until asymptomatic. (5) Clearance by team physician for return to competition.

RTP, return-to-play; NFL, National Football League; NHL, National Hockey League; MLS, Major League Soccer; NBA, National Basketball Association; MLB, Major League Baseball; NCAA, National Collegiate Athletic Association.

Discussion

Concussion awareness is now mainstream with numerous, often polarizing, viewpoints.^{5–8,67} The pathophysiology of concussion remains unknown but robust active research may soon elucidate a mechanism. This unknown, along with the potential link of repeat concussion to CTE, has been the fuel to these polarizing viewpoints. In a recent 2017 study of concussion, researchers suggested the presence of CTE in more than 90% of the studies of deceased National Football League (NFL) players. This link has, however, been convincingly contested by Castellani and colleagues^{68,69} This ambiguity is problematic because there are potentially severe negative consequences related to concussive impacts and repeated concussions on athletes across all levels of play. Without a better understanding of sport-related concussion, the overdiagnosis and underdiagnosis of concussion can be detrimental to athletes and sport as a

whole. For example, overdiagnosis of concussion can cause athletes to unnecessarily miss games, lose training time, and even suffer reduction in pay. On the other hand, underdiagnosis of concussion becomes a public health concern with concussed and sub-concussed athletes continuing to play which may lead to neurodegenerative processes later in life.^{68–70}

The exact pathology of concussion is not known and its sequel, such as CTE, debatable among leaders in the field.^{7,67–69} For this reason, a list of identifiable risk factors is dynamic. Potential risk factors for concussion include prior concussion, participation in full or limited contact sports, the position played in a sport, female sex, older age, occurrence of migraines, alcohol/drug abuse, dementia, motor-neuron disease, depression, and several psychiatric disorders.⁷¹ Further potential consequences of concussion include the onset of headache, cognitive deficits, balance and coordination problems, behavioral

issues, and neurodegenerative disease such as CTE.⁶⁷ If one accepts CTE as a neuropathological entity, then potential links have been made to behavioral and emotional changes such as agitation, apathy, aggressiveness, suicidal ideation, and poor impulse control.^{2,10} CTE may also be associated with disease processes such as depression, Parkinsonism, and dementia.^{2,10,70}

After examining concussion publications, the authors' confirm that the literature is vast and not yet ready for simplification into 1 neuropathological explanation. However, our review of the literature did simplify the existing literature and did surface several findings that may show trends in concussion literature, concussion across sports, concussion pathophysiology, and future direction in concussion research and management: (1) there are potentially 5 primary explanations for concussion pathophysiology amidst the plethora of theories described in the reviewed literature; (2) concussion pathophysiology is a both structural and functional phenomenon; (3) concussion pathophysiology may vary across individual sports; (4) concussion RTP guidelines have a similar structure across different sports.

Theories of concussion pathophysiology

Although sport-related concussion has been loosely defined as neurological symptoms and cognitive problems that result from direct blow to the head, face, neck, or other body parts that transmits force to the head, there are many areas of debate regarding the pathophysiology producing these symptoms on head trauma.⁷⁰ By simplifying the voices in the literature, our first finding indicated the presence of 5 predominant pathophysiological explanations for sport-related concussion. These pathophysiological findings may help elucidate the true mechanism and definition of a concussion. Our survey of the literature identified 53 studies that clearly explained a pathophysiological explanation for sport-related concussion. The studies either looked at concussions across all different sports or focused on concussions in a specific sport (see Tables 1 to 7). Analysis of the individual concussion pathophysiological processes across all sports and within individual sports shows both commonalities and differences. In all of the articles systematically reviewed across all sports, (1) tauopathy, (2) white matter tract changes, and (3) alterations in neuronal connectivity showed up as the top 3 explanations for concussion symptoms, followed by the less common explanations of (4) reduction in cerebral perfusion and (5) gray matter atrophy (see Table 8).

Each of the 5 primary neuropathological explanations for sport-related concussion identified in our systematic literature review has pathophysiological theories: (1) The tauopathy explanation is centered on the abnormal accumulation of abnormally phosphorylated tau protein in the depth of cortical sulci on concussion trauma.⁷² The tau protein normally stabilizes microtubules in axons necessary for neuronal communication. The distribution of hyperphosphorylated tau in the brain

leads to the accumulation of neurofibrillary lesions, a loss in the neuronal network connectivity, and mass cell death and dysfunction.⁶⁸ Tau hyperphosphorylation is characteristic of normal aging, but in athletes who suffer concussions abnormally phosphorylated tau is stained at an early age. (2) The white matter change explanation is centered on the alteration of myelin, a lipid-rich substance that insulates axons and is responsible for coordinating neural communication.⁷³ In this theory, athletes who suffer concussions undergo multifocal axonal varicosities involving the subcortical white matter microstructure, leading to delayed action potentials and signals.⁶⁹ Examples of changes in white matter interfaces include widening of frontal and occipital cortical sulci, damage that leads to release of neural proteins that injure the brain, reduced diffusivity and total brain volume, and diffuse axonal injury.⁷⁴ (3) The neural connectivity explanation relies on neuroimaging to show a reduction in regional and global neuronal networks.⁷⁵ Such a reduction leads to delayed coordinated communication between brain regions, a loss in cognition, and other neurological symptoms. (4) The cerebral perfusion explanation models the vascular hypothesis described previously, emphasizing concussive symptoms resulting from obstruction or trauma resulting in a change in cerebral blood flow and potentially ischemia.⁷¹ (5) Finally, the gray matter atrophy explanation attributes concussion symptoms to trauma leading to overall shrinkage of cortical and subcortical structures, like the thalamus and hippocampus.⁷⁶ These structures are crucial for memory and cognition, processes that are reduced as a result of sport-related concussion.

Concussion: a structural and functional phenomenon

Even though these theories behind concussion may be shared, there is still ongoing debate and constant generation of new theories. One of the major controversies is whether a concussion is caused by functional or structural damage.⁸ Our second finding indicated that concussion pathophysiology cannot be reduced to one or the other, and has both structural and functional mechanisms. Structural damage refers to changes in brain structure that may be related to concussion symptomology. Functional damage refers to changes in cellular or neuronal connections and functions that produce concussion symptoms. The finding that structural and functional deficits results from concussion is no surprise. Concussion is a complicated disease of the most complex organ in the body. Various structural and functional factors go into a concussion. For example, different players have different brain structures and functional situations before a concussion that affect brain pathology. Point of impact may be different. Risk factors are present for different individuals. Factors like the analysis of tissue samples post concussion are constantly prone to error through the variable brain fixation process that can produce both functional and structural changes in brain pathology.⁸

With such factors attributing concussions to both functional and structural origins along with the same finding in the review of literature, further investigation of concussion requires looking at the problem from a heterogeneous and multifactorial lens.

Pathophysiological differences across sport

The third finding in our study is that concussion pathophysiology may vary among sports and have specific trends within individual sports. In football, studies attributing concussions to white matter changes predominated (7 out of 15 studies). In hockey, tauopathy explanations predominated (2 out of 4 studies). In soccer, the neural connectivity alteration (2 out of 4 studies) and tauopathy (2 out of 4 studies) explanations predominated. Basketball and baseball both only had 1 study in the primary literature over the past 5 years that mentioned pathophysiology of concussion. An explanation for the lack of literature may be that, among all sports, baseball and basketball consistently have the lowest incidence of concussion. In articles that looked at the concussion pathophysiology in all sports, tauopathy (7 out of 21 studies), white matter changes (6 out of 21 studies), and neuronal connectivity alterations (6 out of 21 studies) were the most predominant explanations. The presence of these 3 theories in the studies that mention all sports aligns with the overall systematic review identification of these 3 theories as the most common explanations for sport-related concussion pathophysiology.

Concussion RTP guidelines

Finally, we identified that RTP guidelines between sports may share common ground. Despite the differences in concussion by the sports listed above, there may be a clear commonality among theories regarding concussions across all sports. Through collection of concussion RTP guidelines in sports at the professional and collegiate levels (see Table 9), the authors of this study identified different concussion protocols for individual sports. Despite the different number of steps between protocols, a common pattern of rest until concussion symptoms disappear, exert and exercise, and return to full contact play exists among different concussion RTP guidelines. This brings up a major question of whether concussion RTP guidelines should be standardized or adjusted to better manage athlete symptoms and recovery. The mechanism of trauma to the head in sports can differ; however, across all sports, signs and symptoms of concussion remain shared. A further analysis of concussion pathophysiology, prevalence, relapse of symptoms, and long-term consequences across different sports may shed light on this question.

By unifying existing theories on sport-related concussion pathophysiology, summarizing risk factors/comorbidities associated with sport-related concussion, and identifying sport-specific concussion findings, the authors of this review article

simplify sport-related concussion into a more concise and approachable subject. By investigating the consequences of concussion, diagnosis of concussion, and sport RTP guidelines, the authors shed light on the importance of future studies that can help connect concussion management with the concussion mechanism and pathophysiology. Using the simplified information collected in this review article, future studies can further unravel pathophysiology and ideal management of sport-related concussion. As the literature on sport-related concussion becomes clearer, player- and sport-specific guidelines can also be instituted for better management of concussion in sport.

Conclusions

The continued study of concussion pathophysiology across all sports and within individual sports at all levels of play is important and should be continued. The output of new theories and knowledge, the growing medical concern of increasing concussions, the creation of new evaluation rules and protocols, and the overall uncertainty that still exists regarding the topic make this a subject that deserves attention and a focused effort toward improvement. Further investigation, using the information provided in our simplifying literature review of sport-related concussion, will hopefully help shape concussion management and sports in general.

Acknowledgements

The authors would like to give special thanks to Anthony N Pass, Assistant Athletic Director for Sports Medicine and Head Football Athletic Trainer, The University of Texas at Austin and John S Kuo, MD, PhD Chairman Department of Neurosurgery, Dell Medical School, The University of Texas at Austin for their critical review of the article and editorial comments.

Author Contributions

MTK- conception, design, review, and supervision. PS- Acquisition of data, drafting article, critically revising article, drafting revision. All authors- Critically revising article.

ORCID iD

Michael T Koltz  <https://orcid.org/0000-0002-0404-0339>

REFERENCES

1. Facts about concussion and brain injury. *Centers for Disease Control and Prevention*. https://www.cdc.gov/traumaticbraininjury/pdf/Fact_Sheet_ConcussTBI-a.pdf. Published 2017. Accessed June 10, 2018.
2. Manley G, Gardner A, Schneider K, et al. A systematic review of potential long-term effects of sport-related concussion. *Br J Sports Med*. 2017;51:969–977. doi:10.1136/bjsports-2017-097791.
3. Zhang AL, Sing DC, Rugg CM, Feeley BT, Senter C. The rise of concussion in the adolescent population [published online ahead of print August 16, 2016]. *Orthop J Sports Med*. doi:10.1177/2325967116662458.
4. Wenner L. *Media, Sports & Society*. Newbury Park, CA: SAGE; 1991.
5. Workewych AM, Ciuffetelli Muzzi M, Jing R, Zhang S, Topolovec-Vranic J, Cusimano MD; on behalf of the Canadian Brain Injury and Violence Research Team. Twitter and traumatic brain injury: a content and sentiment analysis of tweets pertaining to sport-related brain injury [published online ahead of print August 25, 2017]. *SAGE Open Medicine*. doi:10.1177/2050312117720057.

6. Sullivan SJ, Schneiders AG, Cheang CW, et al. "What's happening?" a content analysis of concussion-related traffic on Twitter. *Br J Sports Med.* 2012;46:258–263. doi:10.1136/bjsm.2010.080341.
7. Sharp DJ, Jenkins PO. Concussion is confusing us all. *Pract Neurol.* 2015;15:172–186. doi:10.1136/practneurol-2015-001087.
8. McCrory P, Meeuwisse WH, Aubry M, et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Br J Sports Med.* 2013;47:250–258. doi:10.1136/bjsports-2013-092313.
9. Mez J, Daneshvar DH, Kiernan PT, et al. Clinicopathological evaluation of chronic traumatic encephalopathy in players of American Football. *JAMA.* 2017;318:360–370. doi:10.1011/jama.2017.8334.
10. Slobounov SM, Walter A, Breiter HC, et al. The effect of repetitive subconcussive collisions on brain integrity in collegiate football players over a single football season: a multi-modal neuroimaging study. *Neuroimage Clin.* 2017;14:708–718. doi:10.1016/j.nicl.2017.03.006
11. Coughlin JM, Wang Y, Minn I, et al. Imaging of glial cell activation and white matter integrity in brains of active and recently retired National Football League Players. *JAMA Neurol.* 2017;74:67–74. doi:10.1001/jamaneurol.2016.3764.
12. Cherry JD, Tripodis Y, Alvarez VE, et al. Microglial neuroinflammation contributes to tau accumulation in chronic traumatic encephalopathy. *Acta Neuropathol Commun.* 2016;4:112. doi:10.1186/s40478-016-0382-8.
13. Bahrami N, Sharma D, Rosenthal S, et al. Subconcussive head impact exposure and white matter tract changes over a single season of youth football. *Radiology.* 2016;281:919–926. doi:10.1148/radiol.2016160564.
14. Strain JF, Didehban N, Spence J, et al. White matter changes and confrontation naming in retired aging national football league athletes. *J Neurotrauma.* 2017;34:372–379. doi:10.1089/neu.2016.4446.
15. Amen DG, Willeumier K, Omalu B, Newberg A, Raghavendra C, Raji CA. Perfusion neuroimaging abnormalities alone distinguish National Football League Players from a healthy population. *JAD.* 2016;53:237–241. doi:10.3233/JAD-160207.
16. Meier TB, Savitz J, Singh R, Teague TK, Bellgowan PSF. Smaller dentate gyrus and CA2 and CA3 volumes are associated with kynurenine metabolites in collegiate football athletes. *J Neurotrauma.* 2016;33:1349–1357. doi:10.1089/neu.2015.4118.
17. Koerte IK, Hufschmidt J, Muehlmann M, et al. Cavum septi pellucidi in symptomatic former professional football players. *J Neurotrauma.* 2016;33:346–353. doi:10.1089/neu.2015.3880.
18. Wang Y, Nelson LD, LaRoche AA, et al. Cerebral blood flow alterations in acute sport-related concussion. *J Neurotrauma.* 2016;33:1227–1236. doi:10.1089/neu.2015.4072.
19. Meier TB, Bellgowan PSF, Bergamino M, Ling JM, Mayer AR. Thinner cortex in collegiate football players with, but not without, a self-reported history of concussion. *J Neurotrauma.* 2016;33:330–338. doi:10.1089/neu.2015.3919.
20. Gardner RC, Hess CP, Brus-Ramer M, et al. Cavum septum pellucidum in retired American pro-football players. *J Neurotrauma.* 2016;33:157–161. doi:10.1089/neu.2014.3805.
21. Poole VN, Breedlove EL, Shenk TE, et al. Sub-concussive hit characteristics predict deviant brain metabolism in football athletes. *Develop Neuropsychol.* 2015;40:12–17. doi:10.1080/87565641.2014.984810
22. Davenport EM, Whitlow CT, Urban JE, et al. Abnormal white matter integrity related to head impact exposure in a season of high school varsity football. *J Neurotrauma.* 2014;31:1617–1624. doi:10.1089/neu.2013.3233.
23. Strain J, Didehban N, Cullum CM, et al. Depressive symptoms and white matter dysfunction in retired NFL players with concussion history. *Neurology.* 2013;81:25–32. doi:10.1212/WNL.0b013e318299ccf8.
24. Shahim P, Tegner Y, Marklund N, Blennow K, Zetterberg H. Neurofilament light and tau as blood biomarkers for sports-related concussion. *Neurology.* 2018;90:e1780–e1788. doi:10.1212/WNL.0000000000005518.
25. Manning KY, Schranz A, Bartha R, et al. Multiparametric MRI changes persist beyond recovery in concussed adolescent hockey players. *Neurology.* 2017;89:2157–2166. doi:10.1212/WNL.0000000000004669.
26. Shahim P, Zetterberg H, Tegner Y, Blennow K. Serum neurofilament light as a biomarker for mild traumatic brain injury in contact sports. *Neurology.* 2017;88:1788–1794. doi:10.1212/WNL.0000000000003912.
27. Wright AD, Jarrett M, Vavasour I, et al. Myelin water fraction is transiently reduced after a single mild traumatic brain injury—a prospective cohort study in collegiate hockey players. *PLoS ONE.* 2016;11:e0150215. doi:10.1371/journal.pone.0150215.
28. Yengo-Kahn AM, Zuckerman SL, Stotts J, et al. Performance following a first professional concussion among National Basketball Association players. *Phys Sportsmed.* 2016;44:297–303.
29. Ling H, Morris HR, Neal JW, et al. Mixed pathologies including chronic traumatic encephalopathy account for dementia in retired association football (soccer) players. *Acta Neuropathol.* 2017;133:337–352. doi:10.1007/s00401-017-1680-3.
30. Di Virgilio TG, Hunter A, Wilson L, et al. Evidence for acute electrophysiological and cognitive changes following routine soccer heading. *EBioMedicine.* 2016;13:66–71. doi:10.1016/j.ebiom.2016.10.029.
31. Grinberg LT, Anghinah R, Nascimento CF, et al. Chronic traumatic encephalopathy presenting as Alzheimer's dementia in a retired soccer player. *JAD.* 2016;54:169–174. doi:10.3233/JAD-160312.
32. Hwang S, Ma L, Kawata K, Tierney R, Jeka JJ. Vestibular dysfunction after subconcussive head impact. *J Neurotrauma.* 2017;34:8–15. doi:10.1089/neu.2015.4238.
33. Lipton ML, Kim N, Zimmerman ME, et al. Soccer heading is associated with white matter microstructural and cognitive abnormalities. *Radiology.* 2013;268:850–857. doi:10.1148/radiol.13130545.
34. Zimmer A, Piccora K, Schuster D, Webbe F. Sport and team differences on baseline measures of sport-related concussion. *J Athlet Train.* 2013;48:659–667. doi:10.4085/1062-6050-48.5.06.
35. Bernick C, Banks SJ, Shin W, et al. Repeated head trauma is associated with smaller thalamic volumes and slower processing speed: the Professional Fighters' Brain Health Study. *Br J Sports Med.* 2015;49:1007–1011. doi:10.1136/bjsports-2014-093877.
36. Johnson B, Neuberger T, Gay M, Hallett M, Slobounov S. Effects of subconcussive head trauma on the default mode network of the brain. *J Neurotrauma.* 2014;31:1907–1913. doi:10.1089/neu.2014.3415.
37. Neselius S, Brisby H, Marcusson J, Zetterberg H, Blennow K, Karlsson T. Neurological assessment and its relationship to CSF biomarkers in amateur boxers. *PLoS ONE.* 2014;9:e99870. doi:10.1371/journal.pone.0099870.
38. Neselius S, Zetterberg H, Blennow K, Marcusson J, Brisby H. Increased CSF levels of phosphorylated neurofilament heavy protein following bout in amateur boxers. *PLoS ONE.* 2013;8:e81249. doi:10.1371/journal.pone.0081249.
39. Di Battista AP, Churchill N, Schweizer TA, et al. Blood biomarkers are associated with brain function and blood flow following sports concussion. *J Neuroimmunol.* 2018;319:1–8. doi: 10.1016/j.jneuroim.2018.03.002
40. Churchill NW, Hutchison MG, Di Battista AP, Graham SJ, Schweizer TA. Structural, functional, and metabolic brain markers differentiate collision versus contact and non-contact athletes. *Front Neurol.* 2017;8:390. doi:10.3389/fneur.2017.00390.
41. Gill J, Merchant-Borna K, Jeromin A, Livingston W, Bazarian J. Acute plasma tau relates to prolonged return to play after concussion. *Neurology.* 2017;88:595–602. doi:10.1212/WNL.0000000000003587.
42. Reches A, Kutcher J, Elbin RJ, et al. Preliminary investigation of brain network activation (BNA) and its clinical utility in sport-related concussion. *Brain Injury.* 2017;31:237–246. doi:10.1080/02699052.2016.1231343.
43. Safinia C, Bershady EM, Clark HB, et al. Chronic traumatic encephalopathy in athletes involved with high-impact sports. *J Vasc Intervention Neurol.* 2016;9:34–48.
44. Papa L. Potential blood-based biomarkers for concussion. *Sports Med Arthroscopy Rev.* 2016;24:108–115. doi:10.1097/JSA.0000000000000117.
45. Babcock L, Zhang N, Leach J, Wade SL. Are UCH-L1 and GFAP promising biomarkers for children with mild traumatic brain injury? *Brain Injury.* 2016;30:1231–1238. doi:10.1080/02699052.2016.1178396.
46. Herrera JJ, Bockhorst K, Kondraganti S, Stertz L, Quevedo J, Narayana PA. Acute white matter tract damage after frontal mild traumatic brain injury. *J Neurotrauma.* 2017;34:291–299. doi:10.1089/neu.2016.4407.
47. Merchant-Borna K, Lee H, Wang D, et al. Genome-wide changes in peripheral gene expression following sports-related concussion. *J Neurotrauma.* 2016;33:1576–1585. doi:10.1089/neu.2015.4191.
48. Kontos AP, Reches A, Elbin RJ, et al. Preliminary evidence of reduced brain network activation in patients with post-traumatic migraine following concussion. *Brain Imag Behav.* 2016;10:594–603. doi:10.1007/s11682-015-9412-6.
49. Militana AR, Donahue MJ, Sills AK, et al. Alterations in default-mode network connectivity may be influenced by cerebrovascular changes within one week of sports related concussion in college varsity athletes: a pilot study. *Brain Imag Behav.* 2016;10:559–568. doi:10.1007/s11682-015-9407-3.
50. Goswami R, Dufort P, Tartaglia MC, et al. Frontotemporal correlates of impulsivity and machine learning in retired professional athletes with a history of multiple concussions. *Brain Struct Funct.* 2016;221:1911–1925. doi:10.1007/s00429-015-1012-0.
51. Pham N, Akonasu H, Shishkin R, Taghibiglou C. Plasma soluble prion protein, a potential biomarker for sport-related concussions: a pilot study. *PLoS ONE.* 2015;10:e0117286. doi:10.1371/journal.pone.0117286.
52. Schulte S, Podlog LW, Hamson-Utley JJ, Strathmann FG, Strüder HK. A systematic review of the biomarker S100B: implications for sport-related concussion management. *J Athlet Train.* 2014;49:830–850. doi:10.4085/1062-6050-49.3.33.
53. Tremblay S, Henry LC, Bedetti C, et al. Diffuse white matter tract abnormalities in clinically normal ageing retired athletes with a history of sports-related concussions. *Brain.* 2014;137:2997–3011. doi:10.1093/brain/awu236.
54. Czerniak SM, Sikoglu EM, Navarro AAL, et al. A resting state functional magnetic resonance imaging study of concussion in collegiate athletes. *Brain Imag Behav.* 2015;9:323–332. doi:10.1007/s11682-014-9312-1.

55. Murugavel M, Cubon V, Putukian M, et al. A longitudinal diffusion tensor imaging study assessing white matter fiber tracts after sports-related concussion. *J Neurotrauma*. 2014;31:1860–1871. doi:10.1089/neu.2014.3368.
56. Bazarian JJ, Zhu T, Zhong J, et al. Persistent, long-term cerebral white matter changes after sports-related repetitive head impacts. *PLoS ONE*. 2014;9:e94734. doi:10.1371/journal.pone.0094734.
57. Ji S, Zhao W, Ford JC, et al. Group-wise evaluation and comparison of white matter fiber strain and maximum principal strain in sports-related concussion. *J Neurotrauma*. 2015;32:441–454. doi:10.1089/neu.2013.3268.
58. Kiechle K, Bazarian JJ, Merchant-Borna K, et al. Subject-specific increases in serum S-100B distinguish sports-related concussion from sports-related exertion. *PLoS ONE*. 2014;9:e84977. doi:10.1371/journal.pone.0084977.
59. De Beaumont L, Tremblay S, Henry LC, Poirier J, Lassonde M, Théoret H. Motor system alterations in retired former athletes: the role of aging and concussion history. *BMC Neurol*. 2013;13:109. doi:10.1186/1471-2377-13-109.
60. Dettwiler A, Murugavel M, Putukian M, Cubon V, Furtado J, Osherson D. Persistent differences in patterns of brain activation after sports-related concussion: a longitudinal functional magnetic resonance imaging study. *J Neurotrauma*. 2014;31:180–188. doi:10.1089/neu.2013.2983.
61. NFL head, neck and spine committee's concussion diagnosis management protocol. <https://www.playsmartplaysafe.com/focus-on-safety/protecting-players/nfl-head-neck-spine-committees-protocols-regarding-diagnosis-management-concussion/>. Published 2017. Accessed June 10, 2018.
62. Concussion evaluation and management protocol. <https://nhl.bamcontent.com/images/assets/binary/282574512/binary-file/file.pdf>. Published 2016. Accessed June 10, 2018.
63. U.S. soccer concussion guidelines. <https://www.ussoccer.com/about/recognize-to-recover/concussion-guidelines>. Published 2016. Accessed June 10, 2018.
64. Explaining the NBA's concussion protocol. <https://www.nbcsports.com/boston/boston-celtics/with-al-horford-out-nba-concussion-protocol-explained>. Published 2016. Accessed June 10, 2018.
65. Analyzing MLB's concussion policy. <http://www.espn.com/fantasy/baseball/fib/story?id=6280590>. Published 2011. Accessed June 10, 2018.
66. Concussion diagnosis and management best practices. <http://www.ncaa.org/sport-science-institute/concussion-diagnosis-and-management-best-practices>. Published 2015. Accessed June 10, 2018.
67. Daneshvar DH, Nowinski CJ, McKee A, Cantu RC. The epidemiology of sport-related concussion. *Clin Sports Med*. 2011;30:1–17. doi:10.1016/j.csm.2010.08.006.
68. Davis GA, Castellani RJ, McCrory P. Neurodegeneration and sport. *Neurosurgery*. 2015; 76:743–655. doi:10.1227/NEU.0000000000000722.
69. Iverson GL, Keene CD, Perry G, Castellani RJ. The need to separate chronic traumatic encephalopathy neuropathology from clinical features. *JAD*. 2017;61:17–28. doi:10.3233/JAD-170654.
70. Guth T, Ketcham CJ, Hall EE. Influence of concussion history and genetics on event-related potentials in athletes: potential use in concussion management. *Sports*. 2018;6:E5. doi:10.3390/sports6010005.
71. Scopaz KA, Hatzenbuehler JR. Risk modifiers for concussion and prolonged recovery. *Sports Health*. 2013;5:537–541. doi:10.1177/1941738112473059.
72. Slobounov SM, Wayne SJ. *Foundations of Sport-Related Brain Injuries*. New York, NY: Springer; 2006.
73. Joseph JR, Swallow J, Willsey K, et al. Elevated markers of brain injury as a result of clinically asymptomatic high-acceleration head impacts in high-school football athletes [published online ahead of print July 3, 2018]. *J Neurosurg*. doi:10.3171/2017.12.JNS172386.
74. Kornguth S, Rutledge N, Perla G, Bray J, Hardin A. A proposed mechanism for development of CTE following concussive events: head impact, water hammer injury, neurofilament release, and autoimmune processes. *Brain Sci*. 2017;7:164. doi:10.3390/brainsci7120164.
75. Carman AJ, Ferguson R, Cantu R, et al. Expert consensus document: mind the gaps—advancing research into short-term and long-term neuropsychological outcomes of youth sport-related concussion. *Nat Rev Neurol*. 2015;11:230–244.
76. Osier N, Dixon CE. Mini review of controlled cortical impact: a well-suited device for concussion research. *Brain Sci*. 2017;7:88. doi:10.3390/brainsci7070088.