



Published in final edited form as:

Mucosal Immunol. 2019 July ; 12(4): 930–944. doi:10.1038/s41385-019-0170-4.

Alcohol induced IL-17A production in Paneth cells amplifies endoplasmic reticulum stress, apoptosis and inflammasome-IL-18 activation in the proximal small intestine in mice

B. Gyongyosi^X, Y. Cho^X, P. Lowe, C.D. Calenda, A. Iracheta-Vellve, A. Satishchandran, A. Ambade, and G. Szabo

University of Massachusetts Medical School, Department of Medicine, Worcester, MA 01605, USA

Abstract

Gut microbial translocation contributes to alcoholic hepatitis. Using a mouse model of alcoholic hepatitis, we investigated the effects of chronic alcohol plus binge and found increased abundance of Paneth cells and IL-17A in the proximal small intestine (PSI). Alcohol increased IL-17A production and pro-apoptotic signaling evidenced by Bax, Bim, caspase-3 and -8 increases via endoplasmic reticulum (ER) stress indicated by C/EBP homologous protein (CHOP) upregulation; this was prevented by the ER stress inhibitor, 4-PBA, in isolated crypts *in vitro* and *in vivo*. Mechanistically, IL-17 augmented alcohol-induced ER stress in isolated crypts. *In vivo* IL-17A blocking antibody administration in alcohol-treated mice attenuated ER stress-mediated apoptosis and IL-18 induction and prevented alcohol-induced impairment of tight junctions in the PSI and LPS translocation to the liver. Acute-on-chronic alcohol resulted in inflammasome activation, caspase-1 cleavage and IL-18 production in the PSI. *In vivo* treatment with antibiotics or 4-PBA prevented CHOP upregulation and inflammasome activation. Our data suggest that alcohol upregulates innate immune mechanisms by increasing Paneth cell numbers and IL-17A release contributing to apoptosis amplification, inflammasome activation and gut leakiness in the PSI. Binge alcohol-induced Paneth cell expansion, ER stress and inflammasome activation in the PSI are modulated by the gut microbiome.

Keywords

alcoholic liver disease; ER stress; inflammasome; Paneth cell; IL-17; unfolded protein response; IL-18; apoptosis; ZO-1

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence: Gyongyi Szabo, MD, PhD, Professor, Department of Medicine, LRB208, University of Massachusetts Medical School, 364 Plantation Street, Worcester, MA 01605, Phone No: 00-1-508-856-6113, Fax No: 00-1-508-856-4770, gyongyi.szabo@umassmed.edu.

^XThese authors contributed equally.

Author contributions

BG, YC and GS conceived and designed the experiments; BG, PL, YC, CDC performed the experiments with assistance from AIV, AS, and AA; BG, YC, PL, CDC, AIV and GS analyzed the data; BG, PL and GS wrote the paper; PL, AIV, AS and GS obtained funding and GS supervised the project.

Competing Interests

The authors declare no competing financial interests.

Introduction

Alcohol-induced intestinal barrier dysfunction contributes to the development of alcoholic liver disease and alcoholic hepatitis.¹ Previous studies have shown that chronic alcohol consumption changes the composition of the gut microbiome, results in inflammatory cytokine expression in the small bowel and disrupts the epithelial tight junction barrier that collectively lead to increased circulating levels of pathogen-associated molecular patterns (PAMPs) including bacterial lipopolysaccharide (LPS).^{2,3} It has been shown that PAMPs and alcohol-induced damage-associated molecules (DAMPs) contribute to tissue inflammation in the liver and other organs.⁴

Paneth cells (PCs), immune-like epithelial cells of the crypts of Lieberkühn, are increasingly recognized as important regulators of intestinal homeostasis that shape gut microbial composition and restrain bacterial translocation.^{5,6} PCs produce antimicrobial peptides (lysozyme, cryptdins, phospholipase A [PLA2g2a]) and inflammatory cytokines that are readily induced and secreted in response to various microbial or inflammatory stimuli. Moreover, PC-derived interleukin (IL)-17 is an important regulator of the tight junction,⁷ and has been implicated in the pathogenesis of various organ injuries.⁸ PCs have a highly secretory phenotype with active endoplasmic reticulum (ER) function that renders them susceptible to perturbations in ER function,⁹ as occurs in other organs after chronic alcohol consumption.¹⁰

ER stress and activation of the unfolded protein response (UPR) have been identified both as physiologic and pathogenic factors in the healthy and diseased intestine. Accumulation of nascent or misfolded proteins in the ER leads to dissociation of the ER chaperone, Bip, from its binding partners, inositol-requiring enzyme 1 (IRE1), activating transcription factor 6 (ATF6), and PKR-like ER kinase (PERK), allowing them to initiate adaptive, pro-apoptotic, and pro-inflammatory signaling cascades.¹¹ When ER stress is prolonged or cannot be resolved, the UPR signaling switches from pro-survival to pro-apoptotic, activating C/EBP homologous protein (CHOP) and growth arrest and DNA damage-inducible gene 34 (GADD34) and transmitting proapoptotic signals from the ER to the mitochondria through pore-forming small proteins such as Bax, Bim, and Bid.¹² Recently, ER stress has also been directly and indirectly associated with inflammatory signaling. Of particular interest, ER stress induced caspase-2 activation and mitochondrial damage have been shown to lead to the activation of the inflammasome,¹³ a multi-protein complex that acts through caspase-1, which cleaves and activates pro-IL-1 β and pro-IL-18.¹⁴

Alcohol was found to induce ER stress in different tissues,¹⁰ however the role of UPR signaling and/or the contribution of PCs is yet to be evaluated in alcohol-induced changes in the intestinal mucosa. Here, we show that alcohol consumption results in intestinal, pro-apoptotic and pro-inflammatory signaling, and that PCs are susceptible to alcohol-related changes in the SI.

Results

Alcohol induces cell death and inflammation in the proximal small intestine and leads to bacterial product translocation

Alcohol consumption leads to many physiologic changes throughout the body, beginning in the gastrointestinal tract. Using a model of chronic alcohol plus binge feeding in mice that results in features of early alcoholic hepatitis,¹⁵ we initially measured bacterial translocation from the intestine and found evidence of bacterial product translocation within the liver. Both bacterial 16S ribosomal DNA (rDNA) and the bacterial wall component, lipopolysaccharide (LPS), were increased in the liver of alcohol-fed mice compared to pair-fed animals (Fig. 1A and B). Since the liver is the first organ encountered by the portal circulation after the intestine, we sought to investigate the effects of alcohol on the small intestine. In the proximal small intestine (PSI), we observed that compared to pair-fed mice, alcohol-fed mice had significantly more cell death detected by measuring the extent of Terminal deoxynucleotidyl transferase dUTP Nick-End Labeling (TUNEL) (Fig. 1C and D). Additionally, we measured the expression of multiple PSI proinflammatory cytokines including *Tnfa*, *Il6*, *Il1 β* , *Mcp1*, *Il23* and *Il17a* and found that alcohol increased the mRNA expression of each of these cytokines (Fig. 1E). Together, these data suggest that alcohol induces significant changes in the PSI, including cell death and proinflammatory signaling, and these changes correlate with translocation of bacterial products from the intestinal lumen to the liver.

Alcohol increases the frequency of Paneth cells in the PSI and results in their degranulation

Paneth cells (PCs) are localized in the intestinal crypts of Lieberkühn and show increased abundance from the proximal to the distal segment of the SI in healthy intestines.¹⁶ We next investigated whether the distribution of PCs changed upon alcohol feeding in mice. We first measured PC frequency using PAS-staining (Fig. 2A) and immunohistochemistry for the PC marker, lysozyme, (Fig. 2B) in SI sections. We found a significant increase in the frequency of PCs in the PSI of alcohol-fed mice compared to control pair-fed mice (Fig. 2C and D). Remarkably, the alcohol-induced increase in PC frequency was restricted to the PSI and alcohol feeding did not change PC numbers in the distal SI (DSI) compared to controls. We also found that alcohol feeding resulted in crypt degranulation, indicated by PAS-positive material in the lumen of crypts, suggesting that alcohol exposure promotes release of antimicrobial substances from the PCs (Fig. 2A inserts and E).

PCs play a role in anti-microbial defense and in intestinal self-renewal.¹⁷ Thus, we tested expression of genes implicated in PC differentiation and function and found increased mRNA expression of differentiation markers, *Sox9* and *Igfbp4*,¹⁸ *Spdef* and *Math1*,¹⁹ and *Lgr5*, while the inhibitory *Klf4*²⁰ was decreased in the PSI of alcohol-fed mice compared to pair-fed mice (Fig. 2F). Although these genes could be expressed by multiple cell types, these findings are consistent with the increased numbers of PCs in the PSI after alcohol feeding. The expression of antimicrobial peptide genes (*Cryptdins*; defensins *Defcr1*, *Defcr4* and *Defa-rs1*; *PLA2g2a*; lysozymes *Lyz1* and *Lyz2*; and *Muc2*) was also significantly higher in the PSI of alcohol-fed compared to control diet-fed mice (Fig. 2G). Indeed, an acute binge

increased *Cryptdins* expression after both 4h and 9h, while mice treated with chronic alcohol and no binge did not show an increase in expression (Suppl. Fig. 2).

Alcohol induces IL-17 in the PSI and in isolated small intestinal crypts

Recent studies support that IL-17 contributes to gut homeostasis,²¹ however the role of IL-17 in alcohol-induced gut permeability is unknown. Immunoblot and immunohistochemistry analyses revealed increased IL-17A levels in the PSI of alcohol-compared to pair-fed mice (Fig. 3A–D). The IL-17A staining was localized to the bottom of the crypts identical to the localization of PCs (Fig. 3A). However, IL-17 could also be released by a subset of T cells, the T helper (Th)-17 cells. Therefore, we isolated resident intestinal immune cells of the PSI lamina propria and quantified Th17 cells by flow cytometry. We found no differences in the number of Th17 cells between alcohol- and pair-fed mice (Fig. 3E). These results suggest that PCs are likely the source of alcohol-induced intestinal IL-17A increase.

Having observed PC expansion and degranulation in alcohol-fed mice, we aimed to explore mechanisms by which ethanol might regulate PC function. Therefore, we isolated crypts that are enriched in PCs from wild-type mice and exposed them to different doses of ethanol *ex vivo*. Ethanol exposure of these isolated PSI crypts resulted in a significant increase in IL-17A release that correlated with the *in vivo* observation of increased IL-17A in PCs in the SI in alcohol-fed mice. The *ex vivo* ethanol-induced IL-17A release was time- and dose-dependent, beginning at 2 hours and peaking at 6 hours (Fig. 3F). The IL-17A increase detected in the supernatants was due to secretion from viable crypts and not a consequence of ethanol-induced cell death because there was no increase in lactate dehydrogenase (LDH) cytotoxicity assay in EtOH-treated crypts compared to controls (Suppl. Fig. 1)

Alcohol-induced IL-17A release is mediated by ER stress

PCs are highly secretory and, consequently, susceptible to ER stress.⁹ We therefore hypothesized that alcohol-induced ER stress might be involved in IL-17 release. To test this hypothesis, we measured the effect of 4-PBA (a FDA-approved ER stress inhibitor) on ethanol-induced IL17A release.²² We found that administration of 4-PBA prevented alcohol-induced IL17A release, in a dose-dependent manner (Fig. 4A). Conversely, IL-17A release was induced by treatment with the ER stress inducer, thapsigargin, in a dose-dependent manner with increasing IL-17A detected in crypt supernatants following 0.3, 1 and 3 μ M thapsigargin (Fig. 4B). These results indicate that alcohol induces IL-17A release in SI crypts via an ER stress-dependent mechanism.

Next, we sought to investigate the role of IL-17A on cell signaling pathways in the SI. SI crypts treated with recombinant IL-17A *ex vivo* had increased cleavage of PARP that was not affected further by alcohol exposure (Fig. 4C and D). Administration of an IL-17A blocking antibody *in vivo* abrogated PARP cleavage following acute-on-chronic alcohol exposure. Together, these data suggest an IL-17-mediated process of PSI apoptosis. Interestingly, *in vivo* administration of the IL-17A blocking antibody also showed a trend of reduction in the cleavage and activation of caspase-8, while control antibody had no effect (Fig. 4E and F).

Acute-on-chronic alcohol administration induces pro-apoptotic signaling and ER stress in the PSI

ER stress and activation of the unfolded protein response (UPR) are key features of secretory cells, such as Paneth cells.²³ Given the important connections between IL-17A and ER stress and that prolonged ER stress initiates cell death signaling by relaying stress signals from the ER to the mitochondria, we next investigated protein levels of the pro-apoptotic molecules, Bax, Bim and Bid. These proteins were reported to be central in ER stress-induced apoptosis pathways acting downstream of CHOP signaling.¹² We found that 10-day alcohol feeding alone (10d EtOH no binge) did not significantly increase expression of the pro-apoptotic molecules Bax, Bim and Bid (Fig. 5A and B). However, the acute binge on chronic alcohol diet significantly upregulated the protein levels of Bax, Bim and Bid in the PSI both 4 and 9 hours after the binge (Fig. 5A and B). These molecules are responsible for mitochondrial outer membrane permeabilization leading to downstream caspase activation. In line with this, we found increased cleavage of PARP (Fig. 5A and B), a well-characterized marker of caspase activation.

ER stress is compensated by activation of the unfolded protein response, a homeostatic signaling mechanism that orchestrates recovery of ER function and upon failure to adapt to ER stress, results in apoptosis. We found a significant and selective upregulation of p-eIF2 α , a marker of the PERK arm of ER stress, and CHOP protein expression in the PSI of mice after 10 days of alcohol feeding compared to controls (Fig. 5A and B). Acute binge in the chronic alcohol-fed mice resulted in a further significant increase in CHOP, phosphorylation of eIF2 α and also increased GADD34 (a downstream target of CHOP) levels at 4 and 9 hours after the acute binge (Fig. 5A and B). Interestingly, these alcohol-induced changes were not observed in the DSI (Suppl. Fig. 3). These data suggest that chronic alcohol induces CHOP, and an additional acute alcohol binge augments CHOP expression and downstream pro-apoptotic events, including translational block (phosphorylation of eIF2 α) and GADD34 induction in the PSI.

Alcohol increases caspase expression and activation through ER stress

Next, we measured the expression and activation of caspases involved in apoptosis pathways. We detected cleavage of the executioner caspase, caspase-3, indicative of ongoing apoptotic signaling at 4 and 9 hours after binge in chronically alcohol-fed mice (Fig. 5C and D). We also found increased levels of total and cleaved Bid (tBid) at 9 hours after the acute alcohol binge (Fig. 5A and B). Bid is a known caspase-8 substrate in the extrinsic apoptotic pathway, and has been described as a mediator of the caspase-2 induced inflammasome activation upon ER stress.¹³ Therefore, we measured these two caspases and found a significant induction of both caspase-8 and caspase-2 protein levels as early as 4 hours after the acute binge (Fig. 5C and D), suggesting that acute-on-chronic alcohol activates both extrinsic and intrinsic proapoptotic mechanisms in the PSI.

In order to determine the role of ER stress in small intestinal caspase activation, we treated mice *in vivo* with the ER stress inhibitor, 4-PBA. Acute-on-chronic alcohol administration led to the induction of CHOP expression (Fig. 5A and B) as well as caspase-8 (Fig. 5C and D). 4-PBA administration inhibited ER stress, determined by decreased CHOP expression

(Fig. 5E and F). Interestingly, 4-PBA also abrogated the alcohol-induced increase in activation of caspase 8 (Fig. 5E and F), indicating that alcohol-induced caspase activation is due to the induction of ER stress.

Alcohol-mediated increase of CHOP is localized to Paneth cells in PSI crypts

To characterize the cell type most affected by alcohol, we next isolated epithelial cell (containing PCs) and immune cell fractions (containing intraepithelial and lamina propria leukocytes) from the PSI. Western blot analysis revealed a significant upregulation of both the ER stress proteins CHOP and Bax in the epithelial cell fraction (Fig. 6A and B, upper panels), but not in the intraepithelial (IEL) or lamina propria (LPL) immune cell fractions (Fig. 6A and B, lower panels), suggesting that alcohol primarily affects epithelial cells. Immunofluorescent (IF) staining revealed increased CHOP fluorescent intensity in PCs of the crypts in the PSI of alcohol-fed mice compared to controls (Fig. 6C and D). Finally, we isolated crypts from the PSI of pair- or alcohol-fed mice in order to differentiate the protein expression localized in this particular epithelial cell layer as the epithelial cell isolation includes both crypts (containing the PCs) and villus epithelial cells. In the isolated crypts, we again detected an increase in ER stress proteins Bax and Bim in alcohol-fed compared with pair-fed mice (Fig. 6E and F). These results identified that *in vivo* alcohol increases CHOP expression in PCs in the crypts of the PSI.

Alcohol-reduced intestinal barrier integrity is restored by anti-IL-17 treatment *in vivo*.

To assess the role of IL-17 in alcohol-induced intestinal barrier dysfunction, we used an IL-17 blocking antibody *in vivo* and found that IL-17 blocking prevented the alcohol-induced increase in liver LPS levels (Fig. 7A), suggesting a connection between IL-17 and the integrity of the intestinal barrier. Therefore, we next evaluated tight junction protein expression in the PSI. Although changes in barrier integrity are likely to occur as a continuum across time of exposure to alcohol, our acute-on-chronic alcohol feeding model dramatically reduced the immunofluorescence of zonula occludens-1 (ZO-1), an important component of the PSI tight junction structure (Fig. 7B). Treatment with IL-17 blocking antibody recovered expression of ZO-1, while ZO-1 staining was unaffected by the isotype antibody (Fig. 7B). Anti-IL-17 also prevented the alcohol-induced decrease in occludin dimer protein expression in the PSI (Fig. 7C and D). We found that IL-17 blocking resulted in a trend of upregulation in the mRNA expression of *claudin-1*, *-5* and *-15* as well as *occludin* and *ZO-1* (Fig. 7E). Interestingly, the acute-on-chronic alcohol showed no reduction at the mRNA levels of these tight junction proteins that is likely due to the binge alcohol effect.²

Inflammasome activation by acute alcohol binge in the small intestine is attenuated by *in vivo* inhibition of ER stress

ER stress is known to induce inflammation and PCs are important for the initiation of intestinal inflammation.²⁴ Therefore, we investigated activation of the central sensor and orchestrator of inflammatory signaling, the inflammasome, in the acute-on-chronic alcohol feeding. We found activation of caspase-1 was significantly increased in the PSI of acute-on-chronic mice compared to PF and chronic alcohol alone (Fig. 8A and B). Inflammasome activation was indicated by cleavage of the inflammasome effector, caspase-1 (p10), and its

downstream target, IL-18 (18 kDa cleaved form) that were increased in mice after acute binge alcohol (Fig. 8A and B).

To investigate whether ER stress could play a role in inflammasome activation, we administered the ER stress inhibitor, 4-PBA, in the liquid diet during alcohol feeding. We found that alcohol-related CHOP expression was attenuated in 4-PBA-treated mice indicating effective ER stress inhibition (Fig. 8C and D). Importantly, *in vivo* administration of 4-PBA in alcohol-fed mice resulted in decreased IL-18 cleavage indicating reduced inflammasome activation (Fig. 8C and D). Additionally, 4-PBA treatment significantly reduced the cleavage of PARP (Fig. 8E and F). These data suggested that alcohol-induced ER stress contributes to inflammasome activation in the PSI. Because we had previously observed that induction of ER stress resulted in the release of IL-17 from PSI crypts (Fig. 4) and that ER stress contributes to inflammasome activation (Fig. 8C and D), we next measured the role of IL-17 in inflammasome activation and found that the induction of active IL-18 by alcohol is inhibited when mice were treated with an IL-17 blocking antibody (Fig. 8G and H).

Alcohol-induced CHOP induction, IL-18 activation and increase in PC number are mediated by the gut microbiota load

Since inflammatory signals from the gut microbiome contribute to the pathogenesis of alcohol-induced organ damage, we investigated whether gut decontamination might affect the alcohol-induced changes in the PSI. We used a combination of antibiotics (ampicillin, neomycin, vancomycin and metronidazole) administered before and simultaneously with the alcohol or pair-fed diet. Successful reduction in gut bacterial load was indicated by enlarged cecum and log-reduction of both aerobic and anaerobic colony forming units (Suppl. Fig. 4). We found that administration of antibiotics significantly reduced the alcohol-induced increase in CHOP and IL-18 cleavage in the SI (Fig. 9A and B). Moreover, antibiotics prevented the alcohol-induced increase in PC numbers measured by immunohistochemistry for lysozyme expression (Fig. 9C and D). These results suggest that signals from the gut microbiome are significant contributors to the development of alcohol-induced PSI pathology and PC functions.

Discussion

The biological effects of alcohol consumption on the GI tract are not limited only to intestinal functions, but extend beyond the GI tract due to alcohol-related changes in gut permeability. It has been shown that increased levels of LPS in the portal and systemic circulation contribute to the end-organ effects of alcohol in the liver, lung, and brain.^{4, 25} Previous studies demonstrated disruption of tight junctions in animal models of alcoholic liver disease, as well as in human alcoholics.²⁶ Furthermore, previous studies demonstrated pro-inflammatory pathway activation and increased levels of pro-inflammatory cytokines in the small bowel after chronic alcohol administration in mice.² In this study, we report that acute-on-chronic alcohol, which induces early alcoholic hepatitis in the liver³, modulates small intestinal homeostasis by increasing the numbers of Paneth cells and IL-17 production in the PSI. Mechanistically, we show that alcohol induces IL-17A production via ER stress

and results in upregulation of the eIF2 α and CHOP arm of UPR by chronic alcohol that was further augmented by additional acute alcohol binge in the PSI. We showed acute-on-chronic alcohol induced pro-apoptotic pathways in the PSI and that IL-17 itself induces ER stress in isolated crypts. *In vivo*, IL-17 blocking antibody prevented the alcohol-induced gut permeability and improved gut barrier functions via normalizing ZO-1 and occludin expression. We also found that alcohol induced ER stress, pro-inflammatory genes upregulation, inflammasome activation and IL-18 production after acute-on-chronic alcohol feeding in mice. Finally, decontamination of the gut with antibiotics prevented the alcohol-induced increase in Paneth cells numbers, CHOP upregulation and inflammasome activation in the PSI. Gut decontamination has also been shown to attenuate alcohol-induced liver inflammation³ and the data presented here provides further mechanistic evidence for this important gut-liver interplay.

The cells of the intestine serve a multitude of functions including the primary role of separating the contents of the intestine from the remainder of the body. Here, we show that acute-on-chronic alcohol exposure reduces the PSI gut barrier integrity leading to bacterial product translocation, although other regions of the intestinal tract may also have altered barrier integrity with increased translocation and these regions are an area of opportunity for future study. Reduced PSI integrity is associated with an increase in cell death in the PSI and an increase in pro-inflammatory cytokine expression. Paneth cells are the main contributors in the intestinal mucosal barrier by secreting large amounts of antimicrobial and inflammation-associated proteins.^{5-7, 26} In the normal GI tract, the number of PCs increases from proximal to distal and expansion of the PC population was found in infection.²⁷ Here, we discovered that alcohol selectively increased the abundance of PCs in the PSI, but not the DSI, in mice.

We also describe increased expression of genes involved in intestinal cell differentiation, such as Sox9, and its downstream target, Igfbp4.¹⁸ Spdef and Math1 are important for maintenance and differentiation of the PC lineage,¹⁹ while Lgr5 is a marker for intestinal stem cells.²⁹ The expression of all of these factors was increased in alcohol-fed animals suggesting expansion of PCs. A recent study from Schmitt et al. demonstrated that intestinal inflammation, similar to that induced by alcohol, led to an increase in Paneth cell stemness and PC³⁰ which may explain in part the paradoxical findings presented here that alcohol induces both cell death in the crypts of the PSI while also increasing Paneth cell numbers and expression of stem-related genes. Interestingly, PERK-eIF2 α dependent signaling plays a major role in intestinal stem cell differentiation³¹ and eIF2 α phosphorylation is required for normal PC function and mucosal homeostasis.³² Therefore, it is tempting to speculate that alcohol-induced eIF2 α phosphorylation contributes to increased PC numbers by inducing their differentiation. Other mechanisms, such as microbiota-mediated PC expansion, are also a possibility.

Strikingly, we found that ethanol stimulated IL-17A release in a dose-dependent manner in isolated crypts, mimicking our observations of increased IL-17 *in vivo* expression in alcohol-fed mice. IL-17A is a homeostatic cytokine in the gut that is constitutively present in PC granules³³ and regulates tight junction barrier integrity. Consistent with this, we found that anti-IL-17 blocking *in vivo* prevented the alcohol-induced increase in LPS in the liver

and restored the decreased expression of the tight junction proteins, ZO-1 and occludin, in the PSI in alcohol-fed mice indicating that mechanistically, IL-17 plays a role in increased gut permeability in alcoholic liver disease. This was also supported by our observation of upregulated pro-apoptotic molecules mediated by alcohol and IL-17 and the absence thereof in IL-17 antibody-treated mice.

PC-derived IL-17A has been implicated in distant organ injury, including hepatic ischemia/reperfusion⁸ and shock.³³ In our experiments, the rapid release of IL-17A from ethanol-treated crypts and the low abundance of IL-17⁺ immune cells suggest PCs, rather than immune cells, as the primary source of secreted IL-17A *in vitro* and *in vivo*. While our *in vitro* paradigm uses high concentrations of alcohol (4–100mM) for multiple hour durations, conditions that may be supraphysiologic and difficult to reproduce in an *in vivo* model, our findings do emphasize the results from 10-day acute-on-chronic mouse model. Different factors including Toll-like receptor ligands, bacteria, TNF α , and immune cell-derived interferon- γ have been shown to induce PC degranulation.³⁴ Here, we show that IL-17A secretion was mimicked by the ER stress inducer thapsigargin and alcohol-induced IL-17A was inhibited by the ER stress inhibitor 4-PBA, implicating alcohol-induced ER stress as a promoter of PC degranulation. In human alcoholic hepatitis, increased circulating IL-17 levels were found to be associated with an increase in Th17 cells in the livers of patients with alcoholic hepatitis.³⁵ Here, we found no significant increase in liver IL-17A levels or in Th17 cells in the acute-on-chronic alcoholic liver disease model (data not shown). We speculate that the increase in IL-17A in PCs may be local mechanism in the PSI rather than a systemic effect promoting liver inflammation.¹⁵ ER stress is a well-characterized pathophysiological factor in intestinal inflammatory diseases^{24,36,37} and alcohol is a potent ER stressor¹⁰. Although the mechanisms of alcohol-induced ER stress remain to be elucidated, we found that alcohol significantly increased eIF2 α phosphorylation and CHOP levels in the PSI. This suggests that acute-on-chronic alcohol may activate CHOP-mediated UPR signaling.³⁸

Here, we show that upregulation of CHOP expression in the proximal (but not in the distal) SI of alcohol-fed mice overlapped with the cellular distribution of PCs. Previous studies identified PCs as a main target of ER stress signaling and described PCs as one of the most susceptible intestinal epithelial cell types for perturbations of ER homeostasis.³⁹ PERK and downstream phosphorylation of eukaryotic translation initiation factor 2 α (eIF2 α) play a physiological role in intestinal stem cell differentiation.³¹ Moreover, the p-eIF2 α arm of the UPR is required for the normal function of PCs and mucosal homeostasis.³² Interestingly, bacteria-derived factors have been shown to induce UPR signaling and subsequent inflammation,⁴⁰ while administration of the ER stress inhibitor, 4-phenylbutyric acid (4-PBA), ameliorated dextran sulfate sodium (DSS)-induced colitis²² similar to the effect we observed with alcohol feeding.

We found that reducing gut bacterial load through antibiotic administration prevented CHOP induction *in vivo*. Chronic alcohol ingestion is associated with both quantitative (small intestinal bacterial overgrowth) and qualitative (dysbiosis) microbiome changes in the SI and PC antimicrobial peptide production is highly inducible by the microbiome.⁴¹ Thus, changes in the PSI microbiome might be an indirect mechanism by which alcohol affects PCs.

Importantly, alcohol induces changes in the microbiome throughout the intestinal tract while our present study focuses only on SI. Interestingly, PCs express ethanol-metabolizing enzymes and are prone to acetaldehyde accumulation due to lowered acetaldehyde oxidizing capacity.⁴² Acetaldehyde is a potent ER stressor through adduct formation and by impairing protein glycosylation. Thus, ethanol metabolites in PCs could directly cause ER stress.

Prolonged and/or unresolved ER stress leads to apoptosis. While the chronic alcohol feeding alone did not induce cell death signaling, the addition of an acute binge increased proapoptotic signaling with phosphorylation of eIF2 α and induction of CHOP-related downstream targets, GADD34, Bax and Bim. Phosphorylation of eIF2 α decreases protein load in the cell by attenuating translation, while allowing expression of certain ER stress-related proteins.¹² Removing p-eIF2 α -mediated translational block by GADD34, a downstream target of CHOP, allows for further synthesis of pro-apoptotic molecules.¹² Indeed, we found increased expression of the small mitochondrial pore-forming proteins, Bim and Bax, in the PSI of alcohol-fed animals. Overexpression of CHOP induces apoptosis by enhancing activation and mitochondrial translocation of Bax⁴³ and induces Bim expression,⁴⁴ eventually leading to mitochondrial membrane destabilization and downstream caspase activation. We show increased cleavage of the executioner caspase, caspase-3, and its downstream target PARP, both well-known markers of ongoing apoptotic signaling. Thus, we conclude that acute-on-chronic alcohol activates the CHOP-mediated mitochondrial apoptosis pathway leading to downstream caspase activation.

PCs are a site of origin for intestinal inflammation,²⁴ therefore we sought to identify possible pro-inflammatory pathways. In line with our previous data in the liver^{45, 46} and intestine,² we found that alcohol induces increased expression of the pro-inflammatory cytokines including *Tnfa*, *Il-6*, *Il1 β* , *Mcp1*, *Il23* and *Il17* in the PSI. Now we report increased activation of the inflammasome upon acute-on-chronic alcohol ingestion indicated by activation of the caspase-1 inflammasome and cleavage of its target, IL-18. Dupaul-Chicoine *et al.* recently showed that caspase-1 knockout mice were more susceptible to acute DSS-induced colitis and exogenous IL-18 administration resulted in protection in this model.⁴⁷ Zaki *et al.* found increased IL-18 production in DSS-induced colitis and suggested that epithelial inflammasome activation was followed by a regenerative proliferative response.⁴⁸ Further, these studies identified epithelial cells as the main source of IL-18 production.

Recently, caspase-2-mediated activation of the pro-apoptotic molecule, Bid, was shown to link ER stress to inflammasome activation. ER stress induced truncation of Bid through caspase-2 activation, which in turn translocated to the mitochondrial membrane leading to the release of mitochondrial content and inflammasome activation.¹³ Further, Upton *et al.* showed that ER stress increased caspase-2 levels by downregulating inhibitory microRNA expression leading to apoptosis.⁴⁹ In our study, we found significant upregulation of both caspase-2 and Bid protein expression along with increased cleaved form of Bid, tBid. In addition, caspase-8, the downstream effector of the extrinsic apoptotic pathway that can cleave Bid,⁵⁰ was induced by acute-on-chronic alcohol in the PSI. This suggests that alcohol might independently induce both extrinsic and intrinsic apoptotic pathways, converging on mitochondrial membrane destabilization.

Importantly, our study focused on the PSI where we noted a significant effect of alcohol on PCs, inflammatory cytokines, ER stress and tight junction protein. However, we did not investigate each of these factors throughout the entire gastrointestinal system, including the DSI (for some markers) and colon. Future investigation of these segments, and the regional effect of alcohol on gut barrier function, immune signaling and cellular response may provide additional information on the role of PC in alcohol-induced gut permeability and ALD.

In conclusion, we describe a novel, alcohol-induced pro-apoptotic signaling and inflammasome activation in the small intestine. We show that alcohol upregulates PC numbers in the PSI and that alcohol primarily affects PCs, which respond with IL-17A release *in vitro*. Our findings underscore the importance of examining the PSI not only as a microbiota-reservoir and tight junction barrier, but also as a complex organ with multiple response mechanisms to various impacts, specifically alcohol-mediated injuries.

Materials and Methods

Animal experiments

This study was conducted according to the regulations of the University of Massachusetts Medical School IACUC (Worcester, MA). Eight- to ten-week-old, female, wild-type C57BL/6 mice (Jackson Laboratory, Bar Harbor, ME) were fed a 5% (vol/vol) ethanol containing Lieber-DeCarli liquid diet or calorie-matched diet for 10 days with or without an oral gavage of 5g/kg ethanol or maltose dextrin solution, respectively (the NIAAA model¹⁵). Mice were sacrificed 4 or 9 hours after the gavage. For depletion of intestinal microbiota, ampicillin (1mg/ml), neomycin (10mg/ml), vancomycin (5mg/ml) and metronidazole (10mg/ml) were administered by gavage twice daily 5 days prior to and throughout the experiment. To verify successful decontamination of the gut, feces was cultured under aerobic and anaerobic conditions. 4-PBA (Calbiochem; 1g/kg mouse/day) was dissolved in water and mixed with the liquid diet. Its concentration in the diet was calculated based on daily monitoring of diet consumption (average 15ml/day). IL-17A blocking antibody (R&D) was injected three times before binge alcohol gavage (50ug/mouse, *i.p.*). IL-17A blocking or isotype antibody (R&D, 50ug/mouse through *i.p.* injection) was injected on 1st and 6th day of 5% alcohol diet. 100ug of anti-IL-17A antibody per mouse was injected on the 10th day of 5% alcohol diet for the last treatment, and it was injected 3hours before binge alcohol gavage.

Sample collection, ELISA, Immunoblotting, qRT-PCR reaction and histology are detailed in the Supplementary Materials and Methods.

Statistical analysis

Data are shown as mean \pm standard error of the mean. Statistical tests included Mann-Whitney or unpaired student t-test and $p < 0.05$ was considered statistically significant. Graphpad prism software was used to perform Grubbs test to remove outlier. (*=0.05).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Karen Kodys and Donna Catalano for their technical assistance. The authors also thank Frances S. Brown, Dr. Arlin B. Rogers, and the Cummings School of Veterinary Medicine at Tufts University for their excellent technical assistance and expert analysis of histology samples. We are especially thankful for assistance of Candice Dufour and Melanie Trombly for their assistance with proofreading, preparation and uploading of the manuscript.

Funding

This work was supported by an NIH grant from the NIAAA; R01 AA017729 (to GS), F30 AA024680 (to PL), F31 AA025545 (to AIV) & F30 AA022283 (to AS).

G.S. has received funding from NIH-NIAAA, Gilead, Genfit, University of Florida, Intercept, Allergan, Novartis, SignaBlok and Shire.

Abbreviations

4-PBA	4-phenylbutyric acid
CHOP	C/EBP homologous protein
DSS	dextran sulfate sodium; eIF2 α - eukaryotic translation initiation factor 2 α
ER	endoplasmic reticulum
EtOH	ethanol
GADD34	growth arrest and DNA damage-inducible gene 34
HBSS	Hank's balanced salt solution
IEL	intraepithelial leukocytes
Igfbp4	insulin-like growth factor binding protein 4
IL	interleukin
IRE1	inositol-requiring enzyme 1
Klf4	Krüppel-like factor 4
Lgr5	leucine rich repeat containing G protein coupled receptor 5
LPL	lamina propria leukocytes
Math1	mouse homolog of the <i>Drosophila</i> proneural gene atonal
MCP1	macrophage chemoattractant protein 1
PAS	periodic acid-Schiff

PARP	poly-ADP ribose polymerase
PBS	phosphate-buffered saline
PC	Paneth cell
PERK	PKR-like ER kinase
PF	pair-fed
PLA2g2a	phospholipase A
Sox9	SRY-box 9
Spdef	SAM pointed domain containing ETS transcription factor
Th	T helper
TNFα	tumor necrosis factor α
UPR	unfolded protein response
ZO-1	zonula occludens-1

References

- Hartmann P, Seebauer CT, Schnabl B. Alcoholic liver disease: the gut microbiome and liver cross talk. *Alcoholism, clinical and experimental research* 2015; 39(5): 763–775.
- Lippai D, Bala S, Catalano D, Kodys K, Szabo G. Micro-RNA-155 deficiency prevents alcohol-induced serum endotoxin increase and small bowel inflammation in mice. *Alcoholism, clinical and experimental research* 2014; 38(8): 2217–2224.
- Lowe PP, Gyongyosi B, Satishchandran A, Iracheta-Vellve A, Ambade A, Kodys K et al. Alcohol-related changes in the intestinal microbiome influence neutrophil infiltration, inflammation and steatosis in early alcoholic hepatitis in mice. *PLoS one* 2017; 12(3): e0174544. [PubMed: 28350851]
- Szabo G, Lippai D. Converging actions of alcohol on liver and brain immune signaling. *Int Rev Neurobiol* 2014; 118: 359–380. [PubMed: 25175869]
- Bevins CL, Salzman NH. Paneth cells, antimicrobial peptides and maintenance of intestinal homeostasis. *Nat Rev Microbiol* 2011; 9(5): 356–368. [PubMed: 21423246]
- Clevers HC, Bevins CL. Paneth cells: maestros of the small intestinal crypts. *Annu Rev Physiol* 2013; 75: 289–311. [PubMed: 23398152]
- Lee JS, Tato CM, Joyce-Shaikh B, Gulen MF, Cayatte C, Chen Y et al. Interleukin-23-Independent IL-17 Production Regulates Intestinal Epithelial Permeability. *Immunity* 2015; 43(4): 727–738. [PubMed: 26431948]
- Park SW, Kim M, Brown KM, D'Agati VD, Lee HT. Paneth cell-derived interleukin-17A causes multiorgan dysfunction after hepatic ischemia and reperfusion injury. *Hepatology* 2011; 53(5): 1662–1675. [PubMed: 21360570]
- Hosomi S, Kaser A, Blumberg RS. Role of endoplasmic reticulum stress and autophagy as interlinking pathways in the pathogenesis of inflammatory bowel disease. *Curr Opin Gastroenterol* 2015; 31(1): 81–88. [PubMed: 25426970]
- Ji C Mechanisms of alcohol-induced endoplasmic reticulum stress and organ injuries. *Biochem Res Int* 2012; 2012: 216450. [PubMed: 22110961]
- Bettigole SE, Glimcher LH. Endoplasmic reticulum stress in immunity. *Annual review of immunology* 2015; 33: 107–138.
- Szegezdi E, Logue SE, Gorman AM, Samali A. Mediators of endoplasmic reticulum stress-induced apoptosis. *EMBO Rep* 2006; 7(9): 880–885. [PubMed: 16953201]

13. Bronner DN, Abuaita BH, Chen X, Fitzgerald KA, Nunez G, He Y et al. Endoplasmic Reticulum Stress Activates the Inflammasome via NLRP3- and Caspase-2-Driven Mitochondrial Damage. *Immunity* 2015; 43(3): 451–462. [PubMed: 26341399]
14. Szabo G, Petrasek J. Inflammasome activation and function in liver disease. *Nature reviews Gastroenterology & hepatology* 2015; 12(7): 387–400. [PubMed: 26055245]
15. Bertola A, Mathews S, Ki SH, Wang H, Gao B. Mouse model of chronic and binge ethanol feeding (the NIAAA model). *Nat Protoc* 2013; 8(3): 627–637. [PubMed: 23449255]
16. Van Dussen KL, Dempsey PJ, Samuelson LC. Chapter 11 - Notch Pathway Regulation of Intestinal Cell Fate A2 - Johnson, Leonard R In: Ghishan FK, Kaunitz JD, Merchant JL, Said HM, Wood JD (eds). *Physiology of the Gastrointestinal Tract (Fifth Edition)*. Academic Press: Boston, 2012, pp 329–357.
17. Gassler N Paneth cells in intestinal physiology and pathophysiology. *World J Gastrointest Pathophysiol* 2017; 8(4): 150–160. [PubMed: 29184701]
18. Bastide P, Darido C, Pannequin J, Kist R, Robine S, Marty-Double C et al. Sox9 regulates cell proliferation and is required for Paneth cell differentiation in the intestinal epithelium. *J Cell Biol* 2007; 178(4): 635–648. [PubMed: 17698607]
19. Gregorieff A, Stange DE, Kujala P, Begthel H, van den Born M, Korving J et al. The ets-domain transcription factor Spdef promotes maturation of goblet and paneth cells in the intestinal epithelium. *Gastroenterology* 2009; 137(4): 1333–1345.e1331–1333. [PubMed: 19549527]
20. Yu T, Chen X, Zhang W, Li J, Xu R, Wang TC et al. Kruppel-like factor 4 regulates intestinal epithelial cell morphology and polarity. *PLoS one* 2012; 7(2): e32492. [PubMed: 22384261]
21. Maloy KJ, Kullberg MC. IL-23 and Th17 cytokines in intestinal homeostasis. *Mucosal Immunol* 2008; 1(5): 339–349. [PubMed: 19079198]
22. Cao SS, Zimmermann EM, Chuang BM, Song B, Nwokoye A, Wilkinson JE et al. The unfolded protein response and chemical chaperones reduce protein misfolding and colitis in mice. *Gastroenterology* 2013; 144(5): 989–1000 e1006. [PubMed: 23336977]
23. Hetz C The unfolded protein response: controlling cell fate decisions under ER stress and beyond. *Nat Rev Mol Cell Biol* 2012; 13(2): 89–102. [PubMed: 22251901]
24. Adolph TE, Tomczak MF, Niederreiter L, Ko HJ, Bock J, Martinez-Naves E et al. Paneth cells as a site of origin for intestinal inflammation. *Nature* 2013; 503(7475): 272–276. [PubMed: 24089213]
25. Lowe PP, Gyongyosi B, Satishchandran A, Iracheta-Vellve A, Cho Y, Ambade A et al. Reduced gut microbiome protects from alcohol-induced neuroinflammation and alters intestinal and brain inflammasome expression. *Journal of neuroinflammation* 2018; 15(1): 298. [PubMed: 30368255]
26. Purohit V, Bode JC, Bode C, Brenner DA, Choudhry MA, Hamilton F et al. Alcohol, intestinal bacterial growth, intestinal permeability to endotoxin, and medical consequences: summary of a symposium. *Alcohol* 2008; 42(5): 349–361. [PubMed: 18504085]
27. Martinez Rodriguez NR, Eloi MD, Huynh A, Dominguez T, Lam AH, Carcamo-Molina D et al. Expansion of Paneth cell population in response to enteric Salmonella enterica serovar Typhimurium infection. *Infect Immun* 2012; 80(1): 266–275. [PubMed: 22006567]
28. Yan AW, Fouts DE, Brandl J, Starkel P, Torralba M, Schott E et al. Enteric dysbiosis associated with a mouse model of alcoholic liver disease. *Hepatology* 2011; 53(1): 96–105. [PubMed: 21254165]
29. Sato T, van Es JH, Snippert HJ, Stange DE, Vries RG, van den Born M et al. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature* 2011; 469(7330): 415–418. [PubMed: 21113151]
30. Schmitt M, Schewe M, Sacchetti A, Feijtel D, van de Geer WS, Teeuwssen M et al. Paneth Cells Respond to Inflammation and Contribute to Tissue Regeneration by Acquiring Stem-like Features through SCF/c-Kit Signaling. *Cell reports* 2018; 24(9): 2312–2328.e2317. [PubMed: 30157426]
31. Heijmans J, van Lidth de Jeude JF, Koo BK, Rosekrans SL, Wielenga MC, van de Wetering M et al. ER stress causes rapid loss of intestinal epithelial stemness through activation of the unfolded protein response. *Cell reports* 2013; 3(4): 1128–1139. [PubMed: 23545496]
32. Cao SS, Wang M, Harrington JC, Chuang BM, Eckmann L, Kaufman RJ. Phosphorylation of eIF2alpha is dispensable for differentiation but required at a posttranscriptional level for paneth

- cell function and intestinal homeostasis in mice. *Inflamm Bowel Dis* 2014; 20(4): 712–722. [PubMed: 24577114]
33. Takahashi N, Vanlaere I, de Rycke R, Cauwels A, Joosten LA, Lubberts E et al. IL-17 produced by Paneth cells drives TNF-induced shock. *J Exp Med* 2008; 205(8): 1755–1761. [PubMed: 18663129]
 34. Farin HF, Karthaus WR, Kujala P, Rakhshandehroo M, Schwank G, Vries RG et al. Paneth cell extrusion and release of antimicrobial products is directly controlled by immune cell-derived IFN-gamma. *J Exp Med* 2014; 211(7): 1393–1405. [PubMed: 24980747]
 35. Lemmers A, Moreno C, Gustot T, Marechal R, Degre D, Demetter P et al. The interleukin-17 pathway is involved in human alcoholic liver disease. *Hepatology* 2009; 49(2): 646–657. [PubMed: 19177575]
 36. Grootjans J, Hodin CM, de Haan JJ, Derikx JP, Rouschop KM, Verheyen FK et al. Level of activation of the unfolded protein response correlates with Paneth cell apoptosis in human small intestine exposed to ischemia/reperfusion. *Gastroenterology* 2011; 140(2): 529–539 e523. [PubMed: 20965186]
 37. Deuring JJ, Fuhler GM, Konstantinov SR, Peppelenbosch MP, Kuipers EJ, de Haar C et al. Genomic ATG16L1 risk allele-restricted Paneth cell ER stress in quiescent Crohn's disease. *Gut* 2014; 63(7): 1081–1091. [PubMed: 23964099]
 38. DuRose JB, Tam AB, Niwa M. Intrinsic capacities of molecular sensors of the unfolded protein response to sense alternate forms of endoplasmic reticulum stress. *Molecular biology of the cell* 2006; 17(7): 3095–3107. [PubMed: 16672378]
 39. Kaser A, Lee AH, Franke A, Glickman JN, Zeissig S, Tilg H et al. XBP1 links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease. *Cell* 2008; 134(5): 743–756. [PubMed: 18775308]
 40. Pillich H, Loose M, Zimmer KP, Chakraborty T. Diverse roles of endoplasmic reticulum stress sensors in bacterial infection. *Molecular and cellular pediatrics* 2016; 3(1): 9. [PubMed: 26883353]
 41. Vaishnava S, Behrendt CL, Ismail AS, Eckmann L, Hooper LV. Paneth cells directly sense gut commensals and maintain homeostasis at the intestinal host-microbial interface. *Proc Natl Acad Sci U S A* 2008; 105(52): 20858–20863. [PubMed: 19075245]
 42. Chiang CP, Wu CW, Lee SP, Ho JL, Lee SL, Nieh S et al. Expression pattern, ethanolmetabolizing activities, and cellular localization of alcohol and aldehyde dehydrogenases in human small intestine. *Alcoholism, clinical and experimental research* 2012; 36(12): 2047–2058.
 43. Gotoh T, Terada K, Oyadomari S, Mori M. hsp70-DnaJ chaperone pair prevents nitric oxide- and CHOP-induced apoptosis by inhibiting translocation of Bax to mitochondria. *Cell death and differentiation* 2004; 11(4): 390–402. [PubMed: 14752510]
 44. Puthalakath H, O'Reilly LA, Gunn P, Lee L, Kelly PN, Huntington ND et al. ER stress triggers apoptosis by activating BH3-only protein Bim. *Cell* 2007; 129(7): 1337–1349. [PubMed: 17604722]
 45. Ambade A, Lowe P, Kodys K, Catalano D, Gyongyosi B, Cho Y et al. Pharmacological inhibition of CCR2/5 signaling prevents and reverses alcohol-induced liver damage, steatosis and inflammation in mice. *Hepatology* 2018.
 46. Petrasek J, Bala S, Csak T, Lippai D, Kodys K, Menashy V et al. IL-1 receptor antagonist ameliorates inflammasome-dependent alcoholic steatohepatitis in mice. *The Journal of clinical investigation* 2012; 122(10): 3476–3489. [PubMed: 22945633]
 47. Dupaul-Chicoine J, Yeretsian G, Doiron K, Bergstrom KS, McIntire CR, LeBlanc PM et al. Control of intestinal homeostasis, colitis, and colitis-associated colorectal cancer by the inflammatory caspases. *Immunity* 2010; 32(3): 367–378. [PubMed: 20226691]
 48. Zaki MH, Boyd KL, Vogel P, Kastan MB, Lamkanfi M, Kanneganti TD. The NLRP3 inflammasome protects against loss of epithelial integrity and mortality during experimental colitis. *Immunity* 2010; 32(3): 379–391. [PubMed: 20303296]
 49. Upton JP, Wang L, Han D, Wang ES, Huskey NE, Lim L et al. IRE1alpha cleaves select microRNAs during ER stress to derepress translation of proapoptotic Caspase-2. *Science (New York, NY)* 2012; 338(6108): 818–822.

50. Kantari C, Walczak H. Caspase-8 and bid: caught in the act between death receptors and mitochondria. *Biochimica et biophysica acta* 2011; 1813(4): 558–563. [PubMed: 21295084]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Alcohol induces cell death and inflammation in the proximal small intestine and leads to bacterial product translocation

(A) Bacterial product translocation to the liver was detected by qPCR of bacterial 16s rDNA and (B) by chromogenic endotoxin quantification of liver LPS levels after calorie-controlled pair-fed diet (PF) or 10 days of 5% alcohol in liquid diet plus binge alcohol (10d EtOH 9h binge) in mice. (C-D) Cell death was measured in the proximal small intestine (PSI) using Terminal deoxynucleotidyl transferase dUTP Nick-End Labeling (TUNEL) staining. (E) mRNA expression levels of inflammatory cytokines and chemokines, *Tnfa*, *Il-6*, *Il1β*, *Mcp1*, *Il23*, *Il17a*, were measured by qPCR. n=5–7 mice/group; * $p < 0.05$.




Figure 2. Alcohol increases the frequency of Paneth cells in the proximal small intestine and results in their degranulation

(A) Representative images and (C) quantification of PAS-stained proximal small intestinal sections. Inserts show individual crypts. (B) Representative images and (D) quantification of lysozyme immunohistochemistry (IHC) in PCs from pair-fed (PF) or 10d EtOH 9h binge alcohol-fed mice. (E) Crypt degranulation (highlighted by magnifications of (A) where PAS-positive material is observed in the crypt lumen in EtOH) was measured in the PSI and the DSI after 10d EtOH 9h binge alcohol exposure. (F) Expression of differentiation and stem markers, including *Sox9*, *Igfbp4*, *Spdef*, *Math1*, *Klf4* and *Lgr5*, were measured by qPCR in the PSI and DSI in alcohol- and pair-fed mice. (G) Expression of antimicrobial peptide genes, including *Cryptidins*, defensins *Defcr1*, *Defcr4* and *Defa-rs1*; *PLA2g2a*; lysozymes *Lyz1* and *Lyz2*; and *Muc2*, were also measured by qPCR. n=5–7 mice/group; * $p < 0.05$.




Figure 3. Alcohol induces IL17A expression in Paneth cells but not Th17 cells in the proximal small intestine

(A) Representative images and (B) quantification of IL-17A immunohistochemistry (IHC) in 10d EtOH 9h binge alcohol- and pair-fed (PF) mice. (C) IL-17A from whole PSI protein was also quantified (D) by Western blot. (E) Flow cytometry of small intestinal lamina propria (SI LP) detected Th17 cells, defined as CD45⁺, TCRb⁺, CD4⁺, IL-17⁺ live single cells. (F) Isolated small intestinal crypts were stimulated *ex vivo* with various concentrations of ethanol (0, 4, 25, 50 and 100mM) for 2, 4, 6 or 8 hours and IL-17A was measured from the supernatants by ELISA. n=5–7 mice/group; * $p < 0.05$.




Figure 4. Ethanol-induced IL-17A release is augmented by ER stress

(A) Isolated small intestinal crypts were treated with ethanol (100mM) to induce IL-17A release and some crypts were also treated with 5 or 10mM of the ER stress inhibitor 4-phenylbutyric acid (4-PBA). Released IL-17A was measured by ELISA from the crypt supernatants. (B) IL17A release ELISA performed on isolated crypts treated with the ER stress inducer thapsigargin (0.3, 1 and 3µM) or ethanol (25, 50 and 100 mM). (C) Cleaved (clv) PARP expression was measured by Western blot and quantified (D) from isolated crypts treated *ex vivo* with 10ng/ml of recombinant IL-17A, 50mM EtOH or IL-17A and EtOH. (E) Cleaved PARP, pro-caspase 8 and cleaved caspase 8 p43 subunit from the PSI of pair-fed (PF) or alcohol-fed mice treated or not with a control isotype antibody or anti-IL17A blocking antibody were measured by Western blot and quantified (F). For A-B, n=3 mice/treatment group; $p < 0.05$ * vs untreated, # vs EtOH 100 mM. For D-E, n=3-4, $p < 0.05$ * vs Untreated. For F-G, n=3-4 mice/group; $p < 0.05$ * vs PF.




Figure 5. Alcohol administration induces pro-apoptotic signaling and ER stress in the proximal small intestine.

(A) Pro-apoptotic proteins Bax, Bim, Bid, truncated Bid (tBid) and cleaved (clv) PARP, as well as ER stress related proteins CHOP, p-eIF2 α , eIF2 α , GADD34 full length (fl) were measured by Western blot and quantified (B) from PSI tissue of pair-fed (PF) mice as well as mice fed a 10-day chronic alcohol, a 10d EtOH plus 4h binge or a 10d EtOH 9h binge diet. (C) Caspase-3 (fl), cleaved (clv) caspase-3, caspase-2 and caspase-8 p43 were measured from the same PSI samples by Western blot and quantified (D). (E) Mice received either calorie matched diet (PF) or 10d EtOH 9h binge and some mice from each group also received *in vivo* administration of the ER stress inhibitor 4-phenylbutyric acid (4-PBA). Expression in the PSI of CHOP, procaspase-8 and clv caspase-8 p43 were measured by Western blot and quantified (F). For A-D, n=5–7 mice/group; $p < 0.05$ * vs pair-fed (PF), # vs 10d EtOH no binge, † vs 10d EtOH 4h binge. For E-F, n=5–7 mice/group; $p < 0.05$ * vs pair-fed, # vs 10d EtOH no binge, † vs 10d EtOH 4h binge,⁰ vs 10d EtOH 9h binge.




Figure 6. Alcohol-induced CHOP expression localizes to Paneth cells

(A) PSI epithelial and immune cell fractions (intraepithelial [IEL] and lamina propria leukocytes [LPL]) were isolated from pair-fed (PF) and 10d EtOH 9h binge mice for Western blot analysis and quantified (B) of CHOP and Bax proteins. (C) Representative CHOP immunofluorescent (IF) images of PSI samples of pair-fed (PF) and 10d EtOH 9h binge mice. Insets show individual crypts and the more intense IF staining observed in the 10-day + 9h binge alcohol samples. (E) Crypts were isolated from PF and 10d EtOH 9h binge mice and Bax and Bim proteins were measured by Western blot and quantified (F). n=3–7 mice/group; * $p < 0.05$ vs pair-fed. Scale bars = 50 μ m.




Figure 7. Tight-junction levels in the proximal small intestine are reduced by alcohol and restored by *in vivo* IL-17 blockade
 (A) The bacterial cell wall component LPS was measured in the livers of pair-fed (PF) and 10d EtOH 9h binge mice as well as EtOH mice treated with anti-IL-17 antibody or an isotype control. (B) ZO-1 immunofluorescent (IF) staining was completed on the PSI of these same mice and quantified. (C) Occludin was measured from the PSI by Western blot and quantified (D). mRNA expression of multiple tight junction proteins including *Claudin-1*, *-5* and *-15*, *Occludin* and *ZO-1* were measured by qPCR from PSI. For A, n=5–8 mice/group. For B, n=6 mice/group and 5 images/mouse were quantified. For C-D, n=6 mice/group. For E, n=7–10 mice/group. * $p < 0.05$ vs pair-fed (PF).




Figure 8. Inflammation activation by acute alcohol binge in the small intestine is attenuated by *in vivo* inhibition of ER stress

(A) Activation of inflammasome components caspase-1 and IL-18, including the active cleaved forms Casp-1 p10 and IL-18 p18, were assessed by Western blot and quantified (B) from the PSI of pair-fed (PF), chronic alcohol without binge, 10d EtOH 4h binge and 10d EtOH 9h binge mice. (C) ER stress inhibition through *in vivo* 4-phenylbutyric acid (4-PBA) administration was used to reduce CHOP expression and inflammasome target IL-18 activation was measured by Western blot and quantified (D) from PF and 10d EtOH 9h binge mice. (E) Western blot analysis and quantification (F) of cleaved (clv) PARP from PF and alcohol-fed mice with and without 4-PBA treatment. (G) Activation of IL-18 was measured by Western blot and quantified (H) in the PSI of PF and 10d EtOH 9h binge mice as well as EtOH mice treated with anti-IL-17 antibody or an isotype control. n=5–7 mice/group; $p < 0.05$ * vs pair-fed, # vs 10d EtOH No binge, † vs 10d EtOH 4h binge, ⁰ vs 10d EtOH 9h binge.




Figure 9. Alcohol-induced CHOP induction, IL-18 activation and PC increases are mediated by the gut microbiota load

(A) Gut decontamination using broad-spectrum antibiotics administration (Abx; ampicillin, neomycin, vancomycin and metronidazole) significantly reduced the gut microbiota load. CHOP and IL-18 activation were measured from PSI of PF and 10d EtOH 9h binge mice by Western blot and quantified (B). (C) Representative images and (D) quantification of lysozyme immunohistochemistry (IHC) from PF and 10d EtOH 9h binge alcohol-fed mice. n=6–10 mice/group; $p < 0.05$ * vs pair-fed, ^o vs 10d EtOH 9h binge.




Figure 10. Increased IL-17 production, Paneth cell activation, inflammasome and ER stress in the proximal small intestine after acute-on-chronic alcohol in mice

Working model based on our data indicates that acute binge on chronic alcohol exposure results in increased abundance of Paneth cells in the PSI, leading to ER stress induction, IL-17A release, apoptosis and inflammasome activation in crypts in the PSI. The mechanistic role of alcohol-induced IL-17 is indicated by experiments where IL-17 blocking antibody prevented alcohol-induced ER stress, apoptosis, inflammasome activation and IL-18 production in isolated crypts *in vitro* and *in vivo* in the PSI. Supplementary Table 1