

Original Research



OPEN ACCESS

Received: Sep 30, 2021

Revised: Jan 9, 2022

Accepted: Mar 30, 2022

Published online: Apr 21, 2022

Correspondence to

Bunty Ramchandani, MD

Congenital Cardiovascular Surgery, Hospital Universitario La Paz, Paseo de la Castellana 261, 28046 Madrid, Spain.

Email: bunty.r@gmail.com

Copyright © 2022. The Korean Society of Cardiology

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Bunty Ramchandani

<https://orcid.org/0000-0002-3304-2317>

Raúl Sánchez

<https://orcid.org/0000-0002-0145-9097>

Luz Polo

<https://orcid.org/0000-0002-2615-5858>

Álvaro Gonzalez

<https://orcid.org/0000-0001-9257-1174>

Meeting presentation

A 34th annual meeting of the European Association of Cardio-Thoracic Surgery (online 8–10 October 2020).

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

<https://e-kcj.org>

A Bicentric Propensity Matched Analysis of 158 Patients Comparing Porcine Versus Bovine Stented Bioprosthetic Valves in Pulmonary Position

Bunty Ramchandani , MD^{1,2}, Raúl Sánchez , PhD¹, Juvenal Rey, MD¹, Luz Polo , PhD¹, Álvaro Gonzalez , PhD¹, María-Jesús Lamas, PhD², Tomasa Centella, PhD², Jesús Díez, PhD³, and Ángel Aroca, PhD¹

¹Congenital Cardiovascular Surgery, Hospital Universitario La Paz, Madrid, Spain

²Congenital Cardiovascular Surgery, Hospital Universitario Ramón y Cajal, Madrid, Spain

³Instituto de Investigación, Hospital Universitario La Paz, Madrid, Spain

AUTHOR'S SUMMARY

Pulmonary valve replacement is the most common operation performed in adults with congenital heart disease (CHD). There is controversy regarding the best bioprosthesis in pulmonary position. The most commonly used xenografts are the porcine and bovine pericardial valves. With our bicentric propensity matched comparison we compared the mid-long term hemodynamic and clinical performance of Carpentier Perimount Magna Ease (Edwards™; Edwards Lifesciences, Irvine, CA, USA) a bovine pericardial bioprosthesis against the Mosaic (Medtronic™; Medtronic Inc., Minneapolis, MN, USA) a porcine bioprosthesis in pulmonary position in patients with CHD. Both bioprosthesis have similar outcomes during the first 5 years. Beyond 5 years porcine bioprosthesis have less structural valve degeneration.

ABSTRACT

Background and Objectives: Pulmonary valve replacement (PVR) is the most common operation in adults with congenital heart disease (CHD). There is controversy regarding the best bioprosthesis. We compare the performance of stented bioprosthetic valves (the Mosaic [Medtronic™] porcine pericardial against Carpentier Perimount Magna Ease [Edwards™] bovine) in pulmonary position in patients with CHD.

Methods: Between January 1999 and December 2019, all the PVRs were identified from hospital databases in 2 congenital heart centres in Spain. Valve performance was evaluated using clinical and echocardiographic criteria. Propensity score matching was used to balance the 2 treatment groups.

Results: Three hundred nineteen patients were retrospectively identified. After statistical adjustment, 79 propensity-matched pairs were available for comparison. Freedom from reintervention for the porcine cohort was 98.3%, 96.1%, and 91.9% at 3, 5, and 10 years and 100%, 98%, and 90.8% for the bovine cohort (p=0.88). Freedom from structural valve degeneration (SVD) for the porcine cohort was 96.9%, 92.8% and 88.7% at 3, 5, and 10 years and 100%, 98%, and 79.1% for the bovine cohort (p=0.38). Bovine prosthesis was associated with a reintervention hazard ratio (HR), 1.12; 95% confidence intervals (CIs), 0.24–5.26;

Conflict of Interest

The authors have no financial conflicts of interest.

Data Sharing Statement

The data generated in this study is available from the corresponding author upon reasonable request.

Author Contributions

Conceptualization: Ramchandani B; Data curation: Ramchandani B, Sánchez R, Rey J, Gonzalez Á, Lamas MJ, Centella T, Polo L; Formal analysis: Ramchandani B, Díez J; Investigation: Ramchandani B; Methodology: Ramchandani B, Díez J; Supervision: Polo L, Aroca Á; Validation: Polo L, Díez J, Aroca Á; Visualization: Ramchandani B; Writing - original draft: Ramchandani B; Writing - review & editing: Sánchez R, Rey J, Gonzalez Á, Lamas MJ, Centella T, Polo L, Díez J, Aroca Á.

p=0.89 and SVD HR, 1.69 (0.52–5.58); p=0.38. In the first 5 years, there was no difference in outcomes. After 5 years, the recipients of the bovine bioprosthesis were at higher risk for SVD (reintervention HR, 2.08 [0.27–16.0]; p=0.49; SVD HR, 6.99 [1.23–39.8]; p=0.03).

Conclusions: Both bioprosthesis have similar outcomes up to 5 years, afterwards, porcine bioprosthesis seem to have less SVD.

Keywords: Pulmonary valve replacement; Congenital heart disease; Bioprosthesis

INTRODUCTION

Pulmonary valve replacement (PVR) is the most common operation performed in adults with congenital heart disease.¹⁾ Pulmonary regurgitation is the main indication for surgery in patients who have undergone previous right ventricular outflow tract (RVOT) reconstruction.¹⁾ A competent pulmonary valve has shown to relieve symptoms and preserve cardiac performance.¹⁾²⁾

However, controversy still remains with regard to the best bioprosthesis in pulmonary position.¹⁻³⁾ The most commonly used xenografts are the porcine and bovine pericardial valves. There is evidence that bovine pericardial valves have a longer durability in aortic position, but this difference has not been showed in pulmonary position.³⁾

In this study we aim to compare the hemodynamic and clinical performance of Mosaic (Medtronic™; Medtronic Inc., Minneapolis, MN, USA) a porcine bioprosthesis against Carpentier Perimount Magna Ease (Edwards™; Edwards Lifesciences, Irvine, CA, USA) a bovine pericardial bioprosthesis in pulmonary position in patients with congenital heart disease.

METHODS

Ethical statement

Current guidelines of the Institution Ethics and Review Board for data protection were followed (IRB approval HULP: PI-5211), due to retrospective nature of the study informed consent was not warranted. All data collected arose from standard of care assessments.

Sample

This was a retrospective review of the medical records of all patients who underwent PVR from 1st January 1999 till 31st December 2019 at 2 congenital cardiac surgery reference centres. Study patients had either a Mosaic (Medtronic™; Medtronic Inc.) or a Carpentier Perimount Magna Ease (Edwards™; Edwards Lifesciences) bioprosthesis implanted in pulmonary position. Indications of PVR as well as preoperative image studies; such as echocardiography, cardiac magnetic resonance imagings and cath-lab data, were reviewed.

End points

Valve performance was evaluated using previously published criteria with a focus on echocardiographic follow-up to monitor hemodynamic valve function.⁴⁾ The primary endpoints were freedom from reintervention (surgical or catheter based) and structural valve degeneration (SVD) with a peak transpulmonary gradient of >50 mm Hg and/or more than moderate pulmonary regurgitation.

Hospital mortality is defined as death in hospital or within 30 days of discharge. All other mortality is considered late.

Patients, families, cardiologists and primary care physicians were contacted when necessary to acquire current follow-up information.

Operative technique

Mosaic bioprosthesis were implanted in Hospital Universitario La Paz (Madrid, Spain), while Carpentier were the prosthesis of choice in Hospital Universitario Ramón y Cajal (Madrid, Spain). The same team of surgeons operated in both hospitals. Due to administrative reasons only one brand of bioprosthesis was available in each institution. Both institutions are tertiary level referral hospitals that give coverage to different geographic areas of the Spanish national territory.

PVR with or without tricuspid repair were performed with beating heart using bicaval cannulation and mild hypothermia (32–34°C). Aortic cross-clamp was necessary for the repair of septal defects or left side valve surgery. Myocardial protection was done by cold single shot Celsior cardioplegia.

Either a failed right ventricle to pulmonary artery conduit, transannular patch or infundibular chamber was opened and residual patch material with native valve tissue was resected.⁵⁾ The posterior one third of the stented valve annulus was sewn to native pulmonary valve annulus with continuous polypropylene suture or interrupted mattress sutures. Anteriorly, a diamond-shaped patch of Contegra™ (formerly Venpro, Medtronic Inc.) or bovine pericardium was used to roof the RVOT over the prosthesis.

Statistical analysis

A descriptive analysis of all the variables was performed. Quantitative variables with normal distribution are expressed as means±SD. Non-normally distributed variables are expressed as medians and interquartile range (IQR). Qualitative variables are reported as absolute values and percentages. Comparison of means was done either with t-student or U Mann-Whitney depending on the Kolmogorov-Smirnov test. Comparison of qualitative variables was done with Fisher's test. Kaplan-Meier (KM) and Log-rank test were used to perform survival analysis. A proportional Cox hazard model was used to evaluate the relation between variables and events in follow-up. Statistical significance was considered when $p < 0.05$.

Due to notable imbalances in baseline covariates between the study groups a propensity score (PS) matching system was used, with a nearest neighbour greedy algorithm.^{6,7)} PS were calculated for each group using multivariate logistic regression according to the following clinical covariates: age, body surface area (BSA), size of the prosthesis, sex, base diagnosis, year of procedure and the presence of concomitant procedures.⁸⁻¹⁰⁾ Common support was subjectively assessed by graphical examination of PS across both groups. Balance of PS across both groups was evaluated in quintiles. If it was not equivalent one or more quintiles were split into smaller blocks. Balance of covariates across both groups were evaluated within blocks of the PS. Balance of covariates after matching was checked comparing standardized differences. The KM curves, stratified log-rank tests and adjusted Cox regression analyses were all then run on the PS matched sample.

Matched subjects have a similar value of PS, thus have a similar distribution of observed baseline covariates. Because baseline covariates are related to outcome, matched subjects are

likely to display a greater similarity in outcome compared with 2 randomly selected subjects. When outcomes are time-to-event in nature, KM survival curves can be estimated for treated and untreated subjects in the PS matched sample.⁷⁾ The KM curve calculates the survival rate each time the event of interest occurs rather than at preset follow-up intervals as in actuarial life tables. KM analysis is thus event dependent and not duration dependent.¹¹⁾

All analysis and graphical representations were done with STATA version 14.2 (Satacorp LP, College Station, TX, USA).

RESULTS

Overall sample

The study included a total of 319 patients with 337 PVR by a bioprosthesis. One hundred and ninety-nine prosthesis were porcine Mosaic (Medtronic™; Medtronic Inc.) and 138 were bovine Carpentier Perimount Magna Ease (Edwards™; Edwards Lifesciences). Due to difficulty for homograft procurement in Spain,¹²⁾ this treatment option was seldom used with few anecdotal cases which are not considered for this study. Baseline information on diagnosis and concomitant procedures are given in **Table 1**. The evolution of PVR and case mix is given in **Supplementary Table 1** and **Supplementary Figure 1**. Median follow-up duration was 3.27 years (IQR, 0.96–6.64) for porcine prosthesis and 5.92 years (IQR, 2.7–8.87) for the bovine group (rank-sum $p < 0.01$). The median age in the porcine group was 33.2 years whereas in the bovine group it was 26.7 years ($p < 0.01$). The median weight in the porcine group was 63.5 kg compared to 64.0 kg in the bovine group. Fifty-five patients (39.9%) were males in the porcine group and 99 (49.7%) in the bovine group. The median prosthesis size was 27.1 mm (SD, 1.81) for the porcine group and 26.8 mm (SD, 1.99) for the bovine group ($p = 0.079$).

In hospital mortality of the global cohort was of 2.5% ($n = 5$) in the porcine group and 5.8% ($n = 8$) in the bovine group, no statistical significance. Five patients in the porcine group were lost to follow-up because they had moved abroad, none in the bovine group. Late mortality was of 3 patients, 2 in the porcine group and one in the bovine group. Freedom from reintervention for patients who received a porcine bioprosthesis ($n = 189$) was 98.3%, 96.1%, and 93.6% at 3, 5, and 10 years and for patients who received a bovine bioprosthesis ($n = 130$) was 98.3%, 96.1%, and 80.9% at 3, 5, and 10 years, retrospectively ($\chi^2 = 5.48$; Log-rank $p = 0.02$). Freedom from SVD for patients with a porcine bioprosthesis was 97.6%, 94.1%, and 89.1% at 3, 5, and 10 years and for patients with bovine bioprosthesis was 97.5%, 94.3%, and 68% at 3, 5, and 10 years, respectively ($\chi^2 = 9.74$; Log-rank $p < 0.01$).

Figure 1A and B show the KM curves for reintervention and SVD for the full sample. For both outcomes, worse performance was noted in the bovine group, which was confirmed by log-rank test (reintervention $p = 0.02$; SVD $p = 0.002$). However, the KM curves indicate a similar performance among both groups until 5 years of follow-up postimplantation. This was confirmed by converging slopes of the residuals for both outcomes, despite the fact that proportional hazards assumption was not violated. We therefore included a time interaction in the Cox regression. Time was split into 2 periods ($t \leq 5$ years and $t > 5$ years) with separate hazard ratio (HR) estimation in each duration. The interactions were significant for both outcomes (reintervention $p = 0.019$; SVD $p = 0.005$). This indicated that in the first 5 years there was no statistical difference in outcomes (HR, 2.11 [0.47–9.50]; $p = 0.33$ and SVD HR, 1.61 [0.52–5.05]; $p = 0.41$). Whereas, after 5 years recipients of bovine bioprosthesis were at higher risk of both

Table 1. Covariate distribution by prosthesis type

Baseline characteristics	Porcine (n=199)	Bovine (n=138)	p value
Sex	99 (49.7)	55 (39.9)	0.073
Age	33.17 (19.15–44.07)	26.74 (20.69–31.40)	<0.001
BSA DuBois formula	1.68 (1.51–1.89)	1.72 (1.55–1.87)	0.35
Weight in kg	63.50 (50.00–77.00)	64.00 (53.00–74.00)	0.73
Height in cm	164.00 (155.00–172.00)	166.00 (159.00–172.00)	0.090
Pump time in mins	128.00 (97.00–188.00)	94.50 (71.00–123.50)	<0.001
Surgical Indication			<0.001
PI	140 (70.7)	118 (85.5)	
DPL	38 (19.2)	3 (2.2)	
PS	20 (10.1)	17 (12.3)	
Diagnosis			0.16
TOF	139 (70.2)	95 (68.8)	
PA+VSD	10 (5.1)	13 (9.4)	
PA+IS	10 (5.1)	2 (1.4)	
TGA+VSD+PS	5 (2.5)	4 (2.9)	
PS	26 (13.1)	23 (16.7)	
Truncus	5 (2.5)	1 (0.7)	
Ross sequel	3 (1.5)	0 (0.0)	
Pulmonary prosthesis diameter			0.25
19	1 (0.5)	1 (0.7)	
21	2 (1.0)	1 (0.7)	
23	5 (2.5)	10 (7.3)	
25	38 (19.1)	30 (21.9)	
26	0 (0.0)	1 (0.7)	
27	81 (40.7)	53 (38.7)	
29	72 (36.2)	41 (29.9)	
Concomitant procedures*	135 (67.8)	44 (32.1)	<0.001
Mean pulmonary prosthesis diameter	27.00 (27.00–29.00)	27.00 (25.00–29.00)	0.079
In-hospital mortality	5 (2.5)	8 (5.8)	0.15
Late mortality	2 (1.0)	1 (0.7)	1.00
Median follow-up in years	3.27 (0.96–6.64)	5.92 (2.70–8.87)	<0.001

Data are shown as mean±SD or number (%).

BSA = body surface area; DPL = double pulmonary lesion; PA+IS = pulmonary atresia with intact septum; PA+VSD = pulmonary atresia with ventricular septal defect; PS = pulmonary stenosis; PI = pulmonary insufficiency; TOF = tetralogy of fallot; TGA+VSD+PS = transposition of great arteries with ventricular septal defect and pulmonary stenosis; Truncus = truncus arteriosus.

*A detailed description of the concomitant procedures is available in the **Supplementary Table 2**.




Figure 1. Kaplan-Meier curves of the overall cohort.

outcomes (reintervention HR, 4.68 [1.29–17.0]; p=0.019 and SVD HR, 6.48 [2.17–19.33]; p=0.01). After accounting for time interaction, we found no evidence of non-proportional hazards.

Propensity-matched analysis

PS matching resulted in a total of 79 pairs of patients (79 porcine and 79 bovine bioprosthesis, n=158). **Table 2** shows the covariate imbalance by bioprosthesis among 79 matched patients. Some imbalance was detected in age and year of procedure as the standardized differences exceeded 0.10.

In hospital mortality of the PS cohort was of 3% (n=2) in the porcine group and 6% (n=5) in the bovine group, no statistical significance. Freedom from reintervention for patients who received a porcine bioprosthesis (n=79) was 98.3%, 96.1%, and 91.9% at 3, 5, and 10 years and for patients who received a bovine bioprosthesis (n=79) was 100%, 98%, and 90.8% at 3, 5, and 10 years, retrospectively ($\chi^2=0.02$; Log-rank p=0.88). Freedom from SVD for patients with a porcine bioprosthesis was 96.9%, 92.8%, and 88.7% at 3, 5, and 10 years and for patients with bovine bioprosthesis was 100%, 98%, and 79.1% at 3, 5, and 10 years, respectively ($\chi^2=0.77$; Log-rank p=0.38). **Figure 2A and B** show the KM curves for reintervention and SVD for the PS sample, no difference was found for both outcomes.

Bovine prosthesis was associated with a reintervention HR, 1.12; 95% confidence intervals, (0.24–5.26); p=0.89 and SVD HR, 1.69 (0.52–5.58); p=0.38. Time was split in 2 periods (t ≤5 years and t >5 years) with separate HR estimation in each duration. During the first 5 years

Table 2. Covariate imbalance by prosthesis type

Covariate balance	Unmatched			Matched		
	Bovine (n=138)	Porcine (n=199)	Stan dif	Bovine (n=79)	Porcine (n=79)	Stan dif
Female (%)	29.9	49.7	0.20	43.0	44.3	0.02
TOF (%)	68.8	70.2	0.04	68	68	0.02
Concomitant procedure (%)	32.1	66.67	0.73	48.1	44.3	0.01
Mean age (years)	26.39	32.1	0.47	29.2	27.78	0.14
Mean BSA (m ²)	1.69	1.65	-0.14	1.71	1.71	0.01
Mean prosthesis size (mm)	26.16	27.14	0.20	27.3	27.35	0.03
Mean year of procedure	2011	2014	0.61	2013	2012	0.19

BSA = body surface area; Stan dif = standardized differences; TOF = tetralogy of fallot.




Figure 2. Kaplan-Meier curves of the propensity matched cohort.

there was no difference in outcomes (reintervention HR, 0.52 [0.05–5.75]; $p=0.59$ and SVD HR, 0.27 [0.03–2.41]; $p=0.24$). Whereas after 5 years recipients of bovine bioprosthesis remained at higher risk of SVD but not reintervention (reintervention HR, 2.08 [0.27–16.0]; $p=0.49$ and SVD HR, 6.99 [1.23–39.8]; $p=0.03$). After accounting for time interaction, we found no evidence of non-proportional hazards.

DISCUSSION

There is enough accumulated evidence to accept the superiority of bovine pericardial valves in aortic position.¹³⁾¹⁴⁾ In vitro studies show different performance of bovine valves under left and right sided pressures. Hence left sided hemodynamic performance of these valves can not be extrapolated to the right side, low pressure, chambers.⁸⁾¹⁵⁾ It is this change in performance which casts doubts of the superior nature of the bovine bioprosthesis in pulmonary position.

Bioprosthesis in pulmonary position is an off-label indication and thus a different scenario. Here the long term data is scarce, with limited reports of over 10 year follow-up.¹⁶⁾¹⁷⁾ Comparisons between bioprosthesis is further hindered by confounding factors.¹⁸⁾ Studies include different timings or “eras” of operation,¹⁸⁾ different ages¹⁹⁾ and different valve sizes.²⁰⁾ Some factors are difficult to account for such as immunological factors²¹⁾ and operative complexity.²²⁾ In order, to make more robust comparisons we used a PS adjusting for age, BSA, size of the prosthesis, sex, base diagnosis, year of procedure and presence of concomitant procedures.

As calculated by Pragt et al.,¹⁷⁾ overall our results fared better than the median freedom from reintervention reported in the literature: 95.2±15.98% (range, 50–98.2%) for 5 year follow-up and 81.2% (66.4–84.7%) for 10 year follow-up. One has to take into account that these studies have a younger age cohort which itself is a risk factor for valve failure. Nomoto et al.,²³⁾ who included a slightly older population (median, 17.5 years) describe better results of freedom of reintervention of 96.7% at 5 and 10 years of follow-up.

Neither the definition nor the echocardiographic follow-up has been uniform between studies. The choice to include moderate regurgitation and/or moderate stenosis as definition for valve failure is arbitrary.¹⁶⁾ There is much debate regarding the best timing for reintervention of malfunctioning pulmonary valve as these are well tolerated.¹⁾³⁾¹⁶⁾¹⁸⁾ Hence, indication for valve reintervention is not solely indicated by the severity of valve malfunction. Our results show a low rate of valve failure: at 10 years of follow-up, 10.9% for porcine and 32% for bovine bioprosthesis. Comparisons with other studies is complicated due to age and valve failure definition confounders.

Although in our study none of the explanted valves were histologically examined, there are reports which identify deterioration patterns of bovine and porcine bioprosthesis.¹⁷⁾ Bovine bioprosthesis have a tendency to cusp stiffening with calcium deposits, whereas, porcine bioprosthesis have a tendency to cusp tearing. These findings have been described when the bioprosthesis are implanted in left side positions. In the right side, pulmonary pressure is lower than systemic pressures even in patients with pulmonary atresia and major aortopulmonary collateral arteries.¹⁵⁾¹⁶⁾²⁴⁾ This rarely has been the cause of acute deterioration of the bioprosthetic valve structure, which could explain the better performance of porcine bioprosthesis.

Our study included a number of limitations due to its retrospective nature. To improve the internal validity of our analysis a robust PS was used to compare both prosthesis. There may be other potential confounders not accounted for in the PS. Our study population was from 2 different centres with echocardiographic evaluations performed by 2 different teams of cardiologists using different machines, this could have introduced bias. Finally, our data included small number of patients beyond 5 years of follow-up especially in the porcine group, thus comparisons of survival beyond this point should be interpreted with caution.

In conclusions, bovine and porcine bioprosthesis have similar outcomes up to 5 years after implantation in pulmonary position. After 5 years, there was no difference in the rate of reintervention, however, porcine bioprosthesis seem to to have less SVD.

SUPPLEMENTARY MATERIALS

Supplementary Table 1

Distribution of base diagnosis

[Click here to view](#)

Supplementary Table 2

Concomitant procedures

[Click here to view](#)

Supplementary Figure 1

Evolution of pulmonary valve replacement.

[Click here to view](#)

REFERENCES

1. Kogon BE, Rosenblum JM, Mori M. Current readings: issues surrounding pulmonary valve replacement in repaired tetralogy of fallot. *Semin Thorac Cardiovasc Surg* 2015;27:57-64.
[PUBMED](#) | [CROSSREF](#)
2. Fiore AC, Rodefeld M, Turrentine M, et al. Pulmonary valve replacement: a comparison of three biological valves. *Ann Thorac Surg* 2008;85:1712-8.
[PUBMED](#) | [CROSSREF](#)
3. Emani SM. Options for prosthetic pulmonary valve replacement. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu* 2012;15:34-7.
[PUBMED](#) | [CROSSREF](#)
4. Akins CW, Miller DC, Turina MI, et al. Guidelines for reporting mortality and morbidity after cardiac valve interventions. *J Thorac Cardiovasc Surg* 2008;135:732-8.
[PUBMED](#) | [CROSSREF](#)
5. Bermudez CA, Dearani JA, Puga FJ, et al. Late results of the peel operation for replacement of failing extracardiac conduits. *Ann Thorac Surg* 2004;77:881-7.
[PUBMED](#) | [CROSSREF](#)
6. Garrido MM, Kelley AS, Paris J, et al. Methods for constructing and assessing propensity scores. *Health Serv Res* 2014;49:1701-20.
[PUBMED](#) | [CROSSREF](#)

7. Austin PC. The use of propensity score methods with survival or time-to-event outcomes: reporting measures of effect similar to those used in randomized experiments. *Stat Med* 2014;33:1242-58.
[PUBMED](#) | [CROSSREF](#)
8. Koziarz A, Makhdoum A, Butany J, Ouzounian M, Chung J. Modes of bioprosthetic valve failure: a narrative review. *Curr Opin Cardiol* 2020;35:123-32.
[PUBMED](#) | [CROSSREF](#)
9. Kostyunin AE, Yuzhalin AE, Rezvova MA, Ovcharenko EA, Glushkova TV, Kutikhin AG. Degeneration of bioprosthetic heart valves: update 2020. *J Am Heart Assoc* 2020;9:e018506.
[PUBMED](#) | [CROSSREF](#)
10. Salaun E, Mahjoub H, Girerd N, et al. Rate, timing, correlates, and outcomes of hemodynamic valve deterioration after bioprosthetic surgical aortic valve replacement. *Circulation* 2018;138:971-85.
[PUBMED](#) | [CROSSREF](#)
11. Barakat A, Mittal A, Ricketts D, Rogers BA. Understanding survival analysis: actuarial life tables and the Kaplan-Meier plot. *Br J Hosp Med (Lond)* 2019;80:642-6.
[PUBMED](#) | [CROSSREF](#)
12. Cuerpo Caballero G, López Menéndez J, Polo López L, et al. Cirugía cardiovascular en España en el año 2019. Registro de intervenciones de la Sociedad Española de Cirugía Cardiovascular y Endovascular. *Cirugía Cardiovasc* 2021;28:162-76.
[CROSSREF](#)
13. Raghav V, Okafor I, Quach M, Dang L, Marquez S, Yoganathan AP. Long-term durability of Carpentier-Edwards magna ease valve: a one billion cycle in vitro study. *Ann Thorac Surg* 2016;101:1759-65.
[PUBMED](#) | [CROSSREF](#)
14. Forcillo J, Pellerin M, Perrault LP, et al. Carpentier-Edwards pericardial valve in the aortic position: 25-years experience. *Ann Thorac Surg* 2013;96:486-93.
[PUBMED](#) | [CROSSREF](#)
15. Pragt H, van Melle JP, Verkerke GJ, Mariani MA, Ebels T. Pulmonary versus aortic pressure behavior of a bovine pericardial valve. *J Thorac Cardiovasc Surg* 2020;159:1051-1059.e1.
[PUBMED](#) | [CROSSREF](#)
16. Kwak JG, Bang JH, Cho S, et al. Long-term durability of bioprosthetic valves in pulmonary position: pericardial versus porcine valves. *J Thorac Cardiovasc Surg* 2020;160:476-84.
[PUBMED](#) | [CROSSREF](#)
17. Pragt H, Schoots MH, Accord RE, et al. A stented bovine pericardial prosthesis in the pulmonary position. *J Thorac Cardiovasc Surg* 2020;159:1063-1071.e1.
[PUBMED](#) | [CROSSREF](#)
18. Abbas JR, Hoschtitzky JA. Which is the best tissue valve used in the pulmonary position, late after previous repair of tetralogy of fallot? *Interact Cardiovasc Thorac Surg* 2013;17:854-60.
[PUBMED](#) | [CROSSREF](#)
19. Dorobantu DM, Sharabiani MT, Taliotis D, et al. Age over 35 years is associated with increased mortality after pulmonary valve replacement in repaired tetralogy of Fallot: results from the UK National Congenital Heart Disease Audit database. *Eur J Cardiothorac Surg* 2020;58:825-31.
[PUBMED](#) | [CROSSREF](#)
20. Aroca Á, Polo L, Pérez-Farinós N, et al. ¿Funcionan igual las prótesis pulmonares porcinas en menores de edad que en adultos? Una llamada a la prudencia. *Cirugía Cardiovasc* 2017;24:135-41.
[CROSSREF](#)
21. Marathe SP, Bell D, Betts K, et al. Homografts versus stentless bioprosthetic valves in the pulmonary position: a multicentre propensity-matched comparison in patients younger than 20 years. *Eur J Cardiothorac Surg* 2019;2019:1-8.
[PUBMED](#) | [CROSSREF](#)
22. Tatewaki H, Shiose A. Pulmonary valve replacement after repaired tetralogy of fallot. *Gen Thorac Cardiovasc Surg* 2018;66:509-15.
[PUBMED](#) | [CROSSREF](#)
23. Nomoto R, Sleeper LA, Borisuk MJ, et al. Outcome and performance of bioprosthetic pulmonary valve replacement in patients with congenital heart disease. *J Thorac Cardiovasc Surg* 2016;152:1333-1342.e3.
[PUBMED](#) | [CROSSREF](#)
24. Bortolotti U, Milano AD, Valente M, Thiene G. The stented porcine bioprosthesis: a 50-year journey through hopes and realities. *Ann Thorac Surg* 2019;108:304-8.
[PUBMED](#) | [CROSSREF](#)