



OPEN

Low fundamental and formant frequencies predict fighting ability among male mixed martial arts fighters

Toe Aung¹, Stefan Goetz², John Adams², Clint McKenna³, Catherine Hess¹, Stiven Roytman², Joey T. Cheng⁴, Samuele Zilioli² & David Puts^{1✉}

Human voice pitch is highly sexually dimorphic and eminently quantifiable, making it an ideal phenotype for studying the influence of sexual selection. In both traditional and industrial populations, lower pitch in men predicts mating success, reproductive success, and social status and shapes social perceptions, especially those related to physical formidability. Due to practical and ethical constraints however, scant evidence tests the central question of whether male voice pitch and other acoustic measures indicate actual fighting ability in humans. To address this, we examined pitch, pitch variability, and formant position of 475 mixed martial arts (MMA) fighters from an elite fighting league, with each fighter's acoustic measures assessed from multiple voice recordings extracted from audio or video interviews available online (YouTube, Google Video, podcasts), totaling 1312 voice recording samples. In four regression models each predicting a separate measure of fighting ability (win percentages, number of fights, Elo ratings, and retirement status), no acoustic measure significantly predicted fighting ability above and beyond covariates. However, after fight statistics, fight history, height, weight, and age were used to extract underlying dimensions of fighting ability via factor analysis, pitch and formant position negatively predicted "Fighting Experience" and "Size" factor scores in a multivariate regression model, explaining 3–8% of the variance. Our findings suggest that lower male pitch and formants may be valid cues of some components of fighting ability in men.

Sexually dimorphic vocalizations appear to have been shaped by sexual selection across a variety of species, including Australian field crickets (*Teleogryllus oceanicus*)¹, túngara frogs (*Physalaemus pustulosus*)², common loons (*Gavia immer*)³, red deer (*Cervus elaphus*)⁴, chacma baboons (*Papio ursinus*)⁵, and other anthropoid primates⁶, including humans (*Homo sapiens*)⁷. Pitch is the most perceptually salient acoustic characteristic of the human voice and is determined primarily by fundamental frequency (f_0), the rate of vocal fold vibration during phonation⁸. Pubertal increases in circulating testosterone and abundant androgen receptors in the vocal folds⁹ cause male vocal folds to grow approximately 60% longer than those of women⁸. Consequently, men's vocal folds vibrate at lower frequencies, and men speak at approximately half the f_0 of women⁶, a difference of about five standard deviations¹⁰.

Cross-species comparison suggests that sexual dimorphism in f_0 is likely to evolve under conditions of intense male-male competition in anthropoid primates⁶. Among humans living in both traditional and industrial populations, lower male f_0 predicts mating success, reproductive success, and social status¹¹ and conveys the impression that the speaker is more physically formidable^{7,12–16}. Because low f_0 reliably distinguishes adult males from adult females and children, it is a valid cue of physical formidability among humans generally¹⁷. More puzzling, however, is the abundance of evidence suggesting that listeners readily use f_0 as a cue for inferring a variety of traits even *within* sex, attributing greater size, strength, and formidability to adult male voices with lower f_0 ^{12,14,18,19}. Yet, scholars debate the accuracy of these perceptions^{11,17,18,20,21}. According to one hypothesis, attention to men's f_0 is the perceptual by-product of a broader tendency to associate low-frequency sounds with physically larger sources. Hence, listeners may perceive low frequencies to be intimidating even when f_0 is unrelated to the size of the sound producer^{18,21}. However, lower f_0 has been found to predict men's status, as well as mating and

¹Department of Anthropology, The Pennsylvania State University, 409 Carpenter Building, University Park, PA 16802, USA. ²Wayne State University, Detroit, USA. ³University of Michigan, Ann Arbor, USA. ⁴York University, Ontario, Canada. ✉email: dap27@psu.edu

reproductive success¹⁷. If these benefits were obtained partly through the influence of f_0 on social impressions, then deferring to men with low f_0 appears to have important costs that could be compensated only if there is an adaptive advantage to doing so^{11,17}. Selection should favor inattention to individual differences in men's f_0 unless the signal is at least partly honest^{22,23}.

Previous studies have investigated the information content of f_0 by exploring relationships with perceived formidability and its correlates such as size, strength, and hormonal concentrations. Participants can rapidly and accurately assess strength from men's voices^{19,24}. Although some studies have reported null findings between voice pitch and measures of strength^{25–28}, testosterone^{27,29,30}, and health³¹ among men, meta-analyses indicate that low f_0 in men predicts greater height ($r \sim -0.2$)³², testosterone concentrations ($r \sim -0.2$)¹⁷, and upper-body strength ($r \sim -0.1$)¹⁷. Although the strengths of these relationships are notably weak, each measure is an imperfect correlate of formidability, and any relationship between f_0 and overall formidability would be attenuated by error in quantifying formidability with a single correlate such as height. In addition, associations between f_0 and testosterone were found to be stronger among men with lower cortisol levels^{6,33}, a pattern of hormonal associations that may reflect greater immunocompetence³⁴. Lower male f_0 has been reported to be associated with elevated mucosal immunity²⁹ and energetic condition³⁵.

Despite the growing interest in the human male voice as a target of sexual selection, scant evidence directly tests the central question raised by the strong effects of f_0 on perceived formidability: Does low f_0 predict actual ability in physical confrontations among men? One means of circumventing the practical and ethical constraints on addressing this question is by utilizing data from men who engage in fighting for sport. In the only previous study of this type of which we are aware²⁸, vocal f_0 did not predict win percentage among 29 male mixed martial arts (MMA) fighters. In this small group of fighters, f_0 was obtained from a standardized passage (i.e., counting from 1 to 10) recorded on the day prior to competing in the IMMAF European Open Championships. This study was underpowered for detecting what is likely to be a small to moderate effect. Difficulties inherent to quantifying small differences in fighting ability among the upper echelons of competitive fighters, as well as naturally occurring within-individual variability in acoustic parameters across contexts^{13,16,36–38}, means that any strong empirical tests and efforts to precisely estimate effect size will require large samples, combined with multiple measures of both fighting ability (i.e., beyond win percentages) and acoustic parameters of interest.

In the current pre-registered study, we obtained 1312 voice samples from interviews of 475 MMA fighters from an elite fighting league. From these samples, we measured f_0 and several other sexually dimorphic acoustic parameters that have previously been linked with dominance, including f_0 variability (f_0 -SD), formant dispersion (D_f), and formant position (P_f)^{10,32,39}. Lower f_0 -SD indicates a more monotone voice, and both D_f and P_f are inversely related to vocal tract length and negatively influence perceptions of vocal timbre. D_f measures the average distance between successive formant (resonant) frequencies (in Hz) for the first n (usually 4) formants, whereas P_f is the average standardized formant value for the first n (usually 4) formants and measures how high or low formants are on average in standard deviations from the sample mean¹⁰. We tested the pre-registered hypotheses that lower, more male-typical f_0 , f_0 -SD, D_f , and P_f predict prevailing in physical confrontations, defined as fighting success indexed by fight outcomes and statistics across multiple physical encounters. We adhered to our pre-registered plans and conducted analyses using both D_f and P_f . However, at the request of a reviewer, we report only P_f as the formant measure in our main manuscript, as D_f is a less precise measure of formant structure^{10,32,40}. Results involving D_f can be found in our result output document available online. In our initial pre-registration, we planned to assess fighting ability via three measures of fighting success outcomes: total number of fights, Elo rating, and retirement status. However the reasons outlined below in our exploratory analyses section (i.e., fighting ability is likely a multidimensional construct), such separate measures are unlikely to fully capture fighting ability. Thus, we went beyond our pre-registered plans to additionally extract distinct dimensions of fighting ability via factor analysis of all fighting-related measures obtained in our study (height, weight, years active, age, total number of fights, Elo rating, retirement status, and win percentage) and explore how these dimensions were associated with f_0 and other acoustic measures.

Results

Pre-registered analyses. In multilevel models with individual acoustic measures as sole predictors (Fig S1), lower f_0 ($b = -0.11$, $p = 0.010$), f_0 -SD ($b = -0.11$, $p = 0.021$), and P_f ($b = -0.15$, $p < 0.001$) significantly predicted total number of fights, only P_f ($b = -0.15$, $p = 0.030$) predicted Elo ratings, and no acoustic measures predicted retirement status (Table S2). Zero-order correlations between best linear unbiased estimates of acoustic parameters and measures of fighting success produced similar results (Table 1). When all four acoustic measures and covariates, such as height and weight, were entered simultaneously as predictors of fighting success (including win percentage, an exploratory, non-pre-registered variable), none reached conventional levels of significance, indicating that acoustic measures did not explain unique variance over and above each other (Table S3). This is perhaps unsurprising given that some acoustic measures are positively correlated and thus share some degree of overlapping variance (see Table 1).

Exploratory analyses. Two key issues likely complicated our initial pre-registered assessments of fighting ability. The first issue is that although single measures of fighting ability, such as Elo ratings and win percentage, are correlated ($r = 0.37$; Table 1) and thus internally consistent, there are limitations to each; for example, fighters with no losses have a win percentage of 100% regardless of their number of fights. To address this issue, we conducted a principal axis factoring to reduce the set of eight fighting ability-related measures (height, weight, years active, age, total number of fights, Elo rating, retirement status, and win percentage) into dimensional factors ($n = 3$) using the “nFactors” package⁴¹, and rotated factors with the Varimax method using the “psych” package⁴². As shown in Table 3, onto Factor 1 (which we label “Fighting Experience”), number of fights, years active, age,

Variable status	f_0	f_0 -SD	P_f	Height	Weight	Active years	Age	Total fights	Elo rating	Retirement
f_0 -SD	0.48***									
P_f	0.24***	0.21***								
Height	-0.18***	-0.10*	-0.24***							
Weight	-0.15**	-0.08	-0.27***	0.74***						
Active years	-0.10*	-0.10*	-0.08*	0.08	0.17***					
Age	-0.17***	-0.09	-0.20***	0.11*	0.30***	0.30***				
Total fights	-0.11*	-0.09*	-0.13**	0.11*	0.21***	0.73***	0.44***			
Elo rating	-0.04	-0.03	-0.11*	0.16**	0.23***	0.47***	0.15**	0.41***		
Retirement status	-0.04	-0.01	-0.03	0.07	0.05	-0.31***	0.20***	0.09	-0.22***	
Win percentage	-0.03	-0.01	-0.05	0.02	0.03	0.31***	0.03	0.20***	0.37***	-0.37***

Table 1. Zero-order correlations among study variables. Acoustic variables are computed as the best linear unbiased measures for multiple recordings measured at the individual level by using micro-macro multi-level modeling⁷³, which is preferred to using simple group means⁷⁴. $p < .05^*$, $p < .01^{**}$, $p < .001^{***}$.

Predictors	Factors		
	Fighting Experience	Fighting Success	Size
Intercept	-0.01 (.903)	0.01 (.976)	-0.01 (.966)
f_0	-0.10 (.059)	0.04 (.405)	-0.13 (.018)
f_0 -SD	-0.04 (.507)	-0.04 (.633)	0.04 (.534)
P_f	-0.20 (.002)	0.02 (.832)	-0.32 (<.001)
R^2	0.04	0.01	0.08

Table 2. Results of micro-macro multi-level models predicting three component measures of fight metrics from acoustic measures. Results are reported as effect size (p-value). Effect sizes are beta-weights, except for retirement status, which is odds ratio. In all models, the variance inflation factor (VIF) of each predictor was < 1.5. f_0 = fundamental frequency; f_0 -SD = Variability in fundamental frequency; P_f = formant position.

and Elo rating loaded most strongly. Onto Factor 2 (“Fighting Success”), retirement status, win percentage, Elo rating, and years active loaded most strongly. Finally, onto Factor 3 (“Size”), height and weight loaded most strongly. Factor scores on each of these 3 factors were then used as dependent variables, in separate regression models, with all acoustic measures as predictors (Table 2). Results from these models indicate that P_f negatively predicted Fighting Experience, whereas f_0 and P_f negatively predicted Size (Fig S2). To strengthen our findings, we also conducted a structural equation model that takes measurement error into account and obtained the same pattern of results and acceptable model fit (Fig. 1).

A second issue, highlighted by our principal axis factoring, is that fighting ability is multidimensional. Although Fighting Experience explained the most variance in fighting ability-related variables (27%), Fighting Success (23%) and Size (22%) also explained substantial proportions and captured unique components of fighting ability. For example, fighters compete only against opponents within the same weight class. Body size is sufficiently decisive in fights to necessitate weight classes; a 52.2 kg strawweight fighter and a 120.2 kg heavy-weight fighter with identical Elo ratings, win percentage, and so forth are not equivalent in fighting ability, for instance. To test whether fighting ability relates to an acoustic parameter such as f_0 , it is important to consider all major components of fighting ability simultaneously. Although the structural equation model reported above considered all three extracted components of fighting ability, structural equation models test association, not significance, as their main function is to evaluate whether the model conforms to the data⁴³. To test whether f_0 and other acoustic parameters predict overall fighting ability, we therefore conducted a multivariate regression model where Fighting Experience, Fighting Success and Size were simultaneously entered as dependent variables and acoustic measures as predictors (Table 4). We observed statistically significant relationships between fighting ability and both f_0 and P_f , but not f_0 -SD. When we compared the multivariate model with all acoustic measures to the model with only f_0 and P_f as predictors, we found that the latter model performed as well as the former. Both f_0 and P_f negatively predicted Fighting Experience and Size but neither predicted Fighting Success. We obtained similar results when f_0 and P_f were entered as single predictors in separate multivariate regression models (Fig. 2).

Discussion

Sexual dimorphism in f_0 likely arose in the common ancestor of the catarrhine primates after their divergence from the New World monkeys⁶ approximately 43.5mya⁴⁴ and appears to have been subsequently elaborated or reduced depending on the form and degree of male mating competition⁶. Relatively low male f_0 may have evolved as a means of exaggerating the appearance of size to same-sex competitors and/or potential mates^{6,18}, but there is considerable debate regarding whether male f_0 is purely deceptive^{20,21,45} or provides any reliable information about formidability in men^{11,17,33}.




Figure 1. Path diagram for a structural equation model highlighting standardized associations among all observed (rectangular) and latent (oval) variables. We first fitted an orthogonal model, but allowing residual correlations between latent variables significantly improved the model ($p < .001$). Note. f_0 = fundamental frequency; f_0 -SD = Variability in fundamental frequency; P_f = formant position; χ^2 = model chi square; CFI = comparative fit index; RMSEA = root mean square error of approximation; SRMR = standardized root mean square residual. $p < .05$, $p < .01$, $p < .001$ ***.

To shed light on this debate, we investigated whether f_0 is associated with fighting ability among a large sample of male MMA fighters. Results of our pre-registered analyses were generally in the direction of lower f_0 predicting greater fighting ability but were mixed in terms of statistical significance. When we addressed the limitations of these analyses by creating more precise measures of fighting ability and accounting for the contributions of the distinct dimensions revealed through principal axis factoring, we found that lower f_0 was associated with greater fighting ability in all analyses. In the statistical model that most precisely measured fighting ability by including principal axis factors related to fighting experience, fighting success, and body size, f_0 predicted fighting ability generally and specifically components of fighting ability related to experience and size, but not within-weight class fighting success. Overall, these results suggest that low f_0 is an honest cue of formidability in men (Table 3).

Effect sizes were small, however. Even when fighting ability was measured most precisely, f_0 explained only 1–3% of the variance (Table 4). On the one hand, the strength of these associations accords with theoretical predictions derived from the fact that signaling is multimodal and multi-component¹⁷. On the other hand, it is important to emphasize that relationships between f_0 and fighting success among MMA fighters may underestimate those in the general population due to range restriction on fighting ability. Among elephant seals, the body length of males who occupied the center of harems correlated with neither maximum harem size nor tenure length on the beach during the breeding season⁴⁶. However, when all males were analyzed together, including those that were peripheral to or outside of the harem, male body length explained 17% of the variation in tenure length on the beach. The degree to which f_0 correlates with fighting ability in the general population has yet to be tested for obvious ethical reasons.

Voice pitch and other acoustic variables are modulated across social contexts, including those related to perceived fighting ability relative to a competitor^{13,47}, relative dominance and prestige⁴⁸, authority³⁶, current aggressive intent⁴⁹, emergent rank¹⁶, and volitional exaggeration^{50,51}. On the one hand, if voice pitch is modulated in relation to self-perceived relative formidability and status, as this prior research indicates, then it is possible that any voice modulation of fighters during interviews could have strengthened relationships between acoustic parameters and measures of fighting ability in the present study. On the other hand, if voice modulation is less patterned or less dependent upon perceived relative formidability, then this would introduce noise in measuring individual differences in voices, which would tend to weaken relationships between acoustic parameters and true fighting ability. In these data based on naturalistic observations, although we cannot rule out that some fighters may have modulated their voice pitch or other vocal parameters during interviews to sound stronger, we sought to strengthen our measures of individual differences in acoustic traits by sampling across multiple interview occasions. Indeed, we found that acoustic measures were consistent across fighters, even between pre- and post-fight conditions in the subset of our sample for which this information was available (see Supplemental Procedures




Figure 2. Relationships between fighting ability components and acoustic measures. The relationship between fundamental frequency (left column) and formant frequency (right column) are plotted against Fighting Experience (top row), Fighting Success (middle row), and Size (bottom row).

Predictors	Factors		
	Factor 1: Fighting Experience	Factor 2: Fighting Success	Factor 3: Size
Number of fights	0.90	0.13	
Years active	0.74	0.48	
Age	0.72	-0.27	0.16
Elo rating	0.44	0.56	0.21
Weight	0.19		0.91
Retirement status	0.19	-0.81	
Win percentage	0.15	0.73	
Height			0.93
Variance explained	0.27	0.23	0.22

Table 3. Component axis analysis for measures related to fighting ability. Loadings that were above $|0.4|$ are bolded. Factors are named descriptively according to the variables that load most strongly onto them. Empty cells indicate factor loadings $<|0.1|$.

for details), and we attempted to capture more stable individual acoustic differences rather than measurements specific to particular recording conditions by using the unbiased linear estimates across recordings for each fighter in all analyses.

Only one previous study of which we are aware²⁸ examined links between fighting success and f_0 or other acoustic measures and found that acoustic measures did not predict win percentage among male MMA fighters. Likewise, in our bivariate correlation analysis, we did not find a significant relationship between win percentage

	Fighting Experience	Fighting Success	Size	Pillai test
Model 1				
f_o	- 0.10 (.903)	0.05 (.429)	- 0.13 (.017)	0.02 (.019)
f_o -SD	- 0.04 (.577)	- 0.04 (.604)	0.04 (.536)	0.01 (.819)
P_f	- 0.20 (.005)	0.02 (.829)	- 0.32 (<.001)	0.06 (<.001)
R^2	0.04	0.01	0.08	
Model 2				
f_o	- 0.11 (.022)	0.03 (.539)	- 0.12 (.018)	0.02 (.008)
P_f	- 0.15 (.016)	- 0.01 (.954)	- 0.31 (<.001)	0.07 (<.001)
R^2	0.03	0.01	0.08	
Model 3				
f_o	- 0.14 (.003)	0.03 (.536)	- 0.17 (<.001)	0.05 (<.001)
R^2	0.02	0.01	0.03	
Model 4				
P_f	- 0.19 (.002)	0.01 (.927)	- 0.35 (<.001)	0.09 (<.001)
R^2	0.02	0.01	0.07	

Table 4. Results of a multi-variate regression model. Results are reported as effect size (p-value). Effect sizes are beta-weights, except for Pillai's trace test statistics. f_o = fundamental frequency; f_o -SD = Variability in fundamental frequency; P_f = formant position.

and vocal f_o (Table 1), although lower f_o predicted a greater number of UFC matches. Our ability in subsequent analyses to detect relationships between f_o and fighting ability where a previous study did not²⁸ may have been due in part to our examination of a larger sample of fighters (474 vs. 29), along with longer total voice samples from each fighter (approximately 85 s vs. 8 s).

Our use of unstandardized, spontaneous speech samples also offered enhanced ecological validity⁴⁷, and although this approach adds noise to the measurement of individual differences in acoustic parameters, previous research shows strong correspondence across speech and/or vocalization types for both fundamental⁵² and formant¹⁰ frequencies. Moreover, the best linear unbiased composite estimates of acoustic measures from multiple voice recordings of each fighter, along with relatively long (> 85 s) average total speech samples from individual fighters, provided reliable estimates of individual differences in acoustic features.

Perhaps most importantly, we extracted component variables via factor analysis to produce more comprehensive measures of fighting ability than can be offered by individual measures such as Elo rating or win percentage. For example, win percentage alone is an imperfect indicator of fighting success because fighters with fewer fights and losses can achieve higher win rates^{53,54} and because win percentage does not account for the strength of opponents as Elo ratings do. Our factor analysis addressed such limitations and produced three unique components of fighting ability, which we termed Fighting Experience, Fighting Success, and Size. Fighting Experience reflects the component of fighting ability most strongly related to total number of fights but also strongly related to years active in the UFC and age. Given that Elo ratings also loaded positively onto this component, it may represent the component of fighting skill attributable to experience. Fighting Success reflects a component of fighting ability related to a history of winning in one's weight class that is relatively unrelated to experience or size, perhaps tapping characteristics such as speed, agility, and strength for one's size. The extracted Size factor reflects the component of fighting ability related to height and weight. Larger size is generally associated with lower f_o across mammals⁵⁵ and is a major determinant of dominance and social status across species⁵⁶⁻⁶⁰. Physical size is such a strong determinant of outcomes in combat sports such as wrestling, boxing, and martial arts that weight classes are needed to prevent dangerous lopsided contests, and fighters are willing to sacrifice energy and hydration by cutting weight to fight smaller opponents. Parallel to findings in previous studies^{32,61}, fighters' f_o predicted their height and weight, as well as their Size factor scores.

Our orthogonal factor analysis extracted independent components of fighting ability, maximizing the variance among items and facilitating interpretation. A structural equation model further confirmed the structure of our proposed components of fighting ability, but allowing residual correlations between latent variables produces a better model fit. These analyses highlight the importance of considering fighting ability across related measures instead of using a single measure of fighting experience, fighting success, or size. Because fighting ability measures (e.g., win percentage and number of fights) reflect weight division-specific measures, results from analyses that do not incorporate the influence of fighter's weight (Table 1) may be less valid than others (Table 4).

Although not the focus of the present paper, other sexually dimorphic acoustic parameters were associated with fighting ability. In correlation analyses, lower f_o -SD and P_f predicted a greater number of UFC fight matches, and lower P_f predicted higher Elo ratings (Table 1). In addition, P_f predicted overall measures of fighting ability in the multivariate regression model (Fig. 2) and independently predicted Fighting Experience and Size (Fig. S2). Our findings extend those from previous studies showing that P_f predicts objective measures of threat potential, such as size³², strength, and physical aggression¹⁰, as well as perceived fighting ability^{14,62}, and suggest that P_f communicates size-independent information about vocalizers' physical formidability.

A potential statistical issue concerns the use of uncorrected p -values associated with multiple tests. However, the adjustment of p -values, such as Bonferroni correction, not only decreases the rates of Type I error, but also increases the rates of Type II error and has been highly criticized⁶³. Additional analyses used in this study, such as multi-level modeling⁶⁴, multi-variate regression, and structural equation modeling, that consider dependent variables simultaneously should reduce concerns associated with multiple comparisons.

In general, our findings are consistent with the broader perspective that men's anatomy, behavior, and psychology have been shaped by an evolutionary history of contest competition^{27,65}, the use of force or threat of force to exclude same-sex competitors from mates⁵⁹. Contest competition should favor psychological mechanisms to attend to and assess the formidability and threat potential of competitors¹⁹. People can accurately assess physical strength from body¹⁹ and face⁶⁶ images. For example, sexually dimorphic facial cues predict fighting ability among MMA fighters^{33,67} but see⁶⁸, and using these cues participants can predict fighting outcomes above chance⁶⁹. The voice appears to be another aspect of the phenotype that indicates formidability^{10,13,19}; we showed that f_0 was associated with both body size and fighting experience (independent of body size) among male MMA fighters. Although associations between measures of fighting ability and f_0 and other sexually dimorphic acoustic parameters were small, they comport with the notion that f_0 is but one of many components of a vocal acoustic signal, and that voices represent one of many signals of men's formidability¹⁷. A deep, resonant voice clearly indicates status as a physically mature or maturing male. Our findings support the further possibility that attention to differences in voice pitch among men may be functional as well, as f_0 and other sexually dimorphic components of men's voices appear to provide information about differences in men's threat potential.

Method

Samples. Out of all fighters in the MMA promotion company Ultimate Fighting Championship (UFC) who were active up to the end of 2013 (UFC event 168, 12/28/2013), we obtained data on 475 fighters for our analyses (see Supplemental Procedures for details). Data were not collected after 2013, as there was a spike in the number of UFC events in 2014 that may have confounded results. Following prior work⁵⁴, we included only experienced fighters, defined as those who have fought at least 10 professional fights, at least one of which was in the UFC.

Fighting ability measurement. We obtained fighters' age, height, weight, total number of fights, and total number of wins from Wikipedia and Sherdog websites. Because, during the sampling period, fighters who experienced three consecutive losses were likely eliminated from the UFC⁷⁰, only relatively successful fighters were likely to remain in the UFC. To measure long-term fighting ability, we obtained the retirement status (whether the fighter retired between 2013 and 2018) and calculated the total number of years a fighter was active (the total number of years between the fighter's first UFC fight and his last UFC fight between 2013 and 2018). In addition, we obtained Elo-equivalent ratings as another measure of fighting ability for each fighter. These are computerized objective rankings generated by the FightMatrix website that utilizes a proprietary engine (CIRRS – Combat Intelli-Rating and Ranking System) and a comprehensive MMA fight database to adjust ratings according to each fighter's wins and losses and the records of opponents. For descriptive statistics on fighting ability measures, see Table S1.

Acoustic measurement. To provide a more precise measure of person-level f_0 , we attempted to collect multiple (up to five) voice samples from each fighter samples (range = 1–5, $M = 2.76$, $SD = 1.15$) from unique audio and video interviews obtained from YouTube, video.google.com, or podcasts from radio shows and downloaded as .wav files. We prioritized pre-fight recordings but also obtained post-fight recordings, identifying them accordingly in our dataset. Secondary sounds (e.g., background noise, interviewers' voices) were removed from downloaded recordings so that only the voice of each fighter was extracted. A mean of 30.99 s (range = 4–594.98, $SD = 55.62$) of a fighter's voice was extracted from each interview, so that the combined length of all voice samples from a fighter averaged 85.32 s (range = 5.43–1099.33, $SD = 133.75$). We then used Praat to measure f_0 , f_0 -SD, D_f , and P_f from each voice sample. For mean f_0 and f_0 -SD, pitch floor was set to 75 Hz, and pitch ceiling was set to 300 Hz in accordance with programmer recommendations. Otherwise, default settings were used. For D_f and P_f calculations¹⁰, f_1 through f_4 were measured at each glottal pulse (automated detection by Praat) and averaged across measurements. Before analyses, we removed one fighter from our sample whose P_f measurement was 7.5 SD above the mean, a decision not specified in our pre-registration document. Our final sample comprised 1311 recordings from 474 fighters (see Supplemental Procedures for details).

Data analysis

Most common multilevel models focus on macro–micro conditions; a mix of both lower micro-level (level 1) and higher macro-level (level 2) explanatory variables are used to predict lower micro-level (level 1) outcomes⁷¹. However, our data diverge from this typical condition, as they exhibit the opposite micro–macro pattern, with our acoustic measures predictors collected multiple times for (that is, nested within) the same fighter (level 1), and our dependent variables (e.g., fighting ability) captured for each fighter (level 2). Given this somewhat anomalous feature of our data, predicting level 2 variables from level 1 variables using standard multilevel techniques would produce statistically biased results⁷². To circumvent these issues, we used the “MicroMacroMultilevel” package⁷³ in R that allows micro–macro multi-level models⁷⁴ by producing the best linear unbiased predictors (BLUPs) for all group aggregates of variables measured at the lowest level. All acoustic measures and covariates, including Elo ratings, were z-scored before analyses (see Table 1 for correlations among study variables), so all results reported below represent standardized effects. All analyses were conducted using R 3.6.1. Three models in our pre-registered tests were estimated, each with a different dependent variable—a given fighter's total number of fights, Elo ratings, and retirement status. For each dependent variable, the BLUP of the group mean

for each acoustic measure was entered as the sole predictor in our multi-level models. In subsequent models, all four BLUPs of the group mean for acoustic measures were simultaneously entered as predictor variables, and fighters' height, weight, age, and number of active years were entered as additional level 2 covariates. Detailed methods for additional tests are described above. Data and scripts for all analyses are made available online (see Supplemental Procedures).

Ethics. Data were obtained from online sources that are freely available to public.

Data availability

The raw data and scripts for all our models are also made available online at https://osf.io/md6wj/?view_only=81cf6446a90448a594e1e1ec6b25ce59.

Received: 16 June 2020; Accepted: 8 December 2020

Published online: 13 January 2021

References

- Tregenza, T., Simmons, L. W., Wedell, N. & Zuk, M. Female preference for male courtship song and its role as a signal of immune function and condition. *Anim. Behav.* **72**, 809–818 (2006).
- Guerra, M. A., Ryan, M. J. & Cannatella, D. C. Ontogeny of sexual dimorphism in the larynx of the túngara frog *Physalaemus pustulosus*. *Copeia* **2014**, 123–129 (2014).
- Mager, J. N. III., Walcott, C. & Piper, W. H. Male common loons, *Gavia immer*, communicate body mass and condition through dominant frequencies of territorial yodels. *Anim. Behav.* **73**, 683–690 (2007).
- Reby, D., Charlton, B. D., Locatelli, Y. & McComb, K. Oestrous red deer hinds prefer male roars with higher fundamental frequencies. *Proc. R. Soc. B Biol. Sci.* **277**, 2747–2753 (2010).
- Kitchen, D. M., Seyfarth, R. M., Fischer, J. & Cheney, D. L. Loud calls as indicators of dominance in male baboons (*Papio cynocephalus ursinus*). *Behav. Ecol. Sociobiol.* **53**, 374–384 (2003).
- Puts, D. A. *et al.* Sexual selection on male vocal fundamental frequency in humans and other anthropoids. *Proc. R. Soc. B Biol. Sci.* **283**, 0–7 (2016).
- Rosenfield, K. A., Sorokowska, A., Sorokowski, P. & Puts, D. A. Sexual selection for low male voice pitch among Amazonian forager-horticulturists. *Evol. Hum. Behav.* **41**, 3–11 (2020).
- Titze, I. R. *Principles of Voice Production* (National Center for Voice and Speech, 2000).
- Markova, D. *et al.* Age- and sex-related variations in vocal-tract morphology and voice acoustics during adolescence. *Horm. Behav.* **81**, 84–96 (2016).
- Puts, D. A., Apicella, C. L. & Cárdenas, R. A. Masculine voices signal men's threat potential in forager and industrial societies. *Proc. R. Soc. B Biol. Sci.* **279**, 601–609 (2012).
- Puts, D. A. *et al.* Does men's voice pitch signal formidability? A reply to Feinberg. *Trends Ecol. Evol.* **34**, 189–190 (2019).
- Feinberg, D. R., Jones, B. C., Little, A. C., Burt, D. M. & Perrett, D. I. Manipulations of fundamental and formant frequencies influence the attractiveness of human male voices. *Anim. Behav.* **69**, 561–568 (2005).
- Puts, D. A., Gaulin, S. J. C. & Verdolini, K. Dominance and the evolution of sexual dimorphism in human voice pitch. *Evol. Hum. Behav.* **27**, 283–296 (2006).
- Puts, D. A., Hodges, C. R., Cárdenas, R. A. & Gaulin, S. J. C. Men's voices as dominance signals: vocal fundamental and formant frequencies influence dominance attributions among men. *Evol. Hum. Behav.* **28**, 340–344 (2007).
- Saxton, T. K., Mackey, L. L., McCarty, K. & Neave, N. A lover or a fighter? Opposing sexual selection pressures on men's vocal pitch and facial hair. *Behav. Ecol.* **27**, 512–519 (2016).
- Cheng, J. T., Tracy, J. L., Ho, S. & Henrich, J. Listen, follow me: dynamic vocal signals of dominance predict emergent social rank in humans. *J. Exp. Psychol. Gen.* **145**, 1–12 (2016).
- Aung, T. & Puts, D. A. Voice pitch: a window into the communication of social power. *Curr. Opin. Psychol.* **33**, 154–161 (2020).
- Rendall, D., Vokey, J. R. & Nemeth, C. Lifting the curtain on the Wizard of Oz: biased voice-based impressions of speaker size. *J. Exp. Psychol. Hum. Percept. Perform.* **33**, 1208–1219 (2007).
- Sell, A. *et al.* Adaptations in humans for assessing physical strength from the voice. *Proc. R. Soc. B Biol. Sci.* **277**, 3509–3518 (2010).
- Feinberg, D. R., Jones, B. C. & Armstrong, M. M. No evidence that men's voice pitch signals formidability. *Trends Ecol. Evol.* **34**, 190–192 (2019).
- Armstrong, M. M., Lee, A. J. & Feinberg, D. R. A house of cards: Bias in perception of body size mediates the relationship between voice pitch and perceptions of dominance. *Anim. Behav.* **147**, 43–51 (2019).
- Dawkins, R. & Krebs, J. R. Animal signals: Information or manipulation? In *Behavioral Ecology: An Evolutionary Approach* (eds Krebs, J. R. & Davies, N. B.) 282–309 (Sinauer Associates, Inc., Sunderland, 1978).
- Searcy, W. A. & Nowicki, S. *The Evolution of Animal Communication: Reliability and Deception in Signaling Systems* (Princeton University Press, Princeton, 2005).
- Raine, J., Id, K. P., Bond, R., Simner, J. & Reby, D. Human roars communicate upper-body strength more effectively than do screams or aggressive and distressed speech. *PLoS ONE* **14**, e0213034 (2019).
- Han, C. *et al.* No clear evidence for correlations between handgrip strength and sexually dimorphic acoustic properties of voices. *Am. J. Hum. Biol.* **30**, 1–4 (2018).
- Sell, A. *et al.* Human adaptations for the visual assessment of strength and fighting ability from the body and face. *Proc. R. Soc. B Biol. Sci.* **276**, 575–584 (2009).
- Kordsmeyer, T. L., Hunt, J., Puts, D. A., Ostner, J. & Penke, L. The relative importance of intra- and intersexual selection on human male sexually dimorphic traits. *Evol. Hum. Behav.* **39**, 424–436 (2018).
- Šebesta, P., Třebický, V., Fialová, J. & Havlíček, J. Roar of a champion: Loudness and voice pitch predict perceived fighting ability in MMA fighters. *Front. Psychol.* **10**, (2019).
- Arnocky, S., Hodges-Simeon, C. R., Ouellette, D. & Albert, G. Do men with more masculine voices have better immunocompetence? *Evol. Hum. Behav.* **39**, 602–610 (2018).
- Skrinda, I. *et al.* Body height, immunity, facial and vocal attractiveness in young men. *Naturwissenschaften* **101**, 1017–1025 (2014).
- Albert, G., Arnocky, S., Puts, D. A. & Hodges-Simeon, C. R. Can listeners assess men's self-reported health from their voice? *Evol. Hum. Behav.* <https://doi.org/10.1016/j.evolhumbehav.2020.08.001> (2020).
- Pisanski, K. *et al.* Vocal indicators of body size in men and women: a meta-analysis. *Anim. Behav.* **95**, 89–99 (2014).
- Schild, C. *et al.* Linking human male vocal parameters to perceptions, body morphology, strength and hormonal profiles in contexts of sexual selection. *Sci. Rep.* <https://doi.org/10.1038/s41598-020-77940-z> (2020).

34. Rantala, M. J. *et al.* Evidence for the stress-linked immunocompetence handicap hypothesis in humans. *Nat. Commun.* **3**, 694–695 (2012).
35. Hodges-Simeon, C. R., Gurven, M. & Gaulin, S. J. C. The low male voice is a costly signal of phenotypic quality among Bolivian adolescents. *Evol. Hum. Behav.* **36**, 294–302 (2015).
36. Sorokowski, P. *et al.* Voice of authority: professionals lower their vocal frequencies when giving expert advice. *J. Nonverbal. Behav.* **43**, 257–269 (2019).
37. Moon, K. R., Chung, S. M., Park, H. S. & Kim, H. S. Materials of acoustic analysis: sustained vowel versus sentence. *J. Voice* **26**, 563–565 (2012).
38. Pisanski, K., Cartei, V., McGettigan, C., Raine, J. & Reby, D. Voice modulation: a window into the origins of human vocal control?. *Trends Cogn. Sci.* **20**, 304–318 (2016).
39. Hodges-Simeon, C. R., Gurven, M., Puts, D. A. & Gaulin, S. J. C. Vocal fundamental and formant frequencies are honest signals of threat potential in peripubertal males. *Behav. Ecol.* **25**, 984–988 (2014).
40. Reby, D. & McComb, K. Anatomical constraints generate honesty: acoustic cues to age and weight in the roars of red deer stags. *Anim. Behav.* **65**, 519e530 (2003).
41. Raiche, G. nFactors: an R package for parallel analysis and non graphical solutions to the Cattell scree test. R package version 2.3.3.1 (2010).
42. Revelle, W. Psych: procedures for personality and psychological research version 1.8.12 (2018).
43. Dattalo, P. *Analysis of Multiple Dependent Variables* (Oxford University Press, Oxford, 2013).
44. Perelman, P. *et al.* A molecular phylogeny of living primates. *PLoS Genet.* **7**, 1–17 (2011).
45. Feinberg, D. R., Jones, B. C. & Armstrong, M. M. Sensory exploitation, sexual dimorphism, and human voice pitch. *Trends Ecol. Evol.* **33**, 901–903 (2018).
46. Modig, A. O. Effects of body size and harem size on male reproductive behaviour in the southern elephant seal. *Anim. Behav.* **51**, 1295–1306 (1996).
47. Hodges-Simeon, C. R., Gaulin, S. J. C. & Puts, D. A. Different vocal parameters predict perceptions of dominance and attractiveness. *Hum. Nat.* **21**, 406–427 (2010).
48. Leongómez, J. D., Mileva, V. R., Little, A. C. & Roberts, S. C. Perceived differences in social status between speaker and listener affect the speaker's vocal characteristics. *PLoS One* **12**, (2017).
49. Zhang, J., Hodges-Simeon, C., Gaulin, S. J. C. & Reid, S. A. Pitch lowering enhances men's perceived aggressive intent, not fighting ability. *Evol. Hum. Behav.* <https://doi.org/10.1016/j.evolhumbehav.2020.07.007> (2020).
50. Fraccaro, P. J. *et al.* Faking it: Deliberately altered voice pitch and vocal attractiveness. *Anim. Behav.* **85**, 127–136 (2013).
51. Pisanski, K. *et al.* Volitional exaggeration of body size through fundamental and formant frequency modulation in humans. *Sci. Rep.* **6**, 1–8 (2016).
52. Pisanski, K., Raine, J. & Reby, D. Individual differences in human voice pitch are preserved from speech to screams, roars and pain cries. *R. Soc. Open Sci.* **7**(2), 191642 (2020).
53. Richardson, T. & Gilman, R. T. Left-handedness is associated with greater fighting success in humans. *Sci. Rep.* **9**, 1–16 (2019).
54. Zilioli, S. *et al.* Face of a fighter: bizygomatic width as a cue of formidability. *Aggress. Behav.* **41**, 322–330 (2015).
55. Bowling, D. L. *et al.* Body size and vocalization in primates and carnivores. *Sci. Rep.* **7**, 1–11 (2017).
56. French, A. R. & Smith, T. B. Importance of body size in determining dominance hierarchies among diverse tropical frugivores. *Biotropica* **37**, 96–101 (2005).
57. Darwin, C. *The Descent of Man, and Selection in Relation to Sex* (Princeton University Press, Princeton, 1871).
58. Clutton-Brock, T. Reproductive competition and sexual selection. *Philos. Trans. R. Soc. B Biol. Sci.* **372**, 20160310 (2017).
59. Andersson, M. *Sexual Selection* (Princeton University Press, Princeton, 1994).
60. Wright, E. *et al.* Male body size, dominance rank and strategic use of aggression in a group-living mammal. *Anim. Behav.* **151**, 87–102 (2019).
61. Evans, S., Neave, N. & Wakelin, D. Relationships between vocal characteristics and body size and shape in human males: an evolutionary explanation for a deep male voice. *Biol. Psychol.* **72**, 160–163 (2006).
62. Wolff, S. E. & Puts, D. A. Vocal masculinity is a robust dominance signal in men. *Behav. Ecol. Sociobiol.* **64**, 1673–1683 (2010).
63. Perneger, T. V. What's wrong with Bonferroni adjustments. *BMJ* **316**, 1236–1238 (1998).
64. Gelman, A., Hill, J. & Yajima, M. Why we (usually) don't have to worry about multiple comparisons. *J. Res. Educ. Eff.* **5**, 189–211 (2012).
65. Hill, A. K., Bailey, D. H. & Puts, D. A. Gorillas in our midst? Human sexual dimorphism and contest competition in men. In *On Human Nature: Biology, Psychology, Ethics, Politics, and Religion* (eds. Tibayrenc, M. & Ayala, F. J.) 235–249 (Elsevier Science, Amsterdam, 2016).
66. Holzleitner, I. J. & Perrett, D. I. Perception of strength from 3D faces is linked to facial cues of physique. *Evol. Hum. Behav.* **37**, 217–229 (2016).
67. Třebický, V., Havlíček, J., Roberts, C. S., Little, A. C. & Kleisner, K. Perceived aggressiveness predicts fighting performance in mixed-martial-arts fighters. *Psychol. Sci.* **24**, 1664–1672 (2013).
68. Třebický, V. *et al.* Predictors of fighting ability inferences based on faces. *Front. Psychol.* **9**, 1–11 (2019).
69. Little, A. C., Roberts, S. C. & Kleisner, K. Human perception of fighting ability: facial cues predict winners and losers in mixed martial arts fights. *Behav. Ecol.* **26**, 1470–1475 (2015).
70. Horne, L. Dana White on Anthony Johnson: “three strikes and you're out”. *Bleacher Rep.* <http://bleacherreport.com/articles/1025522-dana-white-on-anthony-johnson-three-strikes-and-youre-out> (2012).
71. Raudenbush, S. W. & Bryk, A. S. *Hierarchical Linear Models: Applications and Data Analysis Methods* (Sage, Thousand Oaks, 2002).
72. Croon, M. A. & Van Veldhoven, M. J. P. M. Predicting group-level outcome variables from variables measured at the individual level: a latent variable multilevel model. *Psychol. Methods* **12**, 45–57 (2007).
73. Lu, J. G., Page-Gould, E. & Xu, N. R. Micro-macromultilevel Modeling [Computer software] (2017).
74. Akinola, M., Page-Gould, E., Mehta, P. H. & Lu, J. G. Collective hormonal profiles predict group performance. *Proc. Natl. Acad. Sci. USA* **113**, 9774–9779 (2016).

Author contributions

S.G., J.T.C., S.Z., and D.P. conceived the study. T.A., S.G., C.M., C.H., and S.R. collected data. J.T.C., S.Z., D.P., and T.A. conceived analysis plan. D.P. supervised the study, and T.A. analyzed the data. T.A. and D.P. wrote the paper with input from co-authors.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-020-79408-6>.

Correspondence and requests for materials should be addressed to D.P.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021