

S-Benzylthiouronium 3-nitrobenzene-sulfonate

Hoong-Kun Fun,^{a*} Samuel Robinson Jebas,^{a‡}
Ibrahim Abdul Razak,^a E. Deepak D'Silva,^b P. S. Patil^b and
S. M. Dharmaprkash^b

^aX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^bDepartment of Studies in Physics, Mangalore University, Mangalagangotri, Mangalore 574 199, India
Correspondence e-mail: hkfun@usm.my


Received 9 May 2008; accepted 28 May 2008

Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(C-C) = 0.004 \text{ \AA}$; R factor = 0.044; wR factor = 0.113; data-to-parameter ratio = 20.5.

In the title compound, $C_8H_{11}N_2S^+ \cdot C_6H_4NO_5S^-$, the asymmetric unit is composed of two crystallographically independent *S*-benzylthiouronium cations and two independent nitrobenzenesulfonate anions. An intramolecular hydrogen bond generates an *S*(5)*S*(5) ring motif. The crystal packing is stabilized by intramolecular $C-H \cdots O$ and intermolecular $C-H \cdots O$, $N-H \cdots O$ and $N-H \cdots S$ hydrogen bonds which, along with short $S \cdots O$ [3.034 (2) \AA] and $N \cdots O$ [2.796 (3) \AA] contacts, form a two-dimensional network parallel to the *ab* plane.

Related literature

For related literature on nonlinear optical materials, see: Chantrapromma *et al.* (2005, 2006); Fun *et al.* (2006); Patil, Dharmaprakash *et al.* (2007); Patil, Fun *et al.* (2007). For bond-length data, see: Allen *et al.* (1987). For graph-set analysis of hydrogen bonding, see: Bernstein *et al.* (1995).



Experimental

Crystal data

$C_8H_{11}N_2S^+ \cdot C_6H_4NO_5S^-$
 $M_r = 369.41$
Triclinic, $P\bar{1}$

$a = 6.0397 (1) \text{ \AA}$
 $b = 7.7856 (1) \text{ \AA}$
 $c = 17.4680 (2) \text{ \AA}$

‡ Permanent address: Department of Physics, Karunya University, Karunya Nagar, Coimbatore 641 114, India.

$\alpha = 81.366 (1)^\circ$
 $\beta = 89.322 (1)^\circ$
 $\gamma = 87.057 (1)^\circ$
 $V = 811.01 (2) \text{ \AA}^3$
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.36 \text{ mm}^{-1}$
 $T = 100.0 (1) \text{ K}$
 $0.32 \times 0.19 \times 0.04 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2005)
 $T_{\min} = 0.894$, $T_{\max} = 0.986$

19026 measured reflections
8874 independent reflections
7378 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.044$
 $wR(F^2) = 0.112$
 $S = 1.03$
8874 reflections
433 parameters
3 restraints

H-atom parameters constrained
 $\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$
Absolute structure: Flack (1983),
4108 Friedel pairs
Flack parameter: -0.03 (5)

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
N3—H1N3 \cdots O4 ⁱ	0.86	1.97	2.796 (3)	162
N3—H1N3 \cdots S1 ⁱ	0.86	2.84	3.694 (3)	170
N3—H2N3 \cdots O3 ⁱⁱ	0.86	2.05	2.898 (3)	171
N4—H1N4 \cdots O3 ⁱ	0.86	2.26	3.080 (3)	160
N4—H2N4 \cdots O5 ⁱⁱⁱ	0.86	2.37	3.126 (3)	146
N5—H1N5 \cdots O10 ⁱ	0.86	1.91	2.764 (3)	176
N5—H1N5 \cdots S2 ^j	0.86	2.85	3.642 (3)	154
N5—H2N5 \cdots O9 ⁱⁱ	0.86	1.94	2.783 (3)	168
N6—H1N6 \cdots O1 ⁱⁱ	0.86	2.27	3.072 (3)	156
N6—H2N6 \cdots O8 ⁱⁱⁱ	0.86	2.07	2.787 (3)	141
C6—H6A \cdots O4	0.93	2.56	2.900 (3)	102
C8—H8A \cdots O10	0.93	2.57	2.896 (4)	101
C19—H19B \cdots O4 ⁱⁱⁱ	0.97	2.51	3.331 (4)	142
C27—H27B \cdots O8 ⁱⁱⁱ	0.97	2.53	3.259 (4)	132

Symmetry codes: (i) $x + 1, y + 1, z$; (ii) $x, y + 1, z$; (iii) $x + 1, y, z$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *APEX2*; data reduction: *SAINT* (Bruker, 2005); program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2003).

HKF and SRJ thank the Malaysian Government and Universiti Sains Malaysia for Science Fund grant No. 305/PFIZIK/613312. HKF and IAR also thank the Malaysian Government and Universiti Sains Malaysia for FRGS grant No. 203/PFIZIK/671064. SRJ thanks the Universiti Sains Malaysia for a post-doctoral research fellowship. This work was supported by the Department of Science and Technology (DST) and the Government of India (grant No. SR/S2/LOP-17/2006).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SU2055).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bruker (2005). *APEX2, SAINT and SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chantrapromma, S., Jindawong, B., Fun, H.-K., Anjum, S. & Karalai, C. (2005). *Acta Cryst. E61*, o2096–o2098.
- Chantrapromma, S., Ruanwas, P., Jindawong, B., Razak, I. A. & Fun, H.-K. (2006). *Acta Cryst. E62*, o875–o877.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Fun, H.-K., Rodwatcharapiban, P., Jindawong, B. & Chantrapromma, S. (2006). *Acta Cryst. E62*, o2725–o2727.
- Patil, P. S., Dharmaprakash, S. M., Ramakrishna, K., Fun, H.-K., Sai Santosh Kumar, R. & Rao, D. N. (2007). *J. Cryst. Growth*, **303**, 520–524.
- Patil, P. S., Fun, H.-K., Chantrapromma, S. & Dharmaprakash, S. M. (2007). *Acta Cryst. E63*, o2497–o2498.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2008). E64, o1195–o1196 [doi:10.1107/S1600536808016115]

S-Benzylthiouronium 3-nitrobenzenesulfonate

H.-K. Fun, S. R. Jebas, I. A. Razak, E. Deepak D'Silva, P. S. Patil and S. M. Dharmaprakash

Comment

In continuation of our research on nonlinear optical (NLO) materials (Chantrapromma *et al.*, 2005, 2006; Fun *et al.*, 2006; Patil, Dharmaprakash *et al.* (2007); Patil, Fun *et al.* (2007), the crystal structure determination of the title compound, (I), was undertaken in order to obtain detailed information on its three-dimensional structure and crystal packing. Since the title compound crystallizes in a non-centrosymmetric space group, it should exhibit second-order nonlinear optical properties.

In the asymmetric unit of (I) there are two crystallographically independent s-benzylthiuronium cations and two independent m-nitrobenzene sulfonate anions (Fig. 1). Bond lengths and angles are found to have normal values (Allen *et al.*, 1987). The dihedral angle formed by the mean plane of ring (C13–C18) with the mean planes through rings (C21–C26), (C1–C6) and (C7–C12) are 57.6 (2) $^{\circ}$, 49.9 (2) $^{\circ}$ and 46.98 (1) $^{\circ}$, respectively. The dihedral angle formed by the mean planes of rings (C1–C6) and (C7–C12) are 3.07 (1) $^{\circ}$, indicating that they are almost coplanar.

An intramolecular hydrogen bond generates a S(5)S(5) ring motif (Bernstein *et al.*, 1995). The crystal packing (Fig. 2) is stabilized by intramolecular C—H···O, intermolecular C—H···O, N—H···O and N—H···S hydrogen bonds which together with short S···O and N···O contacts, in the range 2.796 (3)–3.282 (3) Å [symmetry code: x, 1+y, z; 1+x, 1+y, z], form a two-dimensional network parallel to the ab plane.

Experimental

Compound (I) was synthesized by mixing solutions of the sodium salt of *m*-nitrobenzene sulfonic acid (0.5 g) in 5 ml of distilled water with 5 drops of 0.1 N HCl and *S*-benzylthiuronium chloride (1 g) in 5 ml of distilled water. The mixing immediately yielded a precipitate when the reaction container was placed in ice cold water. The resulting precipitate was filtered and dried. Crystals suitable for single-crystal X-ray diffraction were grown by slow evaporation of an ethanol solution at room temperature.

Refinement

H atoms were positioned geometrically [C—H = 0.93 Å; N—H = 0.86 Å and CH₂ = 0.97 Å] and refined using a riding model with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C, N})$.

Figures




Fig. 1. The molecular structure of compound (I) showing 50% probability displacement ellipsoids and the atom numbering scheme.

supplementary materials




Fig. 2. The crystal packing of compound (I) viewed down the *c* axis, showing the two-dimensional network. Short intra and intermolecular contacts and hydrogen bonds are shown as dashed lines.

(5-Methylpyrazine-2-carboxylato)diphenyltin(IV)

Crystal data

$C_8H_{11}N_2S^+ \cdot C_6H_4NO_5S^-$	$Z = 2$
$M_r = 369.41$	$F_{000} = 384$
Triclinic, $P\bar{1}$	$D_x = 1.513 \text{ Mg m}^{-3}$
Hall symbol: $P\bar{1}$	Mo $K\alpha$ radiation
$a = 6.0397 (1) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 7.7856 (1) \text{ \AA}$	Cell parameters from 4354 reflections
$c = 17.4680 (2) \text{ \AA}$	$\theta = 2.8\text{--}34.6^\circ$
$\alpha = 81.366 (1)^\circ$	$\mu = 0.36 \text{ mm}^{-1}$
$\beta = 89.322 (1)^\circ$	$T = 100.0 (1) \text{ K}$
$\gamma = 87.057 (1)^\circ$	Plate, colourless
$V = 811.006 (19) \text{ \AA}^3$	$0.32 \times 0.19 \times 0.04 \text{ mm}$

Data collection

Bruker SMART APEXII CCD area-detector diffractometer	8874 independent reflections
Radiation source: fine-focus sealed tube	7378 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.043$
$T = 100.0(1) \text{ K}$	$\theta_{\max} = 30.1^\circ$
φ and ω scans	$\theta_{\min} = 1.2^\circ$
Absorption correction: multi-scan (SADABS; Bruker, 2005)	$h = -7 \rightarrow 8$
$T_{\min} = 0.894$, $T_{\max} = 0.986$	$k = -10 \rightarrow 10$
19026 measured reflections	$l = -24 \rightarrow 24$

Refinement

Refinement on F^2	Hydrogen site location: inferred from neighbouring sites
Least-squares matrix: full	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.0534P)^2]$
$wR(F^2) = 0.112$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.03$	$(\Delta/\sigma)_{\max} < 0.001$
	$\Delta\rho_{\max} = 0.38 \text{ e \AA}^{-3}$

8874 reflections	$\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$
433 parameters	Extinction correction: none
3 restraints	Absolute structure: Flack (1983), with how many Friedel pairs?
Primary atom site location: structure-invariant direct methods	Flack parameter: -0.03 (5)
Secondary atom site location: difference Fourier map	

Special details

Experimental. The data was collected with the Oxford Cyrosystem Cobra low-temperature attachment.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.37546 (10)	0.53723 (8)	0.82205 (4)	0.01444 (14)
O1	0.4811 (4)	0.4607 (3)	0.47221 (12)	0.0252 (5)
O2	0.1931 (4)	0.4114 (3)	0.54531 (12)	0.0241 (5)
O3	0.4577 (3)	0.3635 (3)	0.85663 (11)	0.0180 (4)
O4	0.1377 (3)	0.5430 (3)	0.80849 (11)	0.0184 (4)
O5	0.4448 (3)	0.6737 (3)	0.86311 (12)	0.0206 (4)
N1	0.3820 (4)	0.4617 (3)	0.53419 (14)	0.0193 (5)
C1	0.4988 (4)	0.5723 (3)	0.72854 (15)	0.0147 (5)
C2	0.7020 (4)	0.6478 (4)	0.71677 (16)	0.0167 (5)
H2A	0.7710	0.6903	0.7569	0.020*
C3	0.8017 (5)	0.6590 (4)	0.64337 (18)	0.0207 (6)
H3A	0.9384	0.7085	0.6351	0.025*
C4	0.7004 (5)	0.5977 (4)	0.58323 (17)	0.0184 (6)
H4A	0.7673	0.6044	0.5347	0.022*
C5	0.4953 (5)	0.5255 (4)	0.59719 (16)	0.0163 (5)
C6	0.3912 (5)	0.5112 (3)	0.66886 (15)	0.0154 (5)
H6A	0.2539	0.4623	0.6767	0.018*
S2	0.31950 (11)	-0.08507 (8)	0.41082 (4)	0.01834 (15)
O6	0.5275 (5)	0.1046 (3)	0.73759 (13)	0.0379 (6)
O7	0.2256 (5)	0.0097 (4)	0.69863 (14)	0.0493 (8)
O8	0.1789 (4)	0.0611 (3)	0.37490 (12)	0.0231 (5)
O9	0.4738 (4)	-0.1484 (3)	0.35681 (13)	0.0301 (5)
O10	0.1907 (4)	-0.2203 (3)	0.45482 (13)	0.0282 (5)
N2	0.4156 (5)	0.0558 (4)	0.68796 (15)	0.0285 (6)
C7	0.4798 (5)	-0.0078 (4)	0.48202 (16)	0.0168 (5)

supplementary materials

C8	0.3892 (5)	-0.0087 (4)	0.55576 (17)	0.0185 (6)
H8A	0.2493	-0.0502	0.5677	0.022*
C9	0.5124 (5)	0.0534 (4)	0.61044 (17)	0.0215 (6)
C10	0.7218 (5)	0.1167 (4)	0.59464 (18)	0.0227 (6)
H10A	0.8017	0.1578	0.6327	0.027*
C11	0.8082 (5)	0.1169 (4)	0.52112 (19)	0.0239 (6)
H11A	0.9479	0.1590	0.5094	0.029*
C12	0.6880 (5)	0.0546 (4)	0.46421 (17)	0.0195 (6)
H12A	0.7471	0.0549	0.4148	0.023*
S3	0.77990 (11)	0.92863 (9)	0.89784 (5)	0.02185 (17)
N3	0.9177 (4)	1.2377 (3)	0.85658 (14)	0.0211 (5)
H1N3	1.0112	1.3173	0.8448	0.025*
H2N3	0.7778	1.2643	0.8544	0.025*
N4	1.1997 (4)	1.0283 (3)	0.88215 (14)	0.0199 (5)
H1N4	1.2973	1.1048	0.8708	0.024*
H2N4	1.2408	0.9209	0.8963	0.024*
C13	0.8638 (5)	0.8126 (4)	1.08658 (18)	0.0229 (6)
H13A	0.7231	0.8611	1.0730	0.027*
C14	0.9388 (6)	0.8053 (4)	1.16156 (19)	0.0292 (7)
H14A	0.8479	0.8490	1.1982	0.035*
C15	1.1470 (6)	0.7337 (4)	1.18259 (18)	0.0294 (7)
H15A	1.1961	0.7290	1.2332	0.035*
C16	1.2827 (6)	0.6688 (4)	1.12789 (19)	0.0274 (7)
H16A	1.4234	0.6208	1.1417	0.033*
C17	1.2086 (5)	0.6754 (4)	1.05282 (17)	0.0217 (6)
H17A	1.3000	0.6312	1.0164	0.026*
C18	0.9993 (5)	0.7473 (4)	1.03121 (17)	0.0178 (6)
C19	0.9156 (5)	0.7358 (4)	0.95108 (18)	0.0216 (6)
H19A	0.8127	0.6432	0.9556	0.026*
H19B	1.0406	0.7015	0.9204	0.026*
C20	0.9890 (5)	1.0752 (4)	0.87791 (15)	0.0165 (5)
S4	0.70823 (11)	0.42238 (9)	0.29521 (4)	0.02029 (15)
N5	0.8675 (4)	0.6477 (3)	0.37270 (14)	0.0205 (5)
H1N5	0.9647	0.6932	0.3980	0.025*
H2N5	0.7379	0.6980	0.3650	0.025*
N6	1.1136 (4)	0.4218 (3)	0.35682 (14)	0.0209 (5)
H1N6	1.2132	0.4650	0.3819	0.025*
H2N6	1.1425	0.3262	0.3388	0.025*
C21	0.7614 (5)	0.3251 (4)	0.09917 (17)	0.0211 (6)
H21A	0.8905	0.3852	0.0900	0.025*
C22	0.6228 (5)	0.3107 (4)	0.03818 (17)	0.0241 (6)
H22A	0.6595	0.3609	-0.0118	0.029*
C23	0.4305 (5)	0.2224 (4)	0.05098 (18)	0.0258 (7)
H23A	0.3378	0.2128	0.0099	0.031*
C24	0.3766 (5)	0.1484 (4)	0.12517 (19)	0.0266 (7)
H24A	0.2463	0.0897	0.1341	0.032*
C25	0.5157 (5)	0.1608 (4)	0.18650 (17)	0.0228 (6)
H25A	0.4794	0.1091	0.2362	0.027*
C26	0.7088 (5)	0.2503 (4)	0.17390 (16)	0.0180 (6)

C27	0.8588 (5)	0.2713 (4)	0.23962 (17)	0.0242 (6)
H27A	0.9976	0.3182	0.2200	0.029*
H27B	0.8905	0.1601	0.2718	0.029*
C28	0.9169 (5)	0.5019 (4)	0.34598 (16)	0.0175 (6)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0132 (3)	0.0156 (3)	0.0147 (3)	-0.0010 (2)	-0.0006 (2)	-0.0028 (2)
O1	0.0274 (12)	0.0322 (12)	0.0163 (10)	0.0034 (9)	0.0009 (9)	-0.0061 (9)
O2	0.0218 (11)	0.0285 (12)	0.0236 (11)	-0.0070 (9)	-0.0016 (9)	-0.0066 (9)
O3	0.0197 (10)	0.0164 (10)	0.0169 (9)	0.0018 (8)	-0.0002 (8)	0.0000 (8)
O4	0.0138 (10)	0.0201 (10)	0.0209 (10)	-0.0010 (8)	0.0001 (8)	-0.0019 (8)
O5	0.0202 (10)	0.0223 (10)	0.0222 (10)	-0.0061 (8)	0.0008 (8)	-0.0111 (8)
N1	0.0229 (14)	0.0157 (12)	0.0188 (12)	0.0024 (10)	-0.0025 (10)	-0.0019 (9)
C1	0.0144 (13)	0.0126 (13)	0.0167 (13)	0.0009 (10)	-0.0022 (10)	-0.0018 (10)
C2	0.0130 (13)	0.0167 (13)	0.0199 (13)	-0.0007 (10)	-0.0032 (10)	-0.0012 (11)
C3	0.0128 (13)	0.0199 (14)	0.0273 (15)	-0.0018 (10)	0.0006 (11)	0.0034 (12)
C4	0.0167 (14)	0.0195 (14)	0.0182 (13)	0.0018 (11)	0.0020 (11)	-0.0008 (11)
C5	0.0176 (13)	0.0149 (13)	0.0166 (12)	0.0015 (10)	-0.0048 (10)	-0.0031 (10)
C6	0.0141 (13)	0.0144 (13)	0.0170 (13)	-0.0010 (10)	-0.0017 (10)	-0.0002 (10)
S2	0.0198 (4)	0.0174 (3)	0.0176 (3)	0.0013 (3)	-0.0044 (3)	-0.0023 (3)
O6	0.0591 (18)	0.0348 (14)	0.0216 (12)	-0.0018 (12)	-0.0106 (11)	-0.0094 (10)
O7	0.0337 (15)	0.092 (2)	0.0243 (13)	-0.0050 (15)	0.0061 (11)	-0.0138 (14)
O8	0.0235 (11)	0.0209 (11)	0.0232 (10)	0.0045 (8)	-0.0055 (9)	0.0004 (9)
O9	0.0310 (13)	0.0378 (13)	0.0224 (11)	0.0120 (10)	-0.0038 (9)	-0.0118 (10)
O10	0.0330 (13)	0.0221 (11)	0.0284 (11)	-0.0104 (9)	-0.0110 (10)	0.0035 (9)
N2	0.0390 (18)	0.0263 (14)	0.0199 (13)	0.0067 (12)	-0.0021 (12)	-0.0052 (11)
C7	0.0169 (14)	0.0150 (13)	0.0175 (13)	0.0034 (10)	-0.0031 (10)	-0.0003 (10)
C8	0.0172 (14)	0.0171 (13)	0.0205 (13)	0.0024 (11)	-0.0019 (11)	-0.0019 (11)
C9	0.0232 (16)	0.0201 (15)	0.0206 (14)	0.0059 (11)	-0.0056 (11)	-0.0036 (11)
C10	0.0206 (15)	0.0202 (15)	0.0276 (15)	0.0037 (11)	-0.0091 (12)	-0.0055 (12)
C11	0.0149 (14)	0.0199 (15)	0.0363 (17)	-0.0018 (11)	-0.0032 (12)	-0.0021 (13)
C12	0.0178 (14)	0.0164 (14)	0.0235 (14)	0.0011 (11)	-0.0009 (11)	-0.0013 (11)
S3	0.0150 (4)	0.0199 (4)	0.0292 (4)	-0.0045 (3)	-0.0034 (3)	0.0025 (3)
N3	0.0147 (12)	0.0189 (12)	0.0294 (13)	-0.0045 (9)	-0.0001 (10)	-0.0012 (10)
N4	0.0142 (12)	0.0211 (13)	0.0241 (12)	-0.0025 (9)	0.0002 (10)	-0.0016 (10)
C13	0.0206 (15)	0.0208 (15)	0.0268 (15)	-0.0007 (11)	0.0053 (12)	-0.0024 (12)
C14	0.0363 (19)	0.0219 (16)	0.0287 (16)	-0.0027 (13)	0.0097 (14)	-0.0010 (13)
C15	0.038 (2)	0.0278 (18)	0.0200 (15)	-0.0090 (14)	-0.0013 (13)	0.0053 (13)
C16	0.0228 (16)	0.0265 (17)	0.0298 (17)	-0.0019 (12)	-0.0054 (13)	0.0067 (13)
C17	0.0192 (15)	0.0220 (15)	0.0224 (14)	0.0022 (11)	0.0039 (12)	-0.0002 (12)
C18	0.0176 (14)	0.0130 (13)	0.0224 (14)	-0.0041 (10)	0.0024 (11)	-0.0004 (11)
C19	0.0195 (15)	0.0150 (14)	0.0298 (16)	-0.0016 (11)	0.0014 (12)	-0.0017 (12)
C20	0.0156 (13)	0.0207 (14)	0.0136 (12)	-0.0031 (10)	0.0007 (10)	-0.0027 (10)
S4	0.0170 (4)	0.0252 (4)	0.0202 (3)	-0.0003 (3)	-0.0022 (3)	-0.0082 (3)
N5	0.0177 (12)	0.0222 (13)	0.0228 (12)	-0.0013 (9)	-0.0009 (10)	-0.0067 (10)
N6	0.0187 (13)	0.0233 (13)	0.0212 (12)	0.0006 (10)	-0.0056 (10)	-0.0050 (10)

supplementary materials

C21	0.0222 (15)	0.0188 (14)	0.0224 (14)	-0.0025 (11)	0.0019 (12)	-0.0027 (12)
C22	0.0329 (18)	0.0236 (16)	0.0157 (13)	-0.0003 (13)	-0.0014 (12)	-0.0027 (12)
C23	0.0283 (17)	0.0249 (16)	0.0265 (16)	0.0023 (13)	-0.0125 (13)	-0.0116 (13)
C24	0.0242 (16)	0.0241 (16)	0.0335 (17)	-0.0064 (12)	-0.0009 (13)	-0.0087 (13)
C25	0.0269 (16)	0.0221 (15)	0.0185 (14)	-0.0005 (12)	0.0036 (12)	-0.0001 (12)
C26	0.0215 (15)	0.0182 (14)	0.0149 (12)	0.0013 (11)	-0.0023 (11)	-0.0047 (11)
C27	0.0256 (16)	0.0286 (16)	0.0197 (14)	0.0069 (12)	-0.0035 (12)	-0.0103 (12)
C28	0.0163 (14)	0.0215 (15)	0.0144 (12)	-0.0026 (11)	0.0008 (11)	-0.0011 (11)

Geometric parameters (Å, °)

S1—O5	1.449 (2)	N4—C20	1.306 (4)
S1—O4	1.456 (2)	N4—H1N4	0.8600
S1—O3	1.461 (2)	N4—H2N4	0.8600
S1—C1	1.776 (3)	C13—C14	1.383 (5)
O1—N1	1.232 (3)	C13—C18	1.396 (4)
O2—N1	1.229 (3)	C13—H13A	0.9300
N1—C5	1.464 (4)	C14—C15	1.380 (5)
C1—C2	1.388 (4)	C14—H14A	0.9300
C1—C6	1.390 (4)	C15—C16	1.387 (5)
C2—C3	1.403 (4)	C15—H15A	0.9300
C2—H2A	0.9300	C16—C17	1.383 (4)
C3—C4	1.379 (4)	C16—H16A	0.9300
C3—H3A	0.9300	C17—C18	1.389 (4)
C4—C5	1.390 (4)	C17—H17A	0.9300
C4—H4A	0.9300	C18—C19	1.510 (4)
C5—C6	1.386 (4)	C19—H19A	0.9700
C6—H6A	0.9300	C19—H19B	0.9700
S2—O9	1.440 (2)	S4—C28	1.740 (3)
S2—O8	1.452 (2)	S4—C27	1.835 (3)
S2—O10	1.460 (2)	N5—C28	1.310 (4)
S2—C7	1.778 (3)	N5—H1N5	0.8600
O6—N2	1.222 (4)	N5—H2N5	0.8600
O7—N2	1.224 (4)	N6—C28	1.315 (4)
N2—C9	1.471 (4)	N6—H1N6	0.8600
C7—C12	1.386 (4)	N6—H2N6	0.8600
C7—C8	1.393 (4)	C21—C22	1.384 (4)
C8—C9	1.376 (4)	C21—C26	1.387 (4)
C8—H8A	0.9300	C21—H21A	0.9300
C9—C10	1.390 (4)	C22—C23	1.379 (5)
C10—C11	1.380 (5)	C22—H22A	0.9300
C10—H10A	0.9300	C23—C24	1.379 (4)
C11—C12	1.396 (4)	C23—H23A	0.9300
C11—H11A	0.9300	C24—C25	1.387 (4)
C12—H12A	0.9300	C24—H24A	0.9300
S3—C20	1.742 (3)	C25—C26	1.388 (4)
S3—C19	1.806 (3)	C25—H25A	0.9300
N3—C20	1.316 (4)	C26—C27	1.504 (4)
N3—H1N3	0.8600	C27—H27A	0.9700

N3—H2N3	0.8600	C27—H27B	0.9700
O5—S1—O4	113.68 (12)	C14—C13—C18	120.0 (3)
O5—S1—O3	113.38 (12)	C14—C13—H13A	120.0
O4—S1—O3	111.39 (12)	C18—C13—H13A	120.0
O5—S1—C1	107.07 (12)	C15—C14—C13	120.7 (3)
O4—S1—C1	105.35 (12)	C15—C14—H14A	119.6
O3—S1—C1	105.17 (12)	C13—C14—H14A	119.6
O2—N1—O1	122.9 (2)	C14—C15—C16	119.6 (3)
O2—N1—C5	118.8 (2)	C14—C15—H15A	120.2
O1—N1—C5	118.3 (2)	C16—C15—H15A	120.2
C2—C1—C6	121.3 (3)	C17—C16—C15	120.0 (3)
C2—C1—S1	121.1 (2)	C17—C16—H16A	120.0
C6—C1—S1	117.5 (2)	C15—C16—H16A	120.0
C1—C2—C3	118.9 (3)	C16—C17—C18	120.7 (3)
C1—C2—H2A	120.5	C16—C17—H17A	119.6
C3—C2—H2A	120.5	C18—C17—H17A	119.6
C4—C3—C2	121.1 (3)	C17—C18—C13	119.0 (3)
C4—C3—H3A	119.4	C17—C18—C19	119.0 (3)
C2—C3—H3A	119.4	C13—C18—C19	121.7 (3)
C3—C4—C5	118.1 (3)	C18—C19—S3	117.6 (2)
C3—C4—H4A	121.0	C18—C19—H19A	107.9
C5—C4—H4A	121.0	S3—C19—H19A	107.9
C6—C5—C4	122.7 (3)	C18—C19—H19B	107.9
C6—C5—N1	117.9 (2)	S3—C19—H19B	107.9
C4—C5—N1	119.4 (2)	H19A—C19—H19B	107.2
C5—C6—C1	117.8 (3)	N4—C20—N3	122.3 (3)
C5—C6—H6A	121.1	N4—C20—S3	123.1 (2)
C1—C6—H6A	121.1	N3—C20—S3	114.6 (2)
O9—S2—O8	112.40 (13)	C28—S4—C27	103.46 (14)
O9—S2—O10	113.93 (15)	C28—N5—H1N5	120.0
O8—S2—O10	111.89 (14)	C28—N5—H2N5	120.0
O9—S2—C7	106.75 (14)	H1N5—N5—H2N5	120.0
O8—S2—C7	106.93 (13)	C28—N6—H1N6	120.0
O10—S2—C7	104.21 (13)	C28—N6—H2N6	120.0
O6—N2—O7	123.8 (3)	H1N6—N6—H2N6	120.0
O6—N2—C9	118.6 (3)	C22—C21—C26	120.4 (3)
O7—N2—C9	117.6 (3)	C22—C21—H21A	119.8
C12—C7—C8	120.9 (3)	C26—C21—H21A	119.8
C12—C7—S2	120.8 (2)	C23—C22—C21	120.4 (3)
C8—C7—S2	118.3 (2)	C23—C22—H22A	119.8
C9—C8—C7	118.1 (3)	C21—C22—H22A	119.8
C9—C8—H8A	120.9	C22—C23—C24	119.6 (3)
C7—C8—H8A	120.9	C22—C23—H23A	120.2
C8—C9—C10	122.5 (3)	C24—C23—H23A	120.2
C8—C9—N2	118.3 (3)	C23—C24—C25	120.4 (3)
C10—C9—N2	119.2 (3)	C23—C24—H24A	119.8
C11—C10—C9	118.4 (3)	C25—C24—H24A	119.8
C11—C10—H10A	120.8	C24—C25—C26	120.2 (3)
C9—C10—H10A	120.8	C24—C25—H25A	119.9

supplementary materials

C10—C11—C12	120.6 (3)	C26—C25—H25A	119.9
C10—C11—H11A	119.7	C21—C26—C25	119.0 (3)
C12—C11—H11A	119.7	C21—C26—C27	119.5 (3)
C7—C12—C11	119.4 (3)	C25—C26—C27	121.4 (3)
C7—C12—H12A	120.3	C26—C27—S4	105.9 (2)
C11—C12—H12A	120.3	C26—C27—H27A	110.6
C20—S3—C19	104.77 (14)	S4—C27—H27A	110.6
C20—N3—H1N3	120.0	C26—C27—H27B	110.6
C20—N3—H2N3	120.0	S4—C27—H27B	110.6
H1N3—N3—H2N3	120.0	H27A—C27—H27B	108.7
C20—N4—H1N4	120.0	N5—C28—N6	121.5 (3)
C20—N4—H2N4	120.0	N5—C28—S4	116.1 (2)
H1N4—N4—H2N4	120.0	N6—C28—S4	122.4 (2)
O5—S1—C1—C2	30.8 (3)	O7—N2—C9—C10	175.5 (3)
O4—S1—C1—C2	152.1 (2)	C8—C9—C10—C11	0.1 (4)
O3—S1—C1—C2	−90.1 (2)	N2—C9—C10—C11	−178.6 (3)
O5—S1—C1—C6	−153.3 (2)	C9—C10—C11—C12	−0.2 (4)
O4—S1—C1—C6	−32.0 (2)	C8—C7—C12—C11	0.1 (4)
O3—S1—C1—C6	85.8 (2)	S2—C7—C12—C11	179.0 (2)
C6—C1—C2—C3	−1.3 (4)	C10—C11—C12—C7	0.1 (4)
S1—C1—C2—C3	174.5 (2)	C18—C13—C14—C15	0.0 (5)
C1—C2—C3—C4	0.5 (4)	C13—C14—C15—C16	0.1 (5)
C2—C3—C4—C5	0.5 (4)	C14—C15—C16—C17	−0.2 (5)
C3—C4—C5—C6	−0.8 (4)	C15—C16—C17—C18	0.3 (5)
C3—C4—C5—N1	179.1 (2)	C16—C17—C18—C13	−0.2 (4)
O2—N1—C5—C6	5.8 (4)	C16—C17—C18—C19	−173.9 (3)
O1—N1—C5—C6	−174.1 (2)	C14—C13—C18—C17	0.1 (4)
O2—N1—C5—C4	−174.1 (3)	C14—C13—C18—C19	173.6 (3)
O1—N1—C5—C4	6.0 (4)	C17—C18—C19—S3	−135.9 (2)
C4—C5—C6—C1	0.0 (4)	C13—C18—C19—S3	50.5 (4)
N1—C5—C6—C1	−179.8 (2)	C20—S3—C19—C18	67.8 (2)
C2—C1—C6—C5	1.0 (4)	C19—S3—C20—N4	17.4 (3)
S1—C1—C6—C5	−174.9 (2)	C19—S3—C20—N3	−163.9 (2)
O9—S2—C7—C12	28.0 (3)	C26—C21—C22—C23	0.2 (5)
O8—S2—C7—C12	−92.5 (2)	C21—C22—C23—C24	0.1 (5)
O10—S2—C7—C12	148.9 (2)	C22—C23—C24—C25	−0.7 (5)
O9—S2—C7—C8	−153.1 (2)	C23—C24—C25—C26	1.0 (5)
O8—S2—C7—C8	86.4 (2)	C22—C21—C26—C25	0.1 (4)
O10—S2—C7—C8	−32.2 (3)	C22—C21—C26—C27	−178.5 (3)
C12—C7—C8—C9	−0.2 (4)	C24—C25—C26—C21	−0.7 (4)
S2—C7—C8—C9	−179.1 (2)	C24—C25—C26—C27	177.9 (3)
C7—C8—C9—C10	0.1 (4)	C21—C26—C27—S4	108.2 (3)
C7—C8—C9—N2	178.8 (3)	C25—C26—C27—S4	−70.4 (3)
O6—N2—C9—C8	177.2 (3)	C28—S4—C27—C26	−159.2 (2)
O7—N2—C9—C8	−3.3 (4)	C27—S4—C28—N5	160.8 (2)
O6—N2—C9—C10	−4.0 (4)	C27—S4—C28—N6	−18.8 (3)

Hydrogen-bond geometry (Å, °)

<i>D—H···A</i>	<i>D—H</i>	<i>H···A</i>	<i>D···A</i>	<i>D—H···A</i>
N3—H1N3···O4 ⁱ	0.86	1.97	2.796 (3)	162
N3—H1N3···S1 ⁱ	0.86	2.84	3.694 (3)	170
N3—H2N3···O3 ⁱⁱ	0.86	2.05	2.898 (3)	171
N4—H1N4···O3 ⁱ	0.86	2.26	3.080 (3)	160
N4—H2N4···O5 ⁱⁱⁱ	0.86	2.37	3.126 (3)	146
N5—H1N5···O10 ⁱ	0.86	1.91	2.764 (3)	176
N5—H1N5···S2 ⁱ	0.86	2.85	3.642 (3)	154
N5—H2N5···O9 ⁱⁱ	0.86	1.94	2.783 (3)	168
N6—H1N6···O1 ⁱⁱⁱ	0.86	2.27	3.072 (3)	156
N6—H2N6···O8 ⁱⁱⁱ	0.86	2.07	2.787 (3)	141
C6—H6A···O4	0.93	2.56	2.900 (3)	102
C8—H8A···O10	0.93	2.57	2.896 (4)	101
C19—H19B···O4 ⁱⁱⁱ	0.97	2.51	3.331 (4)	142
C27—H27B···O8 ⁱⁱⁱ	0.97	2.53	3.259 (4)	132

Symmetry codes: (i) $x+1, y+1, z$; (ii) $x, y+1, z$; (iii) $x+1, y, z$.

supplementary materials

Fig. 1




Fig. 2

