

MACROSCOPIC ANALYSES OF THE EFFECTS OF HYALURONATES AND CORTICOSTEROIDS ON INDUCED OSTEOARTHRITIS IN RABBITS' KNEES

Mauro Batista Albano¹, Leandro Vidigal², Marcello Zaia de Oliveira³, Mario Massatomo Namba⁴, João Luiz Vieira da Silva⁵, Francisco de Assis Pereira Filho⁶, Marcio Alves Barbosa⁷, Elias Marcelo Batista da Silva⁸

ABSTRACT

Objective: To evaluate the effects of intra-articular injections of corticosteroids, native hyaluronic acid and branched-chain hyaluronic acid in experimentally-induced osteoarthritis. *Methods:* 44 rabbits underwent anterior cruciate ligament resection and were then divided into four groups of eleven. Group 1: one intra-articular injection of saline solution per week, for three weeks; Group 2: three injections (one per week) of native hyaluronic acid; Group 3: three injections (one per week) of branched-chain hyaluronic acid; Group 4: two injections of betamethasone with an interval of three weeks. The cartilage of the tibial plateaus was evaluated macroscopically twelve weeks after surgery. Changes to the joint surface were graded as follows: Grade 0: smooth joint surface without relief changes; Grade 1: rough surface without any depressions; Grade 2: similar to

grade 1, but with depressions on the joint surface; and Grade 3: subchondral bone exposure. The statistical analysis consisted of the use of Student's *t* test, chi-square test and analysis of variance (ANOVA). The significance level used was 5%. *Results:* A statistical difference was found between the control group and the three study groups (2, 3 and 4) in relation to the development and severity of arthrosis. However, there was no difference between the groups regarding the drugs studied. *Conclusion:* A similar degree of attenuation of the osteoarthritis process in the rabbits' knees was found with the use of intra-articular injections of low-molecular-weight and high-molecular-weight glycosaminoglycans, and the corticosteroid betamethasone, compared with placebo.

Keywords – Hyaluronic acid. Anterior cruciate ligament; Osteoarthritis; Knee; Rabbits

INTRODUCTION

Osteoarthritis is the most common joint disease among the elderly population. Its prevalence is greater than 10% among individuals over the age of 50 years and it may cause significant functional limitations^(1,2). Among the different types of treatment available today, therapy with intra-articular injections of hyaluronic acid has shown beneficial effects in relation to controlling the symptoms of gonarthrosis⁽³⁾.

Hyaluronic acid, which is a polysaccharide in the glycosaminoglycan family, contributes towards homeostasis in

normal joints and presents lower concentrations and diminished molecular weight in the synovial fluid of joints with osteoarthritis⁽⁴⁻⁶⁾. When administered in the form of intra-articular injections, hyaluronic acid may increase the power of the regenerative effects of endogenous hyaluronic acid on the joint cartilage, by restoring the viscoelasticity of the synovial fluid, contributing towards synthesis of hyaluronic acid by synoviocytes and preventing degradation of proteoglycans and collagen fibers in the extracellular matrix. It stimulates the metabolism, prevents apoptosis of chondrocytes and inhibits chondral degradation and inflammatory responses in

1 – Mestre em Clínica Cirúrgica pela UFPR; Professor do Curso de Especialização em Traumatologia Esportiva e Artroscopia da UFPR; Membro do Serviço de Ortopedia e Traumatologia da UFPR.

2 – Médico Ortopedista; Aluno de Pós-graduação Universidade Federal do Paraná (SOT-UFPR).

3 – Médico Ortopedista; Aluno de Pós-graduação Universidade Federal do Paraná (SOT-UFPR).

4 – Mestre em Clínica Cirúrgica pela UFPR; Professor e Coordenador do Curso de Especialização em Traumatologia Esportiva da UFPR; Membro do Serviço de Ortopedia e Traumatologia da UFPR; Membro Titular da SBOT.

5 – Mestre e Doutor em Clínica Cirúrgica pela UFPR; Professor do Curso de Especialização em Traumatologia Esportiva e Artroscopia da UFPR; Membro do Serviço de Ortopedia e Traumatologia da UFPR; Membro Titular da SBOT.

6 – Professor do Curso de Especialização em Traumatologia Esportiva e Artroscopia da UFPR; Membro do Serviço de Ortopedia e Traumatologia da UFPR; Membro Titular da SBOT.

7 – Professor do Curso de Especialização em Traumatologia Esportiva e Artroscopia da UFPR; Membro do Serviço de Ortopedia e Traumatologia da UFPR; Membro Titular da SBOT.

8 – Professor do Curso de Especialização em Traumatologia Esportiva e Artroscopia da UFPR; Membro do Serviço de Ortopedia e Traumatologia da UFPR; Membro Titular da SBOT.

Trabalho realizado no Serviço de Ortopedia e Traumatologia da Universidade Federal do Paraná (SOT-UFPR) – Serviço do Prof. Dr. Luiz Antonio M. Da Cunha.

Correspondência: – Rua Padre Dehon 1119, ap. 410, V. Hauer – Curitiba, PR. E-mail: albano_mauro@yahoo.com.br

We declare that there is no conflict of interests in this article

the joints⁽⁶⁾. These therapeutic effects from hyaluronic acid are attributed not only to its capacity to ease the symptoms relating to osteoarthritis, but also to its interference with the progression of joint degeneration^(4,5).

Another therapeutic option for managing gonarthrosis is intra-articular injections of corticosteroids. Clinical studies have shown favorable results with regard to pain relief and functional improvement following intra-articular injections of betamethasone disodium phosphate, betamethasone acetate or methylprednisolone^(7,8). Intra-articular administration of corticosteroids at low to moderate doses may reduce the progression of cartilage erosion and the formation of osteophytes⁽⁹⁾. In addition to suppressing the production of metalloproteinases, which have been implicated in the genesis of osteoarthritis, it has the capacity to partially prevent destruction of cells and tissues of the cartilaginous matrix by oxygen free radicals^(10,11).

In view of the widespread occurrence and implications of knee osteoarthritis, the importance of diagnosing and treating it at its initial stages, in order to minimize its consequences, is now understood⁽¹²⁾. Because no interventions capable of inhibiting its evolution are currently available, what remain are options that allow its progression to be reduced. Intra-articular injections of different types of hyaluronic acid and corticosteroids may be used for this purpose.

To evaluate the effects of these substances on gonarthrosis, it was proposed in this investigation to use an experimental model for osteoarthritis that would resemble what is observed in humans. Sectioning of the anterior cruciate ligament of rabbits' knees mimics the morphological and biochemical changes that are observed in human osteoarthritis, which enables accurate reproduction of the results obtained⁽¹³⁻¹⁴⁾.

The aim in this study was to evaluate the effect of intra-articular injections of native hyaluronic acid (Polireumin[®], TRB Pharma, São Paulo, Brazil), branched-chain hyaluronic acid (Synvisc[®], Novartis, São Paulo, Brazil) and the corticosteroid betamethasone (Diprospan[®], Schering-Plough, São Paulo, Brazil), separately and in comparison with each other, in osteoarthritis induced by sectioning the anterior cruciate ligament in rabbits' knees.

METHODS

Forty-four male rabbits of the California breed were used. They were kept housed in cages (two animals per

cage) in a vivarium, before and while performing the procedures. They received standard feed and had free access to water. The lighting conditions were controlled, with light-dark cycles of 12 hours each, and the temperature ($22 \pm 1^\circ\text{C}$), humidity and noise levels were kept stable. These conditions were maintained until the rabbits reached a mean weight of 3.5 kg.

All the animals firstly underwent resection of the anterior cruciate ligament.

The surgical procedure consisted of preoperative anesthesia using 10 mg/kg of ketamine hydrochloride (Dopalen[®]) and 50 mg/kg of xylazine hydrochloride (Anasedan[®]), administered in the same syringe, intramuscularly into the belly of the semimembranosus and semitendinosus muscles of the right hind leg. On the same occasion, injections of 14,400 IU of penicillin and 6 mg of streptomycin (Pentabiótico Veterinário Reforçado[®] - Eurofarma) were administered as antibiotic prophylaxis, and Flunamine[®] (Bayer), at a dose of 2.2 mg/kg intramuscularly, was administered for postoperative analgesia.

The right knee was shaved and antisepsis consisting of polyvinylpyrrolidone (Povidone[®]) was applied. After setting up sterile fields, a medial parapatellar incision was made in the skin and subcutaneous tissue, followed by capsulotomy and lateral luxation of the patella.

The knee was then positioned at maximum flexion, which allowed the anterior cruciate ligament to be viewed. This was sectioned using a no. 15 scalpel blade, the joint was irrigated with isotonic saline solution and then capsulorrhaphy and skin suturing were performed using 4-0 mononylon (Figure 1).

The rabbits continued to be housed in their respective cages after the operation, without restriction on placing weight on the operated leg. They were distributed randomly into four groups of 11 animals each. Three weeks after the surgical procedure, the respective intra-articular injections were started.

Group 1: control, injections of isotonic saline solution at one-week intervals, for three weeks.

Group 2: three infiltrations at one-week intervals, using native hyaluronic acid (Polireumin[®]).

Group 3: three infiltrations at one-week intervals, using branched-chain hyaluronic acid (Synvisc[®]).

Group 4: two infiltrations separated by a three-week interval, using betamethasone 0.25 mg as the dipropionate and 0.1 mg as the phosphate (Diprospan[®]).

One rabbit in group 2 and another in group 3 evolved




Figure 1 – Photographs demonstrating the surgical exposure of the anterior cruciate ligament and the closure of the skin after the resection

with infection of the operative site that extended to the joint, and were excluded from the study.

The rabbits were killed 12 weeks after the operation. For this, they were firstly anesthetized as described above and then subjected to intracardiac injection of thiopental (5 ml) and potassium chloride (10 ml).

The tibial plateaus were resected aseptically, and were immersed in a flask containing 10% formol, for subsequent macroscopic evaluation of the joint cartilage.

The changes to the joint surface were graded macroscopically as follows: grade 0 – smooth and shiny joint surface, without abnormalities of relief; grade 1 – rough surface, with loss of shininess, without depressions; grade 2 – similar to grade 1, with depressions on the surface; grade 3 – exposure of the subchondral bone (Figure 2). The lateral and medial tibial plateaus were evaluated separately in each group.

The statistical analysis consisted of using Student's t test to determine whether there were any differences between the different groups in relation to the degrees of osteoarthritis, viewed macroscopically. To determine whether there were any internal differences in the groups regarding the degree of osteoarthritis in the lateral and medial compartments, the chi-square test was performed. To investigate differences between the degrees of osteoarthritis, independent of the drug used, analysis of variance (Anova) was used.

RESULTS

Osteoarthritis induced by resection of the anterior cruciate ligament occurred in the placebo group in both compartments in all the knees. In groups 2, 3 and 4, it occurred in 40%, 65% and 45% of the compartments, respectively (Table 1 and Figure 3).

Table 1 – Frequencies of the different degrees of osteoarthritis in the medial and lateral compartments of the study groups

Disease grade	placebo	%	Polireumin	%	Synvisc	%	corticoid	%
G0	0	0.00	12	60.00	7	35.00	12	54.54
G1	13	59.09	1	5.00	7	35.00	7	31.82
G2	5	22.73	7	35.00	6	30.00	3	13.64
G3	4	18.18	0	0.00	0	0.00	0	0.00
Total	22	100.00	20	100.00	20	100.00	22	100.00

The significance level used for all the analyses was 5%.

Student's t test was used to compare the different drugs and the placebo, regarding the development of osteoarthritis. A statistical difference was observed between each of the groups 2, 3 and 4 and the control group (Table 2), but there were no differences among the drug groups, taking $p < 0.05$.




Figure 2 – Macroscopic classification of the tibial plateaus: A) grade 0; B) grade 1 (arrow: loss of shininess; roughness); C) grade 2 (arrow: surface depression); D) grade 3 (arrow: exposure of subchondral bone)

Table 2 – Student's t test

Drugs	Calculated t
Synvisc/ corticoid	0.8232
Synvisc/ Polireumin	1.0000
Corticoid/ Polireumin	0.8790

The critical t value for all the comparisons was 2.7764
Significance level of 5%

Through Anova, it was seen that there was a difference between different degrees of osteoarthritis, independent of the drug used (Table 3).

Table 3 – Variance test (Anova)

Source of variance	F	P value
Disease grade	6.5890	0.025

Comparison of the degree of osteoarthritis within each group, between the medial and lateral compartments, showed that there was no significant difference in any of the study groups, according to the chi-square test (Table 4).

Table 4 – Chi-square test

Drugs	calculated
placebo	0.8707
Synvisc	0.0631
corticoid	0.1066
Polireumin	0.0040

Chi-square value tabulated with significance of 0.05 and one degree of freedom was 3.84

DISCUSSION

In this experiment, the joint cartilage of the medial and lateral tibial plateaus of the rabbits' knees that were subjected to sectioning of the anterior cruciate ligament and to infiltrations of hyaluronates, corticosteroid and isotonic saline solution (control) were evaluated macroscopically. The definitions of the different degrees of osteoarthritis were based on observations of the appearance of the joint cartilage surface, going from normal (grade 0) to complete destruction of the cartilage and exposure of the subchondral bone (grade 3). Based on this classification, it was observed that, in all the knees infiltrated with isotonic saline solution, there was some degree of chondral lesion and severe joint degeneration (grade 3) in 18%. In the other groups, there were cases that did not evolve to arthrosis, and the maximum that was observed was changes to the joint surface relief (Table 1 and Figure 3).




Figure 3 – Occurrence of osteoarthritis according to the grade in the tibial plateaus in the different groups

The results demonstrated that there was a statistical difference with regard to prevention of chondral lesions, between the placebo group and the groups that received joint injections of different types of hyaluronic acid or corticosteroid (Table 2). There was no predominance of joint lesions between the lateral and medial tibial compartments in any of the groups (Table 4). In the same way as shown by other studies in the literature, the present study demonstrated that the intra-articular injections of low and high molecular weight glycosaminoglycans and corticosteroid had a protective effect on the chondral bone^(5,15-18). This is also in agreement with the study by Karakurum *et al*⁽⁸⁾, who compared intra-articular injections of predniso-

lone with injections of hyaluronic acid, in an experimental osteoarthritis model in rabbits, and observed that the two drugs administered separately prevented joint degeneration to the same extent, while the effect was boosted when the drugs were administered together.

For the infiltrations with corticosteroids, doses proportional to those of human treatments were used, thereby avoiding worsening of the chondral lesions due to excess medication. The protective effect on chondral bone that has been observed in different studies occurred with the use of corticosteroids in low doses. The three-week interval between the applications of corticosteroid was used because corticosteroids are less soluble and may remain inside the joint for this period⁽¹⁹⁾.

Yoshimi *et al*⁽⁴⁾ developed the osteoarthritis model involving sectioning the anterior cruciate ligament in rabbits' knees. In comparison with other models, such as partial medial meniscectomy in rabbits and sectioning the anterior cruciate ligament in dogs or rats, their model is the one that comes closest to the joint degeneration process in humans⁽²⁰⁻²⁵⁾. Published studies in the literature that used this model to evaluate the protective effect on chondral bone of glycosaminoglycans and corticosteroids have mostly demonstrated that the control groups develop arthrosis^(4,5,8-10,14,15,24-26). However, contrary to experimental studies, recent randomized prospective clinical trials have not proven the protective effect on chondral bone of different types of hyaluronic acid and corticosteroid. What is presented as evidence is the improvement in pain levels and functional ability regarding activities of daily living among patients who made use of these medications over short periods⁽²⁷⁻³⁸⁾. Controlled studies with longer follow-up periods are needed in order to clarify the role of these drugs in prevention of degeneration of the joint cartilage.

In a continuation of this investigation, it is intended to use histological analysis and histomorphometric quantification of glycosaminoglycans in the extracellular matrix to assess the influence of these substances on the osteoarthritis model used here.

CONCLUSION

From the present study, it is concluded that attenuation of the process of osteoarthritis takes place to the same extent through the use of intra-articular injections of low molecular weight glycosaminoglycans, high molecular weight glycosaminoglycans and corticosteroid composed of betamethasone dipropionate and betamethasone disodium phosphate, in rabbits' knees.

REFERENCES

- Wang CT, Lin J, Chang CJ, Lin YT, Hou SM. Therapeutic effects of hyaluronic acid on osteoarthritis of the knee. *J Bone Joint Surg Am.* 2004;86(3):538-45.
- Bedson J, Jordan K, Croft P. The prevalence and history of knee osteoarthritis in general practice: a case-control study. *Fam Pract.* 2005;22(1):103-8.
- Altman RD, Moskowitz R. Intraarticular sodium hyaluronate in the treatment of patients with osteoarthritis of the knee: a randomized clinical trial. *J Rheumatol.* 1998;25(11):2203-12.
- Yoshimi T, Kikuchi T, Obara T, Yamaguchi T, Sakakibara Y, Itoh H, et al. Effects of high-molecular-weight sodium hyaluronate on experimental osteoarthritis induced by the resection of rabbit anterior cruciate ligament. *Clin Orthop Relat Res.* 1994;(298):296-304.
- Hulmes DJ, Marsden ME, Strachan RK, Harvey RE, McInnes N, Gardner DL. Intra-articular hyaluronate in experimental rabbit osteoarthritis can prevent changes in cartilage proteoglycan content. *Osteoarthritis Cart.* 2004;12(3):232-8.
- Schiavinato A, Finesso M, Cortivo R, Abatangelo G. Comparison of the effects of intra-articular injections of Hyaluronan and its chemically cross-linked derivative. *Clin Exp Rheumatol.* 2002;20(4):445-54.
- Leopold SS, Redd BB, Warne WJ, Wehrle PA, Pettis PD, Shott S. Corticosteroid compared with hyaluronic acid injections for the treatment of osteoarthritis of the knee. *J Bone Joint Surg Am.* 2003;85(7):1197-203.
- Karakurum G, Karakok M, Tarakcioglu M, Koger NE, Kocabas R, Bagci C. Comparative effect of intra-articular administration of hyaluronan and/or cortisone with evaluation of malondialdehyde on degenerative osteoarthritis of the rabbit's knee. *Tohoku J Exp Med.* 2003;199(3):127-34.
- Pelletier JP, Mineau F, Raynauld JP, Woessner JF, Gunia-Smith Z, Pelletier JM. Intraarticular injections with methylprednisolone acetate reduce osteoarthritic lesions in parallel with chondrocyte stromelysin synthesis in experimental osteoarthritis. *Arthritis Rheum.* 1994;37(3):414-23.
- Pelletier JP, Cloutier JM, Martel-Pelletier J. In vitro effects of tiaprofenic acid, sodium sodium salicylate and hydrocortisone on the proteoglycan metabolism of human osteoarthritic cartilage. *Rheumatol.* 1989;16(5):646-55.
- Cortivo R, Brun P, Cardarelli L, O'regan M, Radice M, Abatangelo G. Antioxidant effects of hyaluronan and its alpha-methyl-prednisolone derivative in chondrocyte and cartilage cultures. *Semin Arthritis Rheum.* 1996;26(1):492-501.
- Cooper C, Snow S, McAlindon TE, Kellingray S, Stuart B, Coggon D, et al. Risk factors for the incidence and progression of radiographic knee osteoarthritis. *Arthritis Rheum.* 2000;43(5):995-1000.
- Moskowitz RW. Experimental models of osteoarthritis. In: Moskowitz RW, Altman RD, Buckwalter JA, Goldberg VM, Hochberg MC. *Osteoarthritis: diagnosis and medical/surgical management.* 2th. Philadelphia: Saunders; 1992. p. 213-52.
- Sah RL, Yang AS, Chen AC, Hant JJ, Halili RB, Yoshioka M, et al. Physical properties of rabbit articular cartilage after transaction of the anterior cruciate ligament. *J Orthop Res.* 1997;15(2):197-203.
- Kobayashi K, Amiel M, Harwood FL, Healey RM, Sonoda M, Moriya H, et al. The long-term effects of hyaluronan during development of osteoarthritis following partial meniscectomy in a rabbit model. *Osteoarthritis Cart.* 2000;8(5):359-65.
- Butler M, Colombo C, Hickman L, O'Byrne E, Steele R, Steinetz B, et al. A new model of osteoarthritis in rabbits. III. Evaluation of anti-osteoarthritis effects of selected drugs administered intraarticularly. *Arthritis Rheum.* 1983;26(11):1380-6.
- Pelletier JP, Mineau F, Raynauld JP, Woessner JF Jr, Gunja-Smith Z, Martel-Pelletier J. Intraarticular injection with methylprednisolone acetate reduces osteoarthritic lesions at the same time as chondrocyte stromelysin synthesis in experimental osteoarthritis. *Arthritis Rheum.* 1994;37(3):414-23.
- Pelletier JP, DiBattista JA, Raynauld JP, Wilhelm S, Martel-Pelletier J. The *in vivo* effects of intraarticular corticosteroid injections on cartilage lesions, stromelysin, interleukin-1, and oncogene protein synthesis in experimental osteoarthritis. *Lab Invest.* 1995;72(5):578-86.
- Caldwell, JR. Intra-articular corticosteroids: guide to selection and indications for use. *Drugs* 1996;52(4):507-14.
- Moskowitz RW, Davis W, Sammarco J, Martens M, Baker J, Mayor M, et al. Experimentally induced degenerative joint lesions following partial meniscectomy in the rabbit. *Arthritis Rheum.* 1973;16(3):397-405.
- Moskowitz RW, Howell DS, Goldberg VM, Muniz O, Pita JC. Cartilage proteoglycan alterations in a experimentally induced model of rabbit osteoarthritis. *Arthritis Rheum.* 1979;22(2):155-63.
- Pond MJ, Nuki G. Experimentally-induced osteoarthritis in the dog. *Ann Rheum Dis.* 1973;32(4):387-8.
- Stoop R, Buma P, van der Kraan PM, Hollander AP, Billingham RC, Meijers TH, et al. Type II collagen degradation in articular cartilage fibrillation after anterior cruciate ligament transection in rats. *Osteoarthritis Cart.* 2001;9(4):308-15.
- Amiel D, Toyoguchi T, Kobayashi K, Bowden K, Amiel M, Healey RM. Long-term effect of sodium hyaluronate (Hyalgan®) on osteoarthritis progression in a rabbit model. *Osteoarthritis Cart.* 2003;11(9):636-43.
- Yoshioka M, Coutts RD, Amiel D, Hacker SA. Characterization of a model of osteoarthritis in the rabbit knee. *Osteoarthritis Cart.* 1996;4(2):87-98.
- Chang DG, Iverson EP, Schinagl RM, Sonoda M, Amiel D, Coutts RD, et al. Quantitation and localization of cartilage degeneration following the induction of osteoarthritis in the rabbit knee. *Osteoarthritis Cart.* 1997;5(5):357-72.
- Divine JG, Zazulak BT, Hewett TE. Viscosupplementation for knee osteoarthritis. A systematic review. *Clin Orthop Relat Res.* 2006;(455):113-22.
- Kemper F, Gebhardt U, Meng T, Murray C. Tolerability and short-term effectiveness of hylan G-F 20 in 4253 patients with osteoarthritis of the knee in clinical practice. *Current Med Res Opin.* 2005;21(8):1261-9.
- Waddell DD, Cefalu CA, Bricker DC. A second course of hylan G-F 20 for the treatment of osteoarthritic knee pain: 12-month patient follow-up. *J Knee Surg.* 2005;18(1):7-15.
- Waddell D, Bricker DC. Clinical experience with the effectiveness and tolerability of hylan G-F 20 in 1047 patients with osteoarthritis of the knee. *J Knee Surg.* 2006;19(1):19-27.
- Raynauld JP, Goldsmith CH, Bellamy N, Torrance GW, Polissos R, Belovich D, et al. Effectiveness and safety of repeat courses of hylan G-F 20 in the patients with knee osteoarthritis. *Osteoarthritis Cart.* 2005;13(2):111-9.
- Pyne D, Ioannou Y, Mootoo R, Bhanji A. Intra-articular steroids in knee osteoarthritis: a comparative study of triancinolone hexacetonide and methylprednisolone acetate. *Clin Rheumatol.* 2004;23(2):116-20.
- Leopold SS, Warne WJ, Pettis PD, Shott S. Increased frequency of acute local reaction to intra-articular hylan GF 20 (Synvisc) in patients receiving more than one course of treatment. *J Bone J Surg Am.* 2002;84(9):1619-23.
- Pagnano M, Westrich G. Successful nonoperative management of chronic osteoarthritis pain of the knee: safety and efficacy of retreatment with intra-articular hyaluronans. *Osteoarthritis Cart.* 2005;13(9):751-61.
- Marino A, Wadell D, Kolomytkin O, Pruetz S, Sadasivan K, Albright J. Assessment of immunologic mechanisms for flare reactions to Synvisc. *Clin Orthop Relat Res.* 2006;(442):187-94.
- Bellamy N, Campbell J, Robinson V, Gee T, Bourne R, Wells G. Viscosupplementation for osteoarthritis of the knee. *Cochrane Database Syst Rev.* 2006:CD005321.
- Kotevoglou N, Ivibozkurt PC, Hiz O, Toktas H, Kuran B. A prospective randomized controlled clinical trial comparing the efficacy of different molecular weight hyaluronan solutions in the treatment of knee osteoarthritis. *Rheumatol Int.* 2006;26(4):325-30.
- Bellamy N, Campbell J, Robinson V, Gee T, Bourne R, Wells G. Intraarticular corticosteroid for treatment of osteoarthritis of the knee. *Cochrane Database Syst Rev.* 2005:CD005328