



## Association between accelerometer-determined physical activity and flavonoid-rich fruit and vegetable consumption among a national sample of U.S. adults

Paul D. Loprinzi \*

Center for Health Behavior Research, Department of Health, Exercise Science, and Recreation Management, The University of Mississippi, 229 Turner Center, University, MS 38677, USA

### ARTICLE INFO

Available online 11 December 2015

#### Keywords:

Accelerometry  
Epidemiology  
Flavonoids  
Fruits  
NHANES  
Vegetables

### ABSTRACT

Few population studies have examined the association between accelerometer-assessed physical activity and dietary behavior. Further, no studies have systematically examined the association between accelerometer-assessed physical activity and flavonoid rich fruits and vegetables in a national sample. As a result, the purpose of this study was to examine the association between objectively measured physical activity and consumption of flavonoid rich fruits and vegetables among a national sample of U.S. adults. Data from the 2003–2006 National Health and Nutrition Examination Survey (NHANES) were used ( $N = 2949$ ). Physical activity was measured via accelerometry and fruit and vegetable consumption was measured from the NHANES Food Frequency Questionnaire. After adjustments, moderate-to-vigorous physical activity was positively associated ( $p < 0.05$ ) with apples ( $\beta = 0.30$ ), grapes ( $\beta = 0.27$ ), strawberries ( $\beta = 0.32$ ), oranges ( $\beta = 0.35$ ), raw greens ( $\beta = 0.19$ ), carrots ( $\beta = 0.23$ ), peppers ( $\beta = 0.29$ ) and an overall flavonoid index variable ( $\beta = 2.34$ ). Future studies employing a longitudinal design are needed to better understand the direction of the observed associations. If future studies do indeed support the possibility that physical activity may help to foster changes in dietary behavior, then this will have strong implications for health behavior interventions, particularly among individuals finding it difficult to change multiple health behaviors concurrently.

© 2015 Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

### Introduction

Emerging research demonstrates that consuming foods rich in flavonoids, such as oranges, grapes, and apples, is associated with improved health, including lower inflammation and reduced incidence of diabetes (Holt et al., 2009; Chun et al., 2008; Zamora-Ros et al., 2013). There is also some encouraging work showing that individuals engaging in higher levels of physical activity tend to eat an ‘overall’ healthier diet than their less active counterparts (Loprinzi et al., 2014; Gillman et al., 2001; Emmons et al., 1994; Eaton et al., 1995; French et al., 1996; Blair et al., 1996; Matthews et al., 1997). However, unlike studies in children and adolescents (Vissers et al., 2013; Jago et al., 2010), these adult studies are limited in the extent that they have exclusively used self-reported physical activity methodology, which is prone to considerable measurement error (Shephard, 2003). Validation studies examining the association between self-report physical activity and some gold-standard (e.g., accelerometry, indirect calorimetry, and doubly labeled water) typically show a poor correlation in the range of 0.3–0.5 (van Poppel et al., 2010; Helmerhorst et al., 2012). Thus, these

‘validated’ self-report questionnaires only account for 9–25% of the variance in the explanatory parameter and are therefore likely to result in considerable misclassification. Further, the few previous studies on this topic did not systematically examine the association between physical activity and foods rich in flavonoids.

As a result, the purpose of this brief study was to examine the association between objectively-measured (accelerometry) physical activity and 15 different food items rich in flavonoids, as identified from the USDA database for the flavonoid content of selected foods. To improve generalizability, data from the 2003–2006 National Health and Nutrition Examination Survey (NHANES) was used. I hypothesize that adults who are more active will consume more food items rich in flavonoids.

### Study design and participants

Data from the 2003–2006 National Health and Nutrition Examination Survey (NHANES) were used (only available cycles with accelerometry data at the time of this writing). All procedures for data collection were approved by the National Center for Health Statistics ethics review board, and all participants provided written informed consent prior to data collection. For the present analyses,

\* Fax: +1 662 915 5525.

E-mail address: [pdloprin@olemiss.edu](mailto:pdloprin@olemiss.edu).

2949 adult participants (20–85 yrs) provided data for all study variables.

#### Assessment of physical activity

2003–2006 NHANES participants were asked to wear an ActiGraph 7164 accelerometer during all activities, except water-based activities and while sleeping. Prior to the participant's examination, accelerometers were initialized to collect data in one minute time periods. The output of an accelerometer is *activity counts*, which are proportional to measured acceleration. The ActiGraph 7164 accelerometer measures accelerations in the vertical axis using a piezoelectric plate. The accelerometer output is digitized using an analog-to-digital converter, and once digitized, the signal passes through a digital filter that detects accelerations ranging from 0.05 to 2.00 g in magnitude with frequency responses ranging from 0.25 to 2.5 Hz to filter motion outside normal human movement. The filtered signal is then rectified and summed over a pre-determined epoch period. After the activity count is sorted into an epoch, it is stored in the internal memory and then the integrator is reset to zero.

Activity counts per minute of  $\geq 2020$  were used to denote moderate-to-vigorous physical activity (MVPA) intensity (Troiano et al., 2008). Nonwear was defined by a period of a minimum of 60 consecutive minutes of zero activity counts, with the allowance of 1–2 min of activity counts between 0 and 100 (Troiano et al., 2008). For the analyses described here, only those participants with at least 4 days with 10 or more hours per day of monitoring data were included in the analyses (Troiano et al., 2008).

#### Assessment of flavonoid-rich fruits and vegetables

Based on the National Cancer Institute Diet History Questionnaire (DHQ) that is widely used in nutritional epidemiology research (Subar et al., 2001), participants completed the NHANES Food Frequency Questionnaire (FFQ) (Subar et al., 2006). Briefly, participants were asked to report the proportion of time certain types of foods were eaten. For the present study, which is consistent with other studies (Loprinzi and Mahoney, 2015; Mahoney and Loprinzi, 2014), 15 [solid] fruits and vegetables rich in flavonoids (i.e., apples, grapes, strawberries, oranges, cooked greens, raw greens, carrots, string beans, peas, broccoli, onions, peppers, cucumbers, tomatoes, and lettuce) were identified by using the USDA flavonoid content of foods (USDA, 2007). For each of the 15 food items, response options ranged from 1 to 11, and included never (1), 1–6 times/yr (2), 7–11 times/yr (3), 1 time/month (4), 2–3 times/month (5), 1 time/wk (6), 2 times/wk (7), 3–4 times/wk (8), 5–6 times/wk (9), 1 time/day (10), and 2 or more times/day (11). To create a flavonoid index variable, responses were summed, with higher values indicating more frequent consumption of flavonoid-rich fruits and vegetables. With 15 items, the possible range for the flavonoid index variable is 15–165.

#### Covariates

Covariates included age (yrs; continuous), gender, race-ethnicity (Mexican American, non-Hispanic white, non-Hispanic black, other), cotinine (ng/dL; continuous), poverty level (range: 0–5), BMI ( $\text{kg}/\text{m}^2$ ; continuous), and comorbidity index (0–1 + comorbidities). Information about age, gender, and race-ethnicity were obtained from a questionnaire. As a measure of socioeconomic status, poverty-to-income ratio (PIR) was assessed, with a PIR value below 1 considered below the poverty threshold. The PIR is calculated by dividing the family income by the poverty guidelines, which is specific to the family size, year assessed, and state of residence. Serum cotinine was measured as a marker of active smoking status or environmental exposure to tobacco (i.e., passive smoking). Serum cotinine was measured by an isotope dilution-high performance liquid chromatography/atmospheric pressure chemical

ionization tandem mass spectrometry. BMI was calculated from measured weight and height (weight in kilograms divided by the square of height in meters). With regard to chronic disease, participants were classified as having 0 or 1 + comorbidities based on self-report of the following chronic diseases/events: arthritis, coronary heart disease, stroke, congestive heart failure, cancer, heart attack, emphysema, chronic bronchitis, asthma, or hypertension.

#### Data analysis

All statistical analyses (STATA, version 12.0, College Station, TX) accounted for the complex survey design used in NHANES by using survey sample weights, clustering, and primary sampling units. Means and standard errors were calculated for continuous variables and proportions were calculated for categorical variables. To examine the association between MVPA and flavonoid-rich fruits and vegetables (outcome variable), multivariable linear regression analysis was employed. Separate models were computed for each of the fruit and vegetable items. A model using the index variable was also computed. In total, 16 regression models (15 fruit and vegetable items plus the index variable) were computed. Models controlled for age, gender, race-ethnicity, cotinine, poverty level, BMI, and comorbidity index. A  $p < 0.05$  denoted statistical significance for all analyses.

#### Results

Weighted characteristics of the analyzed sample are shown in Table 1. Participants were, on average, 50 years of age, 56% were female,

**Table 1**  
Weighted characteristics of the analyzed sample, NHANES 2003–2006 (n = 2949).

Variable	Mean/proportion	(95% CI)
<i>Demographics</i>		
Age, yr	49.9	48.9–50.9
% Female	56.1	54.2–58.0
Race-ethnicity, %		
Mexican American	7.8	5.7–9.9
Other Hispanic	1.9	1.2–2.6
Non-Hispanic White	75.7	71.3–80.1
Non-Hispanic Black	8.3	6.0–10.7
Other Race	6.0	4.3–7.8
Cotinine, ng/mL	47.6	41.4–53.7
Poverty-to-Income Ratio	3.3	3.1–3.4
Body Mass Index, $\text{kg}/\text{m}^2$	28.1	27.7–28.5
Comorbidity Index, %		
0 Comorbidities	44.7	42.4–46.9
1 + Comorbidities	55.2	53.0–57.5
<i>Physical activity</i>		
MVPA, min/day	22.3	21.0–23.5
<i>Diet<sup>a</sup></i>		
Apples	4.9	4.7–5.0
Grapes	4.3	4.2–4.4
Strawberries	4.5	4.4–4.7
Oranges	4.8	4.7–5.0
Cooked Greens	3.5	3.4–3.7
Raw Greens	2.9	2.8–3.1
Carrots	4.9	4.8–5.1
String Beans	5.1	4.9–5.3
Peas	4.1	4.0–4.2
Broccoli	4.6	4.4–4.7
Onions	6.6	6.5–6.8
Peppers	4.6	4.5–4.8
Cucumbers	4.3	4.1–4.4
Tomatoes	6.9	6.8–7.1
Lettuce	6.4	6.2–6.5
Index Variable (sum of each item)	73.1	72.0–74.2

MVPA = Moderate-to-vigorous physical activity.

<sup>a</sup> For each of the 15 food items, response options ranged from 1 to 11, and included never (1), 1–6 times/yr (2), 7–11 times/yr (3), 1 time/month (4), 2–3 times/month (5), 1 time/wk (6), 2 times/wk (7), 3–4 times/wk (8), 5–6 times/wk (9), 1 time/day (10), and 2 or more times/day (11).

75.7% were non-Hispanic white, the mean BMI (28.1 kg/m<sup>2</sup>) was in the 'overweight' category, 55.2% had at least one comorbidity, and on average, participants engaged in 22.3 min/day of MVPA. On average, the mean value for the different fruit and vegetable items was less than 5 (index score was 73.1; 73.1/15 items = 4.8). On the scale of 1–11, a value of 5 indicates that the participant consumed that fruit and vegetable 2–3 times a month.

Table 2 reports the weighted multivariable linear regression analyses examining the association between MVPA and each of the flavonoid-rich fruit and vegetable items. For each model, MVPA was expressed as a 30 unit/min change. At the  $p < 0.05$  level, MVPA was associated with apples ( $\beta = 0.30$ ), grapes ( $\beta = 0.27$ ), strawberries ( $\beta = 0.32$ ), oranges ( $\beta = 0.35$ ), raw greens ( $\beta = 0.19$ ), carrots ( $\beta = 0.23$ ), peppers ( $\beta = 0.29$ ) and the index variable ( $\beta = 2.34$ ). The inclusion of other covariates, including education and total dietary energy intake (kcal) from a 24-h recall, did not alter the findings. For example, MVPA remained significantly associated with the index variable after controlling for the original covariates plus education and energy intake ( $\beta = 2.34$ ; 95% CI: 0.61–3.87;  $P = 0.009$ ). Further, when total saturated fat (gm) and total dietary cholesterol (mg) were added to this model, MVPA remained significantly associated with the index variable ( $\beta = 2.21$ ; 95% CI: 0.55–3.87;  $P = 0.01$ ). Even further, when other chronic diseases, such as diabetes status (self-report of physician diagnosis), was added to this last model, results were unchanged ( $\beta = 2.21$ ; 95% CI: 0.54–3.88;  $P = 0.01$ ).

Although it is plausible to suggest an association between MVPA and each of these items, given the multiple comparisons, interpreting these associations while using a more conservative significance threshold may help to minimize a type 1 research error. After applying a Bonferroni-corrected  $p$ -value (i.e., 0.003; 0.05/16 regression models = 0.003), MVPA was associated with grapes and strawberries.

## Discussion

The main finding of this study was that adults engaging in more MVPA generally consumed more flavonoid rich fruits and vegetables, and in particular, consumed more grapes and strawberries. One potential reason for the null association for some fruits and vegetables may be

a result of some individuals having a dislike or non-preference for certain fruits and vegetables.

The observed association between objectively-measured physical activity and flavonoid rich fruits and vegetables is in support of the few other studies on this topic. For example, Gillman et al. (2001) demonstrated that adults in New England engaging in higher levels of self-reported physical activity, compared to those engaging in less physical activity, consumed greater amount of foods that are considered to be healthy (e.g., fruits, vegetables, fiber, calcium, folate, and vitamins A, C, and E). Similar results were also observed among a different sample of adults in New England (Eaton et al., 1995). Jago et al. (Jago et al., 2005) also showed that, among young adults in the Bogalusa Heart Study (Bogalusa, Louisiana), greater participation in self-reported physical activity was associated with higher fruit intake. Although speculative, the present findings are likely an underestimate of the relationship between MVPA and flavonoid-rich fruit/vegetable intake. In order to assess habitual movement patterns, only those with at least 4 days of 10+ h/day were included in the analysis, and previous work using this data set demonstrates that those with invalid monitoring data tend to be less active and have worse health profiles (Loprinzi et al., 2013), often attenuating associations.

These findings suggest that regular self-reported physical activity participation may be associated with greater consumption of flavonoid-rich fruits and vegetables. The present study extends these findings by employing an objective measure of physical activity and specifically examining flavonoid rich fruits and vegetables that are associated with lower inflammation and reduced incidence of chronic disease (Holt et al., 2009; Chun et al., 2008; Zamora-Ros et al., 2013).

Given the observed cross-sectional association between objectively measured physical activity and flavonoid rich fruits and vegetables, future studies employing a longitudinal design are needed. This will help to better understand the direction of the association between physical activity and fruit and vegetable consumption. It is plausible that consuming a healthier diet may help to facilitate physical activity behavior as individuals who eat 'healthier' may have, for example, more energy to engage in physical activity. It is also plausible to suggest that engaging in physical activity may help to foster changes in dietary behavior. For example, physical activity may increase self-efficacy to change other behaviors (e.g., diet) and physical activity has been shown to increase executive functioning, which may help facilitate purposeful, goal-directed behaviors and inhibit goal inconsistent behaviors (Guiney and Machado, 2013).

In summary, the main finding of this study was that objectively-measured physical activity was associated with flavonoid rich fruits and vegetables among this national sample of U.S. adults. Future studies employing a longitudinal design are needed to better understand the direction of the observed association. Such work should also aim to overcome the subjective assessment of dietary behavior in this study. Further, future studies should also aim to assess the association between MVPA and flavonoid rich fruits and vegetables while considering the ratio of flavonoid intake to overall energy intake. Another careful consideration for future studies is to minimize the potential for residual confounding. Although various covariates were included in the present study's analyses, there is the possibility that unmeasured confounding may be present.

If future studies do indeed support the possibility that physical activity may help to foster changes in dietary behavior, then this will have strong implications for health behavior interventions, particularly among individuals finding it difficult to change multiple health behaviors concurrently.

## Conflict of interest

The author declares no conflict of interest.

**Table 2**

Weighted multivariable linear regression analysis examining the association between MVPA (independent variable) and each flavonoid-rich food item, NHANES 2003–2006 (n = 2949).

Flavonoid-rich food item (dependent variable) Outcome variable	MVPA (independent variable) <sup>a</sup>		
	$\beta$	(95% CI)	P-Value
Apples	0.30	0.06–0.54	0.01
Grapes	0.27	0.10–0.43	0.002
Strawberries	0.32	0.17–0.48	<0.001
Oranges	0.35	0.12–0.58	0.004
Cooked Greens	−0.05	−0.23–0.12	0.52
Raw Greens	0.19	0.01–0.38	0.04
Carrots	0.23	0.03–0.43	0.02
String Beans	−0.12	−0.32–0.08	0.24
Peas	−0.02	−0.17–0.13	0.77
Broccoli	0.11	−0.10–0.32	0.29
Onions	0.05	−0.08–0.19	0.43
Peppers	0.29	0.09–0.48	0.006
Cucumbers	0.13	−0.05–0.32	0.14
Tomatoes	0.12	−0.05–0.30	0.16
Lettuce	0.14	−0.01–0.29	0.07
Index Variable (sum of each item)	2.34	0.65–4.02	0.008

16 separate multivariable linear regression models were computed; one for each flavonoid-rich food item (and the index variable). Each model was adjusted for age, gender, race-ethnicity, cotinine, poverty level, body mass index, and comorbidity index.

MVPA = Moderate-to-vigorous physical activity

<sup>a</sup> Independent variable, MVPA, expressed as a 30 unit/min change.

## Acknowledgments

The author declares no competing interests.

## References

- Blair, S.N., Horton, E., Leon, A.S., et al., 1996. Physical activity, nutrition, and chronic disease. *Med. Sci. Sports Exerc.* 28, 335–349.
- Chun, O.K., Chung, S.J., Claycombe, K.J., et al., 2008. Serum C-reactive protein concentrations are inversely associated with dietary flavonoid intake in U.S. adults. *J. Nutr.* 138, 753–760.
- Eaton, C.B., McPhillips, J.B., Gans, K.M., et al., 1995. Cross-sectional relationship between diet and physical activity in two southeastern New England communities. *Am. J. Prev. Med.* 11, 238–244.
- Emmons, K.M., Marcus, B.H., Linnan, L., et al., 1994. Mechanisms in multiple risk factor interventions: smoking, physical activity, and dietary fat intake among manufacturing workers. Working Well Research Group. *Prev. Med.* 23, 481–489.
- French, S.A., Hennrikus, D.J., Jeffery, R.W., 1996. Smoking status, dietary intake, and physical activity in a sample of working adults. *Health Psychol.* 15, 448–454.
- Gillman, M.W., Pinto, B.M., Tennstedt, S., et al., 2001. Relationships of physical activity with dietary behaviors among adults. *Prev. Med.* 32, 295–301.
- Guiney, H., Machado, L., 2013. Benefits of regular aerobic exercise for executive functioning in healthy populations. *Psychon. Bull. Rev.* 20, 73–86.
- Helmerhorst, H.J., Brage, S., Warren, J., et al., 2012. A systematic review of reliability and objective criterion-related validity of physical activity questionnaires. *Int. J. Behav. Nutr. Phys. Act.* 9, 103.
- Holt, E.M., Steffen, L.M., Moran, A., et al., 2009. Fruit and vegetable consumption and its relation to markers of inflammation and oxidative stress in adolescents. *J. Am. Diet. Assoc.* 109, 414–421.
- Jago, R., Nicklas, T., Yang, S.J., et al., 2005. Physical activity and health enhancing dietary behaviors in young adults: Bogalusa Heart Study. *Prev. Med.* 41, 194–202.
- Jago, R., Ness, A.R., Emmett, P., et al., 2010. Obesogenic diet and physical activity: independent or associated behaviours in adolescents? *Public Health Nutr.* 13, 673–681.
- Loprinzi, P.D., Mahoney, S.E., 2015. Association between flavonoid-rich fruit and vegetable consumption and total serum bilirubin. *Angiology* 66, 286–290.
- Loprinzi, P.D., Cardinal, B.J., Crespo, C.J., et al., 2013. Differences in demographic, behavioral, and biological variables between those with valid and invalid accelerometry data: implications for generalizability. *J. Phys. Act. Health* 10, 79–84.
- Loprinzi, P.D., Smit, E., Mahoney, S., 2014. Physical activity and dietary behavior in US adults and their combined influence on health. *Mayo Clin. Proc.* 89, 190–198.
- Mahoney, S.E., Loprinzi, P.D., 2014. Influence of flavonoid-rich fruit and vegetable intake on diabetic retinopathy and diabetes-related biomarkers. *J. Diabetes Complicat.* 28, 767–771.
- Matthews, C.E., Hebert, J.R., Ockene, I.S., et al., 1997. Relationship between leisure-time physical activity and selected dietary variables in the Worcester Area Trial for Counseling in Hyperlipidemia. *Med. Sci. Sports Exerc.* 29, 1199–1207.
- Shephard, R.J., 2003. Limits to the measurement of habitual physical activity by questionnaires. *Br. J. Sports Med.* 37, 197–206 (discussion 206).
- Subar, A.F., Thompson, F.E., Kipnis, V., et al., 2001. Comparative validation of the Block, Willett, and National Cancer Institute food frequency questionnaires: the Eating at America's Table Study. *Am. J. Epidemiol.* 154, 1089–1099.
- Subar, A.F., Dodd, K.W., Guenther, P.M., et al., 2006. The food propensity questionnaire: concept, development, and validation for use as a covariate in a model to estimate usual food intake. *J. Am. Diet. Assoc.* 106, 1556–1563.
- Troiano, R.P., Berrigan, D., Dodd, K.W., et al., 2008. Physical activity in the United States measured by accelerometer. *Med. Sci. Sports Exerc.* 40, 181–188.
- USDA, 2007. Database for the Flavonoid Content of Selected Foods. Release 2.1. USDA, Washington, DC.
- van Poppel, M.N., Chinapaw, M.J., Mokkink, L.B., et al., 2010. Physical activity questionnaires for adults: a systematic review of measurement properties. *Sports Med.* 40, 565–600.
- Vissers, P.A., Jones, A.P., van Sluijs, E.M., et al., 2013. Association between diet and physical activity and sedentary behaviours in 9–10-year-old British White children. *Public Health* 127, 231–240.
- Zamora-Ros, R., Forouhi, N.G., Sharp, S.J., et al., 2013. The association between dietary flavonoid and lignan intakes and incident type 2 diabetes in European populations: the EPIC-InterAct study. *Diabetes Care* 36, 3961–3970.