

The prognosis of refractory hypotension and severe intraventricular hemorrhage in very low birth weight infants

Seok Hwang-Bo, MD^a, Yu-Mi Seo, MD^a, Moon-Yeon Oh, MD^a, Soo-Ah Im, MD, PhD^b, Young-Ah Youn, MD, PhD^a 

Abstract

Background: The increased survival rate among very low birth weight infants has resulted in a higher risk for developing neuro-complications such as intraventricular hemorrhage (IVH), periventricular leukomalacia (PVL), and adverse neurodevelopmental outcomes.

Purpose: We examined refractory hypotension experienced within a week of life in association with severe IVH (grades 3–4) among very low birth weight infants (VLBWIs).

Method: Between Jan 2014 and Dec 2017, the clinical data of 191 VLBWIs were retrospectively chart reviewed. Of a total of 191 VLBWIs, 71.2% (136/191) had IVH, and 28.7% (55/191) had severe IVH.

Results: The VLBWI with severe IVH group (grade 3–4) presented with a significantly lower gestational age along with higher use of postnatal hydrocortisone for refractory hypotension within a week of life. Resuscitation at delivery, pulmonary hemorrhage, neonatal seizure, and PVL were significantly more frequent in the severe IVH group ($P < .05$). Higher mortality occurred in the VLBWI with severe IVH group ($P < .001$). The multivariable logistic regression analysis consistently showed that refractory hypotension within a week of life and neonatal seizures were significantly associated with severe IVH. Those in the severe IVH and refractory hypotension groups had significantly lower composite cognitive, language, motor scores in Bayley Scales of Infant and Toddler Development III scores at corrected 18–24 months.

Conclusion: Refractory hypotension within a week of life and seizures were consistently associated with severe IVH and developmental delay at corrected 18–24 months. VLBWI who experienced refractory hypotension within a week of life may indicate a more vulnerable clinical setting with a higher risk for developmental delay.

Abbreviations: BPD = bronchopulmonary dysplasia, IVH = intraventricular hemorrhage, NEC = necrotizing enterocolitis, PVL = periventricular leukomalacia, RDS = respiratory distress syndrome, ROP = retinopathy of prematurity, VLBWI = very low birth weight infants

Keywords: intraventricular hemorrhage, outcomes, refractory hypotension, risk factor, very low birth weight infants

1. Introduction

Progressive advances in neonatal practice along with antenatal and postnatal care have led to the increased survival rate of very low birth weight infants (VLBWIs). The increased survival rate in this population has resulted in a higher risk for developing neurocognitive and motor delays such as intraventricular hemorrhage (IVH), periventricular leukomalacia (PVL), and adverse neurodevelopmental outcomes. Neurological sequelae are

found in approximately 50%–75% of preterm survivors with severe IVH, and the finding of IVH invokes considerable anxiety about long-term sequelae when even survivors with mild grades of IVH develop physical impairments, such as cerebral palsy; as such, it remains a significant public health concern worldwide.^[1] Prior studies have shown that IVH is well known to be associated with a lower gestational age and birth weight, while low Apgar score and male sex were not consistently associated with severe IVH^[2,3] Along with these risk factors, refractory

The authors declare that they have no conflicts of interest.

The datasets used during the current study are available from the corresponding author upon request. Medical records are available in the Archive of the Department of Pediatrics of the Seoul St. Mary's Hospital.

For this retrospective study, formal consent was waived due to chart review in this study; any personal data was protected. The study was approved by the Ethics Committee of Seoul St. Mary's Hospital. Informed consent was waived for this study because it was a retrospective chart review, and all methods were performed in accordance with the relevant guidelines and regulations by the Ethics Committee of Seoul St. Mary's Hospital (KC13ONMI0228).

We, each author listed on the manuscript, approved the publication of this version of the manuscript and take full responsibility for the manuscript.

^a Department of Pediatrics, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea, ^b Department of Radiology, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, Republic of Korea.

Correspondence: Young-Ah Youn, Department of Pediatrics, Seoul St. Mary's Hospital, College of Medicine, The Catholic University of Korea, 222 Banpo-daero, Seocho-gu, Seoul, 06591, Republic of Korea (e-mail: lea732@hanmail.net).

Copyright © 2022 the Author(s). Published by Wolters Kluwer Health, Inc. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial License 4.0 (CCBY-NC), where it is permissible to download, share, remix, transform, and buildup the work provided it is properly cited. The work cannot be used commercially without permission from the journal.

How to cite this article: Hwang-Bo S, Seo Y-M, Oh M-Y, Im S-A, Youn Y-A. The prognosis of refractory hypotension and severe intraventricular hemorrhage in very low birth weight infants. *Medicine* 2022;101:30(e29598).

Received: 4 January 2022 / Received in final form: 25 April 2022 / Accepted: 2 May 2022

<http://dx.doi.org/10.1097/MD.00000000000029598>

hypotension is a common problem in the very premature infant population, with an estimated incidence of 20% to 45%,^[4] and there is an association with IVH.^[5] Clinically, hypotension reflects an impairment in cerebral autoregulation which is a protective mechanism to maintain stable cerebral perfusion despite fluctuations in blood flow due to vulnerable premature associated conditions. Refractory hypotension can diminish CO₂ reactivity in the brain due to vasodilated cerebral vessels, which can eventually decrease cerebral blood flow. The immaturity of the germinal matrix as well as fluctuations in cerebral blood flow with pressure changes can also affect premature infants' vulnerable cerebral blood flow.^[6] In very low birth weight infants, hypotension within the first week of life can be the first indication of a circulatory collapse, a condition that may require inotropic therapy or steroid treatment.

Infants with refractory hypotension experienced mostly adrenal insufficiency, shock or RDS, demonstrating the importance of the severity of illness within a week of life and possibly suggesting a causal relationship with severe IVH.

The goal of this retrospective chart review was to explore the risk factors for IVH among those vulnerable VLBWI who experienced refractory hypotension within a week of life and compare IVH outcomes. In our study, we hypothesized that infants who experienced refractory hypotension might be at risk for severe IVH and may indicate a more vulnerable perinatal setting with a higher risk for developmental delay.

2. Methods

2.1. Design and sample

We retrospectively analyzed the medical records of 217 VLBWIs admitted at Seoul St. Mary's Hospital between Jan 2014 and Dec 2017. Our study included VLBWIs born with birth weight <1500g or gestational age <32 weeks at birth. Infants with chromosomal anomalies were excluded (n = 3) for confounding risk factors for IVH. Additionally, 4 infants were excluded for no record of brain sonogram. A total of 217 VLBWI were admitted during the study period. Among these, 7 infants were excluded from this analysis; 2 infants had congenital anomalies and 4 infants did not have brain sonograms (Fig. 1). Out of 210 VLBWI, 19 infants were lost to follow-up at 18–24 months of neurodevelopmental outcome assessment. Eventually, 191 VLBWI were analyzed in this study (Fig. 1). The clinical risk factors influencing the development of IVH were analyzed. A brain sonogram was performed within a week of life for IVH screening among VLBWIs. For infants experiencing systemic hypotension, dopamine infusion was started at a rate of 5 µg/kg/min and increased by 2.5 µg/kg/min until adequate mean blood pressure was achieved after fluid bolus challenges. If 10 µg/kg/min dopamine failed, dobutamine (10 µg/kg/min) was added. If dopamine and dobutamine treatment failed, hydrocortisone was added at a dose of 5 mg/kg/day IV, divided into three doses at 8-hour intervals.

Patients were evaluated for neurodevelopmental outcomes at corrected 18 months based on the Bayley Scales of Infant and Toddler Development III. The study was approved by the Ethics Committee of Seoul St. Mary's Hospital. Informed consent was waived for this study because it was a retrospective chart review, and all methods were performed per the relevant guidelines and regulations by the Ethics Committee of Seoul St. Mary's Hospital (KC13ONMI0228).

2.2. Definitions

Systemic hypotension in very low birth weight infants was defined as a mean blood pressure measured noninvasively as below the 3rd percentile for gestational age or below 30 mmHg^[7] with decreased urine output >12 hours. Refractory




Figure 1. Study flowchart.

hypotension was defined when combination dopamine (10 µg/kg/min) and dobutamine (10 µg/kg/min) treatment failed and hydrocortisone (5 mg/kg/day IV, divided into 3 doses at 8-hour intervals) was needed for refractory hypotension. Pulmonary hemorrhage was defined as massive pulmonary hemorrhage, which affects vital signs as manifested by cardiovascular collapse or acute respiratory failure. Pulmonary hypertension was defined by the need to use nitric oxide or sildenafil or ≤1 week of life. Bronchopulmonary dysplasia (BPD) was diagnosed if oxygen use exceeding 0.21% was still needed at a corrected gestational age of 36 weeks.

Seizures were clinically diagnosed by experienced neonatology or neurologist staff as paroxysmal alterations in motor function and occasional autonomic function; this included clonic, tonic, and “subtle” seizure manifestations [Volpe JJ]. Necrotizing enterocolitis (NEC) was defined as grade II or higher using Bell's classification. IVH > grade II was defined as active bleeding in the ventricles, and the grade designation was based on Drs. Papile and Levene's classification criteria.^[8] At the corrected age of 18 months, patients were considered at risk if scores were >2 SDs below the test mean (scores of <70). The abnormal neurodevelopmental outcome was defined when composite scores of <70 in the areas of cognitive, language, or motor were found.

2.3. Statistical analysis

Continuous variables were compared using the Student *t*-test and are expressed as the mean ± standard deviation. Discrete

variables were compared using a χ^2 test or Fisher's exact test and are expressed as percentages. All the analyses were two-tailed, and clinical significance was defined as a *P* value less than .05. To identify any confounding risk factors for IVH, we used a multivariate logistic regression model. Odds ratios (ORs) and 95% confidence intervals (CIs) were calculated using both a multivariate statistical model that included the following predictors related to severe IVH and a stepwise logistic regression analysis. All statistical analyses were performed using the Statistical Package for the Social Sciences (SPSS), version 15.0 (SPSS-PC Inc., Chicago, IL, USA).

3. Results

Of the 191 VLBWIs who were admitted to the neonatal intensive care unit between January 2014 and December 2017, 71.2% (136/191) of VLBWIs had IVH, and 28.7% (55/191) had no IVH. Table 1 compares the clinical characteristics of VLBWI without and with IVH; VLBWI with IVH had a significantly lower gestational age (27.54 ± 2.37 weeks vs 28.58 ± 2.75 weeks, *P* < .009) and RDS (129 (94.9%) vs 47 (85.5%), *P* = .029). Concerning morbidities, infants who developed IVH had significantly more neonatal seizures, sepsis, packed red blood cell (PRC) transfusions, bronchopulmonary disease (BPD) ≥ moderate, and PVL. Additionally, VLBWI with IVH had a longer duration of total parenteral nutrition (TPN), duration of mechanical ventilation, and hospital stays (*P* < .001) than those without IVH. The hospital mortality rate was significantly different between the two groups.

Table 2 shows the higher severity of IVH characteristics among the VLBWIs with IVH (n = 136). The severe IVH group (> IVH grade II) exhibited a significantly lower gestational age (26.94 ± 2.47 weeks vs 27.94 ± 2.23 weeks, *P* = 0.016) than

those with ≤ IVH grade II. Early hydrocortisone use was significantly higher in the severe IVH group. Resuscitation at delivery, pulmonary hemorrhage, neonatal seizure, and PVL were significantly more frequent in the severe IVH group (*P* < .05). Higher mortality occurred in the VLBWI with severe IVH group (*P* < .001). To explore the influence of any possible confounding factors on IVH and severe IVH in VLBWI, we performed a multivariable logistic regression analysis to identify any confounding factors related to IVH severity. Table 3 provides the risks for severe IVH (those with grades 3 and 4 (n = 55)) in a logistic regression analysis. We included gestational age, early hydrocortisone use for refractory hypotension, pulmonary hemorrhage, and neonatal seizure in a multiple logistic regression analysis. Because the other significant variables in Tables 1 and 2 usually occur after IVH incidence, they were not included in this analysis to observe as risk factors.

Refractory hypotension and seizures were consistently associated with severe IVH.

At the corrected age of 18 months, the VLBWI who had IVH (n = 136) completed the cognitive, language, and motor components of the Bayley Scales of Infant and Toddler Development III. Those with a history of severe IVH (grade 3–4) had significantly lower composite scores for all five areas – cognitive, language, motor, socioemotional, and adaptive behavior on the Bayley Scales of Infant and Toddler Development III (Table 4). In addition, the severe IVH group was at significantly high risk of neurodevelopmental outcomes at corrected 18 months, defined as a score > 2 SDs below the test mean (scores of < 70) (Table 4). Additionally, those experiencing refractory hypotension within a week of life had significantly lower composite scores for all three areas – cognitive, language, and motor – of the Bayley Scales of Infant and Toddler Development III

Table 1

Clinical characteristics and outcomes of VLBWIs (n = 191).

	No IVH (n = 55)	IVH (n = 136)	<i>P</i> value
Gestational age, week	28.58 ± 2.75	27.54 ± 2.37	.009 ^a
Birth weight, kg	1.103 ± 0.302	1.042 ± 0.260	.163
Male, n (%)	24 (43.6%)	76 (55.9%)	.125
Maternal chorioamnionitis	14 (25.5)	51 (37.5)	.112
Antenatal steroid use	36 (65.5)	82 (60.3)	.506
Refractory hypotension ≤ 1 week of life	23 (41.8)	65 (47.8)	.453
Resuscitation at the time of delivery*	50 (90.9)	128 (94.1)	.425
RDS	47 (85.5)	129 (94.9)	.029 ^a
Surfactant > 2 times	15 (27.3)	56 (41.2)	.072
Pneumothorax	5 (9.1)	20 (14.7)	.297
Pulmonary hemorrhage	7 (12.7)	33 (24.3)	.076
Pulmonary hypertension†	6 (10.9)	23 (16.9)	.295
Neonatal seizure	13 (23.6)	86 (63.2)	<.001 ^a
Sepsis	15 (27.3)	60 (44.1)	<.031 ^a
PDA ligation	4 (7.3)	24 (17.6)	.066
NEC operation	0 (0)	7 (5.1)	.196
ROP operation	1 (1.8)	10 (7.4)	.182
PRC transfusion	47 (85.5)	132 (97.1)	.003 ^a
BPD ≥ moderate	14 (25.5)	82 (60.3)	<.001 ^a
PVL	5 (9.1)	67 (49.3)	<.001 ^a
TPN duration‡	25.11 ± 25.11	43.36 ± 31.72	<.001 ^a
Mechanical ventilation, days‡	12.85 ± 23.82	27.17 ± 31.62	.001 ^a
Hospital stay, days‡	35.11 ± 27.96	62.59 ± 38.41	<.001 ^a
Mortality	16 (29.1)	26 (19.1)	.132

BPD = bronchopulmonary dysplasia, IVH = intraventricular hemorrhage, NEC = necrotizing enterocolitis, PDA = patent ductus arteriosus, PRC = packed red cells, PVL = periventricular leukomalacia, RDS = respiratory distress syndrome, ROP, retinopathy of prematurity, TPN = total parenteral nutrition, VLBWI = very low birth weight infants.

*Resuscitation included oxygen use or positive pressure ventilation or intubation.

†Use of nitric oxide and sildenafil within 1 week of birth.

‡Expired patients were excluded.

^a*P* < .05.

Table 2

Clinical characteristics and outcomes of VLBWIs with IVH (n = 136).

	Mild IVH (Grades 1–2) (n = 81)	Severe IVH (grades 3–4) (n = 55)	<i>P</i> value
Gestational age, week	27.94 ± 2.23	26.95 ± 2.47	.016 ^a
Birth weight, kg	1.07 ± 0.24	1.01 ± 0.29	.224
Male, n (%)	40 (49.4)	36 (65.5)	.064
Antenatal steroid use	42 (51.9)	40 (72.7)	.015 ^a
Refractory hypotension ≤ 1 week of life	27 (33.3)	38 (69.1)	<.001 ^a
Maternal chorioamnionitis	26 (32.1)	25 (45.5)	.114
Resuscitation at delivery*	73 (90.1)	55 (100.0)	.021 ^a
RDS	76 (93.8)	53 (96.4)	.701
Surfactant > 2 times	32 (39.5)	24 (43.6)	.631
Pneumothorax	9 (11.1)	11 (20.0)	.151
Pulmonary hemorrhage	13 (16.0)	20 (36.4)	.007 ^a
Pulmonary hypertension†	10 (12.3)	13 (23.6)	.085
Neonatal seizure	40 (49.4)	46 (83.6)	<.001 ^a
Sepsis	32 (39.5)	28 (50.9)	.189
PDA ligation	17 (21.0)	7 (12.7)	.215
NEC operation	5 (6.2)	2 (3.6)	.701
ROP operation	4 (4.9)	6 (10.9)	.190
BPD ≥ moderate	49 (60.5)	33 (60.0)	.954
PVL	31 (38.3)	36 (65.5)	.002
PRC transfusion	77 (95.1)	55 (100)	.147
TPN duration‡	43.25 ± 27.0	43.53 ± 37.9	.960
Mechanical ventilation, days‡	23.84 ± 25.9	32.07 ± 38.25	.137
Hospital stay, days‡	60.98 ± 27.54	64.96 ± 50.56	.595
Mortality	7 (8.3)	19 (34.5)	<.001 ^a

BPD = bronchopulmonary dysplasia, IVH = intraventricular hemorrhage, NEC = necrotizing enterocolitis, PDA = patent ductus arteriosus, PRC = packed red cells, PVL = periventricular leukomalacia, RDS = respiratory distress syndrome, ROP, retinopathy of prematurity, TPN = total parenteral nutrition, VLBWI = very low birth weight infants.

*Resuscitation included oxygen use or positive pressure ventilation or intubation.

†Use of nitric oxide, sildenafil, or iloprost within 1 week of birth.

‡Expired patients were excluded.

^a*P* < .05.

Table 3

Severe IVH (adjusted for gestational age, refractory hypotension, pulmonary hemorrhage and neonatal seizure) in a multiple logistic regression analysis (n = 55).

	P	OR	95% CI
Gestational age, week	.487	0.937	0.780–1.126
Refractory hypotension	.025	2.746	1.133–6.652
Pulmonary hemorrhage	.571	1.323	0.502–3.488
Neonatal seizure	.042	2.689	1.038–6.968

IVH, intraventricular hemorrhage.

Table 4

Neurodevelopmental outcomes of severe IVH on Bayley Scales of Infant and Toddler Development III at corrected 18 months (n = 136).

	Severe IVH (grade 3–4) (n = 55)	Mild IVH (grades 1–2) (n = 81)	P value
Score, mean ±SD			
Cognitive	74.89 ± 27.24	88.61 ± 23.01	.023
Language	73.64 ± 22.38	85.43 ± 20.81	.025
Motor	68.78 ± 26.10	81.52 ± 23.84	.035
Socio-emotional score	79.26 ± 18.95	88.95 ± 20.64	.049
Adaptive behavior score	73.68 ± 20.80	85.31 ± 16.12	.043
At risk, n (%) [*]			
Cognitive score	13 (37.1)	8 (13.6)	.008 [*]
Language score	13 (37.1)	10 (16.9)	.028 [*]
Motor score	15 (42.9)	13 (21.7)	.029 [*]
Socio-emotional score	10 (28.6)	5 (8.5)	.010 [*]
Adaptive behavior score	16 (45.7)	10 (16.9)	.003 [*]

IVH = intraventricular hemorrhage, VLBWI = very low birth weight infants.

^{*}Children were considered to be at risk if their scores were >2 standard deviations below the test mean (scores of <70).

(Table 5). They were also at high risk of neurodevelopmental delay, defined as a score >2 SDs below the test mean (scores of <70) (Table 5).

4. Discussion

In our study of VLBWI, we observed a 28.7% incidence of severe IVH among 191 very preterm infants, which was higher than the reported severe IVH incidence in the National Institute of Child Health and Human Development (NICHD) Neonatal Research Network study in 2016, which was only 7.5%.^[9] However, the mean gestational age in our study group was 27–28 weeks of age, weighing 1–1.1 instead of comparing VLBWI weighing a mean of 1.5 kg. A prior study investigating the IVH prevalence in premature Israeli infants according to weight and gestational age similarly reports IVH incidence of 20.9%,^[6] which was similar to our IVH rate of 28.7%. Among the multifactorial pathogenetic risks for IVH, abnormalities of the hemodynamic systems certainly play a role.

We found that VLBWIs experiencing refractory hypotension within the first week of life who ultimately needed hydrocortisone to maintain appropriate blood pressure were at high risk for severe IVH (stages 3–4). Further, refractory hypotension and seizures were associated with severe IVH and developmental delay at corrected 18–24 months.

Refractory hypotension can be a risk factor in the very preterm population, with an estimated incidence of 20% to 45%^[10] which is associated with IVH.^[5] Other studies also indicate that hypotension independently predicts adverse outcomes and inotropic use was an independent predictor of IVH and death in premature infants.^[11] Faust et. al reported

Table 5

Outcomes of refractory hypotension ≤1 week on Bayley Scales of Infant and Toddler Development III at 18 months (n = 92).

	Hypotension ≤ 1 week (n = 52)	No hypotension (n = 40)	P value
Score, mean ±SD			
Cognitive	73.98 ± 21.77	89.50 ± 25.57	.018
Language	76.69 ± 19.80	87.50 ± 23.09	.022
Motor	78.62 ± 23.23	86.09 ± 25.03	.036
At risk, n (%) [*]			
Cognitive score	17 (32.7)	4 (10.0)	.021
Language score	17 (32.7)	4 (10.0)	.021
Motor score	23 (44.2)	7 (17.5)	.028

^{*}Children were considered to be at risk if their scores were >2 standard deviations below the test mean (scores of <70).

that emergency cesarean section, low Apgar scores, mechanical ventilation, and multiple births were associated with hypotension.^[12]

The more severe hospital morbidities in the VLBWI group exposed to early hydrocortisone use in our study are directly related to increased severity of illness in VLBWI, especially when it occurred in the first 72 hours of life, as reported by Jonathan et al.^[13] Similar to our study, refractory hypotension is reported to be associated with severe IVH, poor long-term neurodevelopment (permanent neurological injuries/deficits or cerebral palsy), and even death.^[14,15] Our study also reported higher mortality in the severe IVH group (34.5%) and 38 (69.1%) VLBWIs in the severe IVH group. Overall, VLBWI experiencing refractory hypotension within the first week of life indicates hemodynamic instability and changes in cerebral blood flow, which increases the incidence of severe IVH.

Some studies have highlighted the negative effects of vasoconstrictors in compromising cerebral autoregulation.^[16] The vasoconstrictive effects of inotropic drugs could worsen the hypoxia-hypoperfusion status of the periventricular areas of the immature brain and result in glutamate- and free radical-induced damage to preoligodendrocytes, which are the precursors of oligodendrocytes that eventually form the white matter.^[17] In fact, studies reported that the use of inotropes was associated with increased severe IVH in infants born before 32 weeks gestation.^[18,19] This may be because hemodynamic instability and changes in cerebral blood flow are increased risks of IVH.^[20]

In our study, our center only used hydrocortisone as an alternative to dexamethasone to minimize the negative effect on neurodevelopmental outcomes. Until the early 2000s, the postnatal steroid dexamethasone was used widely as clinical trials reported that dexamethasone reduced the risk of death or BPD in VLBWI; however, this benefit was outweighed by a possible increased risk of neurodevelopmental impairment.^[21] While many randomized and control trials (RCTs) have also shown adverse neurodevelopmental outcomes after postnatal dexamethasone treatment for BPD, no multicenter RCT studies have demonstrated adverse effects on the long-term outcomes after hydrocortisone therapy.^[22] This is because dexamethasone is 40 to 50 times more potent than hydrocortisone and longer-acting at 18 to 24 hours and may cause a less harmful effect on neurodevelopmental outcomes.^[23]

Since hydrocortisone targets both glucocorticoid and mineralocorticoid receptors, it has less potent toxicity than dexamethasone, suggesting that appropriate use of hydrocortisone may not develop complications such as brain toxicity. A current practice survey showed that hydrocortisone treatment has already been implemented in many neonatal units across the world.^[24] Hydrocortisone use can also counter the untoward effects of cytokines and hemodynamic shock on cellular

toxicity^[25] and may improve the inflammatory process and effectively reduce cytokine- and hypofusion-induced cellular and white matter injury. Our study has several limitations in its observational nature: (i) the retrospective study design, which might not be appropriate for confirming the examined relationships; (ii) the relatively small sample size of the study group; (iii) hidden disabilities may subsequently have become apparent later, and many infants might have important developmental lags that were not classified as impairments; and (iv) many clinical conditions that may originate from prematurity itself. However, the strength of our study is the observation of neurodevelopmental outcomes by the Bayley Scales of Infant and Toddler Development III at a corrected age of 18 months as a long-term study.

In summary, we concluded that refractory hypotension within a week of life and seizures may be predictors of the severity of IVH. Severe IVH (grades 3–4) was also significantly associated with an increased risk of neurodevelopmental outcome at corrected 18 months.

Author contributions

YYA wrote the first draft of the manuscript. All Authors have made a substantial contribution: YYA designed the patient study; HBS, SYM, OMY, and YYA conducted the research. YYA wrote the paper and had primary responsibility for the final content. OMY did the statistical analysis and ISA performed the brain imaging interpretation. Each author listed on the manuscript has seen and approved the submission of this version of the manuscript and takes full responsibility for the manuscript.

References

- [1] Sherlock RL, Anderson PJ, Doyle LW; Victorian Infant Collaborative Study Group. Neurodevelopmental sequelae of intraventricular hemorrhage at 8 years of age in a regional cohort of ELBW/very preterm infants. *Early Hum Dev.* 2005;81:909–16.
- [2] Sarkar S, Dechert R, Schumacher RE, et al. Is refractory hypotension in preterm infants a manifestation of early ductal shunting? *J Perinatol.* 2007;27:353–8.
- [3] Heuchan AM, Evans N, Smart DH, et al. Perinatal risk factors for major intraventricular hemorrhage in the Australian and New Zealand Neonatal Network, 1995–97. *Arch Dis Child-Fetal.* 2002;86:F86e–90.
- [4] Efrid MM, Heerens AT, Gordon PV, et al. A randomized-controlled trial of prophylactic hydrocortisone supplementation for the prevention of hypotension in extremely low birth weight infants. *J Perinatol.* 2004;25:119–24.
- [5] Watkins AM, West CR, Cooke RW. Blood pressure and cerebral hemorrhage and ischemia in very low birthweight infants. *Early Hum Dev.* 1989;19:103–10.
- [6] Kenet G, Kuperman AA, Strauss T, et al. Neonatal IVH—mechanisms and management. *Thromb Res.* 2011;127(Suppl 3):S120–2.
- [7] Zubrow AB, Hulman S, Kushner H, et al. Determinants of blood pressure in infants admitted to neonatal intensive care units: a prospective multicenter study. *J Perinatol.* 1995;15:470–9.
- [8] Papile L, Burstein J, Burstein R, et al. Incidence and evolution of subependymal and intraventricular hemorrhage in premature infants: a study of infants. *J Pediatr.* 1978;92:529–34.
- [9] Wei JC, Catalano R, Profit J, et al. Impact of antenatal steroids on intraventricular hemorrhage in very-low-birth weight infants. *J Perinatol.* 2016 ;36:352–6.
- [10] Efrid MM, Heerens AT, Gordon PV, et al. A randomized-controlled trial of prophylactic hydrocortisone supplementation for the prevention of hypotension in extremely low birth weight infants. *J Perinatol.* 2004;25:119–24.
- [11] Peter D, Gandy C, Hoffman SB. Hypotension and adverse outcomes in prematurity. *Neonatology.* 2017;111:228–33.
- [12] Faust K, Härtel C, Preuß M, et al. Short-term outcome of very-low-birthweight infants with arterial hypotension in the first 24h of life. *Arch Dis Child Fetal Neonatal Ed.* 2015;100:F388–92.
- [13] Fanaroff JM, Wilson-Costello DE, Newman NS. Treated hypotension is associated with neonatal morbidity and hearing loss in extremely low birth weight infants. *Pediatrics.* 2006;117:1131–5.
- [14] Abdul Aziz AN, Thomas S, Murthy P, et al. Early inotropes use is associated with higher risk of death and/or severe brain injury in extremely premature infants. *J Matern Fetal Neonatal Med.* 2019;22:1–8.
- [15] Dempsey EM. Challenges in treating low blood pressure in preterm infants. *Children (Basel).* 2015;2:272–88.
- [16] Hahn GH, Hyttel-Sorensen S, Petersen SM, et al. Cerebral effects of commonly used vasopressor-inotropes: a study in newborn piglets. *PLoS One.* 2013;8:e63069e 63069.
- [17] Falahati S, Breu M, Waickman AT, et al. Ischemia-induced neuroinflammation is associated with disrupted development of oligodendrocyte progenitors in a model of periventricular leukomalacia. *Dev Neurosci.* 2013;35:182–96.
- [18] Alotaibi WSM, Alsaif NS, Ahmed IA, et al. Reduction of severe intraventricular hemorrhage, a tertiary single-center experience: incidence trends, associated risk factors, and hospital policy. *Childs Nerv Syst* 2020;36:2971–9.
- [19] Szpecht D, Szymankiewicz M, Nowak I, et al. Intraventricular hemorrhage in neonates born before 32 weeks of gestation—retrospective analysis of risk factors. *Childs Nerv Syst.* 2016;32:1399–404.
- [20] Noori S, McCoy M, Anderson MP, et al. Changes in cardiac function and cerebral blood flow in relation to peri/intraventricular hemorrhage in extremely preterm infants. *J Pediatr.* 2014;164:264–70.e1.
- [21] Doyle LW, Cheong JL, Ehrenkranz RA, et al. Late (>7 days) systemic postnatal corticosteroids for prevention of bronchopulmonary dysplasia in preterm infants. *Cochrane Database Syst Rev.* 2017;10:CD001145.
- [22] Bartholomew J, Kovacs L, Papageorgiou A. Review of the antenatal and postnatal use of steroids. *Indian J Pediatr.* 2014;81:466–72.
- [23] Huang CC, Lin HR, Liang YC, et al. Effects of neonatal corticosteroid treatment on hippocampal synaptic function. *Pediatr Res.* 2007;62:267–70.
- [24] Virkud YV, Hornik CP, Benjamin DK, et al. Respiratory support for very low birth weight infants receiving dexamethasone. *J Pediatr.* 2017;183:26–30.
- [25] Fantuzzi G, Ghezzi P. Glucocorticoids as cytokine inhibitors: role in neuroendocrine control and therapy of inflammatory diseases. *Mediators Inflamm.* 1993;2:263–70.