



Short Communication

First report of the bat fly species *Basilia italica* in Romania

Áron Péter[‡], Andrei Daniel Mihalca[‡], Attila D Sándor^{‡,§}

[‡] University of Agricultural Sciences and Veterinary Medicine, Cluj-Napoca, Romania

[§] University of Veterinary Medicine, Budapest, Hungary

Corresponding author: Áron Péter (aronpeter92@gmail.com)

Academic editor: Pierfilippo Cerretti

Received: 17 Aug 2020 | Accepted: 18 Dec 2020 | Published: 18 Jan 2021

Citation: Péter Á, Mihalca AD, Sándor AD (2021) First report of the bat fly species *Basilia italica* in Romania.

Biodiversity Data Journal 9: e57680. <https://doi.org/10.3897/BDJ.9.e57680>

Abstract

Bat flies are haematophagous ectoparasites, highly specialised to bats and are also considered to have vectorial potential for several pathogens like *Bartonella* spp. or *Polychromophilus* spp. In Romania, past studies mostly focused on the ectoparasitic fauna of cave-dwelling bats, listing the occurrence of 10 bat fly species in the country, with only scarce information on bat flies infesting crevice-roosting bat species. Here we report the occurrence of *Basilia italica*, a rare nycteribiid species infesting primarily forest-dwelling bats. This is the first country-record for Romania and also represents the easternmost occurrence of this species. Further studies are needed to evaluate the vectorial potential of *B. italica*.

Keywords

Chiroptera, distribution, host-parasite relationships, *Myotis mystacinus*, Nycteribiidae

Introduction

Bat flies (Diptera, Hippoboscoidea) are highly specialised ectoparasites of bats representing two families in the superfamily Hippoboscoidea (Streblidae and Nycteribiidae;

Reeves and Lloyd 2018). Species belonging to Nycteribiidae are wingless, while most streblids have functional wings (with some exceptions). These dipterans are exclusive parasites of bats and are highly adapted to a parasitic life style. Bat flies are haematophagous and have highly specialised reproductive organs. These parasites reproduce by viviparous puparity and only females leave the host, when larvadeposit on the roost-substrate (Dick and Patterson 2006). Most bat flies show high host-specificity (monoxenous), being exclusive parasites of one or a few morphologically similar and phylogenetically closely related bat species (Dick 2007). Bat flies are increasingly recognised as vectors for several pathogen groups like bacteria of the genus *Bartonella* spp. or parasitic protozoa, like *Polychromophilus* spp. (Megali et al. 2010, Dick and Dittmar 2014, Sándor et al. 2018). Although the research interest is growing worldwide, most papers focus on parasites of cave-roosting bats in Romania, listing 10 species (Burghel-Balacescu 1966), namely: *Basilisa nana*, *B. nattereri*, *Nycteribia latreillii*, *N. kolenatii*, *N. pedicularia*, *N. schmidlii*, *N. vexata*, *Penicillidia duforii*, *P. conspicua* and *Phthiridium biarticulatum*). Gheorghiu (2006) reported the presence of several species at the Piatra Craiului forest and Sándor et al. (2018) described several new parasite-host species associations, while focusing on the epidemiological aspects of nycteribiid-infestation. Information on the biology, ecology or distribution of bat flies is still scarce to support further conclusions about their epidemiological risk contribution. Most of all, we lack this information on bat flies infesting crevice-roosting bat species. Recent studies of bat-related parasites in Romania provided new information for a checklists of bat ticks (Sándor et al. 2019) and their vectorial importance (Hornok et al. 2019, McKee et al. 2019), which also led to the study of bat flies (Sándor et al. 2018). Here we report the first records of a rare bat fly (*Basilisa italica*) in Romania, these being the easternmost reports of this species in Europe.

Material and methods

(Methods used for this study were published at protocols.io with the following DOI: [dx.doi.org/10.17504/protocols.io.bi2ykgfw](https://doi.org/10.17504/protocols.io.bi2ykgfw)).

Bats were trapped using mist nets erected close to the roosting site of an all-male colony of multiple species in an abandoned building close to Ic Ponor (Munții Apuseni, 46.629842N, 22.806450E, Fig. 1), at an altitude of 1044 m on 17.06.2020 and 01.07.2020. Upon capture, bats were examined, species were identified, based on morphological characters (Dietz et al. 2009) and all external parasites were collected using tweezers and were stored in 70% ethanol (single tube/host). In the laboratory, bat flies were identified under stereomicroscope, using morphological keys (Theodor 1967).

Results

On both trapping occasions, a single bat fly individual was collected from a single adult male *Myotis mystacinus*. None of the other examined individuals (all males, *Vespertilio murinus* – 106, *Pipistrellus pipistrellus* – 17, *M. mystacinus* – 16 and *M. brandtii* - 2) hosted

bat flies (Suppl. material 1). Bat flies were identified as a male and a female *Basilia italica*, based on the following morphological characteristics: a small-sized fly with visible eyes, but a body shape similar to the genus of Nycteribiae, a group of spines at the end of sternite 5. The surface of tergites are more or less bare, but on the marginal ends, there are some longer hairs. Lastly, there are long setae on the tibiae and the overall form of sternite 5 is unique (Fig. 2; see Theodor 1967, page 202).



Figure 1. [doi](#)

Location of the newly-reported bat fly species *B. italica*.

Discussion

The main host of *B. italica* is *M. mystacinus*, with occasional records on *M. alcahoe* and *M. brandtii* (Aellen 1963, Hürka 1964, Beaucournu and Noblet 1966, Czuppon and Molnár 2001, Krištofik and Danko 2012, Szentivanyi et al. 2016). In addition, five more bat species were listed as hosts, each with a single record (*Barbastella barbastellus*, *Eptesicus serotinus*, *M. nattereri* - Czuppon and Molnár 2001, *M. emarginatus*, *M. myotis* – Theodor 1967). It is a rare species, with a handful of observations, with a mainly Central European distribution (Hürka and Soós 1986).

The finds reported here are the first records of *B. italica* in Romania, hence increasing to 11 the number of known bat fly species in this country. This is a novel and a geographically-distant record, therefore considerably extending the range towards the East. (Fig. 3). The geographically closest occurrence of this fly species is in Hungary, where it was reported from six different species (Czuppon and Molnár 2001, Haelewaters et al. 2017).



Figure 2. [doi](#)

Dorsal view of the male *Basilica italica* caught on *Myotis mystacinus* in Ic Ponor, Munții Apuseni, Romania.




Figure 3. [doi](#)

Current geographical distribution of *B. italica* in Europe. Green coloured countries with data from literature by Szentivanyi et al. 2016 and our first record for Romania with red colour.

Although bat flies are usually abundant on cave dwelling species (like *Miniopterus schreibersii* or *M. myotis*) and their flies are relatively well-known (Zahn and Rupp 2004);, however, the host species of *B. italica* tend to roost in crevices (primarily tree-holes, but also abandoned buildings) and they are captured in smaller numbers (Dietz et al. 2009). After examining almost 500 individuals from the potential host species of *B. italica* in the past five years, these are the first observed individuals of this species in Romania.

Acknowledgements

Hereby, we would like to acknowledge the help provided by numerous volunteers in the fieldwork throughout the last five years, with special thanks for the assistance of B. Sándor and E. Sándor in data collection in 2020. ÁP was supported by the Collegium Talentum programme of Hungary. ADS was the beneficiary of the János Bolyai Research Scholarship of Hungarian Academy of Science and the ÚNKP 19-4-ÁTE-10 New National Excellence Program of the MIT.

Funding program

The NKFIH 132794 (National Research, Development and Innovation Office of Hungary) provided financial resources to ADS.

Hosting institution

University of Agricultural Sciences and Veterinary Medicine, Cluj-Napoca, Romania

Ethics and security

Bat capture permits were issued by the Underground Heritage Commission (Romania).

Author contributions

ÁP contributed to the bat capture and sample collection, did the identification and wrote the early version of the manuscript.

ADM made the photos and corrected the manuscript.

SDA organised the field trip, revised and corrected the manuscript.

Conflicts of interest

We (all authors) declare that there are no conflict of interest.

References

- Aellen V (1963) Les nycteribiidés de la Suisse, diptères parasites de chauves-souris. Bulletin de la Société Neuchâteloise des Sciences Naturelles 86: 143-154.
- Beaucournu JC, Noblet JF (1966) Les Diptères pupipares parasites de chauves-souris dans les Alpes et les Préalpes françaises (Diptera, Streblidae et Nycteribiidae). Bulletin de la Société entomologique de France 101: 235-240.
- Burghel-Balacescu A (1966) Date noi asupra raspândirii nycteribiidelor (Diptera, Pupipara) in România. Travaux de l'Institut de Spéologie "Émile Racovitza 5: 115-123.
- Czuppon B, Molnár V (2001) Bat fly genus and species new to Hungary (Diptera: Nycteribiidae). Folia Entomologica Hungarica 62: 313-317.
- Dick C, Patterson B (2006) Bat flies: Obligate ectoparasites of bats. Micromammals and Macroparasites 179-194. https://doi.org/10.1007/978-4-431-36025-4_11
- Dick C (2007) High host specificity of obligate ectoparasites. Ecological Entomology 32 (5): 446-450. <https://doi.org/10.1111/j.1365-2311.2007.00836.x>
- Dick CW, Dittmar K (2014) Parasitic bat flies (Diptera: Streblidae and Nycteribiidae): Host specificity and potential as vectors. In: Klimpel S, Mehlhorn H (Eds) Bats (Chiroptera) as vectors of diseases and parasites. URL: <https://link.springer.com/book/10.1007/978-3-642-39333-4>
- Dietz C, von Helversen O, Nill D (2009) Bats of Britain, Europe and northwest Africa. A & C Black [ISBN 1408105314]
- Gheorghiu V (2006) Contribution to the knowledge of the family Nycteribiidae (Diptera: Pupipara) of Piatra Craiului National Park. Researches in Piatra Craiului National Park 218-222.
- Haelewaters D, Pfliegler WP, Szentivnyi T, Fldvri M, Sndor AD, Barti L, Camacho JJ, Gort G, Estk P, Hiller T, Dick CW (2017) Parasites of parasites of bats: Laboulbeniales (Fungi: Ascomycota) on bat flies (Diptera: Nycteribiidae) in central Europe. Parasites & Vectors 1 (10): 96. [In English]. <https://doi.org/10.1186/s13071-017-2022-y>
- Hornok S, Szóke K, Meli M, Sándor A, Görföl T, Estók P, Wang Y, Tu VT, Kováts D, Boldogh S, Corduneanu A, Sulyok K, Gyuranecz M, Kontschán J, Takács N, Halajian A, Epis S, Hofmann-Lehmann R (2019) Molecular detection of vector-borne bacteria in bat ticks (Acari: Ixodidae, Argasidae) from eight countries of the Old and New Worlds. Parasites & Vectors 12 (1). <https://doi.org/10.1186/s13071-019-3303-4>
- Húrka K (1964) Distribution, bionomy and ecology of the European bat flies with special regard to the Czechoslovak fauna (Diptera, Nycteribiidae). Acta Universitatis Carolinae Biologica (3): 167-231.
- Húrka K, Soós Á (1986) Nycteribiidae. In: Soós Á, Papp L (Eds) Catalogue of Palaearctic Diptera . 11. Akadémiai Kiadó, Budapest, 226–233 pp.
- Krištofik J, Danko S (2012) Arthropod ectoparasites (Acarina, Heteroptera, Diptera, Siphonaptera) of bats in Slovakia. Vespertilio 16: 167-189.
- McKee C, Krawczyk A, Sándor A, Görföl T, Földvári M, Földvári G, Dekeukeleire D, Haarsma A, Kosoy M, Webb C, Sprong H (2019) Host Phylogeny, Geographic Overlap, and Roost Sharing Shape Parasite Communities in European Bats. Frontiers in Ecology and Evolution 7 <https://doi.org/10.3389/fevo.2019.00069>
- Megali A, Yannic G, Christe P (2010) Disease in the dark: molecular characterization of *Polychromophilus murinus* in temperate zone bats revealed a worldwide distribution of

this malaria-like disease. *Molecular Ecology* 20 (5): 1039-1048. <https://doi.org/10.1111/j.1365-294x.2010.04905.x>

- Reeves W, Lloyd J (2018) Louse flies, keds, and bat flies (Hippoboscoidea). In: Mullen G, Durden L (Eds) *Medical and Veterinary Entomology*. 3. Elsevier, 17 pp. [ISBN 9780128140437]. <https://doi.org/10.1016/B978-0-12-814043-7.00020-0>
- Sándor A, Földvári M, Krawczyk A, Sprong H, Corduneanu A, Barti L, Görföl T, Estók P, Kováts D, Szekeres S, László Z, Hornok S, Földvári G (2018) Eco-epidemiology of novel *Bartonella* genotypes from parasitic flies of insectivorous bats. *Microbial Ecology* 76 (4): 1076-1088. <https://doi.org/10.1007/s00248-018-1195-z>
- Sándor A, Corduneanu A, Péter Á, Mihalca AD, Barti L, Csősz I, Szőke K, Hornok S (2019) Bats and ticks: host selection and seasonality of bat-specialist ticks in eastern Europe. *Parasites & Vectors* 12 (1). <https://doi.org/10.1186/s13071-019-3861-5>
- Szentivanyi T, Estók P, Földvári M (2016) Checklist of host associations of European bat flies (Diptera: Nycteribiidae, Streblidae). *Zootaxa* 4205 (2). <https://doi.org/10.11646/zootaxa.4205.2.1>
- Theodor O (1967) *An illustrated catalogue of the Rothschild Collection of Nycteribiidae in the British Museum (Nat. Hist.)*. The British Museum, London.
- Zahn A, Rupp D (2004) Ectoparasite load in European vespertilionid bats. *Journal of Zoology* 262 (4): 383-391. <https://doi.org/10.1017/s0952836903004722>

Supplementary material

Suppl. material 1: first occurrence of *B. italica* in Romania [doi](#)

Authors: Áron Péter, Sándor D. Attila

Data type: species occurrence in DwC

Brief description: Collection data of the bat fly *B. italica* for the first time in Romania

[Download file](#) (24.77 kb)