

Efficacy and safety of sodium cantharidinate and vitamin B6 injection for the treatment of digestive system neoplasms: a meta-analysis of randomized controlled trials

This article was published in the following Dove Medical Press journal:
Drug Design, Development and Therapy

Meirong Liu¹
Chunhong Xu²
Yingying Sun³

¹Department of Oncology, Liaocheng People's Hospital, Liaocheng Clinical School of Taishan Medical University, Liaocheng 252000, Shandong Province, China; ²Department of Gastroenterology, Liaocheng People's Hospital, Liaocheng Clinical School of Taishan Medical University, Liaocheng 252000, Shandong Province, China; ³Department of Radiotherapy, Liaocheng People's Hospital, Liaocheng Clinical School of Taishan Medical University, Liaocheng 252000, Shandong Province, China

Objective: To systematically evaluate the efficacy and safety of sodium cantharidinate and vitamin B6 (SC/B6) combined with conventional medical treatment (CMT) for the treatment of patients with advanced digestive system neoplasms (DSNs).

Methods: The Cochrane Library, Embase, PubMed, Web of Science, Chinese Scientific Journal Database (VIP), China National Knowledge Infrastructure, and Wanfang databases were searched for clinical trials using SC/B6 for DSNs. Outcome measures, including therapeutic efficacy, quality of life (QoL), and adverse events, were extracted and systematically evaluated.

Results: Data from 24 trials including 1,825 advanced DSN patients were included. Compared with CMT alone, its combination with SC/B6 significantly improved the patients' overall response rate (OR =2.25, 95% CI =1.83–2.76, $P<0.00001$), disease control rate (OR =2.41, 95% CI =1.85–3.15, $P<0.00001$), and QoL improvement rate (OR =2.75, 95% CI =2.13–3.55, $P<0.00001$). Moreover, adverse events caused by chemotherapy, including leukopenia, nausea and vomiting, gastrointestinal side effects, hepatotoxicity, diarrhea, transaminase disorder, myelosuppression, anorexia, and anemia, were significantly alleviated ($P<0.05$) when SC/B6 was applied to DSN patients. Nephrotoxicity, thrombocytopenia, hand-foot syndrome, and oral mucositis were not significantly alleviated in patients receiving combination therapy ($P>0.05$).

Conclusion: The combination of SC/B6 and CMT is more effective in treating DSNs than CMT alone. This combination alleviates the adverse effects associated with chemotherapy and improves the QoL of DSN patients, and its application in the clinic is worth promoting.

Keywords: sodium cantharidinate and vitamin B6, conventional medical treatment, digestive system neoplasms, meta-analysis

Introduction

Digestive system neoplasms (DSNs) are the leading cause of cancer-related death worldwide, and cause 3,056,412 deaths in 2018, which accounts for 32% of all cancer deaths worldwide.^{1–3} This category comprises colorectal cancer, gastric cancer, liver cancer, esophageal cancer, and pancreatic cancer, which are the fourth, sixth, seventh, ninth, and fourteenth most common cancers, respectively.¹ Despite improvements in diagnostic and therapeutic methods in the past decades,⁴ the prognosis of DSNs is still poor, because they are mostly diagnosed at advanced stages, which may be accompanied by extensive invasion and distant metastasis.^{4–6} Therefore, effective therapeutic approaches should be developed.

Correspondence: Yingying Sun
Department of Radiotherapy, Liaocheng People's Hospital, Liaocheng Clinical School of Taishan Medical University, Dongchang West Road, No 67, Liaocheng 252000, Shandong Province, China
Tel +86 152 7583 9331
Email yingsunyy@163.com

In recent years, traditional Chinese medicine has been more widely used as auxiliary treatment in tumor therapy and has shown promising therapeutic effects in many clinical studies.⁷⁻⁹ Sodium cantharidinate/vitamin B6 (SC/B6) is a combination of sodium cantharidinate (SC) and vitamin B6, and has the pharmacologic characteristics of both.^{7,8} SC is a derivative of cantharidin, which is extracted from the body of meloidae insects such as *Mylabris phalerata pallas* and *Mylabris cichorii linnaeus*.¹⁰ SC preserves the unique anticancer activity of cantharidin and has lower toxicity and fewer adverse effects.^{7,10} Its combination with vitamin B6 can even further lower the side effects.⁷ In recent years, SC has been used as a safe auxiliary antitumor drug for malignancies such as gastric cancer, liver cancer, and non-small-cell lung cancer.^{7-9,11} Tao et al¹² indicated that SC induces HepG2 cells to undergo apoptosis through the LC3 autophagy pathway. Liang et al¹³ showed that SC can inhibit tumor growth by downregulating vascular endothelial growth factor expression and blocking tumor angiogenesis. In addition, SC can also have an anticancer effect by blocking progression through the cell cycle, inhibiting invasion/metastasis, and improving the immunity of cancer patients.¹⁴⁻¹⁸

Several clinical studies^{8,19-41} have revealed the prominent therapeutic effects of SC/B6 and conventional medical treatment (CMT, including chemotherapy, symptomatic, and supporting therapy) for advanced DSNs but clinical efficacy and safety have not been systematically evaluated. In this study, we performed a meta-analysis to evaluate the efficacy and safety of SC/B6 for DSN treatment, with a comparison between SC/B6 and CMT combined therapy and CMT alone, in order to provide scientific reference for the design of future clinical trials.

Materials and methods

Search strategy and selection criteria

Publications were searched across the Cochrane Library, Embase, Pubmed, Web of Science, Chinese Scientific Journal Database (VIP), China National Knowledge Infrastructure, and Wanfang databases, using the search terms “sodium cantharidinate” or “disodium cantharidinate” and “vitamin B6” combined with “gastric cancer” or “colorectal cancer” or “gastrointestinal cancer” or “liver cancer” or “esophageal cancer” or “pancreatic cancer” or “digestive system neoplasms” without restriction on the language. The retrieval was initiated in May 2018 and updated in August 2018.

All of the clinical trials brought into this analysis were randomized controlled trials with reference to advanced DSNs, in which patients in the experimental groups were

treated by SC/B6 and CMT combined therapy, and patients in the control groups were treated by CMT alone.

Data extraction and quality assessment

Literature screening and data extraction were carried out by two independent investigators (Meirong Liu and Chunhong Xu) and verified by a third reviewer (Yingying Sun). All included studies were summarized as follows: first author name, year of publication, study location, Karnofsky Performance Score (KPS), number of cases, patient ages, study parameter type, treatment regimen and enrollment period, and administration route and dosage of SC/B6. The quality of the included trials was evaluated as described in the Cochrane Handbook.⁴²

Outcome definition

Clinical responses, including therapeutic effects, quality of life (QoL), and adverse events, were analyzed. Therapeutic effects were evaluated by overall survival (OS) rate, complete response (CR) rate, partial response (PR) rate, stable disease (SD) rate, progressive disease (PD) rate, overall response rate (ORR, $ORR = CR + PR$), and disease control rate (DCR = $CR + PR + SD$). OS was defined as the length of time from the start of treatment to death from any cause; QoL was assessed using KPS scales and the European Organization for Research and Treatment of Cancer core quality-of-life questionnaire. The QoL improvement rate (QIR) was defined as the improvement in QoL after treatment. Adverse events, including leukopenia, nausea and vomiting, gastrointestinal side effects, hepatotoxicity, nephrotoxicity, diarrhea, thrombocytopenia, transaminase disorder, myelosuppression, hand-foot syndrome, oral mucositis, anorexia, and anemia, were also assessed.

Statistical analysis

Review Manager 5.3 (Nordic Cochrane Centre, Copenhagen, Denmark) and Stata 13.0 (Stata Corp., College Station, TX, USA) were the main statistical analysis tools in this study. $P < 0.05$ indicated statistically significant differences. Cochran's Q test was used to determine heterogeneity among studies,⁴³ and publication bias was analyzed by Begg's and Egger's regression asymmetry tests and presented by funnel plots.⁴⁴ $I^2 < 50\%$ or $P > 0.1$ indicated study homogeneity. Therapeutic effects were mainly represented by HRs and ORs presented with 95% CIs. HRs were collected for survival data. If HRs can neither be collected directly nor calculated, survival curve plots were extracted by Engauge Digitizer software and then transformed by specialized form.⁴⁵⁻⁴⁷

Pooled analysis with publication bias determined that the trim-and-fill method would be applied to coordinate the estimates of unpublished studies, and the adjusted results were compared with the original pooled OR.⁴⁸ Sensitivity analysis (subgroup analyses) was conducted to evaluate the impact of different cancer types, SC/B6 dosages, therapeutic regimens, sample sizes, and study types on clinical efficacy.

Results

Search results

A total of 974 articles were identified with the initial search, and 602 papers were excluded due to duplication. After title and abstract review, 269 articles were further excluded because they did not include clinical trials (n=209), were reviews or meta-analyses (n=6), were unrelated studies (n=43), or were case reports (n=11), leaving 103 studies as potentially relevant. After detailed assessment of full texts,

articles without a control group (n=11), studies with inappropriate criteria in the experimental or control group (n=16), studies with insufficient data (n=5), and studies including patients with non-digestive system tumors (n=47) were excluded. Finally, data from 24 trials^{8,19-41} (gastric cancer, n=7; colorectal cancer, n=5; gastrointestinal cancer, n=3; liver cancer, n=7; esophageal cancer, n=1; and pancreatic cancer, n=1) including 1,825 advanced DSN patients were included in the present analysis (Figure 1).

Patient characteristics

All studies involved in this analysis were carried out in different hospitals in China. These trials include 1,825 patients with advanced DSNs; of these, 933 were treated by combined SC/B6 and CMT, and 892 were treated by CMT alone. Detailed information on the included trials and patients is presented in Tables 1 and 2.




Figure 1 Flow diagram of the selection process.

Table 1 Clinical information from the eligible trials in the meta-analysis

Included studies	Nation	KPS	Patients Con/Exp	Age (years)		Parameter types
				Con	Exp	
Chen Y 2016 ¹⁹	China	ND	25/25	61.27±1.46 (mean)	61.25±1.44 (mean)	ORR, DCR, QIR, AE
Fan LJ 2009 ²⁰	China	KPS ≥60	42/42	51.5 (mean)	52.3 (mean)	ORR, DCR
Fan QL 2013 ²¹	China	KPS >60	19/23	ND	ND	ORR, DCR, QIR, AE
Fang XH 2016 ²²	China	KPS >50	37/37	64.3±10.3 (mean)	66.3±9.3 (mean)	ORR, DCR
Guan LY 2015 ²³	China	KPS >60	27/27	ND	ND	ORR, DCR, QIR, AE
Jia JM 2013 ²⁴	China	KPS ≥60	18/18	ND	ND	ORR, DCR, QIR, AE
Li GP 2010 ²⁵	China	KPS >60	25/25	40–58	42–65	AE
Liu GW 2017 ²⁶	China	KPS ≥60	20/20	35–76 (mean)	37–74 (mean)	ORR, DCR, QIR, AE
Liu SH 2008 ²⁷	China	60–90 (KPS)	32/32	54.7 (mean)	52.2 (mean)	ORR, DCR, QIR, AE
Mao WD 2016 ²⁸	China	KPS ≥70	32/33	56.3±15.5 (mean)	55.7±17.2 (mean)	ORR, DCR, AE
Shao H 2014 ⁸	China	ND	41/63	41.71±8.55 (mean)	38.74±11.06 (mean)	ORR, DCR
Shi XY 2017 ²⁹	China	KPS >60	48/48	62.14±11.23 (mean)	61.59±11.02 (mean)	ORR, DCR, QIR, AE
Tian XL 2006 ³⁰	China	KPS ≥70	36/36	52.5±9.6 (mean)	53.4±10.5 (mean)	ORR, DCR, QIR, AE
Wang JH 2010 ³¹	China	50–90 (KPS)	26/26	51.79 (mean)	53.26 (mean)	ORR, DCR, QIR, AE
Wang YW 2017 ³²	China	KPS ≥70	42/42	62.1±10.2 (mean)	61.2±9.7 (mean)	ORR, DCR, QIR, AE
Wei YF 2015 ³³	China	KPS >70	44/48	ND	ND	ORR, DCR, AE
Wu ZM 2013 ³⁴	China	ND	32/32	ND	ND	ORR, DCR, AE
Xie ZX 2016 ³⁵	China	ND	32/32	58.1±3.2 (mean)	57.3±2.8 (mean)	ORR, DCR, QIR, AE
You ZY 2015 ³⁶	China	KPS ≥60	85/85	ND	ND	ORR, DCR, QIR
Zeng L 2009 ³⁷	China	60–80 (KPS)	63/63	ND	ND	ORR, DCR, QIR
Zhang MJ 2011 ³⁸	China	KPS ≥60	38/38	55.0±2.2 (mean)	54.0±2.4 (mean)	ORR, DCR, QIR, AE
Zhang W 2012 ³⁹	China	KPS ≥70	42/42	61.2 (mean)	62.1 (mean)	ORR, DCR
Zhang W 2015 ⁴⁰	China	KPS ≥70	36/48	59.6 (median)	54.2 (median)	ORR, DCR, QIR, AE
Zhu WQ 2014 ⁴¹	China	ND	50/48	ND	ND	ORR, DCR, AE

Abbreviations: AE, adverse events; CMT, conventional medical treatment; Con, control group (CMT alone group); DCR, disease control rate; Exp, experimental group (SC/B6 plus CMT combined group); KPS, Karnofsky Performance Score; ND, nondetermined; ORR, overall response rate; QIR, quality-of-life improved rate; SC/B6, sodium cantharidinate and vitamin B6 injection.

Quality assessment

The evaluation of bias risk is presented in Figure 2. Twenty-two studies had low risk, and the other two articles did not have a clear description of the randomization process. None of the included trials provided a clear description of the performance and detection risks. Two studies were regarded as high-risk due to the absence of follow-up and seven trials were considered as unclear risk owing to selective reporting.

Therapeutic efficacy assessments

As shown in Figures 3 and 4, Table 3, and Figure S1, patients who underwent combined therapy had a significantly improved CR rate (OR = 2.06, 95% CI = 1.41–3.00, $P=0.0002$), PR rate (OR = 1.85, 95% CI = 1.50–2.29, $P<0.00001$), ORR (OR = 2.25, 95% CI = 1.83–2.76, $P<0.00001$), and DCR (OR = 2.41, 95% CI = 1.85–3.15, $P<0.00001$), and

significantly decreased SD and PD rates (SD, OR = 0.77, 95% CI = 0.63–0.93, $P=0.009$; PD, OR = 0.45, 95% CI = 0.35–0.59, $P<0.00001$) compared to patients receiving CMT alone. The OS rates of patients who received combination treatment (HR = 0.74, 95% CI = 0.47–1.17, $P=0.20$) did not differ significantly from those in patients who received CMT alone.

QoL assessment

QoL evaluation demonstrated that SC/B6 and CMT combined therapy-treated DSN patients had improved QoL compared to those treated by CMT alone (Figure 5A, OR = 2.75, 95% CI = 2.13–3.55, $P<0.00001$).

Adverse events assessment

As shown in Table 4 and Figure S2, patients treated by SC/B6 and CMT combined therapy had lower incidences

Table 2 Information of SC/B6 combined with conventional medical treatment

Included studies	Therapeutic regimen		Enrollment period	Dosage of apatinib
	Experimental group	Control group		
Chen Y 2016 ¹⁹	CMT + SC/B6	CMT (raltitrexed, oxaliplatin)	2,013.4–2,016.4	30 mL/time (0.1 mg/10 mL, IV), 1 time/day
Fan LJ 2009 ²⁰	CMT + SC/B6	CMT (calcium folinate, 5-Fu)	2,005.2–2,009.7	30 mL/time (0.1 mg/10 mL, IV), 1 time/day
Fan QL 2013 ²¹	CMT + SC/B6	CMT (S-1)	ND	20 mL/time (0.1 mg/10 mL, IV), 1 time/day
Fang XH 2016 ²²	CMT + SC/B6	CMT (ND)	2,012.1–2,014.8	40 mL/time (0.1 mg/10 mL, IV), 1 time/day
Guan LY 2015 ²³	CMT + SC/B6	CMT (S-1)	2,012.10–2,014.10	50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Jia JM 2013 ²⁴	CMT + SC/B6	CMT (oxaliplatin, paclitaxel)	2,011.1–2,012.10	20 mL/time (0.1 mg/10 mL, IV), 1 time/day
Li GP 2010 ²⁵	CMT + SC/B6	CMT (FOLFOX4)	2,008.3–2,009.9	40 mL/time (0.1 mg/10 mL, IV), 1 time/day
Liu GW 2017 ²⁶	CMT + SC/B6	CMT (capecitabine)	2,014.1–2,016.1	40 mL/time (0.1 mg/10 mL, IV), 1 time/day
Liu SH 2008 ²⁷	CMT + SC/B6	CMT (leucovorin, oxaliplatin)	2,005.1–2,007.1	30 mL/time (0.1 mg/10 mL, IV), 1 time/day
Mao WD 2016 ²⁸	CMT + SC/B6	CMT (capecitabine)	2,012.6–2,013.12	30 mL/time (0.1 mg/10 mL, IV), 1 time/day
Shao H 2014 ⁸	CMT + SC/B6	CMT (ND)	2,011.1–2,012.11	50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Shi XY 2017 ²⁹	CMT + SC/B6	CMT (XELOX)	2,013.12–2,015.12	20 mL/time (0.1 mg/10 mL, IV), 1 time/day
Tian XL 2006 ³⁰	CMT + SC/B6	CMT (mitomycin, adriamycin/5-Fu, cisplatin)	2,001.9–2,003.9	50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Wang JH 2010 ³¹	CMT + SC/B6	CMT (FOLFOX4)	2,008.1–2,009.10	50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Wang YW 2017 ³²	CMT + SC/B6	CMT (capecitabine)	2,016.6–2,017.6	20 mL/time (0.1 mg/10 mL, IV), 1 time/day
Wei YF 2015 ³³	CMT + SC/B6	CMT (5-Fu, epirubicin, mitomycin)	2,010.1–2,011.9	80 mL/time (0.1 mg/10 mL, IV), 1 time/day
Wu ZM 2013 ³⁴	CMT + SC/B6	CMT (FOLFIRI)	2,008.5–2,011.1	50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Xie ZX 2016 ³⁵	CMT + SC/B6	CMT (oxaliplatin, S-1)	2,013.4–2,015.4	40 mL/time (0.1 mg/10 mL, IV), 1 time/day
You ZY 2015 ³⁶	CMT + SC/B6	CMT (cisplatin, 5-Fu)	2,010.4–2,012.6	ND
Zeng Li 2009 ³⁷	CMT + SC/B6	CMT (ND)	2,005.3–2,008.6	30–50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Zhang MJ 2011 ³⁸	CMT + SC/B6	CMT (mitomycin, adriamycin)	ND	50 mL/time (0.1 mg/10 mL, IV), 1 time/day
Zhang W 2012 ³⁹	CMT + SC/B6	CMT (capecitabine)	2,007.2–2,011.7	30 mL/time (0.1 mg/10 mL, IV), 1 time/day
Zhang W 2015 ⁴⁰	CMT + SC/B6	CMT (XELOX)	2,012.3–2,014.12	30 mL/time (0.1 mg/10 mL, IV), 1 time/day
Zhu WQ 2014 ⁴¹	CMT + SC/B6	CMT (ND)	2,008.3–2,012.3	50 mL/time (0.1 mg/10 mL, IV), 1 time/day

Abbreviations: 5-Fu, 5-fluorouracil; CMT, conventional medical treatment; Con, control group (CMT alone group); Exp, experimental group (SC/B6 plus CMT combined group); FOLFOX, oxaliplatin + calcium folinate + 5-fluorouracil; FOLFIRI, calcium folinate + irinotecan + 5-fluorouracil; IV, intravenous; S-1, gimeracil and oteracil potassium capsules; ND, nondetermined; SC/B6, sodium cantharidinate and vitamin B6 injection; XELOX, oxaliplatin + capecitabine.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Chen Y 2016 ¹⁹	●	●	?	?	+	+	?
Fang XH 2016 ²²	+	+	?	?	+	?	?
Fan LJ 2009 ²⁰	+	+	?	?	+	?	+
Fan QL 2013 ²¹	+	+	?	?	+	+	?
Guan LY 2015 ²³	+	+	?	?	+	+	?
Jia JM 2013 ²⁴	+	+	?	?	+	+	?
Li GP 2010 ²⁵	+	+	?	?	+	?	+
Liu GW 2017 ²⁶	+	+	?	?	+	+	+
Liu SH 2008 ²⁷	+	+	?	?	+	+	+
Mao WD 2016 ²⁸	+	+	?	?	+	+	+
Shao H 2014 ⁸	+	+	?	?	●	?	?
Shi XY 2017 ²⁹	●	●	?	?	+	+	+
Tian XL 2006 ³⁰	+	+	?	?	●	+	+
Wang JH 2010 ³¹	+	+	?	?	+	+	+
Wang YW 2017 ³²	+	+	?	?	+	+	+
Wei YF 2015 ³³	+	+	?	?	+	+	?
Wu ZM 2013 ³⁴	+	+	?	?	+	+	?
Xie ZX 2016 ³⁵	+	+	?	?	+	+	+
You ZY 2015 ³⁶	+	+	?	?	+	?	?
Zeng Li 2009 ³⁷	+	+	?	?	+	?	?
Zhang MJ 2011 ³⁸	+	+	?	?	+	+	?
Zhang W 2012 ³⁹	+	+	?	?	+	?	+
Zhang W 2015 ⁴⁰	+	+	?	?	+	+	+
Zhu WQ 2014 ⁴¹	+	+	?	?	+	+	?

Figure 2 Risk of bias summary: review of authors' judgments about each risk of bias item for included studies.
Note: Each color represents a different level of bias: red for high risk, green for low risk, and yellow for unclear risk of bias.

of leukopenia, nausea and vomiting, gastrointestinal side effects, hepatotoxicity, diarrhea, transaminase disorder, myelosuppression, anorexia, and anemia than those treated with CMT alone (leukopenia: OR =0.29, 95% CI=0.21–0.39, $P<0.00001$; nausea and vomiting: OR =0.30, 95% CI =0.22–0.40, $P<0.00001$; gastrointestinal side effects: OR=0.42, 95% CI=0.29–0.62, $P<0.00001$; hepatotoxicity: OR=0.49, 95% CI=0.30–0.78, $P=0.003$; diarrhea: OR =0.37, 95% CI =0.23–0.60, $P<0.0001$; transaminase disorder: OR =0.23, 95% CI =0.09–0.62, $P=0.003$; myelosuppression: OR =0.33, 95% CI =0.18–0.60, $P=0.0003$; anorexia: OR=0.37, 95% CI=0.20–0.68, $P=0.001$; anemia: OR=0.54, 95% CI=0.32–0.91, $P=0.02$). No significant difference was found in the occurrence of nephrotoxicity, thrombocytopenia, hand-foot syndrome, and oral mucositis (nephrotoxicity: OR =0.70, 95% CI=0.38–1.30, $P=0.26$; thrombocytopenia: OR =0.77, 95% CI=0.31–1.92, $P=0.57$; hand-foot syndrome: OR =0.75, 95% CI =0.40–1.40, $P=0.36$; oral mucositis: OR =0.45, 95% CI =0.13–1.62, $P=0.22$) between patients receiving combination treatment and those receiving CMT alone.

Publication bias

Publication bias of primary outcomes (CR, PR, SD, PD, ORR, DCR, QIR, and adverse events) was evaluated and presented by funnel plots. All plots were approximately symmetrical, indicating generally controlled publication bias (Figures 6 and S3).

We also assessed the publication bias by Begg's and Egger's regression asymmetry tests, and SD and leukopenia were found to have bias (SD, Egger: 0.024, Begg: 0.039; leukopenia, Egger: 0.041, Begg: 0.080; Table 5). To determine whether the bias affected the pooled risk, we conducted trim-and-fill analysis. The adjusted OR indicated the same trend as the primary analysis (SD, before: $P=0.010$, after: $P<0.0001$; leukopenia, before: $P<0.0001$, after: $P<0.0001$), reflecting the reliability of our primary conclusions, except those based on a small number of trials.

Sensitivity analysis

Subgroup analysis was performed for ORR and DCR heterogeneity assessment concerning cancer types, SC/B6 dosages, therapeutic regimens, sample sizes, and study types of involved trials. No significant difference was observed in the sample sizes, study types, or SC/B6 dosages (Table 6). SC/B6 combined with CMT was more effective in treating gastric cancer, colorectal cancer, and liver cancer. Moreover, SC/B6




Figure 3 Forest plot of the comparison of overall survival between the experimental and control groups.
Notes: Control group, CMT-alone group; Experimental group, sodium cantharidinate and vitamin B6 injection (SC/B6) + CMT. The fixed-effects meta-analysis model (inverse variance method) was used.
Abbreviations: CMT, conventional medical treatment; IV, intravenous.

combined with oxaliplatin and capecitabine (XELOX) or capecitabine regimens was more effective for DSN treatment.

Discussion

The chemotherapeutic regimens commonly used to treat DSNs cause serious side effects, such as myelosuppression, hepatotoxicity, and gastrointestinal side effects, which severely affect the QoL of DSN patients.^{7,9} Therefore, seeking a therapy that can improve treatment outcomes and decrease

the adverse effects of chemotherapy is a major direction in the development of tumor treatment. Traditional Chinese medicine plays a unique role in improving host immunity and lowering the toxic effects of chemotherapy.^{7,9,49-52} In recent decades, SC/B6 has been clinically applied as an adjuvant therapy for malignancies and has been beneficial for advanced DSN patients in several trials.^{7-9,11} Despite the published reports on clinical trials using SC/B6, its therapeutic effects have not been systematically demonstrated.

A




Figure 4 (Continued)

B




Figure 4 Forest plot of the comparison of overall response rate (A) and disease control rate (B) between the experimental and control groups.

Notes: Control group, CMT-alone group; Experimental group, sodium cantharidinate and vitamin B6 injection (SC/B6) + CMT. The fixed-effects meta-analysis model (M-H method) was used.

Abbreviations: CMT, conventional medical treatment; M-H, Mantel-Haenszel.

In the present study, we performed an extensive literature search followed by rigorous contrasting and combining data analysis for categorization to provide clear and systematic conclusions.

Our meta-analysis revealed that SC/B6 and CMT combined therapy for DSN patients achieved more beneficial effects than CMT alone. Combined therapy-treated patients

exhibited markedly improved ORR and DCR ($P<0.05$ for all) and also significantly improved QoL. These results indicated that intravenous infusion of SC/B6 improved the curative effects of CMT for advanced DSNs.

Our analysis indicates that most of the adverse events caused by chemotherapy, including leukopenia, nausea and vomiting, gastrointestinal side effects, and hepatotoxicity,

Table 3 Comparison of CR, PR, SD, PD, ORR, and DCR between the SC/B6 + CMT and SC/B6 group

Parameter	SC/B6 + CMT group	CMT group	Analysis method	Heterogeneity		OR	95% CI	P-value
	No of patients (n)	No of patients (n)		I^2 (%)	P-value			
CR	889	840	Fixed	0	0.99	2.06	1.41–3.00	0.0002
PR	889	840	Fixed	0	0.89	1.85	1.50–2.29	<0.00001
SD	889	840	Fixed	43	0.01	0.77	0.63–0.93	0.009
PD	889	840	Fixed	0	0.91	0.45	0.35–0.59	<0.00001
ORR	889	840	Fixed	0	0.56	2.25	1.83–2.76	<0.00001
DCR	889	840	Fixed	0	0.93	2.41	1.85–3.15	<0.00001

Abbreviations: CMT, conventional medical treatment; CR, complete response rates; DCR, disease control rate; ORR, overall response rate; PD, progressive disease rates; PR, partial response rates; SC/B6, sodium cantharidinate and vitamin B6 injection; SD, stable disease rates.




Figure 5 Forest plot of the comparison of quality-of-life improved rate between the experimental and control groups.

Notes: Control group, CMT-alone group; Experimental group, SC/B6 + CMT. The fixed-effects meta-analysis model (M-H method) was used.

Abbreviations: CMT, conventional medical treatment; M-H, Mantel-Haenszel; SC/B6, sodium cantharidinate and vitamin B6 injection.

Table 4 Comparison of adverse events between the SC/B6 + CMT and SC/B6 group

Adverse events	SC/B6 + CMT group No patients (n)	CMT group No patients (n)	Analysis method	Heterogeneity		OR	95% CI	P-value
				I^2 (%)	P-value			
Leucopenia	449	427	Fixed	0	0.72	0.29	0.21–0.39	<0.00001
Leucopenia I + II	364	344	Fixed	0	0.92	0.39	0.28–0.54	<0.00001
Leucopenia III + IV	399	377	Fixed	0	0.99	0.36	0.21–0.63	0.0003
Nausea, vomiting	407	393	Fixed	0	0.93	0.30	0.22–0.40	<0.00001
Nausea, vomiting I + II	242	226	Fixed	0	0.97	0.28	0.19–0.43	<0.00001
Nausea, vomiting III + IV	242	226	Fixed	0	1.00	0.59	0.23–1.51	0.27
Gastrointestinal side effects	278	271	Fixed	0	0.97	0.42	0.29–0.62	<0.00001
Gastrointestinal side effects I + II	167	162	Fixed	0	0.81	0.49	0.30–0.80	0.004
Gastrointestinal side effects III + IV	190	182	Fixed	0	0.54	0.37	0.17–0.79	0.01
Hepatotoxicity	262	257	Fixed	0	0.67	0.49	0.30–0.78	0.003
Hypertension I + II	206	201	Fixed	0	0.69	0.54	0.31–0.94	0.03
Hypertension III + IV	206	201	Fixed	0	0.79	0.44	0.12–1.61	0.22
Nephrotoxicity	277	272	Fixed	0	0.95	0.70	0.38–1.30	0.26
Nephrotoxicity I + II	154	149	Fixed	0	1.00	0.89	0.39–2.08	0.80
Nephrotoxicity III + IV	154	149	Fixed	Not applicable		1.00	0.14–7.40	1.00
Diarrhea	192	176	Fixed	0	0.61	0.37	0.23–0.60	<0.0001
Diarrhea I + II	192	176	Fixed	0	0.74	0.38	0.23–0.62	<0.0001
Diarrhea III + IV	192	176	Fixed	0	0.81	0.58	0.15–2.30	0.44
Thrombocytopenia	143	169	Random	63	0.03	0.77	0.31–1.92	0.57
Thrombocytopenia I + II	141	137	Fixed	0	0.69	0.50	0.27–0.92	0.03
Thrombocytopenia III + IV	141	137	Fixed	0	0.98	0.43	0.09–1.95	0.27

(Continued)

Table 4 (Continued)

Adverse events	SC/B6 + CMT group	CMT group	Analysis method	Heterogeneity		OR	95% CI	P-value
	No patients (n)	No patients (n)		I ² (%)	P-value			
Transaminase disorder	149	145	Random	55	0.07	0.23	0.09–0.62	0.003
Transaminase disorder I + II	117	113	Fixed	0	0.40	0.33	0.15–0.69	0.004
Transaminase disorder III + IV	117	113	Fixed	0	0.80	0.46	0.08–2.57	0.38
Myelosuppression	151	152	Fixed	0	0.90	0.33	0.18–0.60	0.0003
Myelosuppression I + II	151	152	Random	79	0.003	0.70	0.23–2.08	0.52
Myelosuppression III + IV	113	114	Random	0	0.81	0.28	0.11–0.73	0.009
Hand-foot syndrome	116	104	Fixed	0	0.39	0.75	0.40–1.40	0.36
Hand-foot syndrome I + II	116	104	Fixed	0	0.70	0.83	0.44–1.57	0.56
Hand-foot syndrome III + IV	116	104	Fixed	0	0.51	0.49	0.10–2.41	0.38
Oral mucositis	45	45	Fixed	0	0.98	0.45	0.13–1.62	0.22
Oral mucositis I + II	45	45	Fixed	0	0.63	0.34	0.07–1.59	0.17
Oral mucositis III + IV	45	45	Fixed	Not applicable		1.00	0.13–7.72	1.00
Anorexia	92	88	Fixed	39	0.20	0.37	0.20–0.68	0.001
Anorexia I + II	92	88	Fixed	39	0.20	0.37	0.20–0.68	0.001
Anorexia III + IV	92	88	Fixed	Not applicable				
Anemia	162	162	Fixed	0	0.73	0.54	0.32–0.91	0.02
Anemia I + II	77	77	Fixed	0	0.49	0.60	0.31–1.16	0.13
Anemia III + IV	77	77	Fixed	0	0.84	0.41	0.06–2.90	0.37

Abbreviations: CMT, conventional medical treatment; SC/B6, sodium cantharidinate and vitamin B6 injection.




Figure 6 Funnel plot of percentage of overall response rate (A), disease control rate (B), quality-of-life improved rate (C), leukopenia (D), nausea and vomiting (E), gastrointestinal side effects (F), and hepatotoxicity (G).

Note: Parameters discussed in over eight papers were conducted bias analyses.

Table 5 Publication bias on therapeutic efficacy and adverse events

Publication bias	Therapeutic efficacy							Adverse events			
	CR	PR	SD	PD	ORR	DCR	QIR	Leukopenia	Nausea and vomiting	Hepatotoxicity	Gastrointestinal side effects
Begg	0.058	0.154	0.039	0.195	0.369	0.612	1.000	0.080	0.213	0.386	0.711
Egger	0.078	0.259	0.024	0.149	0.489	0.425	0.808	0.041	0.697	0.198	0.581

Note: Parameters discussed in over eight papers were conducted bias analyses.

Abbreviations: CR, complete response rates; DCR, disease control rate; ORR, overall response rate; PD, progressive disease rates; PR, partial response rates; QIR, quality-of-life improved rate; SD, stable disease rates.

Table 6 Subgroup analyses of ORR and DCR between the SC/B6 + CMT and SC/B6 groups

Parameter	Factors at study level	Exp group	Con group	Analysis method	Heterogeneity		OR	95% CI	P-value
		No patients (n)	No patients (n)		I ² (%)	P-value			
ORR	Type of cancer								
	Gastric cancer	219	206	Fixed	0	0.60	1.78	1.20–2.66	0.005
	Colorectal cancer	161	161	Fixed	0	0.79	2.60	1.59–4.26	0.0001
	Gastrointestinal cancer	150	146	Fixed	0	0.96	2.48	1.53–4.02	0.0002
	Liver cancer	314	282	Fixed	54	0.04	2.42	1.70–3.43	<0.00001
	Esophageal cancer	18	18	Fixed			2.00	0.52–7.69	0.31
	Pancreatic cancer	27	27	Fixed			1.56	0.24–10.19	0.64
	Dosage of SC/B6								
	20 mL/day	131	127	Fixed	0	0.93	2.16	1.28–3.66	0.004
	30 mL/day	222	209	Fixed	0	0.81	2.37	1.56–3.58	<0.0001
	40 mL/day	89	89	Fixed	0	0.61	2.19	1.17–4.09	0.01
	50 mL/day	251	223	Fixed	0	0.54	1.68	1.12–2.52	0.01
	Therapeutic regimen								
	SC/B6 + XELOX	96	84	Fixed	0	0.96	1.83	0.97–3.45	0.06
	SC/B6 + S-1	50	46	Fixed	0	0.76	2.00	0.71–5.63	0.19
	SC/B6 + capecitabine	137	136	Fixed	0	0.94	2.91	1.70–4.97	<0.0001
	Study sample size								
	>80	511	469	Fixed	10	0.35	2.70	2.04–3.57	<0.00001
	<80	378	371	Fixed	0	0.84	1.80	1.32–2.44	0.0002
	Type of control trials								
RCT	816	767	Fixed	0	0.50	2.24	1.81–2.78	<0.00001	
Overall	889	840	Fixed	0	0.56	2.25	1.83–2.76	<0.00001	
DCR	Type of cancer								
	Gastric cancer	219	206	Fixed	0	0.67	2.32	1.43–3.76	0.0006
	Colorectal cancer	161	161	Fixed	0	0.91	2.41	1.37–4.26	0.002
	Gastrointestinal cancer	150	146	Fixed	0	0.61	2.32	0.90–6.02	0.08
	Liver cancer	314	282	Fixed	0	0.69	2.65	1.64–4.27	<0.0001
	Esophageal cancer	18	18	Fixed			10.82	1.17–100.44	0.04
	Pancreatic cancer	27	27	Fixed			1.16	0.40–3.43	0.78
	Dosage of SC/B6								
	20 mL/day	131	127	Fixed	0	0.59	2.87	1.54–5.35	0.0009
	30 mL/day	222	209	Fixed	0	0.79	2.39	1.45–3.93	0.0006
	40 mL/day	89	89	Fixed	46	0.17	2.22	0.91–5.45	0.08
	50 mL/day	251	223	Fixed	0	0.55	2.45	1.53–3.93	0.0002

(Continued)

Table 6 (Continued)

Parameter	Factors at study level	Exp group	Con group	Analysis method	Heterogeneity		OR	95% CI	P-value
		No patients (n)	No patients (n)		I ² (%)	P-value			
Therapeutic regimen									
	SC/B6 + XELOX	96	84	Fixed	0	0.39	2.76	1.48–5.14	0.001
	SC/B6 + S-1	50	46	Fixed	18	0.27	1.63	0.66–4.01	0.29
	SC/B6 + capecitabine	137	136	Fixed	0	0.88	2.08	0.97–4.45	0.06
Study sample size									
	>80	511	469	Fixed	0	0.97	2.40	1.64–3.51	<0.00001
	<80	378	371	Fixed	0	0.59	2.42	1.67–3.52	<0.00001
Type of control trials									
	RCT	816	767	Fixed	0	0.89	2.40	1.80–3.20	<0.00001
	Overall	889	840	Fixed	0	0.93	2.41	1.85–3.15	<0.00001

Abbreviations: CMT, conventional medical treatment; Con, control group (CMT alone group); DCR, disease control rate; Exp, experimental group (SC/B6 plus CMT combined group); RCT, randomized controlled trial; ORR, overall response rate; S-1, gimeracil and oteracil porassium capsules; SC/B6, sodium cantharidinate and vitamin B6 injection; XELOX, oxaliplatin + capecitabine.

were alleviated with SC/B6 combination therapy ($P < 0.05$). Therefore, SC/B6 is a safe auxiliary antitumor medicine for DSN and can effectively alleviate the adverse events associated with chemotherapy.

The analysis of therapeutic effects may be influenced by several factors. In our study, no difference was found between sample sizes, study types, and SC/B6 dosages. SC/B6 combined with CMT was more effective in treating gastric cancer, colorectal cancer, and liver cancer than it was in treating esophageal cancer and pancreatic cancer. Moreover, our subgroup analysis showed that SC/B6 combined with XELOX/capecitabine was more effective for DSN treatment. However, recent studies on the impact of these factors on the curative effect of SC/B6 adjuvant therapy remain insufficient, and further investigations should be performed.

There are some limitations in our analysis. First, the follow-up durations of the included studies were not long enough. Second, as a traditional medicine, SC/B6 was mainly applied in China, which may bring an unavoidable regional bias and subsequently influence the clinical application of SC/B6 worldwide. Furthermore, treatment/medical history is very important for evaluating the efficacy of SC/B6-mediated therapy. However, our data were extracted from published papers rather than from the original patient records; therefore, analytical bias may possibly exist. More original data would be valuable to achieve a higher reliability of statistical analysis on SC/B6 for DSN treatment.

In summary, this meta-analysis indicated that SC/B6 and CMT combined therapy was effective in treating advanced DSNs. Intravenous infusion of SC/B6 not only greatly

improved the therapeutic effects of CMT but also effectively alleviated the toxicity and most of the side effects associated with chemotherapy. Therefore, SC/B6 has potential for development as a new adjuvant therapy for the treatment of DSN.

Author contributions

All authors contributed to data analysis, drafting or revising the article, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2018;68(6):394–424.
- Fidler MM, Bray F, Soerjomataram I. The global cancer burden and human development: a review. *Scand J Public Health*. 2018;46(1):27–36.
- Gong J, Lv L, Huo J. Roles of F-box proteins in human digestive system tumors (review). *Int J Oncol*. 2014;45(6):2199–2207.
- Li MX, Bi XY, Huang Z, et al. Prognostic role of phospho-STAT3 in patients with cancers of the digestive system: a systematic review and meta-analysis. *PLoS One*. 2015;10(5):e0127356.
- Dai C, Wang M, Lu J, et al. Prognostic and predictive values of PD-L1 expression in patients with digestive system cancer: a meta-analysis. *Onco Targets Ther*. 2017;10:3625–3634.
- Fu M, Zou C, Pan L, et al. Long noncoding RNAs in digestive system cancers: Functional roles, molecular mechanisms, and clinical implications (review). *Oncol Rep*. 2016;36(3):1207–1218.
- Wang B, Cui J. Treatment of mid-late stage NSCLC using sodium cantharidinate/vitamin B6/GP regimen in clinic. *J Cancer Res Ther*. 2014;10(Suppl 1):C79–C81.
- Shao H, Hong G, Luo X. Evaluation of sodium cantharidinate/vitamin B6 in the treatment of primary liver cancer. *J Cancer Res Ther*. 2014;10(Suppl 1):75–78.

9. Shi Z, Song T, Wan Y, et al. A systematic review and meta-analysis of traditional insect Chinese medicines combined chemotherapy for non-surgical hepatocellular carcinoma therapy. *Sci Rep*. 2017;7(1):4355.
10. Wen SQ, Chen Q, Hu M. Experimental study on the inhibitory effect of sodium cantharidinate on human hepatoma HepG2 cells. *Afr J Tradit Complement Altern Med*. 2014;11(1):131–134.
11. Zhang D, Wu J, Wang K, Duan X, Liu S, Zhang B. Which are the best Chinese herbal injections combined with XELOX regimen for gastric cancer?: A PRISMA-compliant network meta-analysis. *Medicine*. 2018; 97(12):e0127.
12. Tao R, Sun WY, Yu DH, et al. Sodium cantharidinate induces HepG2 cell apoptosis through LC3 autophagy pathway. *Oncol Rep*. 2017;38(2): 1233–1239.
13. Liang F, Wang MY, Huang WB, Li AJ. Effect of sodium cantharidinate on the angiogenesis of nude mice with human gastric cancer. *Zhong Yao Cai*. 2011;34(3):343–346.
14. Zhao RL, Chen MJ, Zhao FM, Xu DQ, Zhang X, Zhou KF. Study on synergistic effect of sodium cantharidinate combined with chemotherapeutic drugs on hepatic carcinoma and its effective mechanism. *Zhong Yao Cai*. 2014;37(11):1938–1946.
15. Tao R, Wang ZF, Qiu W, et al. Role of S100A3 in human hepatocellular carcinoma and the anticancer effect of sodium cantharidinate. *Exp Ther Med*. 2017;13(6):2812–2818.
16. Zang GH, Li R, Zhou RS, et al. Effects of disodium cantharidinate on dendritic cells of patients with bladder carcinoma. *Oncol Lett*. 2018; 15(2):2273–2277.
17. Verma AK, Prasad SB. Changes in glutathione, oxidative stress and mitochondrial membrane potential in apoptosis involving the anticancer activity of cantharidin isolated from redheaded blister beetles, epicauta hirticornis. *Anticancer Agents Med Chem*. 2013;13(7):1096–1114.
18. Li W, Xie L, Chen Z, et al. Cantharidin, a potent and selective PP2A inhibitor, induces an oxidative stress-independent growth inhibition of pancreatic cancer cells through G2/M cell-cycle arrest and apoptosis. *Cancer Sci*. 2010;101(5):1226–1233.
19. Chen Y, Zhu WM, Zhou DX, Shan YF. Clinical study on disodium cantharidinate and vitamin B6 combined with raltitrexed and oxaliplatin in treatment of advanced colorectal cancer. *Drugs Clinic*. 2016; 31(11):1784–1787.
20. Fan LJ. Efficacy of sodium cantharidinate and vitamin B6 combined with 5-fluorouracil and calcium folinate in treatment of advanced colorectal cancer. *Mod Med J China*. 2009;11(10):64–65.
21. Fan QL, Wang Y, Zhao KX, Kao J. Clinical observation on effects of sodium cantharidinate vitamin B6 injection combined with S-1 treatment of advanced gastrointestinal tract tumor patients. *Guangming J Chin Med*. 2013;28(4):786–788.
22. Fang XH, Xu J. Cantharidin sodium vitamin B6 injection as assistant therapy for treating liver cancer in 37 cases. *China Pharm*. 2016;25(1): 25–27.
23. Guan LY, Ma J, Yuan SF, Qu J. Disodium cantharidinate and vitamin B6 combined with tegafur, gimeracil and oteracil potassium for treating stage IV pancreatic cancer in 27 cases. *Tradit Med*. 2015;24(20):124–125.
24. Jia JM, Li ZH. Efficacy of sodium cantharidinate and vitamin B6 combined with intravenous chemotherapy for advanced esophageal cancer. *Med Inf*. 2013;26(9):188–189.
25. Li GP, Li XR, Wang JH. Clinical observation on sodium cantharidate vitamin B6 injection combined with postoperative chemotherapy in treatment of gastric cancer. *Pract Prev Med*. 2010;17(1):105–106.
26. Liu GW, Ren WD. Cantharis acid sodium vitamin B6 injection combined with maintain capecitabine synchronization for the treatment of advanced colorectal cancer. *World Latest Med Inf*. 2017;17(26):12–14.
27. Liu SH, Feng W, Zhu RX, Feng ZC. Comparative study of disodium cantharidinate and vitamin B6 injection combined with oxaliplatin and tegafur in the treatment of advanced gastric cancer. *Pract Clin J Integ Tradit Chinese West Med*. 2008;8(4):3–4.
28. Mao WD. Clinical study of sodium cantharidinate and vitamin B6 combined with capecitabine in the treatment of advanced gastric cancer. *J Clin Med*. 2016;3(45):9034–9036.
29. Shi XY, Zhang J, Meng W, Ma F, Li ZL, Zhao JF. Clinical observation of sodium cantharidinate and vitamin B6 injection combined with XELOX protocol in treatment of patients with advanced colorectal carcinoma. *Med & Pharm J Chin PLA*. 2017;29(6):36–39.
30. Tian XL, Yang P. Clinical efficacy of treatment of liver cancer with combined sodium cantharidate vitamin B6 injection and TACE therapy. *Chin J Clin Oncol Rehabil*. 2006;13(4):351–353.
31. Wang JH, Li GP, Wu JH. Application of sodium cantharidinate and vitamin B6 injection in patients with advanced colorectal cancer. *J Mod Med Health*. 2010;26(6):844–845.
32. Wang YW, Yang RL. The effect observation of sodium cantharidate vitamin B6 combined with capecitabine in the treatment of advanced stomach and colorectal carcinoma. *Med Innov China*. 2017; 14(36):23–26.
33. Wei YF, Wu JS, Huo ZG, Jia ZY, Qi SS, Wang XX. Efficacy evaluation of sodium cantharidinate and vitamin B6 injection combined with TACE in the treatment of 48 patients with intermediate and advanced primary liver cancer. *China Pharm*. 2015;24(5):72–74.
34. Wu ZM, Liu Q, Shi SM, Yu SY. Clinical study of ai yishu combined with FOLFIRI regimen in the treatment of recurrent gastric cancer. *Med Inf*. 2013;26(6):256.
35. Xie ZX, Wang JS, Liu YB. Short-term efficacy and adverse effects of sodium cantharidinate and vitamin B6 injection combined with chemotherapy in the treatment of advanced gastric cancer. *Hebei Med J*. 2016;38(10):1554–1559.
36. You ZY, Liu Y, Shu XH, Liu BN, Liu XF. Malignant ascites of gastrointestinal tumor treated with sodium cantharidinate and vitamin B6 injection and hyperthermic intraperitoneal chemotherapy. *World J Integ Tradit West Med*. 2015;10(9):1249–1255.
37. Zeng L, Liu YL, Liu XT, Yang ZF. Clinical observation of sodium cantharidinate and vitamin B6 injection in the treatment of liver cancer. *J Hebei Med Univ*. 2009;30(6):596–597.
38. Zhang MJ, Zuo CF. Clinical efficacy of sodium cantharidate vitamin B6 injection combined with chemotherapy in treatment of liver cancer. *J Pract Oncol*. 2011;26(1):50–52.
39. Zhang W, Yang J, Zhong CS. Efficacy of sodium cantharidinate vitamin B6 combined with xeloda in the treatment of advanced gastric cancer. *J Huaihai Med*. 2012;30(5):446–447.
40. Zhang W, Lao WW. Efficacy of sodium cantharidinate vitamin B6 combined with CapeOX in the treatment of advanced gastric cancer. *Med Front*. 2015;5(16):108–110.
41. Zhu WQ. Sodium cantharidinate and vitamin B6 injection in combination with TACE for primary liver cancer. *Int Med Health Guid News*. 2014;20(14):2130–2132.
42. Zeng X, Zhang Y, Kwong JSW, et al. The methodological quality assessment tools for preclinical and clinical studies, systematic review and meta-analysis, and clinical practice guideline: a systematic review. *J Evid Based Med*. 2015;8(1):2–10.
43. Jackson D, White IR, Riley RD. Quantifying the impact of between-study heterogeneity in multivariate meta-analyses. *Stat Med*. 2012;31(29): 3805–3820.
44. Zhang L, Mu Y, Zhang A, et al. Cytokine-induced killer cells/dendritic cells-cytokine induced killer cells immunotherapy combined with chemotherapy for treatment of colorectal cancer in China: a meta-analysis of 29 trials involving 2,610 patients. *Oncotarget*. 2017;8(28):45164–45177.
45. Williamson PR, Smith CT, Hutton JL, Marson AG. Aggregate data meta-analysis with time-to-event outcomes. *Stat Med*. 2002;21(22): 3337–3351.
46. Zhou L, Wang XL, Deng QL, Du YQ, Zhao NQ. The efficacy and safety of immunotherapy in patients with advanced NSCLC: a systematic review and meta-analysis. *Sci Rep*. 2016;6(1):32020.
47. Tierney JF, Stewart LA, Ghersi D, Burdett S, Sydes MR. Practical methods for incorporating summary time-to-event data into meta-analysis. *Trials*. 2007;8(1):16.
48. Liang M, Chen Q, Zhang Y, et al. Impact of diabetes on the risk of bed sore in patients undergoing surgery: an updated quantitative analysis of cohort studies. *Oncotarget*. 2017;8(9):14516–14524.

49. Gu Y, Jiang L, Miao JH, Liang TS, Kan QC, Yang DK. Clinical effects of thermotherapy in combination with intracavitary infusion of traditional Chinese medicine in the treatment of malignant pleural effusion. *J Biol Regul Homeost Agents*. 2016;30(4):1023–1028.
50. Yan Z, Lai Z, Lin J. Anticancer properties of traditional Chinese medicine. *Comb Chem High Throughput Screen*. 2017;20(5):423–429.
51. Xu J, Song Z, Guo Q, Li J. Synergistic effect and molecular mechanisms of traditional Chinese medicine on regulating tumor microenvironment and cancer cells. *Biomed Res Int*. 2016;2016(5):1–14.
52. Liao YH, Li CI, Lin CC, Lin JG, Chiang JH, Li TC. Traditional Chinese medicine as adjunctive therapy improves the long-term survival of lung cancer patients. *J Cancer Res Clin Oncol*. 2017;143(12):2425–2435.

Supplementary materials




Figure S1 (Continued)




Figure S1 Forest plot of the comparison of complete response rates (A), partial response rates (B), stable disease rates (C), and progressive disease rates (D) between the experimental and control groups. Control group, CMT alone group; Experimental group, sodium cantharidinate and vitamin B6 injection (SC/B6) + CMT. The fixed-effects meta-analysis model (M-H method) was used.

Abbreviations: CMT, conventional medical treatment; M-H, Mantel-Haenszel.




Figure S2 (Continued)




Figure S2 (Continued)




Figure S2 (Continued)




Figure S2 Forest plot of the comparison of adverse effects including leukopenia (A), nausea and vomiting (B), gastrointestinal side effects (C), hepatotoxicity (D), nephrotoxicity (E), diarrhea (F), thrombocytopenia (G), transaminase disorder (H), myelosuppression (I), hand foot syndrome (J), oral mucositis (K), anorexia (L), and anemia (M) between the experimental and control groups. Control group, CMT-alone group; Experimental group, sodium cantharidinate and vitamin B6 injection (SC/B6) + CMT. **Abbreviation:** CMT, conventional medical treatment.




Figure S3 Funnel plot of percentage of complete response rates (A), partial response rates (B), stable disease rates (C), and progressive disease rates (D).

References

- Chen Y, Zhu WM, Zhou DX, Shan YF. Clinical study on disodium cantharidinate and vitamin B6 combined with raltitrexed and oxaliplatin in treatment of advanced colorectal cancer. *Drugs Clinic*. 2016; 31(11):1784–1787.
- Fan LJ. Efficacy of sodium cantharidinate and vitamin B6 combined with 5-fluorouracil and calcium folinate in treatment of advanced colorectal cancer. *Mod Med J China*. 2009;11(10):64–65.
- Fan QL, Wang Y, Zhao KX, Kao J. Clinical observation on effects of sodium cantharidinate vitamin B6 injection combined with S-1 treatment of advanced gastrointestinal tract tumor patients. *Guangming J Chin Med*. 2013;28(4):786–788.
- Fang XH, Xu J. Cantharidin sodium vitamin B6 injection as assistant therapy for treating liver cancer in 37 cases. *China Pharm*. 2016;25(1): 25–27.
- Guan LY, Ma J, Yuan SF, Qu J. Disodium cantharidinate and vitamin B6 combined with tegafur, gimeracil and oteracil potassium for treating stage IV pancreatic cancer in 27 cases. *Tradit Med*. 2015;24(20):124–125.
- Jia JM, Zh L. Efficacy of sodium cantharidinate and vitamin B6 combined with intravenous chemotherapy for advanced esophageal cancer. *Med Inf*. 2013;26(9):188–189.
- Liu GW, Ren WD. Cantharis acid sodium vitamin B6 injection combined with capecitabine synchronization for the treatment of advanced colorectal cancer. *World Latest Med Inf*. 2017;17(26):12–14.

8. Liu SH, Feng W, Zhu RX, Feng ZC. Comparative study of disodium cantharidinate and vitamin B6 injection combined with oxaliplatin and tegafur in the treatment of advanced gastric cancer. *Pract Clin J Integr Tradit Chinese West Med*. 2008;8(4):3–4.
9. Mao WD. Clinical study of sodium cantharidinate and vitamin B6 combined with capecitabine in the treatment of advanced gastric cancer. *J Clin Med*. 2016;3(45):9034–9036.
10. Shao H, Hong G, Luo X. Evaluation of sodium cantharidinate/vitamin B6 in the treatment of primary liver cancer. *J Cancer Res Ther*. 2014;10(Suppl 1):75–78.
11. Shi XY, Zhang J, Meng W, Ma F, Li ZL, Zhao JF. Clinical observation of sodium cantharidinate and vitamin B6 injection combined with XELOX protocol in treatment of patients with advanced colorectal carcinoma. *Med & Pharm J Chin PLA*. 2017;29(6):36–39.
12. Tian XL, Yang P. Clinical efficacy of treatment of liver cancer with combined sodium cantharidate vitamin B6 injection and TACE therapy. *Chin J Clin Oncol Rehabil*. 2006;13(4):351–353.
13. Wang JH, Li GP, Wu JH. Application of sodium cantharidinate and vitamin B6 injection in patients with advanced colorectal cancer. *J Mod Med Health*. 2010;26(6):844–845.
14. Wang YW, Yang RL. The effect observation of sodium cantharivitamin B6 combined with capecitabine in the treatment of advanced stomach and colorectal carcinoma. *Med Innov China*. 2017;14(36):23–26.
15. Wei YF, Wu JS, Huo ZG, Jia ZY, Qi SS, Wang XX. Efficacy evaluation of sodium cantharidinate and vitamin B6 injection combined with TACE in the treatment of 48 patients with intermediate and advanced primary liver cancer. *China Pharm*. 2015;24(5):72–74.
16. Wu ZM, Liu Q, Shi SM, Yu SY. Clinical study of ai yishu combined with FOLFIRI regimen in the treatment of recurrent gastric cancer. *Med Inf*. 2013;26(6):256.
17. Xie ZX, Wang JS, Liu YB. Short-term efficacy and adverse effects of sodium cantharidinate and vitamin B6 injection combined with chein the treatment of advanced gastric cancer. *Hebei Med J*. 2016;38(10):1554–1559.
18. You ZY, Liu Y, Shu XH, Liu BN, Liu XF. Malignant ascites of gas-tumor treated with sodium cantharidinate and vitamin B6 injection and hyperthermic intraperitoneal chemotherapy. *World J Integr Tradit West Med*. 2015;10(9):1249–1255.
19. Zeng L, Liu YL, Liu XT, Yang ZF. Clinical observation of sodium cantharidinate and vitamin B6 injection in the treatment of liver cancer. *J Hebei Med Univ*. 2009;30(6):596–597.
20. Zhang MJ, Zuo CF. Clinical efficacy of sodium cantharidate vitamin B6 injection combined with chemotherapy in treatment of liver cancer. *J Pract Oncol*. 2011;26(1):50–52.
21. Zhang W, Yang J, Zhong CS. Efficacy of sodium cantharidinate vitamin B6 combined with xeloda in the treatment of advanced gastric cancer. *J Huaihai Med*. 2012;30(5):446–447.
22. Zhang W, Lao WW. Efficacy of sodium cantharidinate vitamin B6 combined with CapeOX in the treatment of advanced gastric cancer. *Med Front*. 2015;5(16):108–110.
23. Zhu WQ. Sodium cantharidinate and vitamin B6 injection in combination with TACE for primary liver cancer. *Int Med Health Guid News*. 2014;20(14):2130–2132.
24. Li GP, Li XR, Wang JH. Clinical observation on sodium cantharidate vitamin B6 injection combined with postoperative chemotherapy in treatment of gastric cancer. *Pract Prev Med*. 2010;17(1):105–106.

Drug Design, Development and Therapy

Dovepress

Publish your work in this journal

Drug Design, Development and Therapy is an international, peer-reviewed open-access journal that spans the spectrum of drug design and development through to clinical applications. Clinical outcomes, patient safety, and programs for the development and effective, safe, and sustained use of medicines are the features of the journal, which

has also been accepted for indexing on PubMed Central. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <http://www.dovepress.com/drug-design-development-and-therapy-journal>