

# The Impact of COVID-19 on the Tuberculosis Features in a Romanian Pneumology Hospital

George-Cosmin Popovici<sup>1,2,\*</sup>, Costinela-Valerica Georgescu<sup>3,4,\*</sup>, Claudiu-Ionut Vasile<sup>5,6,\*</sup>, Oana-Mariana Mihailov<sup>2,5,\*</sup>, Mihaela-Camelia Vasile<sup>5,7,\*</sup>, Manuela Arbune<sup>5,8,\*</sup>

<sup>1</sup>School for Doctoral Studies in Biomedical Sciences, “Dunarea de Jos” University, Galati, Romania; <sup>2</sup>Pneumology Department II, Pneumophthisiology Hospital, Galati, Romania; <sup>3</sup>Pharmaceutical Sciences Department, “Dunarea de Jos” University, Galati, Romania; <sup>4</sup>Public Health & Management Department, Gynecology and Obstetrics Clinic Hospital “Buna Vestire”, Galati, Romania; <sup>5</sup>Medical Clinic Department, “Dunarea de Jos” University, Galati, Romania; <sup>6</sup>Psychiatry Clinic Department I, “Elisabeta Doamna” Psychiatric Hospital, Galati, Romania; <sup>7</sup>Infectious Diseases Clinic Department II, Clinic Hospital for Infectious Diseases, Galati, Romania; <sup>8</sup>Infectious Diseases Clinic Department I, Clinic Hospital for Infectious Diseases, Galati, Romania

\*These authors contributed equally to this work

Correspondence: Costinela-Valerica Georgescu; Claudiu-Ionut Vasile, Email [costinela.georgescu@ugal.ro](mailto:costinela.georgescu@ugal.ro); [ionut.vasile@ugal.ro](mailto:ionut.vasile@ugal.ro)

**Introduction:** The COVID-19 pandemic and tuberculosis have epidemiological similarities, being transmitted airborne, favored by direct contact, crowded environments, and vulnerable biological status.

**Methods:** We performed a retrospective study of 45 cases of pulmonary tuberculosis associated with COVID-19 (TB+COV+) compared to 45 cases with tuberculous monoinfection (TB+COV-), hospitalized during 2021–2022.

**Results:** The demographic characteristics were similar in the two groups, predominating men, a median age of 51 years, living in rural areas, medium level of education and smoking. Common symptoms of the two groups were cough, weight loss, profuse sweating, loss of appetite and hemoptysis, while fever, headache, myo-arthralgias, and digestive symptoms characterized the TB+COV+ forms. The scores of radiological lesions in the TB+COV+ compared to TB+COV- group were significantly higher and persistent, revealing more frequent bilateral extensive lung lesions. There were no significant differences in the biological parameters between the two groups. Mortality was 2.2%, regardless of the association of COVID-19. The frequency of infections with *Clostridioides difficile* was higher in TB+COV+ cases.

**Conclusion:** The co-infection of COVID-19 had a mild impact on the clinical and biological expression of tuberculosis diagnosed in a pandemic context.

**Keywords:** tuberculosis, COVID-19 pandemic, co-infection, Romania

## Introduction

Tuberculosis was one of the leading causes of death in human history and remains one of the leading infectious causes of death in the world.<sup>1</sup> Globally, the number of new cases of tuberculosis exceeds 10 million, although the incidence of tuberculosis has fallen slowly over the past decade, and mortality has fallen by almost a third.<sup>1</sup>

Although there has been a downward trend in the incidence of tuberculosis, the achievements are below the indicators established by the World Health Organization [WHO]. The aim of the “END TB” strategy, is to reduce the global TB incidence and mortality rates by 90% and 95%, respectively, in 2035 when compared to 2015. The COVID-19 pandemic has interfered with the diagnosis and care program for tuberculosis, negatively impacting the progress achieved in combating the disease by 2019. The reported number of people diagnosed with TB in 2020 and 2021 has decreased suggesting growing the number of undiagnosed and untreated people, with an estimated 10.6 million reported TB cases in 2021, compared to 10.1 million in 2020. Over more, the number of deaths increased to 1.6 million in 2021 compared to 1.5 million in 2020.<sup>1</sup>

COVID-19 has epidemiological similarities with tuberculosis, both infections being airborne, favored by direct contact, crowded environments, and vulnerable biological status. Both tuberculosis and the COVID-19 infection predominantly affect the lung, although extrapulmonary manifestations are also known. The incubation period differs,

with variations between 2 and 14 days for COVID-19 and 3–9 weeks for tuberculosis, although they may start with common manifestations, such as cough, chest pain, and asthenia.<sup>2</sup> Tuberculosis is an airborne bacterial infection caused by *Mycobacterium tuberculosis* that can cause extensive destruction of lung tissue, with the formation of cavities, while COVID-19 is a viral infection primarily transmitted person-to-person, caused by SARS-Cov-2 that alters lung cells, triggering inflammation and acute respiratory dysfunction.<sup>3,4</sup> The infection of COVID-19 can be diagnosed both before, during active tuberculosis, and post-tuberculosis, but the influence of COVID-19 in the reactivation or worsening of active tuberculosis is still unclear.<sup>5,6</sup> Morbidity and mortality risks for COVID-19 and tuberculosis can be influenced by smoking and co-morbidities, such as chronic lung disease, diabetes, or liver failure.<sup>7–11</sup> The convergence of these infections constitutes major challenges for public health globally.<sup>12</sup>

In 2020, of the 33,148 cases of tuberculosis reported by 29 of 30 EU/EEA countries, 23.2% came from Romania. Regarding the number of TB new cases, Romania is at the top ranking of EU countries, though decreasing trend was noted in the last years.<sup>13</sup>

There are some reports regarding the role of COVID-19 to increase the severity and mortality of tuberculosis.<sup>14–16</sup> The purpose of our study is to identify the particularities of the clinical, biological, radiological, and evolutive characteristics on hospitalized patients with tuberculosis–COVID-19 co-infection, from a special hospital of pneumology. To our knowledge, this is the first study on COVID-19 and tuberculosis co-infection from South-East of Romania.

## Materials and Methods

We have conducted a descriptive retrospective study on the cases of tuberculosis hospitalized during the COVID-19 pandemic period, from 1.07.2020 to 30.06.2022, from the Clinical Hospital of Pneumophthisiology Galati, located in the South-East of Romania. We specify that the hospital does not have an intensive care unit, and the complicated cases with organ dysfunctions are admitted/transferred to emergency hospitals.

## Selection of the Patients

The diagnosis of tuberculosis (TB) was considered according to the case definitions established by the national tuberculosis management guideline, classified into new cases and relapses, according to the TB history and treatment.<sup>13</sup> The etiological diagnosis was based on the classical methods of sputum analysis through culture on Loewenstein-Jensen medium and/or microscopic examination of Ziehl-Nielsen colored smears. The laboratory diagnosis of TB was completed by molecular tests by quantitative polymerization reaction in real-time DNA chain - RT-PCR (GeneXpert MTB-Rif-Cepheid). All hospitalized patients with TB were tested for COVID-19 upon admission and during hospitalization, due to clinical or epidemiological suspicion for this infection. The diagnosis of COVID-19 was confirmed by RT-PCR SARS-CoV-2 test. Considering that the average incubation interval of COVID-19 is 7 days, we have classified the cases as community or health care associated infection, depending on the diagnostic data, in the first 7 days of hospitalization or after 7 days, respective. The study was based on the analysis of data from the hospital's archive of the clinical records, that were selected based on the diagnostic codes ICD-10-CM version 37.1 R1, specified A15 (respiratory tuberculosis), associated or not with U07.1 (COVID-19), for each case of co-infection. The control group included cases of TB mono-infection (A15), selecting a pair with the most proximate hospitalization date of each co-infection case. Two independent researchers, not involved in the patient care or data collection, have selected the patients of the control group.

## Data Collection

Demographic, clinical, biological, radiological, and therapeutic data were collected, used according to current hospital procedures for the diagnosis and treatment of TB and COVID-19, according to the recommendations of national protocols.<sup>17–19</sup>

The radiological changes were evaluated by three independent radiologists and were validated by the pneumologist physician. The cavitary, ulcerated, caseous, nodular, infiltrative, fibrous, pleural/extrapulmonary forms of TB have been notified. Chest CT examination was available in isolated cases. The BRIXIA radiological scores for severity were evaluated on the date of COVID-19 diagnosis (baseline) and at discharge (end of study). Extent of lung parenchyma damage evaluation by BRIXIA score, consists of division into six sections of the lung radiologic image, in which each section achieves a damage score of 0, 1, 2 or 3, while total score could sum from 0 to 18.<sup>20–22</sup>

The inclusion criteria were age over 20 years, signing the informed consent for the use of personal data for the purpose of medical statistical studies, according to the standardized form from the clinical observation sheet.

They were excluded the patients vaccinated against COVID-19, patients with HIV infection, with neurocognitive dysfunctions who cannot express their consent, those re-hospitalized for the same episode of the disease of COVID-19 or tuberculosis, as well as cases with incomplete CXR examination of the lung (less than 3 radiological opinions).

## Statistical Analysis

The statistical analysis used the XLSTAT.2022.4.5 program. Continuous variables with normal distribution were compared based on t-tests, and categorical variables were compared with chi-squared tests or Fisher exact tests. We calculated the frequency of the variables and the mean  $\pm$ SD for the data with normal distribution or the median in the case of non-normally distributed data. We used the nonparametric Mann–Whitney *U*-test for data that did not correspond to the standard distribution.

This non-interventional study complies with the Declaration of Helsinki. The institutional approval of was endorsed by the Medical Council of the Pneumology Hospital from Galati no 1546/15.01.2024.

## Results

We analyzed 90 cases hospitalized with tuberculosis during the 2020–2022 pandemic period, of which 45 had the diagnosis of co-infection with COVID-19 and 45 had TB mono-infection.

## Demographic Data

The age of the patients varied between 20 and 86 years, with an average of  $52.1 \pm 14.69$  years, without significant statistical differences between the two groups (Mann Whitney test:  $p = 0.272$ ). The demographic characteristics indicated the predominance of male patients (78.8%), from rural areas (62.2%), with medium/higher education level (53.3%), professionally active (43.33%). The frequency of patients with a low level of education was significantly higher in the group with co-infection with COVID-19 compared to mono-infection with TB ( $p < 0.001$ ), without other demographic differences (Table 1).

**Table 1** Characteristics of Hospitalized Patients with TB Only Vs COVID-19 Co-Infection

		All	TB+COV+	TB+COV-	OR	CI 0.95	X <sup>2</sup>
Gender	Female	19	6	13	0.378	0.132;1.084	0.070
	Male	71	39	32			
Living area	Urban	34	15	19	0.684	0.290;1.609	0.384
	Rural	56	30	26			
Education*	Low	42	29	13	4.461	1.874;10.617	<0.001
	Medium/High	48	16	32			
Employment	Employed	39	22	17	3.364	1.032;10.968	0.044
	Non-employed	18	5	13			
	Retired	33	18	15			
Smoking	Yes	64	33	31	1.241	0.498;3.094	0.641
	No	26	12	14			
Alcohol use	Yes	67	33	30	2.312	0.875;6.108	0.090
	No	23	8	15			
Underweight (<20 kg/m <sup>2</sup> )	Yes	21	7	14	0.407	0.147;1.117	0.081
	No	69	38	31			
Overweight ( $\geq 26$ kg/m <sup>2</sup> )	Yes	12	9	3	3.5	0.935;13.098	0.0628
	No	78	36	42			

**Notes:** \*Low education: less than 8 years; Medium/High education: over 8 years.

Smoking was reported in 71.11% of the patients, with an average lifetime smoking exposure of  $29.48 \pm 11.74$  packs of cigarettes/year. The alcohol use was reported by 74.33% of patients, with average of  $4164 \pm 1.82$  IU/day. The health risk factors, such as smoking, alcohol consumption or altered nutritional status, did not influence the association with COVID-19.

### Associated Chronic Co-Morbidities

Chronic co-morbidities have been associated in 57.77% of patients (Figure 1), mainly chronic lung diseases (41.11%), liver chronic disease (18.88%), hypertension (16.66%), heart chronic diseases (11.11%) (Figure 1).

### Clinical Presentation and Diagnostic

The inventory of clinical symptoms identified manifestations exclusively associated with COVID-19, such as headache (44.4%), myalgias (35.5%), arthralgias (24.4%), as well as manifestations common to the two groups of patients, but with a higher frequency in the case of co-infection COVID-19/TB: fever (71.1% vs 22.2%), chest pain (82.2% vs 28.8%), shortness of breath (73.3% vs 44.4%), decreased appetite (77.7% vs 60%). Other manifestations had equal frequencies, for example, cough (100%), sweating (88.8%), weight loss (73–74%). Hemoptysis was reported in 22.2%, but the frequency was higher in patients with TB only (Figure 2).

The source of COVID-19 was community acquired in 62.2% of cases and healthcare-associated in 37.7%.

Recurrences constituted 30% of TB cases, more frequent in patients with COVID-19 co-infections compared to TB mono-infections (38% vs 22%). The microscopic examination of acid-alcohol-resistant bacilli (AARB) was positive in 91.1% of the cases.

The majority of cases (86.66%) had complex radiological images, highlighting 46.66% cavitary forms, 40% caseous, 26.66% ulcerated, 21.11% infiltrative, 36.66% fibrous, 43.33% nodular, and 7.7% pleurisy or associated extrapulmonary locations (Table 2).

The biological picture evidenced higher values of inflammation markers and of the Neutrophils/Lymphocytes ratio in patients with co-infection with COVID-19. Overmore, significantly higher differences of values in creatinine, ALT and Brixia scores, both at baseline and during the follow-up examination, were found in COVID-19 group (Table 3).

Non-TB bacterial examination of sputum was performed in 17.77% of patients, with 43.75% showing negative results. Isolates from sputum included Streptococcus spp (3), Moraxella catharalis (1), Klebsiella spp (3), Pseudomonas spp (1), and





Figure 1 The frequency of TB related chronic co-morbidities.

Abbreviations: COPD, chronic obstructive pulmonary disease; HTN, hypertension; HCDs, heart chronic diseases; NCDs, neurological chronic disease; CTD, connective tissue disease; CLD, chronic liver disease.



**Figure 2** Frequency of Symptoms on patients with TB+ COV- comparative with TB+COV+.

*E. coli* (1). A higher number of Gram-negative bacteria (BGN) isolates were observed in cases with COVID-19 (3:1), while TB mono-infection had a higher proportion of Gram-positive bacteria (CGP) (3:2).

### Drug Therapy and Outcomes

Treatment with the combination of Isoniazid, Rifampicin, Pyrazinamide, and Ethambutol, was administered to 96.6% of TB patients. Individualized antituberculous regimens were provided for 2 patients with TB-MDR and for one case of TB rifampicin resistance detected in the GeneXPERT test.

**Table 2** Classification Criteria of in TB Only in Comparison with TB-COVID-19 Co-Infection

		N-All	%	n1 TB+COV+	n2 TB+COV-	p	CI
Evolution	First presentation	63	70%	28	35	0.107	-0.344;0.033
	Recurrence	27	30%	17	10	0.107	-0.033;0.344
Microscopy	Positive AFB	82		40	42	0.458	-0.074;0.163
Radiologic features	Cavitary	43	47.77	21	22	0.996	-0.212;0.211
	Caseous	36	40	15	21	0.223	-0.336;0.079
	Ulcerate	24	26.66	12	12	0.947	-0.184;0.196
	Infiltrative	19	21.11	10	9	0.710	-0.207;0.141
	Fibrous	33		15	18	0.562	-0.146;0.266
	Nodular	39		20	19	0.685	-0.253;0.167
	Pleural/Extrapulmonary	7		6	1	0.049	0.021;0.992
Localization of the lesions	Bilateral	61	67.7	37	24	0.003	0.105;0.475
	Right	15		6	9	0.396	-0.222;0.088
	Left	12		1	11	0.001	-0.358;-0.086
	Miliaria	2		1	1	1	-0.061;0.617

**Abbreviations:** N-All, number of all patients; n1 TB+COV+, number of patients with co-infection COVID-19 and Tuberculosis; n2 TB+COV-, number of patients with tuberculosis only; CI, 95% confidence interval; AFB, acid fast bacteria.

**Table 3** Comparative Features of Biological Data and Radiologic Scores in TB Only and TB-COVID-19 Co-Infection


	NRV	Average ±SD		Median		2t-test	CI 0.95
		COV+	COV-	COV+	COV-		
<b>Leucocytes *10<sup>3</sup>/mm<sup>3</sup></b>	4.5–11	13,270±8947	12,300±922	11,400	11,400	0.504	–1.920;3.860
<b>NLR</b>	<3	7.648±8.346	5.468±4.502	4.764	4.229	0.127	–0.641;5.002
<b>CRP [mg/L]</b>	<3	14.448±21.208	8.611±5.63	7	6.7	0.080	–0.732;12.408
<b>ESR [mm/h]</b>	0–22	62.71±31.430	73.2±32.460	55	65	0.123	–23.877;2.898
<b>Creatinine [mg/dL]</b>	0.7–1.3	1.295±1.924	0.678±0.172	0.77	0.66	0.037	0.036;1.197
<b>ALT [U/L]</b>	7–55	72.71±86.69	33.77±27.93	38	22	0.005	11.700;66.166
<b>AST [U/L]</b>	8–48	66.711±73.621	47.667	43	30	0.126	–5.520;43.609
<b>BRIXIA Score-B</b>	0	9.288±3.307	7.688±2.556	9	8	0.012	0.360;2.839
<b>BRIXIA Score C</b>	0	7.266±3.326	3.911±2.172	7	3	<0.001	2.175;4.535

**Abbreviations:** NRV, Normal Reference Values; N/Ly, Neutrophils/Lymphocytes Ratio; CRP, C-Reactive Protein; ESR, Erythrocytes Sedimentation Rate; ALT, Alanine-amino transferase; AST, Aspartate amino transferase; BRIXIA Score -B, Severity Radiologic Score baseline; BRIXIA Score -C, Severity Radiologic Score Control (on hospital discharge).

Antibiotics for non-tuberculous infections were used in 33.3% of patients, including ceftriaxone (14.4%), amoxicillin (2.2%), tienamicin (3.3%), moxifloxacin (3.3%), levofloxacin (2.2%), linezolid (1.1%), gentamicin (1.1%), sulfamethoxazole (1.1%). Vancomycin (14.4%) and metronidazole (6.6%) were used for the treatment of patients with *Clostridioides difficile*-associated diarrhea.

Oxygen support via nasal cannula or face mask was provided to 31.1% of patients with COVID-19, compared to 6.6% in patients with TB monoinfection. Patients with COVID-19 received antivirals (77.7%) with favipiravir or remdesivir, dexamethasone (51.1%), and anticoagulants (62.2%).

Complications associated with the severe evolution of COVID-19 were observed in 8.8% of cases (coma, sepsis, and pleuropneumothorax), and medication toxicity was reported in 20% of patients (hyperuricemia, renal failure, toxic hepatitis, and anemia). *Clostridioides difficile* associated diarrhea occurred in 27.7% of cases, more frequently in the group with COVID-19 (16/9 vs.), but the difference was not statistically significant (Figure 3).



**Figure 3** Comparative Complications in Patients Hospitalized with Co-infection TB-COVID-19 and TB only. **Abbreviations:** PL-PN, pleuropneumonia; AKF, acute kidney failure; CDD, *Clostridioides difficile* diarrhea.

The outcomes were favorable in 92.2% of patients who were discharged with improved conditions. However, 5.5% of patients with COVID-19 deteriorated, requiring transfer to intensive care units in emergency hospitals, and 2.2% of patients died (1 case of TB-COVID co-infection and 1 case of mono-infection).

## Discussion

### Impact of COVID-19 on Tuberculosis Outcomes

*Mycobacterium tuberculosis*, a well-adapted pathogen to humans, has coexisted throughout human history. Approximately 2 billion people worldwide carry latent tuberculosis infections, with an estimated 10% risk of developing active disease during their lifetime, varying geographically and with comorbidities.<sup>23</sup>

COVID-19 induces pulmonary lesions but also affects other compartments of the immune system. In this context, tuberculosis reactivation can be attributed to various mechanisms, such as the immunosuppressive effects of corticosteroid therapy, cytokine storms, and T lymphocyte depletion.<sup>24</sup> An observational case-control study in China highlighted that both active and latent tuberculosis are risk factors for the progression of COVID-19.<sup>25</sup>

Conversely, studies from the early pandemic period provided evidence that COVID-19 increases the frequency and severity of tuberculosis.<sup>18,21,22</sup> The mortality rate in our study for patients with co-infection was 2.2%, comparable to the mono-infection TB group. This hospital mortality rate was lower than reported in some multicenter studies, where rates ranged from 9.2% to 14.2%.<sup>26–28</sup> It should be counted that 5.5% cases with severe evolution were transferred to intensive care units and probably died, increasing the mortality reported per case, and consequent decreasing discrepancy versus other studies. Most radiological lesions observed in our study were cavitory, evenly distributed between the co-infection and mono-infection TB groups, constituting 46.6%. Over 60% of these were new cases of tuberculosis, suggesting that the lesions had been present for at least one month before the onset of the COVID infection, whose incubation period is short.<sup>28</sup>

In view of these observations, our study results in line with the patterns observed in countries with a high prevalence of tuberculosis. In such settings, the diagnosis of COVID-19 might be more of a coincidental revelation of tuberculosis rather than a causative factor for reactivating latent tuberculosis.<sup>23,27,29,30</sup>

### Considerations of COVID-19 Impact on Tuberculosis Appearance in Romanian Patients

In comparison with a report from Serbia, patients with tuberculosis-COVID-19 co-infection in our study exhibited more intense symptoms, with a higher frequency of cough (100% vs 69.8%), fever (71.1% vs 30.2%), and shortness of breath (73.3% vs 30.2%). However, the frequency of cases diagnosed with COVID-19 following tuberculosis was lower in our study (37.7% vs 45%).<sup>31</sup> A noteworthy proportion of patients in both the mono-infection TB and co-infection COVID-19 groups received non-tuberculosis antibiotics. Despite the global alert on excessive antibiotic use for COVID-19 without clear evidence of bacterial infections, our study reported lower antibiotic utilization rates compared to other studies.<sup>32</sup>

Demographic characteristics and risk factors such as alcohol and smoking did not influence the occurrence of COVID-19 co-infection. Nevertheless, lower adherence to COVID-19 prevention rules could be related to lower education level that was found more prevalent in co-infected group. Chronic comorbidities, especially chronic obstructive pulmonary disease (COPD), hypertension, chronic liver diseases, or neurological chronic diseases (NCDs), were more frequent in patients with COVID-19 co-infection, consistent with previous findings.<sup>33,34</sup> Most statistics on tuberculosis has reported more frequent the males, according to our findings.<sup>35,36</sup> A slightly higher rate of male/ female was found in co-infection group, probably related to additional risks of comorbidities and smoking that are considered in COVID-19, but are as well in generally observed between males and females.<sup>37–39</sup>

Clinical symptoms highlighted an overlap between manifestations typical of both COVID-19 and tuberculosis, such as cough, sweating, and weight loss. Fever, chest pain, shortness of breath, and loss of appetite were more common in patients with COVID-19 co-infection, whereas hemoptysis was more frequent in TB alone. The pain syndrome was characterized by headache, myalgia, and arthralgia, and has been mentioned exclusively in patients with COVID-19 co-infection. However, the specificity of clinical manifestations for COVID-19 is limited, with none of the symptoms having

the accuracy to confirm or rule out this infection.<sup>40</sup> Moreover, the clinical spectrum of COVID-19 varies from asymptomatic to severe forms, correlated with epidemiological factors and host characteristics.<sup>41</sup> Hemoptysis is an uncommon manifestation of COVID-19, reported with frequencies between 0.9% and 3.3%.<sup>41,42</sup> Hemoptysis in tuberculosis can be caused by bronchiectasis, bronchial artery erosions, and, more rarely, pulmonary artery erosions.<sup>43</sup> The proportion of hemoptysis revealing the onset of tuberculosis and/or the unfavorable evolution of fibrocavitary secondary TB was 20.71%, in line with recent studies in Romania.<sup>44</sup>

Bilateral lung lesions were significantly more frequent in the COVID-19 group compared to mono-infection TB ( $p = 0.003$ ), consistent with a higher BRIXIA radiologic score. Although the BRIXIA score decreased in both groups upon discharge, the difference in values accentuated, likely due to the slow regression of COVID-19 pulmonary lesions. Prospective evaluation of these cases may be useful to clarify whether there is an additional risk for post-COVID fibrosis and respiratory dysfunction in the case of tuberculosis co-infection.

The average NLR was elevated in most patients in our study, but the differences between the COVID-19 co-infection and mono-infection TB groups were not significant, possibly because critical COVID-19 patterns were not evaluated in the study.<sup>45,46</sup>

*Clostridioides difficile* infection was reported in 27.7% of tuberculosis patients, possibly favored by rifampicin use, listed as one of the antibiotics that can induce diarrhea with this etiology, especially in elderly individuals with chronic comorbidities (underlying conditions).<sup>29</sup> The frequency of *Clostridioides difficile*-associated diarrhea was higher in the COVID-19 co-infection group (35.5% vs 20%), likely explained by the twice as frequent use of non-tuberculosis antibiotics and the occurrence of this type of diarrhea. It could be favored by rifampicin or broad-spectrum antibiotics use, as well as by changes in the intestinal microbiome during COVID-19 infection (fecal microbiota disruption by SARS-CoV-2 infection).<sup>47</sup>

## Limitations of the Study

The retrospective design of the study could have limited the accuracy of collecting the data. The number of cases is small, limiting the statistical significance of the results. There were only hospitalized TB cases, with symptomatic COVID-19 during the first 4 to 8 weeks from the admission. Symptomatic or asymptomatic COVID-19 cases preceding tuberculosis were not analyzed. The potential impact of prior asymptomatic COVID-19 infections on tuberculosis, with potential effects on the biological and radiological changes found in the mono-infection TB group, could not be assessed. Intensive care unit was not available and some patients with severe evolution were transferred to other hospitals, with unknown state, that could distort the rate of mortality per case.

## Conclusion

Symptomatic non-severe infection COVID-19 had a mild impact on the clinical and biological expression of tuberculosis diagnosed in the pandemic context. The hospital mortality of patients with tuberculosis was not influenced by COVID-19 co-infection in a country with a high tuberculosis prevalence. Persistent structural lung damage is suggested by radiological lesions according to the BRIXIA score and could be an indicator for COVID-19 additional risk on tuberculosis sequelae. Co-infection COVID-19 - tuberculosis should be cared by multidisciplinary teams, covering at least specialties as pneumology, infectious diseases, intensive care, microbiology, imaging, epidemiology. Tuberculosis control strategies should consider intensifying efforts to evaluate tuberculosis in COVID-19 patients, as common clinical manifestations and some pathogenic mechanisms continue to pose challenges for medical research and public health.

## Abbreviations

AFB, acid fast bacteria; ALT, Alanine-amino transferase; AKF, acute kidney failure; AST, Aspartate amino transferase; CI, confidence interval; CDD; *Clostridioides difficile* diarrhea; COPD, chronic obstructive pulmonary disease; HCDs, heart chronic diseases; CLD, chronic liver disease; CRP, C-Reactive Protein; CTD, connective tissue disease; ESR, Erythrocytes Sedimentation Rate; GNB, Gram-negative bacilli; GPC, Gram-positive cocci; HTN – hypertension; NCDs, neurological chronic disease; NRV, Normal Reference Values; N/Ly, Neutrophils/Lymphocytes Ratio; OR, Odds Ratio; PL-PN, pleuropneumonia; TB-tuberculosis; WHO, World Health Organization.



## Acknowledgments

We kindly acknowledge the board of “Dunarea de Jos” University from Galati for the financial support for publication. This paper has been uploaded to <https://www.preprints.org> as a preprint: <https://doi.org/10.20944/preprints202401.1145.v1>

## Disclosure

The authors report no conflicts of interest in this work.

## References

1. World Health Organization. Global tuberculosis report 2022; 2022. Available from: <https://iris.who.int/bitstream/handle/10665/363752/9789240061729-eng.pdf?sequence=1>. Accessed 11, December 2023.
2. Cioboata R, Biciusca V, Olteanu M, Vasile CM. COVID-19 and Tuberculosis: unveiling the Dual Threat and Shared Solutions Perspective. *J Clin Med*. 2023;12:4784. doi:10.3390/jcm12144784
3. Gibson PG, Qin L, Pua SH. COVID-19 acute respiratory distress syndrome (ARDS): clinical features and differences from typical pre-COVID-19 ARDS. *Med J Aust*. 2020;213(2):54–56.e51. doi:10.5694/mja2.50674
4. Mocanu AI, Mocanu H, Moldovan C, et al. Some Manifestations of Tuberculosis in Otorhinolaryngology - Case Series and a Short Review of Related Data from South-Eastern Europe. *Infect Drug Resist*. 2022;15:2753–2762. doi:10.2147/IDR.S367885
5. Brosnahan SB, Jonkman AH, Kugler MC, Munger JS, Kaufman DA. COVID-19 and Respiratory System Disorders: current Knowledge, Future Clinical and Translational Research Questions. *Arterioscler Thromb Vasc Biol*. 2020;40(11):2586–2597. doi:10.1161/ATVBAHA.120.314515
6. Visca D, Ong CWM, Tiberi S, et al. Tuberculosis and COVID-19 interaction: a review of biological, clinical and public health effects. *Pulmonology*. 2021;27(2):151–165. doi:10.1016/j.pulmoe.2020.12.012
7. Al Maqbal M, Al Badi K, Al Sinani M, Madkhali N, Dickens GL. Clinical Features of COVID-19 Patients in the First Year of Pandemic: a Systematic Review and Meta-Analysis. *Biol Res Nurs*. 2022;24(2):172–185. doi:10.1177/10998004211055866
8. Biswas SS, Awal SS, Awal SK. COVID-19 and pulmonary tuberculosis - A diagnostic dilemma. *Radiol Case Rep*. 2021;16(11):3255–3259. doi:10.1016/j.radcr.2021.07.079
9. Vasile MC, Arbune AA, Lupasteanu G, Vlase CM, Popovici GC, Arbune M. Epidemiologic and Clinic Characteristics of the First Wave of the COVID-19 Pandemic in Hospitalized Patients from Galați County. *J Clin Med*. 2021;10:18. doi:10.3390/jcm10184210
10. Kumar R, Rai AK, Phukan MM, et al. Accumulating Impact of Smoking and Co-morbidities on Severity and Mortality of COVID-19 Infection: a Systematic Review and Meta-analysis. *Curr Genomics*. 2021;22(5):339–352. doi:10.2174/1389202922666210921101728
11. Vidyasagan AL, Readshaw A, Boeckmann M, et al. Is Tobacco Use Associated With Risk of Recurrence and Mortality Among People With TB?: a Systematic Review and Meta-Analysis. *Chest*. 2024;165(1):22–47. doi:10.1016/j.chest.2023.08.021
12. Herrera MT, Guzmán-Beltrán S, Bobadilla K, et al. Human Pulmonary Tuberculosis: understanding the Immune Response in the Bronchoalveolar System. *Biomolecules*. 2022;12(8). doi:10.3390/biom12081148
13. European Centre for Disease Prevention and Control, WHO Regional Office for Europe. *Tuberculosis Surveillance and Monitoring in Europe 2022 – 2020 Data*. Copenhagen: WHO Regional Office for Europe and Stockholm: European Centre for Disease Prevention and Control; 2022.
14. Patra K, Batabyal S, Mandal K, Ghose D, Sarkar J. Tuberculosis and COVID-19: a combined global threat to human civilization. *Clin Epidemiol Glob Health*. 2022;15:101031. doi:10.1016/j.cegh.2022.101031
15. Gong W, Mao Y, Li Y, Qi Y. BCG Vaccination: a potential tool against COVID-19 and COVID-19-like Black Swan incidents. *Int Immunopharmacol*. 2022;108:108870. doi:10.1016/j.intimp.2022.108870
16. Harries AD, Kumar AMV, Satyanarayana S. TB and COVID-19: measuring key risk factors that affect treatment outcomes. *Int J Tubercul Lung Dis*. 2021;25(4):329–331. doi:10.5588/IJTL.21.0061
17. MINISTERUL SĂNĂTĂȚII ORDIN nr. 956 din 31 martie 2022 privind modificarea Ordinului ministrului sănătății nr. 1.171/2015 pentru aprobarea unor reglementări privind controlul tuberculozei: GHID METODOLOGIC DE IMPLEMENTARE A PROGRAMULUI NAȚIONAL DE PREVENIRE, SUPRAVEGHERE ȘI CONTROL AL TUBERCULOZEI ȘI ALTOR MICOBACTERIOZE. MONITORUL OFICIAL AL ROMÂNIEI, PARTEA I, Nr. 326 bis 4.IV.2022. Available from: <https://legislatie.just.ro/Public/DetaliiDocument/253597?msclcid=09174c29c24711ec95524134f8dfd6f2>. Accessed October 21, 2023.
18. Ministerul Sănătății. ORDIN nr. 487 din 23 martie 2020 pentru aprobarea protocolului de tratament al infecției cu virusul SARS-Cov-2. MONITORUL OFICIAL nr. 242 din 24 martie 2020. Available from: <https://legislatie.just.ro/Public/DetaliiDocument/224341>. Accessed October 21, 2023.
19. Ministerul Sănătății. ORDIN nr. 533 din 22 aprilie 2021 privind modificarea anexei la Ordinul ministrului sănătății nr. 487/2020 pentru aprobarea protocolului de tratament al infecției cu virusul SARS-CoV-2. MONITORUL OFICIAL nr. 434 din 23 aprilie 2021. Available from: <https://legislatie.just.ro/Public/DetaliiDocumentAfis/241318>. Accessed October 21, 2023.
20. Nukovic JA, Opancina V, Zdravkovic N, et al. Brixia Chest X-ray Score, Laboratory Parameters and Vaccination Status for Prediction of Mortality in COVID-19 Hospitalized Patients. *Diagnostics (Basel)*. 2023;13:12. doi:10.3390/diagnostics13122122
21. Au-Yong I, Higashi Y, Giannotti E, et al. Chest Radiograph Scoring Alone or Combined with Other Risk Scores for Predicting Outcomes in COVID-19. *Radiology*. 2022;302(2):460–469. doi:10.1148/radiol.2021210986
22. Little BP. Disease Severity Scoring for COVID-19: a Welcome Semiquantitative Role for Chest Radiography. *Radiology*. 2022;302(2):470–472. doi:10.1148/radiol.2021212212
23. Mousquer GT, Peres A, Fiegenbaum M. Pathology of TB/COVID-19 Co-Infection: the phantom menace. *Tuberculosis (Edinb)*. 2021;126:102020. doi:10.1016/j.tube.2020.102020
24. Shariq M, Sheikh JA, Quadir N, Sharma N, Hasnain SE, Ehtesham NZ. COVID-19 and tuberculosis: the double whammy of respiratory pathogens. *Eur Respir Rev*. 2022;31:164. doi:10.1183/16000617.0264-2021
25. Feldman C, Anderson C. The role of co-infections and secondary infections in patients with COVID-19. *Pneumonia (Nathan)*. 2021;13(1):5. doi:10.1186/s41479-021-00083-w

26. Tadolini M, Codecasa LR, García-García JM, et al. Active tuberculosis, sequelae and COVID-19 co-infection: first cohort of 49 cases. *Eur Respir J.* 2020;56:1. doi:10.1183/13993003.01398-2020
27. Motta I, Centis R, D'Ambrosio L, et al. Tuberculosis, COVID-19 and migrants: preliminary analysis of deaths occurring in 69 patients from two cohorts. *Pulmonology.* 2020;26(4):233–240. doi:10.1016/j.pulmoe.2020.05.002
28. Group TC-GS. Tuberculosis and COVID-19 co-infection: description of the global cohort. *Eur Respir J.* 2022;59:3. doi:10.1183/13993003.02538-2021
29. Khurana AK, Aggarwal D. The (in)significance of TB and COVID-19 co-infection. *Eur Respir J.* 2020;56(2). doi:10.1183/13993003.02105-2020
30. Stochino C, Villa S, Zucchi P, Parravicini P, Gori A, Raviglione MC. Clinical characteristics of COVID-19 and active tuberculosis co-infection in an Italian reference hospital. *Eur Respir J.* 2020;56:1. doi:10.1183/13993003.01708-2020
31. Adzic-Vukicevic T, Stosic M, Antonijevic G, Jevtic M, Radovanovic-Spurnic A, Velickovic J. Tuberculosis and COVID-19 co-infection in Serbia: pandemic challenge in a low-burden country. *Front Med Lausanne.* 2022;9:971008. doi:10.3389/fmed.2022.971008
32. Vaughn VM, Gandhi TN, Petty LA, et al. Empiric Antibacterial Therapy and Community-onset Bacterial Coinfection in Patients Hospitalized With Coronavirus Disease 2019 (COVID-19): a Multi-hospital Cohort Study. *Clin Infect Dis.* 2021;72(10):e533–e541. doi:10.1093/cid/ciaa1239
33. Parolina L, Pshenichnaya N, Vasilyeva I, et al. Clinical characteristics of COVID-19 in patients with tuberculosis and factors associated with the disease severity. *Int J Infect Dis.* 2022;124(Suppl 1):S82–S89. doi:10.1016/j.ijid.2022.04.041
34. Song B, Sheng X, Justice JL, et al. Intercellular communication within the virus microenvironment affects the susceptibility of cells to secondary viral infections. *Sci Adv.* 2023;9(19):eadg3433. doi:10.1126/sciadv.adg3433
35. European Centre for Disease Prevention and Control, WHO Regional Office for Europe. *Tuberculosis Surveillance and Monitoring in Europe 2021 – 2019 Data.* Copenhagen: WHO Regional Office for Europe; 2021.
36. Shaweno D, Horton KC, Hayes RJ, Dodd PJ. Assortative social mixing and sex disparities in tuberculosis burden. *Sci Rep.* 2021;11(1):7530. doi:10.1038/s41598-021-86869-w
37. Bwire GM. Coronavirus: why Men are More Vulnerable to Covid-19 Than Women? *SN Compr Clin. Med.* 2020;2(7):874–876. doi:10.1007/s42399-020-00341-w
38. Kharroubi SA, Diab-El-Harake M. Sex-differences in COVID-19 diagnosis, risk factors and disease comorbidities: a large US-based cohort study. *Front Public Health.* 2022;10:1029190. doi:10.3389/fpubh.2022.1029190
39. Peer V, Schwartz N, Green MS. Gender differences in tuberculosis incidence rates-A pooled analysis of data from seven high-income countries by age group and time period. *Front Public Health.* 2022;10:997025. doi:10.3389/fpubh.2022.997025
40. Martínez Orozco JA, Sánchez Tinajero Á, Becerril Vargas E, et al. COVID-19 and Tuberculosis Coinfection in a 51-Year-Old Taxi Driver in Mexico City. *Am J Case Rep.* 2020;21:e927628. doi:10.12659/AJCR.927628
41. Baj J, Karakuła-Juchnowicz H, Teresiński G, et al. COVID-19: specific and Non-Specific Clinical Manifestations and Symptoms: the Current State of Knowledge. *J Clin Med.* 2020;9:6. doi:10.3390/jcm9061753
42. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med.* 2020;382(18):1708–1720. doi:10.1056/NEJMoa2002032
43. Quigley N, Gagnon S, Fortin M. Aetiology, diagnosis and treatment of moderate-to-severe haemoptysis in a North American academic centre. *ERJ Open Res.* 2020;6(4). doi:10.1183/23120541.00204-2020
44. Chirana C, Vlase I. Hemoptysis in Patients with Lung Tuberculosis. *ARS Medica Tomitana.* 2021;27(3). doi:10.2478/arsm-2021-0025
45. Kim MA, Park YE, Chong YP, Shim TS, Jo KW. Neutrophil-Lymphocyte Ratio and Monocyte-Lymphocyte Ratio According to the Radiologic Severity of. *J Korean Med Sci.* 2022;37(40):e292. doi:10.3346/jkms.2022.37.e292
46. Sarkar S, Khanna P, Singh AK. The Impact of Neutrophil-Lymphocyte Count Ratio in COVID-19: a Systematic Review and Meta-Analysis. *J Intensive Care Med.* 2022;37(7):857–869. doi:10.1177/08850666211045626
47. Lewandowski K, Rosołowski M, Kaniewska M, et al. Clostridioides difficile infection in coronavirus disease 2019 (COVID-19): an underestimated problem? *Pol Arch Intern Med.* 2021;131(2):121–127. doi:10.20452/pamw.15715

Journal of Multidisciplinary Healthcare

Dovepress

## Publish your work in this journal

The Journal of Multidisciplinary Healthcare is an international, peer-reviewed open-access journal that aims to represent and publish research in healthcare areas delivered by practitioners of different disciplines. This includes studies and reviews conducted by multidisciplinary teams as well as research which evaluates the results or conduct of such teams or healthcare processes in general. The journal covers a very wide range of areas and welcomes submissions from practitioners at all levels, from all over the world. The manuscript management system is completely online and includes a very quick and fair peer-review system. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-multidisciplinary-healthcare-journal>