

Poly[[bis(μ_3 -2-[(3,5-dimethyl-1*H*-pyrazol-1-yl)(phenyl)methyl]propanedioato)tetrasodium(I)] 7.5-hydrate]

Ihssan Meskini,^a Maria Daoudi,^{a*} Jean-Claude Daran,^b Taibi Ben Hadda^c and Hafid Zouhri^d

^aLaboratoire de Chimie Organique, Faculté des Sciences Dhar el Mahraz, Université Sidi Mohammed Ben Abdellah, Fès, Morocco, ^bLaboratoire de Chimie de Coordination, 205 Route de Narbonne, 31077 Toulouse Cedex, France, ^cLaboratoire de Chimie des Matériaux, Université Mohammed Ier, Oujda, Morocco, and ^dLaboratoires de Diffraction des Rayons X, Division UATRS, Centre National pour la Recherche Scientifique et Technique, Rabat, Morocco
Correspondence e-mail: daoudimaria@yahoo.fr


Received 6 July 2010; accepted 16 July 2010

Key indicators: single-crystal X-ray study; $T = 293$ K, $P = 0.0$ kPa; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.036; wR factor = 0.130; data-to-parameter ratio = 18.5.

The asymmetric unit of the title polymer, $\{[Na_4(C_{15}H_{14}N_2O_4)_2] \cdot 7.5H_2O\}_n$, contains two 2-[(3,5-dimethyl-1*H*-pyrazol-1-yl)(phenyl)methyl]propanedioate (ppmp) anions, eight water molecules (one located on a twofold rotation axis) and five sodium cations (one located on an inversion center and the other one located on a twofold rotation axis). The carboxylate groups of the ppmp anions and the water molecules bridge the Na cations, forming a two-dimensional polymeric structure. In the structure there are two types of coordination environment around the metal cations: one Na cation is coordinated by five O atoms in a distorted square-pyramidal geometry while the other four Na cations are coordinated by six O atoms in a distorted octahedral geometry. Extensive O—H...O and O—H...N hydrogen bonding is present in the crystal structure. The H atoms of one methyl group of the ppmp anion are disordered equally over two positions.

Related literature

For related compounds displaying biological activity, see: Dayam *et al.* (2007); Patil *et al.* (2007); Ramkumar *et al.* (2008); Sechi *et al.* (2009); Zeng *et al.* (2008). For the synthetic procedure, see: Pommier & Neamati (2006).



Experimental

Crystal data

$[Na_4(C_{15}H_{14}N_2O_4)_2] \cdot 7.5H_2O$
 $M_r = 799.64$
Monoclinic, $C2/c$
 $a = 31.8211$ (11) Å
 $b = 14.4951$ (4) Å
 $c = 16.1113$ (5) Å
 $\beta = 102.139$ (3)°

$V = 7265.2$ (4) Å³
 $Z = 8$
Mo $K\alpha$ radiation
 $\mu = 0.16$ mm⁻¹
 $T = 293$ K
 $0.45 \times 0.38 \times 0.19$ mm

Data collection

Bruker X8 APEXII CCD area-detector diffractometer
76380 measured reflections

9034 independent reflections
6490 reflections with $I > 2\sigma(I)$
 $R_{int} = 0.049$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.036$
 $wR(F^2) = 0.130$
 $S = 1.02$
9034 reflections

488 parameters
H-atom parameters constrained
 $\Delta\rho_{max} = 0.53$ e Å⁻³
 $\Delta\rho_{min} = -0.44$ e Å⁻³

Table 1

Selected bond lengths (Å).

Na1—O3	2.3113 (11)	Na4—O6	2.6565 (14)
Na1—O11	2.4177 (11)	Na4—O11 ^{iv}	2.3956 (14)
Na1—O12	2.4028 (13)	Na4—O12 ⁱⁱⁱ	2.4434 (15)
Na2—O5 ⁱ	2.3507 (13)	Na4—O41	2.4634 (11)
Na2—O5 ⁱⁱ	2.3417 (13)	Na4—O42	2.3423 (15)
Na2—O8	2.2818 (13)	Na5—O4	2.3952 (13)
Na2—O21	2.4546 (13)	Na5—O5	2.5592 (13)
Na2—O22	2.3253 (13)	Na5—O21 ^v	2.3176 (13)
Na3—O2	2.4259 (14)	Na5—O22 ⁱ	2.4275 (14)
Na3—O31	2.4430 (14)	Na5—O31	2.5011 (14)
Na3—O4	2.5862 (13)	Na5—O51	2.4478 (14)
Na4—O3 ⁱⁱⁱ	2.6812 (15)		

Symmetry codes: (i) $-x, y, -z + \frac{1}{2}$; (ii) $x, -y, z + \frac{1}{2}$; (iii) $x, y - 1, z$; (iv) $-x, -y, -z$; (v) $x, -y, z - \frac{1}{2}$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O11—H111 ⁱ ...O2 ⁱ	0.86	1.97	2.7404 (16)	149
O11—H11B...O4	0.85	1.88	2.7067 (16)	165
O12—H121...O1 ⁱ	0.84	1.90	2.7391 (17)	174
O21—H211...O6 ⁱ	0.85	1.92	2.7566 (17)	167
O21—H212...O7	0.85	1.97	2.8007 (16)	165
O22—H221...O31	0.85	2.10	2.8381 (17)	145
O22—H222...O11 ⁱ	0.85	2.31	2.9707 (16)	135
O22—H222...O6 ⁱⁱ	0.85	2.45	3.0532 (18)	128
O31—H31A...O8	0.85	1.87	2.7128 (16)	169
O31—H31B...O51	0.84	2.09	2.8085 (17)	142
O41—H411...O7	0.84	2.27	3.1061 (16)	172

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O42–H421 \cdots O1 ⁱⁱⁱ	0.85	1.92	2.7582 (17)	175
O42–H422 \cdots O7	0.85	2.03	2.8639 (17)	168
O51–H511 \cdots N4	0.86	1.99	2.8424 (18)	171
O51–H512 \cdots N2	0.85	1.94	2.7799 (18)	169

Symmetry codes: (i) $-x, y, -z + \frac{1}{2}$; (ii) $x, -y, z + \frac{1}{2}$; (iii) $x, y - 1, z$.

Data collection: *APEX2* (Bruker, 2005); cell refinement: *SAINTE* (Bruker, 2005); data reduction: *SAINTE*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *publCIF* (Westrip, 2010).

This work was supported by grants from Project PGR-Ump-BH-2005, the Centre National de Recherche Scientifique, CNRS (France), the Centre National pour la Recherche Scientifique et Technique, CNRST (Morocco) and the CURI (Morocco).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2795).

References

- Bruker (2005). *APEX2* and *SAINTE*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dayam, R., Al-Mawsawi, L. O. & Neamati, N. (2007). *Bioorg. Med. Chem. Lett.* **17**, 6155–6159.
- Patil, S., Kamath, S., Sanchez, T., Neamati, N., Schinazi, R. F. & Buolamwini, J. K. (2007). *Bioorg. Med. Chem.* **15**, 1212–1228.
- Pommier, Y. & Neamati, N. (2006). *Bioorg. Med. Chem.* **14**, 3785–3792.
- Ramkumar, K., Tambov, K. V., Gundla, R., Manaev, A. V., Yarovenko, V., Traven, V. F. & Neamati, N. (2008). *Bioorg. Med. Chem.* **16**, 8988–8998.
- Sechi, M., Carta, F., Sannia, L., Dallochio, R., Dessi, A., Al-Safi, R. I. & Neamati, N. (2009). *Antivir. Res.* **81**, 267–276.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Zeng, L.-F., Zhang, H.-S., Wang, Y.-H., Sanchez, T., Zheng, Y.-T., Neamati, N. & Long, Y.-Q. (2008). *Bioorg. Med. Chem. Lett.* **18**, 4521–4524.

supplementary materials

Acta Cryst. (2010). E66, m1009-m1010 [doi:10.1107/S1600536810028515]

Poly[[bis{ μ_3 -2-[(3,5-dimethyl-1*H*-pyrazol-1-yl)(phenyl)methyl]propanedioato}tetrasodium(I)] 7.5-hydrate]

I. Meskini, M. Daoudi, J.-C. Daran, T. Ben Hadda and H. Zouihri

Comment

The rational design of new HIV-1 Integrase (H-I) inhibitors, one validated target for chemotherapeutic intervention (Dayam *et al.*, 2007), is fundamentally based on intermolecular coordination between H—I / chemical inhibitor / metals (Mg^{+2} and Mn^{+2} , co-factors of the enzyme), leading in the formation of bimetallic complexes (Zeng *et al.*, 2008; Sechi *et al.*, 2009). Thereby, several bimetallic metal complexes, in many cases exploring the known-well polydentate ligands, appear in this scenario as the most promising concept to employ in either enzyme / drug interaction or electron transfer process, in the last case involving the biological oxygen transfer (Sechi *et al.*, 2009; Ramkumar *et al.*, 2008). Another exciting example of application for such polydentate ligand involves the synergic water activation, that occurs *via* the so-called -remote metallic atoms-. Such organometallic compounds are structurally deemed to promote or block the H—I activity (Zeng *et al.*, 2008).

The asymmetric unit of the title polymer contains two (3,5-dimethyl-1*H*-pyrazol-1-yl)(phenyl)methylpropanedioate (ppmp) anions, eight water molecules (O41 atom located at a twofold rotation axis) and five sodium cations (Na1 located on an inversion center and Na3 located at a twofold rotation axis). The carboxyl groups of ppmp anions and water molecules bridge the Na cations to form the two-dimensional polymeric structure. In the structure there are two types of coordination environment around the metal cations. The Na2 cation is coordinated by five oxygen atoms with a distorted square-pyramidal geometry; the other four Na cations are coordinated by six oxygen atoms with the distorted octahedral geometry. The Na—O bond distances are ranged from 2.3113 (11) to 2.6812 (15) Å (Table 1). The extensive O—H \cdots O and O—H \cdots N hydrogen bonding is present in the crystal structure (Table 2).

Experimental

A mixture of the sodium salt of 2-[(phenyl)-3,5-dimethyl-pyrazol-1-yl]-malonic acid (Pommier & Neamati, 2006) (0.2 g, 0.61 mmol) and (0.13, 1.22 mmol) of sodium dicarbonate in water (5 ml) was stirred at room temperature, then (0.047 g, 0.305 mmol) of (VO_2SO_4) was added. The mixture was allowed to stand to ambient temperature. Single crystals suitable for X-ray diffraction were obtained a few days later. Yield: 37%.

Refinement

All H atoms attached to C were fixed geometrically and treated as riding with C—H = 0.96 Å (methyl), 0.98 Å (methine) or 0.93 Å (aromatic) with $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ or $1.5U_{\text{eq}}(\text{methyl})$. H atoms of water molecule were located in difference Fourier maps and included in the subsequent refinement using restraints (O—H = 0.85 (1) Å and H \cdots H = 1.39 (2) Å) with $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$. In the last stage of refinement, they were treated as riding on their parent O atoms. The H atoms of the one methyl group in the ligand are disordered equally over two positions.

Figures




Fig. 1. Molecular structure of the title compound with the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are represented as small spheres of arbitrary radii.




Fig. 2. Partial packing view showing the chain generated by C—H...O hydrogen bonds shown as dashed lines.

Poly[[bis{ μ_3 -2-[(3,5-dimethyl-1*H*-pyrazol-1-yl)(phenyl)methyl]propanedioato}tetrasodium(I)] 7.5-hydrate]

Crystal data

[Na₄(C₁₅H₁₄N₂O₄)₂]·7.5H₂O

$M_r = 799.64$

Monoclinic, *C2/c*

Hall symbol: -C 2yc

$a = 31.8211$ (11) Å

$b = 14.4951$ (4) Å

$c = 16.1113$ (5) Å

$\beta = 102.139$ (3)°

$V = 7265.2$ (4) Å³

$Z = 8$

$F(000) = 3352$

$D_x = 1.462$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2648 reflections

$\theta = 1.5$ – 26.3 °

$\mu = 0.16$ mm⁻¹

$T = 293$ K

Block, colourless

$0.45 \times 0.38 \times 0.19$ mm

Data collection

Bruker X8 APEXII CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

φ and ω scans

76380 measured reflections

9034 independent reflections

6490 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.049$

$\theta_{\text{max}} = 28.3$ °, $\theta_{\text{min}} = 1.3$ °

$h = -42$ → 42

$k = -19$ → 19

$l = -21$ → 21

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.036$

$wR(F^2) = 0.130$

$S = 1.02$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0783P)^2 + 2.736P]$

9034 reflections
488 parameters
0 restraints

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.006$$

$$\Delta\rho_{\max} = 0.53 \text{ e } \text{Å}^{-3}$$

$$\Delta\rho_{\min} = -0.44 \text{ e } \text{Å}^{-3}$$

Special details

Experimental. IR (KBr, [UTF-8]^{1/2} cm⁻¹): 1592.46 (C=O), 3248 (OH).

The data collection nominally covered a sphere of reciprocal space, by a combination of seven sets of exposures; each set had a different ϕ angle for the crystal and each exposure covered 0.5° in ω and 20 s in time. The crystal-to-detector distance was 37.5 mm.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
C1	0.13673 (5)	0.43818 (11)	0.21255 (10)	0.0157 (3)	
H1	0.1288	0.5033	0.2036	0.019*	
C2	0.09753 (5)	0.38687 (10)	0.23022 (10)	0.0146 (3)	
H2	0.1049	0.3215	0.2390	0.018*	
N1	0.14699 (4)	0.40316 (9)	0.13308 (8)	0.0171 (3)	
N2	0.15635 (5)	0.31192 (9)	0.12560 (9)	0.0217 (3)	
C11	0.14801 (5)	0.45027 (12)	0.06112 (11)	0.0211 (3)	
C111	0.17568 (5)	0.43350 (12)	0.28481 (10)	0.0189 (3)	
C112	0.18973 (7)	0.35192 (14)	0.32677 (13)	0.0351 (5)	
H112	0.1747	0.2975	0.3109	0.042*	
C113	0.22592 (7)	0.35053 (16)	0.39193 (15)	0.0451 (6)	
H113	0.2350	0.2953	0.4193	0.054*	
C114	0.24851 (6)	0.43032 (17)	0.41628 (14)	0.0420 (5)	
H114	0.2728	0.4292	0.4599	0.050*	
C115	0.23495 (7)	0.51153 (16)	0.37580 (16)	0.0450 (6)	
H115	0.2501	0.5657	0.3920	0.054*	
C116	0.19862 (6)	0.51313 (14)	0.31076 (14)	0.0328 (4)	
H116	0.1895	0.5687	0.2842	0.039*	
C12	0.15814 (6)	0.38751 (13)	0.00471 (11)	0.0268 (4)	
H12	0.1610	0.3990	-0.0506	0.032*	
C13	0.16321 (6)	0.30294 (12)	0.04696 (12)	0.0249 (4)	
C14	0.17451 (8)	0.21067 (14)	0.01556 (14)	0.0417 (5)	
H14A	0.1753	0.1653	0.0593	0.063*	
H14B	0.2022	0.2141	0.0009	0.063*	

supplementary materials

H14C	0.1533	0.1935	-0.0337	0.063*	
C15	0.13940 (7)	0.55134 (13)	0.05108 (13)	0.0308 (4)	
H15A	0.1111	0.5643	0.0598	0.046*	
H15B	0.1414	0.5701	-0.0051	0.046*	
H15C	0.1602	0.5846	0.0920	0.046*	
C21	0.08148 (5)	0.42241 (11)	0.30808 (10)	0.0177 (3)	
C22	0.05932 (5)	0.39513 (11)	0.15432 (10)	0.0163 (3)	
O1	0.09050 (4)	0.50303 (8)	0.33362 (8)	0.0245 (3)	
O2	0.05884 (4)	0.36630 (9)	0.33884 (8)	0.0292 (3)	
O3	0.04974 (4)	0.47344 (8)	0.12450 (8)	0.0246 (3)	
O4	0.03947 (4)	0.32106 (8)	0.12934 (8)	0.0227 (3)	
C3	0.12948 (5)	-0.12031 (11)	0.19213 (10)	0.0152 (3)	
H3	0.1257	-0.1848	0.2073	0.018*	
C4	0.08996 (5)	-0.06637 (11)	0.20531 (10)	0.0141 (3)	
H4	0.0968	-0.0005	0.2054	0.017*	
N3	0.16800 (4)	-0.08512 (9)	0.25020 (9)	0.0177 (3)	
N4	0.17591 (4)	0.00742 (9)	0.25847 (9)	0.0203 (3)	
C31	0.20142 (6)	-0.13530 (13)	0.29413 (12)	0.0249 (4)	
C311	0.13456 (5)	-0.11851 (11)	0.10044 (10)	0.0175 (3)	
C312	0.13502 (6)	-0.20186 (12)	0.05790 (12)	0.0274 (4)	
H312	0.1330	-0.2570	0.0863	0.033*	
C313	0.13850 (7)	-0.20368 (14)	-0.02656 (13)	0.0345 (5)	
H313	0.1389	-0.2598	-0.0543	0.041*	
C314	0.14133 (6)	-0.12223 (14)	-0.06929 (12)	0.0296 (4)	
H314	0.1435	-0.1234	-0.1260	0.035*	
C315	0.14097 (6)	-0.03861 (13)	-0.02781 (11)	0.0255 (4)	
H315	0.1431	0.0163	-0.0565	0.031*	
C316	0.13747 (5)	-0.03674 (12)	0.05677 (11)	0.0222 (3)	
H316	0.1371	0.0195	0.0843	0.027*	
C32	0.23146 (6)	-0.07307 (13)	0.33379 (13)	0.0299 (4)	
H32	0.2578	-0.0862	0.3695	0.036*	
C33	0.21456 (5)	0.01393 (12)	0.30977 (12)	0.0243 (4)	
C34	0.23424 (6)	0.10641 (14)	0.33425 (14)	0.0369 (5)	
H34A	0.2150	0.1539	0.3078	0.055*	0.50
H34B	0.2610	0.1113	0.3159	0.055*	0.50
H34C	0.2394	0.1133	0.3948	0.055*	0.50
H34D	0.2619	0.0984	0.3712	0.055*	0.50
H34E	0.2159	0.1411	0.3631	0.055*	0.50
H34F	0.2375	0.1391	0.2842	0.055*	0.50
C35	0.20217 (7)	-0.23784 (14)	0.29361 (15)	0.0399 (5)	
H35A	0.1782	-0.2610	0.3149	0.060*	
H35B	0.2285	-0.2593	0.3290	0.060*	
H35C	0.2003	-0.2594	0.2366	0.060*	
C41	0.05116 (5)	-0.08420 (11)	0.13273 (10)	0.0154 (3)	
C42	0.07637 (5)	-0.08894 (11)	0.28921 (10)	0.0156 (3)	
O5	0.03524 (4)	-0.01421 (8)	0.09023 (7)	0.0191 (2)	
O6	0.03787 (4)	-0.16533 (8)	0.11998 (8)	0.0230 (3)	
O7	0.08345 (4)	-0.16799 (8)	0.32102 (8)	0.0239 (3)	
O8	0.05735 (4)	-0.02532 (8)	0.31923 (7)	0.0229 (3)	

Na1	0.0000	0.5000	0.0000	0.0202 (2)
Na2	0.03995 (2)	0.00005 (4)	0.44768 (4)	0.01928 (15)
Na3	0.0000	0.28300 (6)	0.2500	0.0230 (2)
Na4	0.03267 (3)	-0.34679 (6)	0.13857 (5)	0.0347 (2)
Na5	0.04893 (2)	0.15926 (4)	0.10917 (4)	0.02081 (15)
O11	-0.03287 (4)	0.35116 (8)	0.00998 (7)	0.0201 (2)
H111	-0.0480	0.3692	0.0450	0.030*
H11B	-0.0086	0.3365	0.0400	0.030*
O12	-0.03957 (4)	0.59680 (9)	0.07746 (8)	0.0276 (3)
H121	-0.0542	0.5642	0.1043	0.041*
H122	-0.0554	0.6369	0.0489	0.041*
O21	0.04694 (4)	-0.16789 (8)	0.46475 (7)	0.0231 (3)
H211	0.0208	-0.1762	0.4404	0.035*
H212	0.0611	-0.1749	0.4263	0.035*
O22	0.02907 (4)	0.15783 (8)	0.42770 (8)	0.0231 (3)
H221	0.0435	0.1722	0.3908	0.035*
H222	0.0368	0.1942	0.4697	0.035*
O31	0.04933 (4)	0.15204 (8)	0.26444 (8)	0.0225 (3)
H31A	0.0500	0.0942	0.2749	0.034*
H31B	0.0750	0.1673	0.2648	0.034*
O41	0.0000	-0.27849 (13)	0.2500	0.0307 (4)
H411	0.0213	-0.2470	0.2736	0.046*
O42	0.09668 (4)	-0.33909 (8)	0.24137 (8)	0.0279 (3)
H421	0.0932	-0.3870	0.2691	0.042*
H422	0.0940	-0.2925	0.2718	0.042*
O51	0.12258 (4)	0.15927 (8)	0.19318 (8)	0.0242 (3)
H511	0.1410	0.1169	0.2115	0.036*
H512	0.1361	0.2027	0.1744	0.036*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0160 (7)	0.0154 (7)	0.0155 (8)	-0.0014 (6)	0.0030 (6)	-0.0002 (6)
C2	0.0146 (7)	0.0150 (7)	0.0137 (7)	-0.0007 (5)	0.0018 (6)	0.0003 (6)
N1	0.0197 (7)	0.0161 (6)	0.0162 (7)	-0.0001 (5)	0.0056 (5)	0.0010 (5)
N2	0.0266 (7)	0.0171 (7)	0.0229 (8)	0.0015 (6)	0.0086 (6)	-0.0003 (6)
C11	0.0212 (8)	0.0232 (9)	0.0191 (8)	0.0007 (6)	0.0051 (7)	0.0050 (7)
C111	0.0150 (7)	0.0245 (8)	0.0174 (8)	-0.0013 (6)	0.0039 (6)	-0.0019 (6)
C112	0.0344 (11)	0.0277 (10)	0.0349 (11)	-0.0031 (8)	-0.0113 (9)	0.0033 (8)
C113	0.0409 (12)	0.0434 (13)	0.0409 (13)	0.0032 (10)	-0.0145 (10)	0.0094 (10)
C114	0.0238 (10)	0.0651 (15)	0.0314 (12)	-0.0069 (10)	-0.0074 (8)	-0.0027 (10)
C115	0.0347 (11)	0.0490 (14)	0.0448 (14)	-0.0203 (10)	-0.0066 (10)	-0.0067 (11)
C116	0.0308 (10)	0.0290 (10)	0.0352 (11)	-0.0083 (8)	-0.0006 (8)	0.0001 (8)
C12	0.0344 (10)	0.0299 (10)	0.0178 (9)	0.0019 (7)	0.0090 (7)	0.0022 (7)
C13	0.0286 (9)	0.0248 (9)	0.0234 (9)	-0.0002 (7)	0.0099 (7)	-0.0022 (7)
C14	0.0624 (15)	0.0312 (11)	0.0370 (12)	0.0035 (10)	0.0229 (11)	-0.0055 (9)
C15	0.0402 (11)	0.0237 (9)	0.0294 (10)	0.0042 (8)	0.0093 (8)	0.0096 (8)
C21	0.0153 (7)	0.0232 (8)	0.0133 (8)	-0.0004 (6)	0.0004 (6)	-0.0002 (6)

supplementary materials

C22	0.0152 (7)	0.0207 (8)	0.0136 (7)	-0.0004 (6)	0.0044 (6)	-0.0012 (6)
O1	0.0314 (7)	0.0219 (6)	0.0219 (6)	-0.0028 (5)	0.0097 (5)	-0.0056 (5)
O2	0.0354 (7)	0.0332 (7)	0.0227 (7)	-0.0136 (6)	0.0143 (6)	-0.0036 (5)
O3	0.0253 (6)	0.0210 (6)	0.0235 (7)	-0.0020 (5)	-0.0040 (5)	0.0064 (5)
O4	0.0211 (6)	0.0196 (6)	0.0237 (6)	-0.0021 (5)	-0.0035 (5)	-0.0039 (5)
C3	0.0161 (7)	0.0152 (7)	0.0144 (7)	0.0005 (6)	0.0034 (6)	-0.0009 (6)
C4	0.0147 (7)	0.0145 (7)	0.0132 (7)	0.0006 (5)	0.0031 (6)	0.0006 (6)
N3	0.0169 (6)	0.0181 (7)	0.0174 (7)	0.0021 (5)	0.0020 (5)	0.0003 (5)
N4	0.0195 (7)	0.0176 (7)	0.0232 (8)	-0.0007 (5)	0.0031 (6)	-0.0005 (6)
C31	0.0201 (8)	0.0272 (9)	0.0264 (9)	0.0069 (7)	0.0024 (7)	0.0052 (7)
C311	0.0146 (7)	0.0219 (8)	0.0167 (8)	0.0012 (6)	0.0053 (6)	-0.0013 (6)
C312	0.0383 (10)	0.0216 (9)	0.0245 (9)	0.0015 (7)	0.0113 (8)	-0.0018 (7)
C313	0.0460 (12)	0.0328 (11)	0.0262 (10)	0.0016 (9)	0.0110 (9)	-0.0115 (8)
C314	0.0245 (9)	0.0476 (12)	0.0175 (9)	-0.0009 (8)	0.0066 (7)	-0.0037 (8)
C315	0.0236 (9)	0.0345 (10)	0.0184 (9)	-0.0029 (7)	0.0044 (7)	0.0041 (7)
C316	0.0243 (8)	0.0232 (9)	0.0198 (9)	0.0001 (7)	0.0059 (7)	-0.0007 (7)
C32	0.0182 (8)	0.0349 (11)	0.0327 (11)	0.0033 (7)	-0.0033 (7)	0.0026 (8)
C33	0.0180 (8)	0.0295 (10)	0.0251 (9)	-0.0021 (7)	0.0038 (7)	-0.0013 (7)
C34	0.0271 (10)	0.0353 (11)	0.0457 (13)	-0.0096 (8)	0.0022 (9)	-0.0067 (9)
C35	0.0343 (11)	0.0265 (10)	0.0536 (14)	0.0121 (8)	-0.0028 (10)	0.0041 (9)
C41	0.0142 (7)	0.0218 (8)	0.0114 (7)	0.0017 (6)	0.0056 (6)	-0.0003 (6)
C42	0.0168 (7)	0.0176 (7)	0.0118 (7)	-0.0008 (6)	0.0016 (6)	-0.0007 (6)
O5	0.0207 (6)	0.0224 (6)	0.0138 (6)	0.0041 (4)	0.0025 (4)	0.0036 (4)
O6	0.0229 (6)	0.0218 (6)	0.0219 (6)	-0.0030 (5)	-0.0007 (5)	-0.0010 (5)
O7	0.0366 (7)	0.0202 (6)	0.0177 (6)	0.0055 (5)	0.0118 (5)	0.0047 (5)
O8	0.0326 (7)	0.0204 (6)	0.0185 (6)	0.0062 (5)	0.0119 (5)	0.0012 (5)
Na1	0.0243 (5)	0.0168 (4)	0.0175 (5)	-0.0013 (3)	-0.0003 (4)	0.0014 (4)
Na2	0.0231 (3)	0.0201 (3)	0.0154 (3)	0.0022 (2)	0.0059 (3)	-0.0001 (2)
Na3	0.0219 (5)	0.0212 (5)	0.0263 (5)	0.000	0.0060 (4)	0.000
Na4	0.0301 (4)	0.0529 (5)	0.0206 (4)	-0.0048 (3)	0.0045 (3)	-0.0084 (3)
Na5	0.0230 (3)	0.0220 (3)	0.0168 (3)	0.0017 (3)	0.0027 (3)	-0.0014 (3)
O11	0.0199 (6)	0.0241 (6)	0.0160 (6)	0.0008 (4)	0.0033 (5)	0.0002 (5)
O12	0.0253 (6)	0.0326 (7)	0.0259 (7)	0.0014 (5)	0.0080 (5)	0.0062 (5)
O21	0.0230 (6)	0.0302 (7)	0.0165 (6)	0.0021 (5)	0.0049 (5)	0.0016 (5)
O22	0.0264 (6)	0.0201 (6)	0.0236 (6)	0.0001 (5)	0.0073 (5)	-0.0035 (5)
O31	0.0241 (6)	0.0185 (6)	0.0260 (7)	0.0012 (5)	0.0078 (5)	0.0035 (5)
O41	0.0288 (10)	0.0320 (10)	0.0328 (11)	0.000	0.0099 (8)	0.000
O42	0.0353 (7)	0.0197 (6)	0.0292 (7)	-0.0003 (5)	0.0080 (6)	0.0001 (5)
O51	0.0214 (6)	0.0181 (6)	0.0337 (7)	0.0005 (5)	0.0073 (5)	0.0057 (5)

Geometric parameters (Å, °)

Na1—O3	2.3113 (11)	C13—C14	1.500 (3)
Na1—O3 ⁱ	2.3113 (11)	C14—H14A	0.9600
Na1—O11	2.4177 (11)	C14—H14B	0.9600
Na1—O11 ⁱ	2.4177 (11)	C14—H14C	0.9600
Na1—O12	2.4028 (13)	C15—H15A	0.9600
Na1—O12 ⁱ	2.4028 (13)	C15—H15B	0.9600

Na2—O5 ⁱⁱ	2.3507 (13)	C15—H15C	0.9600
Na2—O5 ⁱⁱⁱ	2.3417 (13)	C21—O1	1.252 (2)
Na2—O8	2.2818 (13)	C21—O2	1.256 (2)
Na2—O21	2.4546 (13)	C22—O3	1.245 (2)
Na2—O22	2.3253 (13)	C22—O4	1.2676 (19)
Na3—O2	2.4259 (14)	C3—N3	1.468 (2)
Na3—O2 ⁱⁱ	2.4259 (14)	C3—C311	1.520 (2)
Na3—O31	2.4430 (14)	C3—C4	1.533 (2)
Na3—O31 ⁱⁱ	2.4430 (14)	C3—H3	0.9800
Na3—O4	2.5862 (13)	C4—C41	1.533 (2)
Na3—O4 ⁱⁱ	2.5862 (13)	C4—C42	1.538 (2)
Na4—O3 ^{iv}	2.6812 (15)	C4—H4	0.9800
Na4—O6	2.6565 (14)	N3—C31	1.358 (2)
Na4—O11 ^v	2.3956 (14)	N3—N4	1.3661 (19)
Na4—O12 ^{iv}	2.4434 (15)	N4—C33	1.333 (2)
Na4—O41	2.4634 (11)	C31—C32	1.371 (3)
Na4—O42	2.3423 (15)	C31—C35	1.487 (3)
Na5—O4	2.3952 (13)	C311—C312	1.391 (2)
Na5—O5	2.5592 (13)	C311—C316	1.391 (2)
Na5—O21 ^{vi}	2.3176 (13)	C312—C313	1.389 (3)
Na5—O22 ⁱⁱ	2.4275 (14)	C312—H312	0.9300
Na5—O31	2.5011 (14)	C313—C314	1.379 (3)
Na5—O51	2.4478 (14)	C313—H313	0.9300
Na1—Na4 ^{vii}	3.1655 (8)	C314—C315	1.385 (3)
Na1—Na4 ^v	3.1655 (8)	C314—H314	0.9300
Na1—H111	2.6295	C315—C316	1.391 (2)
Na1—H11B	2.4858	C315—H315	0.9300
Na2—Na2 ^{viii}	3.3331 (13)	C316—H316	0.9300
Na2—Na5 ⁱⁱⁱ	3.4453 (9)	C32—C33	1.393 (3)
Na2—Na5 ⁱⁱ	3.6158 (9)	C32—H32	0.9300
Na2—H211	2.6231	C33—C34	1.497 (3)
Na2—H212	2.6649	C34—H34A	0.9600
Na2—H221	2.6683	C34—H34B	0.9600
Na3—Na5	3.5036 (8)	C34—H34C	0.9600
Na3—Na5 ⁱⁱ	3.5036 (8)	C34—H34D	0.9600
Na4—Na1 ^{iv}	3.1655 (8)	C34—H34E	0.9600
Na4—H421	2.6015	C34—H34F	0.9600
Na5—Na2 ^{vi}	3.4453 (9)	C35—H35A	0.9600
Na5—Na2 ⁱⁱ	3.6158 (9)	C35—H35B	0.9600
Na5—H31B	2.4723	C35—H35C	0.9600
C1—N1	1.476 (2)	C41—O6	1.252 (2)
C1—C111	1.513 (2)	C41—O5	1.2684 (19)
C1—C2	1.530 (2)	C42—O8	1.2544 (19)
C1—H1	0.9800	C42—O7	1.2562 (19)
C2—C22	1.537 (2)	O11—Na4 ^v	2.3956 (14)

supplementary materials

C2—C21	1.539 (2)	O11—H111	0.8553
C2—H2	0.9800	O11—H11B	0.8487
N1—C11	1.352 (2)	O12—Na4 ^{vii}	2.4434 (15)
N1—N2	1.3665 (19)	O12—H121	0.8438
N2—C13	1.337 (2)	O12—H122	0.8399
C11—C12	1.371 (2)	O21—Na5 ⁱⁱⁱ	2.3176 (13)
C11—C15	1.493 (2)	O21—H211	0.8507
C111—C116	1.383 (2)	O21—H212	0.8458
C111—C112	1.389 (3)	O22—Na5 ⁱⁱ	2.4275 (14)
C112—C113	1.386 (3)	O22—H221	0.8489
C112—H112	0.9300	O22—H222	0.8522
C113—C114	1.375 (3)	O31—H31A	0.8540
C113—H113	0.9300	O31—H31B	0.8440
C114—C115	1.371 (3)	O41—Na4 ⁱⁱ	2.4634 (11)
C114—H114	0.9300	O41—H411	0.8398
C115—C116	1.388 (3)	O42—H421	0.8453
C115—H115	0.9300	O42—H422	0.8494
C116—H116	0.9300	O51—H511	0.8572
C12—C13	1.395 (3)	O51—H512	0.8533
C12—H12	0.9300		
N1—C1—C111	110.93 (13)	H111—Na1—H11B	30.6
N1—C1—C2	109.38 (12)	O8—Na2—O22	95.19 (5)
C111—C1—C2	113.78 (13)	O8—Na2—O5 ⁱⁱⁱ	168.93 (5)
N1—C1—H1	107.5	O22—Na2—O5 ⁱⁱⁱ	90.64 (5)
C111—C1—H1	107.5	O8—Na2—O5 ⁱⁱ	100.26 (5)
C2—C1—H1	107.5	O22—Na2—O5 ⁱⁱ	86.37 (5)
C1—C2—C22	110.71 (12)	O5 ⁱⁱⁱ —Na2—O5 ⁱⁱ	89.48 (5)
C1—C2—C21	114.35 (13)	O8—Na2—O21	84.74 (4)
C22—C2—C21	106.25 (12)	O22—Na2—O21	176.60 (5)
C1—C2—H2	108.5	O5 ⁱⁱⁱ —Na2—O21	90.00 (4)
C22—C2—H2	108.5	O5 ⁱⁱ —Na2—O21	90.30 (4)
C21—C2—H2	108.5	O8—Na2—Na2 ^{viii}	144.59 (5)
C11—N1—N2	111.61 (13)	O22—Na2—Na2 ^{viii}	87.89 (4)
C11—N1—C1	128.29 (14)	O5 ⁱⁱⁱ —Na2—Na2 ^{viii}	44.85 (3)
N2—N1—C1	120.09 (12)	O5 ⁱⁱ —Na2—Na2 ^{viii}	44.63 (3)
C13—N2—N1	105.07 (13)	O21—Na2—Na2 ^{viii}	90.21 (4)
N1—C11—C12	106.54 (15)	O8—Na2—Na5 ⁱⁱⁱ	125.41 (4)
N1—C11—C15	123.48 (15)	O22—Na2—Na5 ⁱⁱⁱ	138.59 (4)
C12—C11—C15	129.98 (16)	O5 ⁱⁱⁱ —Na2—Na5 ⁱⁱⁱ	47.97 (3)
C116—C111—C112	117.92 (17)	O5 ⁱⁱ —Na2—Na5 ⁱⁱⁱ	93.18 (4)
C116—C111—C1	119.36 (16)	O21—Na2—Na5 ⁱⁱⁱ	42.23 (3)
C112—C111—C1	122.72 (15)	Na2 ^{viii} —Na2—Na5 ⁱⁱⁱ	64.45 (2)
C113—C112—C111	120.82 (18)	O8—Na2—Na5 ⁱⁱ	101.49 (4)
C113—C112—H112	119.6	O22—Na2—Na5 ⁱⁱ	41.53 (3)

C111—C112—H112	119.6	O5 ⁱⁱⁱ —Na2—Na5 ⁱⁱ	89.11 (4)
C114—C113—C112	120.4 (2)	O5 ⁱⁱ —Na2—Na5 ⁱⁱ	44.86 (3)
C114—C113—H113	119.8	O21—Na2—Na5 ⁱⁱ	135.15 (4)
C112—C113—H113	119.8	Na2 ^{viii} —Na2—Na5 ⁱⁱ	59.28 (2)
C115—C114—C113	119.57 (19)	Na5 ⁱⁱⁱ —Na2—Na5 ⁱⁱ	123.73 (2)
C115—C114—H114	120.2	O8—Na2—H211	84.3
C113—C114—H114	120.2	O22—Na2—H211	157.7
C114—C115—C116	120.17 (19)	O5 ⁱⁱⁱ —Na2—H211	93.8
C114—C115—H115	119.9	O5 ⁱⁱ —Na2—H211	71.8
C116—C115—H115	119.9	O21—Na2—H211	18.9
C111—C116—C115	121.16 (19)	Na2 ^{viii} —Na2—H211	80.0
C111—C116—H116	119.4	Na5 ⁱⁱⁱ —Na2—H211	50.4
C115—C116—H116	119.4	Na5 ⁱⁱ —Na2—H211	116.6
C11—C12—C13	106.19 (16)	O8—Na2—H212	67.3
C11—C12—H12	126.9	O22—Na2—H212	162.3
C13—C12—H12	126.9	O5 ⁱⁱⁱ —Na2—H212	106.3
N2—C13—C12	110.59 (15)	O5 ⁱⁱ —Na2—H212	98.9
N2—C13—C14	120.33 (16)	O21—Na2—H212	18.4
C12—C13—C14	129.07 (17)	Na2 ^{viii} —Na2—H212	107.8
C13—C14—H14A	109.5	Na5 ⁱⁱⁱ —Na2—H212	58.4
C13—C14—H14B	109.5	Na5 ⁱⁱ —Na2—H212	141.3
H14A—C14—H14B	109.5	H211—Na2—H212	29.6
C13—C14—H14C	109.5	O8—Na2—H221	78.7
H14A—C14—H14C	109.5	O22—Na2—H221	17.9
H14B—C14—H14C	109.5	O5 ⁱⁱⁱ —Na2—H221	105.7
C11—C15—H15A	109.5	O5 ⁱⁱ —Na2—H221	96.1
C11—C15—H15B	109.5	O21—Na2—H221	163.0
H15A—C15—H15B	109.5	Na2 ^{viii} —Na2—H221	105.4
C11—C15—H15C	109.5	Na5 ⁱⁱⁱ —Na2—H221	152.0
H15A—C15—H15C	109.5	Na5 ⁱⁱ —Na2—H221	53.4
H15B—C15—H15C	109.5	H211—Na2—H221	157.1
O1—C21—O2	125.86 (15)	H212—Na2—H221	144.6
O1—C21—C2	119.19 (14)	O2—Na3—O2 ⁱⁱ	120.30 (8)
O2—C21—C2	114.91 (14)	O2—Na3—O31 ⁱⁱ	147.53 (5)
O3—C22—O4	125.76 (15)	O2 ⁱⁱ —Na3—O31 ⁱⁱ	85.97 (4)
O3—C22—C2	117.81 (14)	O2—Na3—O31	85.97 (4)
O4—C22—C2	116.40 (14)	O2 ⁱⁱ —Na3—O31	147.53 (5)
C21—O2—Na3	121.68 (11)	O31 ⁱⁱ —Na3—O31	78.02 (6)
C22—O3—Na1	123.65 (10)	O2—Na3—O4	84.31 (4)
C22—O3—Na4 ^{vii}	152.98 (11)	O2 ⁱⁱ —Na3—O4	83.51 (4)
Na1—O3—Na4 ^{vii}	78.32 (4)	O31 ⁱⁱ —Na3—O4	119.87 (4)
C22—O4—Na5	143.51 (11)	O31—Na3—O4	80.41 (4)
C22—O4—Na3	103.32 (10)	O2—Na3—O4 ⁱⁱ	83.51 (4)

supplementary materials

Na5—O4—Na3	89.30 (5)	O2 ⁱⁱ —Na3—O4 ⁱⁱ	84.31 (4)
N3—C3—C311	111.60 (12)	O31 ⁱⁱ —Na3—O4 ⁱⁱ	80.41 (4)
N3—C3—C4	109.50 (12)	O31—Na3—O4 ⁱⁱ	119.87 (4)
C311—C3—C4	112.58 (13)	O4—Na3—O4 ⁱⁱ	155.37 (7)
N3—C3—H3	107.6	O2—Na3—Na5	104.12 (3)
C311—C3—H3	107.6	O2 ⁱⁱ —Na3—Na5	105.40 (3)
C4—C3—H3	107.6	O31 ⁱⁱ —Na3—Na5	84.53 (4)
C3—C4—C41	111.15 (12)	O31—Na3—Na5	45.55 (3)
C3—C4—C42	113.80 (12)	O4—Na3—Na5	43.13 (3)
C41—C4—C42	107.72 (12)	O4 ⁱⁱ —Na3—Na5	161.48 (4)
C3—C4—H4	108.0	O2—Na3—Na5 ⁱⁱ	105.40 (3)
C41—C4—H4	108.0	O2 ⁱⁱ —Na3—Na5 ⁱⁱ	104.12 (3)
C42—C4—H4	108.0	O31 ⁱⁱ —Na3—Na5 ⁱⁱ	45.55 (3)
C31—N3—N4	111.53 (14)	O31—Na3—Na5 ⁱⁱ	84.53 (4)
C31—N3—C3	127.11 (14)	O4—Na3—Na5 ⁱⁱ	161.48 (4)
N4—N3—C3	121.02 (13)	O4 ⁱⁱ —Na3—Na5 ⁱⁱ	43.13 (3)
C33—N4—N3	104.89 (13)	Na5—Na3—Na5 ⁱⁱ	118.41 (3)
N3—C31—C32	106.47 (16)	O42—Na4—O11 ^v	121.53 (5)
N3—C31—C35	122.97 (17)	O42—Na4—O12 ^{iv}	154.89 (6)
C32—C31—C35	130.56 (17)	O11 ^v —Na4—O12 ^{iv}	77.79 (5)
C312—C311—C316	118.83 (16)	O42—Na4—O41	84.44 (4)
C312—C311—C3	118.58 (15)	O11 ^v —Na4—O41	145.99 (5)
C316—C311—C3	122.56 (14)	O12 ^{iv} —Na4—O41	85.57 (4)
C313—C312—C311	120.71 (17)	O42—Na4—O6	88.10 (5)
C313—C312—H312	119.6	O11 ^v —Na4—O6	84.27 (5)
C311—C312—H312	119.6	O12 ^{iv} —Na4—O6	111.20 (5)
C314—C313—C312	120.00 (18)	O41—Na4—O6	74.34 (5)
C314—C313—H313	120.0	O42—Na4—O3 ^{iv}	86.82 (5)
C312—C313—H313	120.0	O11 ^v —Na4—O3 ^{iv}	81.18 (4)
C313—C314—C315	120.01 (17)	O12 ^{iv} —Na4—O3 ^{iv}	80.27 (5)
C313—C314—H314	120.0	O41—Na4—O3 ^{iv}	125.19 (6)
C315—C314—H314	120.0	O6—Na4—O3 ^{iv}	159.10 (5)
C314—C315—C316	120.01 (17)	O42—Na4—Na1 ^{iv}	129.79 (4)
C314—C315—H315	120.0	O11 ^v —Na4—Na1 ^{iv}	49.18 (3)
C316—C315—H315	120.0	O12 ^{iv} —Na4—Na1 ^{iv}	48.66 (3)
C315—C316—C311	120.43 (16)	O41—Na4—Na1 ^{iv}	132.04 (4)
C315—C316—H316	119.8	O6—Na4—Na1 ^{iv}	129.35 (4)
C311—C316—H316	119.8	O3 ^{iv} —Na4—Na1 ^{iv}	45.64 (3)
C31—C32—C33	105.99 (16)	O42—Na4—H421	18.8
C31—C32—H32	127.0	O11 ^v —Na4—H421	130.1
C33—C32—H32	127.0	O12 ^{iv} —Na4—H421	136.4
N4—C33—C32	111.11 (15)	O41—Na4—H421	81.7

N4—C33—C34	120.47 (16)	O6—Na4—H421	105.1
C32—C33—C34	128.42 (17)	O3 ^{iv} —Na4—H421	73.9
C33—C34—H34A	109.5	Na1 ^{iv} —Na4—H421	119.4
C33—C34—H34B	109.5	O21 ^{vi} —Na5—O4	96.07 (5)
H34A—C34—H34B	109.5	O21 ^{vi} —Na5—O22 ⁱⁱ	86.80 (5)
C33—C34—H34C	109.5	O4—Na5—O22 ⁱⁱ	83.71 (4)
H34A—C34—H34C	109.5	O21 ^{vi} —Na5—O51	112.08 (5)
H34B—C34—H34C	109.5	O4—Na5—O51	93.32 (4)
C33—C34—H34D	109.5	O22 ⁱⁱ —Na5—O51	161.11 (5)
H34A—C34—H34D	141.1	O21 ^{vi} —Na5—O31	178.58 (5)
H34B—C34—H34D	56.3	O4—Na5—O31	83.09 (4)
H34C—C34—H34D	56.3	O22 ⁱⁱ —Na5—O31	91.97 (5)
C33—C34—H34E	109.5	O51—Na5—O31	69.14 (4)
H34A—C34—H34E	56.3	O21 ^{vi} —Na5—O5	88.01 (4)
H34B—C34—H34E	141.1	O4—Na5—O5	162.77 (5)
H34C—C34—H34E	56.3	O22 ⁱⁱ —Na5—O5	79.79 (4)
H34D—C34—H34E	109.5	O51—Na5—O5	100.67 (4)
C33—C34—H34F	109.5	O31—Na5—O5	92.47 (4)
H34A—C34—H34F	56.3	O21 ^{vi} —Na5—Na2 ^{vi}	45.38 (3)
H34B—C34—H34F	56.3	O4—Na5—Na2 ^{vi}	139.98 (4)
H34C—C34—H34F	141.1	O22 ⁱⁱ —Na5—Na2 ^{vi}	83.76 (4)
H34D—C34—H34F	109.5	O51—Na5—Na2 ^{vi}	109.59 (4)
H34E—C34—H34F	109.5	O31—Na5—Na2 ^{vi}	135.19 (4)
C31—C35—H35A	109.5	O5—Na5—Na2 ^{vi}	42.81 (3)
C31—C35—H35B	109.5	O21 ^{vi} —Na5—Na3	134.47 (4)
H35A—C35—H35B	109.5	O4—Na5—Na3	47.57 (4)
C31—C35—H35C	109.5	O22 ⁱⁱ —Na5—Na3	65.93 (3)
H35A—C35—H35C	109.5	O51—Na5—Na3	98.40 (4)
H35B—C35—H35C	109.5	O31—Na5—Na3	44.21 (3)
O6—C41—O5	125.44 (15)	O5—Na5—Na3	119.51 (4)
O6—C41—C4	118.26 (13)	Na2 ^{vi} —Na5—Na3	148.72 (2)
O5—C41—C4	116.30 (14)	O21 ^{vi} —Na5—Na2 ⁱⁱ	85.72 (4)
O8—C42—O7	124.95 (15)	O4—Na5—Na2 ⁱⁱ	123.06 (4)
O8—C42—C4	115.47 (13)	O22 ⁱⁱ —Na5—Na2 ⁱⁱ	39.43 (3)
O7—C42—C4	119.54 (13)	O51—Na5—Na2 ⁱⁱ	138.34 (4)
C41—O5—Na2 ^{vi}	119.93 (10)	O31—Na5—Na2 ⁱⁱ	93.78 (3)
C41—O5—Na2 ⁱⁱ	118.67 (10)	O5—Na5—Na2 ⁱⁱ	40.38 (3)
Na2 ^{vi} —O5—Na2 ⁱⁱ	90.52 (5)	Na2 ^{vi} —Na5—Na2 ⁱⁱ	56.27 (2)
C41—O5—Na5	133.06 (10)	Na3—Na5—Na2 ⁱⁱ	93.46 (2)
Na2 ^{vi} —O5—Na5	89.22 (4)	O21 ^{vi} —Na5—H31B	161.5
Na2 ⁱⁱ —O5—Na5	94.76 (4)	O4—Na5—H31B	80.8
C41—O6—Na4	159.80 (11)	O22 ⁱⁱ —Na5—H31B	110.8
C42—O8—Na2	134.19 (11)	O51—Na5—H31B	50.3

supplementary materials

O3—Na1—O3 ⁱ	180.00 (8)	O31—Na5—H31B	19.5
O3—Na1—O12 ⁱ	90.88 (4)	O5—Na5—H31B	100.4
O3 ⁱ —Na1—O12 ⁱ	89.12 (4)	Na2 ^{vi} —Na5—H31B	139.0
O3—Na1—O12	89.12 (4)	Na3—Na5—H31B	54.5
O3 ⁱ —Na1—O12	90.88 (4)	Na2 ⁱⁱ —Na5—H31B	111.4
O12 ⁱ —Na1—O12	180.00 (5)	Na4 ^v —O11—Na1	82.24 (4)
O3—Na1—O11 ⁱ	88.82 (4)	Na4 ^v —O11—H111	141.9
O3 ⁱ —Na1—O11 ⁱ	91.18 (4)	Na1—O11—H111	94.7
O12 ⁱ —Na1—O11 ⁱ	101.84 (4)	Na4 ^v —O11—H11B	111.7
O12—Na1—O11 ⁱ	78.16 (4)	Na1—O11—H11B	84.6
O3—Na1—O11	91.18 (4)	H111—O11—H11B	105.7
O3 ⁱ —Na1—O11	88.82 (4)	Na1—O12—Na4 ^{vii}	81.56 (4)
O12 ⁱ —Na1—O11	78.16 (4)	Na1—O12—H121	110.1
O12—Na1—O11	101.84 (4)	Na4 ^{vii} —O12—H121	123.7
O11 ⁱ —Na1—O11	180.00 (7)	Na1—O12—H122	116.0
O3—Na1—Na4 ^{vii}	56.04 (3)	Na4 ^{vii} —O12—H122	113.2
O3 ⁱ —Na1—Na4 ^{vii}	123.96 (3)	H121—O12—H122	109.7
O12 ⁱ —Na1—Na4 ^{vii}	130.22 (3)	Na5 ⁱⁱⁱ —O21—Na2	92.39 (5)
O12—Na1—Na4 ^{vii}	49.78 (3)	Na5 ⁱⁱⁱ —O21—H211	106.7
O11 ⁱ —Na1—Na4 ^{vii}	48.58 (3)	Na2—O21—H211	91.8
O11—Na1—Na4 ^{vii}	131.42 (3)	Na5 ⁱⁱⁱ —O21—H212	146.6
O3—Na1—Na4 ^v	123.96 (3)	Na2—O21—H212	95.0
O3 ⁱ —Na1—Na4 ^v	56.04 (3)	H211—O21—H212	105.5
O12 ⁱ —Na1—Na4 ^v	49.78 (3)	Na2—O22—Na5 ⁱⁱ	99.04 (5)
O12—Na1—Na4 ^v	130.22 (3)	Na2—O22—H221	104.5
O11 ⁱ —Na1—Na4 ^v	131.42 (3)	Na5 ⁱⁱ —O22—H221	120.1
O11—Na1—Na4 ^v	48.58 (3)	Na2—O22—H222	119.2
Na4 ^{vii} —Na1—Na4 ^v	180.00 (2)	Na5 ⁱⁱ —O22—H222	107.5
O3—Na1—H111	88.4	H221—O22—H222	107.1
O3 ⁱ —Na1—H111	91.6	Na3—O31—Na5	90.24 (4)
O12 ⁱ —Na1—H111	96.9	Na3—O31—H31A	140.3
O12—Na1—H111	83.1	Na5—O31—H31A	103.5
O11 ⁱ —Na1—H111	161.1	Na3—O31—H31B	113.4
O11—Na1—H111	18.9	Na5—O31—H31B	78.3
Na4 ^{vii} —Na1—H111	115.8	H31A—O31—H31B	105.8
Na4 ^v —Na1—H111	64.2	Na4—O41—Na4 ⁱⁱ	132.60 (9)
O3—Na1—H11B	73.0	Na4—O41—H411	97.1
O3 ⁱ —Na1—H11B	107.0	Na4 ⁱⁱ —O41—H411	108.3
O12 ⁱ —Na1—H11B	71.1	Na4—O42—H421	98.2
O12—Na1—H11B	108.9	Na4—O42—H422	105.5
O11 ⁱ —Na1—H11B	160.1	H421—O42—H422	107.9
O11—Na1—H11B	19.9	Na5—O51—H511	134.1

Na4 ^{vii} —Na1—H11B	121.6	Na5—O51—H512	107.4
Na4 ^v —Na1—H11B	58.4	H511—O51—H512	107.0

Symmetry codes: (i) $-x, -y+1, -z$; (ii) $-x, y, -z+1/2$; (iii) $x, -y, z+1/2$; (iv) $x, y-1, z$; (v) $-x, -y, -z$; (vi) $x, -y, z-1/2$; (vii) $x, y+1, z$; (viii) $-x, -y, -z+1$.

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O11—H111 \cdots O2 ⁱⁱ	0.86	1.97	2.7404 (16)	149
O11—H11B \cdots O4	0.85	1.88	2.7067 (16)	165
O12—H121 \cdots O1 ⁱⁱ	0.84	1.90	2.7391 (17)	174
O21—H211 \cdots O6 ⁱⁱ	0.85	1.92	2.7566 (17)	167
O21—H212 \cdots O7	0.85	1.97	2.8007 (16)	165
O22—H221 \cdots O31	0.85	2.10	2.8381 (17)	145
O22—H222 \cdots O11 ⁱⁱ	0.85	2.31	2.9707 (16)	135
O22—H222 \cdots O6 ⁱⁱⁱ	0.85	2.45	3.0532 (18)	128
O31—H31A \cdots O8	0.85	1.87	2.7128 (16)	169
O31—H31B \cdots O51	0.84	2.09	2.8085 (17)	142
O41—H411 \cdots O7	0.84	2.27	3.1061 (16)	172
O42—H421 \cdots O1 ^{iv}	0.85	1.92	2.7582 (17)	175
O42—H422 \cdots O7	0.85	2.03	2.8639 (17)	168
O51—H511 \cdots N4	0.86	1.99	2.8424 (18)	171
O51—H512 \cdots N2	0.85	1.94	2.7799 (18)	169

Symmetry codes: (ii) $-x, y, -z+1/2$; (iii) $x, -y, z+1/2$; (iv) $x, y-1, z$.

Fig. 1




Fig. 2

