

CRITICAL REVIEW

Drug-resistant epilepsy and the hypothesis of intrinsic severity: What about the high-frequency oscillations?

Cesar E. Santana-Gomez¹  | Jerome Engel Jr.^{1,2,3,4} | Richard Staba¹

¹Department of Neurology, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

²Brain Research Institute, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

³Department of Neurobiology, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

⁴Department of Psychiatry and Biobehavioral Sciences, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

Correspondence

Cesar E. Santana-Gomez, Department of Neurology, David Geffen School of Medicine, Reed Neurological Research Center, Suite 2155, 710 Westwood Plaza, Los Angeles, CA 90095, USA.
Email: csantanagomez@mednet.ucla.edu

Funding information

National Institute of Neurological Disorders and Stroke, Grant/Award Number: NS033310 and R01 NS106957

Abstract

Drug-resistant epilepsy (DRE) affects approximately one-third of the patients with epilepsy. Based on experimental findings from animal models and brain tissue from patients with DRE, different hypotheses have been proposed to explain the cause(s) of drug resistance. One is the intrinsic severity hypothesis that posits that drug resistance is an inherent property of epilepsy related to disease severity. Seizure frequency is one measure of epilepsy severity, but frequency alone is an incomplete measure of severity and does not fully explain basic research and clinical studies on drug resistance; thus, other measures of epilepsy severity are needed. One such measure could be pathological high-frequency oscillations (HFOs), which are believed to reflect the neuronal disturbances responsible for the development of epilepsy and the generation of spontaneous seizures. In this manuscript, we will briefly review the intrinsic severity hypothesis, describe basic and clinical research on HFOs in the epileptic brain, and based on this evidence discuss whether HFOs could be a clinical measure of epilepsy severity. Understanding the mechanisms of DRE is critical for producing breakthroughs in the development and testing of novel strategies for treatment.

KEYWORDS

drug-resistant epilepsy, EEG, epilepsy, epilepsy severity

1 | DRUG-RESISTANT EPILEPSY

Epilepsy is a brain disorder characterized by the enduring predisposition to generate spontaneous and recurrent seizures.¹ The World Health Organization reports epilepsy affects more than 50 million people around the world,² though others estimate there are 70 million people with epilepsy worldwide.³ Approximately 30% of persons diagnosed with epilepsy continue to have seizures despite taking one or

more antiseizure medication(s) (ASM), and in these cases, the individuals could be considered to have drug-resistant epilepsy (DRE). Though there is no precise definition of drug resistance, one definition considers DRE as a failure of two adequate trials of a well-tolerated and selected ASM regimen to produce sustained seizure freedom.⁴⁻⁶

The current research on drug resistance has focused on cellular and molecular alterations that lead to the reduction in drug sensitivity. Several decades of research

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. *Epilepsia Open* published by Wiley Periodicals LLC on behalf of International League Against Epilepsy.

from animals and resected tissue from surgical patients with DRE have led to two major hypotheses to explain ASM resistance.⁷⁻⁹ The first is the “multidrug transporter hypothesis,” which posits that DRE is caused by the overexpression or gain of function of multidrug transporters (such as P-glycoprotein) and members of multidrug resistance-associated proteins in the blood-brain barrier. This condition leads to a decrease in the concentrations of ASMs in the brain parenchyma and consequently a reduction in drug efficacy.⁷ The second is the “target hypothesis,” which postulates inherited or acquired changes in the structure, function, or localization of the drug targets lead to reduced pharmacodynamic effects of the ASM.⁸

1.1 | The hypothesis of intrinsic severity

Proving the two aforementioned hypotheses has been difficult. Some have noted the basis of these hypotheses, that is, drug resistance develops independently of epilepsy itself, might be constraining research on the mechanisms of drug resistance.¹⁰

An alternative explanation considers a clinical observation, that is, in treatment studies of patients with newly diagnosed epilepsy, those with a higher frequency of seizures before treatment often have lower likelihood for remission of seizures.¹¹ This observation is consistent with the progressive aspects of seizures as described by Gowers in the late 19th century,¹² and later by others who observed more seizures or longer duration of epilepsy is associated with lower probability of remission.¹³ These data suggest there is a continuum in severity of epilepsy and higher frequency of seizures at the early stage of epilepsy diagnosis indicates future drug resistance. In 2008 and again in 2013, Rogawski organized clinical data and the concept of epilepsy severity into the “intrinsic severity hypothesis,” which states “pharmacoresistance is an inherent property of epilepsy related to disease severity.”^{10,14}

The term severity defines the impact of the illness in the biological, physical, or psychosocial function of the patient and is an often-used descriptive feature for most medical symptoms and conditions, such as in the case of traumatic brain injury where injury can be mild, moderate, or severe.¹⁵ Severity is not commonly used to describe clinical epilepsy, and thus, there is little consensus on metrics to assess severity of epilepsy, and as noted by Rogawski in 2008, there is a need to identify measures of epilepsy severity. Nonetheless, as noted above, seizure frequency could be a measure of epilepsy severity associated with drug non-responsiveness. In a study of 780 newly diagnosed patients with epilepsy followed in a single center over 20 years, a higher seizure frequency before treatment with ASM was associated with drug resistance.¹¹ In this same study, the

Key Points

- The hypothesis of intrinsic severity theorizes that DRE is an inherent property of epilepsy related to the severity of the disease.
- Seizure frequency is a relevant marker of the severity, but not the only measure of severity, to predict DRE.
- PIN clusters are proposed to generate pathological HFOs and believed to reflect neuronal disturbances responsible for epilepsy.
- Pathological HFOs correlate with rate of epileptogenesis and, in some types of epilepsy, epileptogenicity of the lesion and propensity for seizures.
- HFOs could be used as an additional clinical marker to evaluate the severity of the disease and the possible development of DRE.

presence of psychiatric comorbidities such as depression, anxiety, and psychosis, as well as recreational drug use, also was associated with DRE.¹¹ Sillampää and colleagues observed that pediatric patients who had three or more seizures in 24 hours, or a seizure cluster, had a lower likelihood for seizure remission than patients without a seizure cluster during the same period.¹⁶ Other work found patients with 2 or fewer seizures in the first 6 months after epilepsy diagnosis had a 77% probability of being seizure-free for 1 year within 2 years of diagnosis. However, patients with 10 seizures during the first 6 months had a 50% chance of being seizure-free during the same period.¹⁷ Based on this evidence, the authors concluded that the rate of epileptic seizures during the first 6 months after the disease diagnosis could be an important and predictive marker for the development of DRE.¹⁷ Likewise, clinical trials have been shown that the delay in adequate treatment with ASM gradually increases the risk of DRE.¹⁸ However, other studies found patients with a low number of seizures at the beginning of the disease could develop DRE.¹⁷

Some of these clinical observations are also seen in pre-clinical studies of epileptogenesis.^{19,20} In these rat studies, self-sustained status epilepticus was induced by electrical stimulation of the basolateral amygdala, and then, rats that developed subsequent spontaneous seizures were treated with phenobarbital. Results show rats with a low seizure frequency before initial treatment were more responsive to phenobarbital and rats with a high seizure frequency were less responsive to the same treatment.^{19,20} However, some of the unresponsive rats had a low seizure

rate similar to responsive rats.²¹ These data indicate seizure frequency is a good predictor of DRE, but it does not explain drug response in all cases and thus unlikely to be the only measure of epilepsy severity. Other measures could include occurrence of precipitating injury such as status epilepticus or traumatic brain injury, the type of seizure, a lesion on MRI verified by electrophysiology as epileptogenic and potentially unresponsiveness to medication, and other functional disturbances in the EEG, like high-frequency oscillations (HFOs).

1.2 | Pathologically interconnected neuron clusters in the epileptic brain

Studies in the epileptic brain show an increase in basal concentrations of glutamate in epileptogenic and nonepileptogenic brain areas.^{22–26} Once a seizure begins, extracellular glutamate levels increase and remain elevated after the seizure has ended.^{22–25} A consequence of this sustained high glutamate level is continuous excitation of neurons, producing abnormally high levels of intracellular calcium, deregulation of signaling pathways and organelle dysfunction, and ultimately cell death.²⁷ Cell loss changes the tissue architecture and among surviving neurons could induce axonal sprouting and activity-dependent synaptic reorganization.^{28,29} These morphological alterations could strengthen excitatory connections between neurons, leading to the formation of pathologically interconnected neuron (PIN) clusters.³⁰ PIN clusters are hypothesized to be the mechanism for generating pathological synchronous bursts of population spikes, which are recorded as HFOs, and could represent the neuronal disturbances responsible for epilepsy.³¹

1.2.1 | High-frequency oscillations

High-frequency oscillations are local field potentials that correspond to an increase in synchronous neuronal spike bursts.³² Based on their spectral frequency properties, HFOs are classified into “ripples” (80–200 Hz) and “fast ripples” (FRs, 200–800 Hz).³² Ripples have been recorded in the normal hippocampus and parahippocampal structures of mammals, including rodents, nonhuman primates, and humans,^{33–37} as well as in eloquent cortex of humans.³⁸ Normal ripples correspond to summated inhibitory postsynaptic potentials (IPSPs) of rhythmically discharging interneurons, which regulate pyramidal cell firing,^{33,35} and in the hippocampus, ripples play a key role in the memory formation, reactivation of previous experiences, and information processing (eg, during somatosensory evoked potentials).³⁹ On the contrary, FRs have been recorded in patients and many animal models

of epilepsy.^{40–42} FRs reflect as a burst of population spikes generated from the out-of-phase firing of distinct cell clusters consisting primarily of principal cells.^{42–50}

In animals and patients with epilepsy, both ripples and FRs have been associated with the seizure-onset zone (SOZ), suggesting some ripple frequency oscillations could also be pathological HFOs.^{42,51–53} For example, in the normal dentate gyrus, oscillatory activity is typically lower than 100 Hz.^{54,55} However, Bragin and colleagues found ripples >100 Hz in the epileptogenic dentate gyrus of kainic acid-treated rats.^{56,57} Several patient studies of epilepsy surgery indicate resection of pathological HFO-generating areas is important for a seizure-free outcome.^{58–61}

1.2.2 | High-frequency oscillations and epileptogenesis

In 2000, Bragin and colleagues hypothesized the development of PIN clusters generating pathological HFOs as a mechanism of epileptogenesis (ie, development of epilepsy).³⁰ Much of the evidence for this hypothesis was derived from the intrahippocampal kainic acid rat model of human temporal lobe epilepsy where rats receive a unilateral injection of kainic acid in posterior hippocampus and status epilepticus begins within 20 minutes of injection and lasts for up to 6 hours. Not all rats that have status epilepticus later develop epilepsy, but between 50% and 75% of them do, indicating there are differences in susceptibility between different strains and between rats within the same strain. In one study, using kainic acid-treated rats, HFOs were found in the dentate gyrus, CA1, and entorhinal cortex ipsilateral to the injection site of rats that later developed epilepsy, but HFOs were not found in rats that did not develop epilepsy.⁵⁷ In this same study, the sooner HFOs appeared after status epilepticus, the sooner the first spontaneous seizure occurred, the shorter the interval between first HFOs and first seizure, and the higher the rate of seizures per month.⁵⁷ However, the rate of HFOs, when they first appeared, which could be as early as one day after status epilepticus, was unrelated to the rate of subsequent spontaneous seizures. In other studies, higher rates of pathological HFOs were found inside and outside the hippocampus in animals that developed epilepsy than those that did not develop epilepsy.^{62,63} It appears pathological HFOs in the epileptic hippocampus can also trigger the occurrence of remote HFOs, as well as spindles,⁶⁴ and likely through cross-frequency coupling with slow waves could contribute to epileptogenesis.⁶⁵ Recording from epileptic rats shows a greater number of electrodes with FRs correlate with a higher number of seizures per day.⁴¹ Interestingly, an intracranial EEG study involving presurgical patients with focal epilepsy found a greater number

of electrode contacts recording more than 20 FRs per minute correlated with a higher number of seizures per month, especially in cases of temporal lobe epilepsy.⁶⁶ In addition to the intrahippocampal kainic acid rat model, pathological HFOs can be recorded in the fluid percussion injury rat model of human traumatic brain injury and posttraumatic epilepsy. Studies recorded pathological HFOs alone or superimposed on EEG spikes within the first two weeks after fluid percussion injury only in rats that later developed focal spontaneous seizures.^{67,68} Results from the intrahippocampal kainic acid and fluid percussion injury models suggest pathological HFOs could be an electrophysiological biomarker of epileptogenesis. The neuronal mechanisms of epileptogenesis could be due to the formation of PIN clusters.⁶⁹ After an episode of status epilepticus or a traumatic brain injury, PIN clusters could form and, through a kindling-like process, lead to the formation of widespread PIN clusters (Figure 1). A greater number of PIN clusters lead the increase in amplitude and duration of HFOs leading up to seizure onset, and correlating with higher density of the PIN clusters with more frequent seizures.^{30,62,70}

1.2.3 | High-frequency oscillations and epilepsy severity

Several studies have identified links between HFOs and aspects of epilepsy that could be associated with severity, such as frequency and type of seizures, MRI pathology,

and, in some types of epilepsy, the propensity to generate seizures. In addition to the studies on interictal FRs and seizure frequency in rats and patients with epilepsy noted in Section 1.2.2, a study in children with drug-resistant focal epilepsy showed higher rates of HFOs in preoperative scalp EEG correlate with higher seizure frequency.⁷¹ Results like this are also observed between interictal ripples and number of seizures in cases of childhood epilepsy with centrottemporal spikes (CECTS).⁷² With respect to the type of seizures in rats, rates of ripples and FRs are higher during seizures with clonus, rearing and falling or with wild running than rates during seizures with rearing only (no falling) or movements limited to forelimbs, face, mouth, or eyes.⁷³ A similar analysis performed in patients showed ripple activity in the SOZ is higher during seizures that subsequently evolve with bilateral tonic-clonic movements than during focal seizures without such movements.⁷⁴ These latter data are consistent with a previous patient study that showed, during low voltage fast (LVF) seizures, a common ictal EEG pattern associated with regional-onset seizures,⁷⁵ that ripples increase in the SOZ and sites of spread.⁷⁶ The LVF activity is associated with an increase in inhibitory cell firing and a reduction in excitatory cell firing that subsequently rebounds.⁷⁷ One interpretation of these data is that a widespread HFO network generates seizures and can facilitate seizure spread. Alternatively, the ripples could be a mechanism to prevent seizure spread that ultimately fails, which might explain the pattern of inhibitory and excitatory cell firing.




FIGURE 1 High-frequency oscillations (HFOs) as a potential marker of epilepsy severity. After an epileptogenic insult, the appearance of HFOs precedes and contributes to the development of clinical seizures (denoted by outer arc). In some cases, HFOs could appear immediately after injury, and lead to the formation of a more extensive HFO network, and soon thereafter the appearance of clinical seizures (inner arc). Once epilepsy is established, the rate of HFOs or the number of HFO-generating sites or both increases to generate more frequent seizures (denoted by black dashed arrow). If the intrinsic severity hypothesis is correct and HFOs are a measure of epilepsy severity, then a high burden of HFOs would predict a severe form of epilepsy with high seizure frequency (solid blue arrow) that could be unresponsive to medication, that is, drug-resistant epilepsy (DRE) (blue dashed arrow)

In mesial temporal lobe epilepsy, a higher ratio of FR to ripples correlates with reduced hippocampal volume and cell loss associated with hippocampal sclerosis.⁷⁸ Similar results can be found in pilocarpine-treated epileptic rats.⁴⁶ In a separate patient study, higher rates of FR alone, but not ripples, correlated with local areas of atrophy in hippocampi ipsilateral and contralateral to the SOZ.⁷⁹ This latter study included patients who had drug-resistant seizures, and results suggest the extent of atrophy and the size or number of the brain areas generating FR are greater in patients with many years of poorly controlled seizures. This interpretation is consistent with the positive correlation between the number of electrodes with FR and seizure frequency in rats with chronic epilepsy.⁴¹ In cases with focal cortical dysplasia (FCD), patients with FCD type 2 have significantly more seizures and higher rates of HFOs than patients with FCD type 1.⁸⁰ Results from these studies suggest HFOs could correspond to the epileptogenicity of the lesion and a measure of epilepsy severity.

Several studies have found HFOs, particularly ripples, either alone or superimposed on spikes (spike ripples) in cases of CECTS. The highest number of ripples is associated with frequent seizures in patients with atypical CECTS and the fewest or absence of ripples in classical CECTS or those patients without seizures.^{72,81} Ripples are better than spikes alone for predicting seizures and can help distinguish patients with atypical CECTS from those with few or no seizures. In other work of CECTS, spike ripples are associated with more recent seizures,⁸² and rates of spike ripples are higher in patients who have at least one seizure within one year than in patients who are seizure-free for more than one year.⁸³ Also, a longer duration without a seizure is associated with a lower spike ripple rate. Patients with CECTS and scalp-recorded HFOs have more cognitive deficits than patients without HFOs.⁸⁴ These data suggest in some types of epilepsy, HFOs correspond to active epilepsy. In a syndrome such as CECTS with a spectrum of severity of focal seizures and neurocognitive impairments, HFOs indicate a greater propensity for seizures and cognitive deficits.

In surgical pediatric patients with diverse etiologies and focal seizures, high rates of scalp-recorded HFOs over the affected hemisphere are associated with recurrent seizures and rates of HFOs decrease after surgery in patients whose seizures are reduced or eliminated.⁷¹ Another study found in surgical patients, HFOs in pre- and postoperative electrocorticography are associated with recurrent seizures.⁸⁵ Interestingly, in some patients, preoperative HFOs outside the resection margins disappear on postoperative recordings, and in a few patients, sites with “new” HFOs could be found in postoperative recordings that were not in preoperative recordings. Of these latter patients, one had recurrent seizures postoperatively, and the other was

seizure-free. There could exist HFO “hub” sites that maintain a high number of direct (possibly synaptic) connections with remote HFO sites, and these interconnected sites could coordinate activity through preferred phases of low-frequency EEG rhythms (see Section 1.2.2).⁸⁶ In this scenario, severe forms of epilepsy could have several HFO hubs that have a strong modulatory effect on remote sites,⁸⁷ and resection of HFO hubs renders remote HFOs sites inactive and abolishes seizures.

1.2.4 | The effect of antiseizure drugs on high-frequency oscillations

A study of the effects of seizures and ASM on spikes and HFOs showed there are higher rates of spikes, but not HFOs, after than before a seizure.⁸⁸ The number of electrode contacts with spikes or HFOs is also higher after a seizure. During reduction in ASM, there is no change in the rates of spikes, but there is an increase in the rate and duration of HFOs. In other work on West syndrome, scalp-recorded interictal HFOs were higher in patients with seizures before the start of treatment than in controls without seizures.⁸⁹ During treatment, the decrease in hypsarrhythmia and seizures is accompanied by a decrease in number of HFOs. A similar decrease is found in HFOs after treatment in patients with atypical benign partial epilepsy.⁸¹ More recently, a study of interictal HFOs after treatment for epileptic spasms showed a higher incidence of HFOs coupling with the trough-to-peak transition of 2-3 Hz slow waves in patients unresponsive to treatment than those who responded to treatment.⁹⁰ Previous studies also found a strong association between HFOs coupling with the trough-to-peak transition of the slow wave in epileptogenic tissue in patients with drug-resistant seizures.⁹¹⁻⁹³ The mechanism(s) for HFO coupling with slow waves described in these cited studies are not known, but one explanation is that the trough-to-peak transition of the slow wave produces a powerful depolarizing volley that excites PIN clusters and generates pathological HFOs. HFOs correspond to abnormal neuronal activity involved with generating seizures, and lowering the threshold for seizures as in the case of ASM withdrawal increases HFOs and likelihood for seizures.

1.3 | Summary and conclusions

The intrinsic severity hypothesis proposes there is a continuum in severity of epilepsy that corresponds to a likelihood the epilepsy will respond to ASM. A high seizure frequency is a risk factor for refractoriness, though the hypothesis does not explain why patients with infrequent

seizures do not respond to ASM or why seizure control on ASMs can vary over time. Seizure frequency should not be the only measure of severity, and other measures such as pathological HFOs appear to reflect other aspects of epilepsy severity.

PIN clusters are proposed to be the mechanism for generating pathological HFOs and may contribute to the generation of spontaneous epileptic seizures (Figure 1). During the development of epilepsy, the greater the extent of pathological HFO-generating sites in terms of their number and spatial distribution, the sooner seizures appear with greater frequency, indicating pathological HFOs could be a measure of the rate and strength of epileptogenesis.

In patients with medication-resistant epilepsy and some childhood epilepsies, increased irritability within PIN clusters as measured by the rate of HFO discharges corresponds with greater propensity to generate seizures. In CECTS, for example, HFOs (particularly ripples) or spikes with HFOs are better than spikes alone to predict active epilepsy that require ASM treatment.

Pathological HFOs are associated with some types of FCD and histological cell loss and MRI atrophy, especially in cases of hippocampal sclerosis, which could correspond to the extent of structural abnormality and epileptogenicity of these lesions. It appears widespread pathological HFOs indicate a large area of brain capable of generating seizures and could be a mechanism for seizure spread with greater motor and behavioral impairment. Incomplete resection of pathological HFOs reduces the likelihood for seizure-free outcome in many, but not all, surgical patients, which implies there could be hub-like sites generating HFOs. Severe forms of epilepsy could have a greater number of HFO hubs that need to be removed to eliminate seizures.

Invasive EEG is best for recording pathological HFOs, especially in deeper brain areas. Scalp EEG can record HFOs and is used more frequently, carries less risk, and has greater spatial sampling than invasive EEG. In clinical epilepsy, recording HFOs could provide a measure of the risk for developing epilepsy and propensity for spontaneous seizures, and, in some types of epilepsy, the epileptogenicity of a lesion and severity of the disease.

ACKNOWLEDGMENT

This study was supported by NIH R01 NS106957 (RS) and NS033310 (JE).

CONFLICT OF INTEREST

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. We confirm that we have read the Journal's position on

issues involved in ethical publication and affirm that this report is consistent with those guidelines.

ORCID

Cesar E. Santana-Gomez  <https://orcid.org/0000-0001-9163-5615>

REFERENCES

1. Fisher RS, Acevedo C, Arzimanoglou A, Bogacz A, Cross JH, Elger CE, et al. ILAE official report: a practical clinical definition of epilepsy. *Epilepsia*. 2014;55(4):475–82.
2. World Health Organization 2021. <https://www.who.int/news-room/fact-sheets/detail/epilepsy#:~:text=Globally%2C%20an%20estimated%20five%20million,as%20139%20per%20100%20000>
3. Thijs RD, Surges R, O'Brien TJ, Sander JW. Epilepsy in adults. *Lancet*. 2019;393(10172):689–701. [https://doi.org/10.1016/S0140-6736\(18\)32596-0](https://doi.org/10.1016/S0140-6736(18)32596-0)
4. Chen Z, Brodie MJ, Liew D, Kwan P. Treatment outcomes in patients with newly diagnosed epilepsy treated with established and new antiepileptic drugs a 30-year longitudinal cohort study. *JAMA Neurol*. 2018;75(3):279–86.
5. Kalilani L, Sun X, Pelgrims B, Noack-Rink M, Villanueva V. The epidemiology of drug-resistant epilepsy: a systematic review and meta-analysis. *Epilepsia*. 2018;59(12):2179–93.
6. Kwan P, Arzimanoglou A, Berg AT, Brodie MJ, Hauser WA, Mathern G, et al. Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia*. 2010;51(6):1069–77.
7. Löscher W, Potschka H. Role of multidrug transporters in pharmacoresistance to antiepileptic drugs. *J Pharmacol Exp Ther*. 2002;301(1):7–14.
8. Remy S, Beck H. Molecular and cellular mechanisms of pharmacoresistance in epilepsy. *Brain*. 2006;129(1):18–35.
9. Löscher W, Potschka H. Drug resistance in brain diseases and the role of drug efflux transporters. *Nat Rev Neurosci*. 2005;6(8):591–602.
10. Rogawski MA, Johnson MR. Intrinsic severity as a determinant of antiepileptic drug refractoriness. *Epilepsy Curr*. 2008;8(5):127–30.
11. Hitiris N, Mohanraj R, Norrie J, Sills GJ, Brodie MJ. Predictors of pharmacoresistant epilepsy. *Epilepsy Res*. 2007;75(2–3):192–6.
12. Gowers W. *Epilepsy and other convulsive disorders: their causes, symptoms and treatment*. London: Churchill; 1881.
13. Reynolds EH, Elwes RD. Why does epilepsy become intractable? Prevention of chronic epilepsy. *Lancet*. 1983;322(8356):952–4.
14. Rogawski MA. The intrinsic severity hypothesis of pharmacoresistance to antiepileptic drugs. *Epilepsia*. 2013;54(Suppl 2):33–40.
15. Mckee AC, Daneshvar DH. The neuropathology of traumatic brain injury. *Handb. Clin. Neurol*. 2015;45–66.
16. Sillanpää M, Schmidt D. Seizure clustering during drug treatment affects seizure outcome and mortality of childhood-onset epilepsy. *Brain*. 2008;131(4):938–44.
17. MacDonald BK, Johnson AL, Goodridge DM, Cockerell OC, Sander JWAS, Shorvon SD. Factors predicting prognosis of epilepsy after presentation with seizures. *Ann Neurol*. 2000;48(6):833–41.

18. Sillanpää M, Schmidt D. Delayed time to first remission identifies poor long-term drug response of childhood-onset epilepsy: a prospective population-based study. *Epilepsy Behav.* 2009;16(3):507–11. <https://doi.org/10.1016/j.yebeh.2009.08.029>
19. Volk HA, Arabadzisz D, Fritschy JM, Brandt C, Bethmann K, Löscher W. Antiepileptic drug-resistant rats differ from drug-responsive rats in hippocampal neurodegeneration and GABAA receptor ligand binding in a model of temporal lobe epilepsy. *Neurobiol Dis.* 2006;21(3):633–46.
20. Brandt C, Glien M, Potschka H, Volk H, Löscher W. Epileptogenesis and neuropathology after different types of status epilepticus induced by prolonged electrical stimulation of the basolateral amygdala in rats. *Epilepsy Res.* 2003;55(1–2):83–103.
21. Löscher W, Brandt C. High seizure frequency prior to antiepileptic treatment is a predictor of pharmacoresistant epilepsy in a rat model of temporal lobe epilepsy. *Epilepsia.* 2010;51(1):89–97.
22. During MJ, Spencer SS. Extracellular hippocampal glutamate and spontaneous seizure in the conscious human brain. *Lancet.* 1993;341:1607–10.
23. Soukupova M, Binaschi A, Falcicchia C, Palma E, Roncon P, Zucchini S, et al. Increased extracellular levels of glutamate in the hippocampus of chronically epileptic rats. *Neuroscience.* 2015;301:246–53. <http://www.ncbi.nlm.nih.gov/pubmed/26073699>
24. Ronne-Engström E, Hillered L, Flink R, Spännare B, Ungerstedt U, Carlson H. Intracerebral microdialysis of extracellular amino acids in the human epileptic focus. *J Cereb Blood Flow Metab.* 1992;12(5):873–6.
25. Carlson H, Ronne-Engström E, Ungerstedt U, Hillered L. Seizure related elevations of extracellular amino acids in human focal epilepsy. *Neurosci Lett.* 1992;140(1):30–2.
26. Santana-Gómez CE, Orozco-Suárez SA, Talevi A, Bruno-Blanch L, Magdaleno-Madrigal VM, Fernández-Mas R, et al. Propylparaben applied after pilocarpine-induced status epilepticus modifies hippocampal excitability and glutamate release in rats. *Neurotoxicology.* 2017;59:110–20. <http://linkinghub.elsevier.com/retrieve/pii/S0161813X17300207>
27. Tapia R, Medina-Ceja L, Peña F. On the relationship between extracellular glutamate, hyperexcitation and neurodegeneration, in vivo. *Neurochem Int.* 1999;34(1):23–31.
28. Covolan L, Mello LE. Assessment of the progressive nature of cell damage in the pilocarpine model of epilepsy. *Braz J Med Biol Res.* 2006;39(7):915–24. <http://www.ncbi.nlm.nih.gov/pubmed/16862283>
29. Fujikawa DG. Prolonged seizures and cellular injury: understanding the connection. *Epilepsy Behav.* 2005;7:S3–11. <http://www.ncbi.nlm.nih.gov/pubmed/16278099>
30. Bragin A, Wilson CL, Engel J. Chronic epileptogenesis requires development of a network of pathologically interconnected neuron clusters: a hypothesis. *Epilepsia.* 2000;41(s6):S144–52. <https://doi.org/10.1111/j.1528-1157.2000.tb01573.x>
31. Jefferys JGR, Menendez de la Prida L, Wendling F, Bragin A, Avoli M, Timofeev I, et al. Mechanisms of physiological and epileptic HFO generation. *Prog Neurobiol.* 2012;98(3):250–64. <http://linkinghub.elsevier.com/retrieve/pii/S0301008212000317>
32. Engel J, Bragin A, Staba R, Mody I. High-frequency oscillations: what is normal and what is not? *Epilepsia.* 2009;50(4):598–604.
33. Buzsáki G, Horvath Z, Urioste R, Wise KD. High-frequency network oscillation in the hippocampus. *Science.* 1992;256(5059):1025–7.
34. Chrobak JJ, Buzsáki G. High-frequency oscillations in the output networks of the hippocampal-entorhinal axis of the freely behaving rat. *J Neurosci.* 1996;16(9):3056–66.
35. Ylinen A, Bragin A, Nádasdy Z, Jandó G, Szabó I, Sik A, et al. Sharp wave-associated high-frequency oscillation (200 Hz) in the intact hippocampus: network and intracellular mechanisms. *J Neurosci.* 1995;15(1 Pt 1):30–46. <http://www.ncbi.nlm.nih.gov/pubmed/7823136>
36. Staba RJ, Wilson CL, Bragin A, Fried I, Engel J. Quantitative analysis of high-frequency oscillations (80–500 Hz) recorded in human epileptic hippocampus and entorhinal cortex. *J Neurophysiol.* 2002;88(4):1743–52. <http://www.ncbi.nlm.nih.gov/pubmed/12364503>
37. Schönberger J, Draguhn A, Both M. Lamina-specific contribution of glutamatergic and GABAergic potentials to hippocampal sharp wave-ripple complexes. *Front Neural Circuits.* 2014;8:1–11.
38. Frauscher B, von Ellenrieder N, Zemann R, Rogers C, Nguyen DK, Kahane P, et al. High-frequency oscillations in the normal human brain. *Ann Neurol.* 2018;84(3):374–85.
39. Buzsáki G, editor. *Rhythms of the brain.* New York: Oxford University Press; 2006. 448 pp.
40. Staba RJ, Bragin A. High-frequency oscillations and other electrophysiological biomarkers of epilepsy: underlying mechanisms. *Biomark Med.* 2011;5(5):545–56.
41. Bragin A, Wilson CL, Engel J. Spatial stability over time of brain areas generating fast ripples in the epileptic rat. *Epilepsia.* 2003;44(9):1233–7.
42. Jirsch JD, Urrestarazu E, LeVan P, Olivier A, Dubeau F, Gotman J. High-frequency oscillations during human focal seizures. *Brain.* 2006;129(6):1593–608.
43. Bragin A, Engel J, Wilson CL, Fried I, Mathern GW. Hippocampal and entorhinal cortex high-frequency oscillations (100–500 Hz) in human epileptic brain and in kainic acid-treated rats with chronic seizures. *Epilepsia.* 1999;40(2):127–37.
44. Bragin A, Engel J, Wilson CL, Fried I, Buzsáki G. High-frequency oscillations in human brain. *Hippocampus.* 1999;9(2):137–42.
45. Dzhala VI, Staley KJ. Mechanisms of fast ripples in the hippocampus. *J Neurosci.* 2004;24(40):8896–906.
46. Foffani G, Uzcategui YG, Gal B, Menendez de la Prida L. Reduced spike-timing reliability correlates with the emergence of fast ripples in the rat epileptic hippocampus. *Neuron.* 2007;55(6):930–41.
47. Ochi A, Otsubo H, Donner EJ, Elliott I, Iwata R, Funaki T, et al. Dynamic changes of ictal high-frequency oscillations in neocortical epilepsy: Using multiple band frequency analysis. *Epilepsia.* 2007;48(2):286–96.
48. Spampinato J, Mody I. Spike timing of lacunosom-molecular targeting interneurons and CA3 pyramidal cells during high-frequency network oscillations in vitro. *J Neurophysiol.* 2007;98(1):96–104.
49. Demont-Guignard S, Benquet P, Gerber U, Biraben A, Martin B, Wendling F. Distinct hyperexcitability mechanisms underlie fast ripples and epileptic spikes. *Ann Neurol.* 2012;71(3):342–52.
50. Bragin A, Benassi SK, Kheiri F, Engel J. Further evidence that pathologic high-frequency oscillations are bursts of population

- spikes derived from recordings of identified cells in dentate gyrus. *Epilepsia*. 2011;52(1):45–52.
51. Fisher R, Webber W, Lesser R, Arroyo S, Uematsu S. High frequency EEG activity at the start of seizures. *J Clin Neurophysiol*. 1992;9:441–8.
 52. Staba RJ, Wilson CL, Bragin A, Jhung D, Fried I, Engel J. High-frequency oscillations recorded in human medial temporal lobe during sleep. *Ann Neurol*. 2004;56(1):108–15.
 53. Grenier F, Timofeev I, Steriade M. Neocortical very fast oscillations (ripples, 80–200 Hz) during seizures: intracellular correlates. *J Neurophysiol*. 2003;89(2):841–52.
 54. Bragin A, Jando G, Nadasy Z, Hetke J, Wise K, Buzsáki G. Gamma (40–100 Hz) oscillation in the hippocampus of the behaving rat. *J Neurosci*. 1995;15(1):47–60. <https://doi.org/10.1523/JNEUROSCI.15-01-00047.1995>
 55. Csicsvari J, Jamieson B, Wise KD, Buzsáki G. Mechanisms of gamma oscillations in the hippocampus of the behaving rat. *Neuron*. 2003;37(2):311–22.
 56. Bragin A, Wilson CL, Engel J. Voltage depth profiles of high-frequency oscillations after kainic acid-induced status epilepticus. *Epilepsia*. 2007;48(Suppl. 5):35–40.
 57. Bragin A, Wilson CL, Almajano J, Mody I, Engel J. High-frequency oscillations after status epilepticus: epileptogenesis and seizure genesis. *Epilepsia*. 2004;45(9):1017–23.
 58. Jacobs J, Zijlmans M, Zelmann R, Chatillon CÉ, Hall J, Olivier A, et al. High-frequency electroencephalographic oscillations correlate with outcome of epilepsy surgery. *Ann Neurol*. 2010;67(2):209–20.
 59. Wu JY, Sankar R, Lerner JT, Matsumoto JH, Vinters HV, Mathern GW. Removing interictal fast ripples on electrocorticography linked with seizure freedom in children. *Neurology*. 2010;75(19):1686–94.
 60. Höller Y, Kutil R, Klaffenböck L, Thomschewski A, Höller PM, Bathke AC, et al. High-frequency oscillations in epilepsy and surgical outcome. A meta-analysis. *Front Hum Neurosci*. 2015;9:1–14.
 61. Jacobs J, Wu JY, Perucca P, Zelmann R, Mader M, Dubeau F, et al. Removing high-frequency oscillations: a prospective multicenter study on seizure outcome. *Neurology*. 2018;91(11):e1040–52.
 62. Li L, Patel M, Almajano J, Engel J, Bragin A. Extrahippocampal high-frequency oscillations during epileptogenesis. *Epilepsia*. 2018;59(4):e51–5.
 63. Sheybani L, Birot G, Contestabile A, Seeck M, Kiss JZ, Schaller K, et al. Electrophysiological evidence for the development of a self-sustained large-scale epileptic network in the kainate mouse model of temporal lobe epilepsy. *J Neurosci*. 2018;38(15):3776–91.
 64. Gelinás JN, Khodagholy D, Thesen T, Devinsky O, Buzsáki G. Interictal epileptiform discharges induce hippocampal-cortical coupling in temporal lobe epilepsy. *Nat Med*. 2016;22(6):641–8.
 65. Sheybani L, Van Mierlo P, Birot G, Michel CM, Quairiaux C. Large-scale 3–5 Hz oscillation constrains the expression of neocortical fast ripples in a mouse model of Mesial temporal lobe epilepsy. *eNeuro*. 2019;6(1):1–13.
 66. Zijlmans M, Jacobs J, Zelmann R, Dubeau F, Gotman J. High frequency oscillations and seizure frequency in patients with focal epilepsy. *Clin Neurophysiol*. 2009;122(4):664–71.
 67. Reid AY, Bragin A, Giza CC, Staba RJ, Engel J. The progression of electrophysiologic abnormalities during epileptogenesis after experimental traumatic brain injury. *Epilepsia*. 2016;57(10):1558–67.
 68. Bragin A, Li L, Almajano J, Alvarado-Rojas C, Reid AY, Staba RJ, et al. Pathologic electrographic changes after experimental traumatic brain injury. *Epilepsia*. 2016;57(5):735–45.
 69. Li L, Bragin A, Staba R, Engel J. Unit firing and oscillations at seizure onset in epileptic rodents. *Neurobiol Dis*. 2019;127:382–9. <https://doi.org/10.1016/j.nbd.2019.03.027>
 70. Weiss SA, Alvarado-Rojas C, Bragin A, Behnke E, Fields T, Fried I, et al. Ictal onset patterns of local field potentials, high frequency oscillations, and unit activity in human mesial temporal lobe epilepsy. *Epilepsia*. 2016;57(1):111–21.
 71. Boran E, Sarnthein J, Krayenbühl N, Ramantani G, Fedele T. High-frequency oscillations in scalp EEG mirror seizure frequency in pediatric focal epilepsy. *Sci Rep*. 2019;9(1):1–10.
 72. van Klink NEC, van 't Klooster MA, Leijten FSS, Jacobs J, Braun KPJ, Zijlmans M. Ripples on rolandic spikes: a marker of epilepsy severity. *Epilepsia*. 2016;57(7):1179–89.
 73. Birk N, Schönberger J, Somerlik-Fuchs KH, Schulze-Bonhage A, Jacobs J. Ictal occurrence of high-frequency oscillations correlates with seizure severity in a rat model of temporal lobe epilepsy. *Front Hum Neurosci*. 2021;15:1–8.
 74. Schönberger J, Birk N, Lachner-Piza D, Dümpelmann M, Schulze-Bonhage A, Jacobs J. High-frequency oscillations mirror severity of human temporal lobe seizures. *Ann Clin Transl Neurol*. 2019;6(12):2479–88.
 75. Memarian N, Madsen SK, Macey PM, Fried I, Engel J, Thompson PM, et al. Ictal depth EEG and MRI structural evidence for two different epileptogenic networks in mesial temporal lobe epilepsy. *PLoS One*. 2015;10(4):1–16.
 76. Weiss SA, Orosz I, Salamon N, Stephanie Moy, Wei L, van 't Klooster MA, et al. Ripples on spikes show increased phase-amplitude coupling in mesial temporal lobe epilepsy seizure onset zones. *Epilepsia*. 2016;57(11):1916–30.
 77. Elahian B, Lado NE, Mankin E, Vangala S, Misra A, Moxon K, et al. Low-voltage fast seizures in humans begin with increased interneuron firing. *Ann Neurol*. 2018;84(4):588–600.
 78. Staba RJ, Frighetto L, Behnke EJ, Mathern GW, Fields T, Bragin A, et al. Increased fast ripple to ripple ratios correlate with reduced hippocampal volumes and neuron loss in temporal lobe epilepsy patients. *Epilepsia*. 2007;48(11):2130–8.
 79. Ogren JA, Wilson CL, Bragin A, Lin JJ, Salamon N, Dutton RA, et al. 3D surface maps link local atrophy & fast ripples in human epileptic hippocampus. *Ann Neurol*. 2009;66(6):783–91.
 80. Kerber K, Levan P, Dümpelmann M, Fauser S, Korinthenberg R, Schulze-Bonhage A, et al. High frequency oscillations mirror disease activity in patients with focal cortical dysplasia. *Epilepsia*. 2013;54(8):1428–36.
 81. Qian P, Li H, Xue J, Yang Z. Scalp-recorded high-frequency oscillations in atypical benign partial epilepsy. *Clin Neurophysiol*. 2016;127(10):3306–13. <https://doi.org/10.1016/j.clinph.2016.07.013>
 82. Kobayashi K, Yoshinaga H, Toda Y, Inoue T, Oka M, Ohtsuka Y. High-frequency oscillations in idiopathic partial epilepsy of childhood. *Epilepsia*. 2011;52(10):1812–9.
 83. Kramer MA, Ostrowski LM, Song DY, Thorn EL, Stoyell SM, Parnes M, et al. Scalp recorded spike ripples predict seizure risk in childhood epilepsy better than spikes. *Brain*. 2019;142(5):1296–309.

84. Zhang J, Yang H, Wu D, Yang C, Yang Y, Zhou W, et al. Electroencephalographic abnormalities are correlated with cognitive deficits in children with benign childhood epilepsy with centrotemporal spikes: a clinical study of 61 cases. *Epilepsy Behav.* 2020;106:107012. <https://doi.org/10.1016/j.yebeh.2020.107012>
85. van 't Klooster MA, van Klink NEC, Zweiphenning WJEM, Leijten FSS, Zelmann R, Ferrier CH, et al. Tailoring epilepsy surgery with fast ripples in the intraoperative electrocorticogram. *Ann Neurol.* 2017;81(5):664–76.
86. Van Diessen E, Diederer SJH, Braun KPJ, Jansen FE, Stam CJ. Functional and structural brain networks in epilepsy: what have we learned? *Epilepsia.* 2013;54(11):1855–65.
87. Zweiphenning WJEM, Keijzer HM, van Diessen E, van 't Klooster MA, van Klink NEC, Leijten FSS, et al. Increased gamma and decreased fast ripple connections of epileptic tissue: a high-frequency directed network approach. *Epilepsia.* 2019;60(9):1908–20.
88. Zijlmans M, Jacobs J, Zelmann R, Debeau F, Gotman J. High-frequency oscillations mirror disease activity in patients with epilepsy. *Neurology.* 2009;72(11):979–86.
89. Kobayashi K, Akiyama T, Oka M, Endoh F, Yoshinaga H. A storm of fast (40–150Hz) oscillations hypsarhythmia in West syndrome. *Ann Neurol.* 2015;77(1):58–67.
90. Bernardo D, Nariai H, Hussain SA, Sankar R, Wu JY. Interictal scalp fast ripple occurrence and high frequency oscillation slow wave coupling in epileptic spasms. *Clin Neurophysiol.* 2020;131(7):1433–43. <https://doi.org/10.1016/j.clinph.2020.03.025>
91. Song I, Orosz I, Chervoneva I, Waldman ZJ, Fried I, Wu C, et al. Bimodal coupling of ripples and slower oscillations during sleep in patients with focal epilepsy. *Epilepsia.* 2017;58(11):1972–84.
92. Weiss SA, Song I, Leng M, Pastore T, Slezak D, Waldman Z, et al. Ripples have distinct spectral properties and phase-amplitude coupling with slow waves, but indistinct unit firing, in human epileptogenic hippocampus. *Front Neurol.* 2020;11:1–8.
93. Frauscher B, Von Ellenrieder N, Ferrari-Marinho T, Avoli M, Dubeau F, Gotman J. Facilitation of epileptic activity during sleep is mediated by high amplitude slow waves. *Brain.* 2015;138(6):1629–41.

How to cite this article: Santana-Gomez CE, Engel J Jr., Staba R. Drug-resistant epilepsy and the hypothesis of intrinsic severity: What about the high-frequency oscillations? *Epilepsia Open.* 2022;7(Suppl. 1):S59–S67. <https://doi.org/10.1002/epi4.12565>