

ORIGINAL RESEARCH:  
EMPIRICAL RESEARCH - MIXED METHODS

# Factors contributing to reported medication administration incidents in patients' homes – A text mining analysis

Marja Härkänen<sup>1</sup>   | Bryony Dean Franklin<sup>2,3</sup>   | Trevor Murrells<sup>4</sup>   |  
Anne Marie Rafferty<sup>4</sup>   | Katri Vehviläinen-Julkunen<sup>1,5</sup>  

<sup>1</sup>Department of Nursing Science, University of Eastern Finland, Kuopio, Finland

<sup>2</sup>Centre for Medication Safety and Service Quality, Imperial College London Healthcare NHS Trust, London, UK

<sup>3</sup>UCL School of Pharmacy, London, UK

<sup>4</sup>Florence Nightingale Faculty of Nursing, Midwifery & Palliative Care, King's College London, London, UK

<sup>5</sup>Kuopio University Hospital, Kuopio, Finland

## Correspondence

Marja Härkänen, Nursing Science, University of Eastern Finland Faculty of Health Sciences, Kuopio, Finland.

Email: marja.harkanen@uef.fi

## Funding information

Academy of Finland, Grant/Award Number: 310708 This work was financially supported by the post-doctoral research funding for the first author of this paper by the Academy of Finland. The second and the fourth authors are supported by the National Institute for Health Research (NIHR) Imperial Patient Safety Translational Research Centre, and the second author by the NIHR Health Protection Research Unit in Healthcare Associated Infections and Antimicrobial Resistance at Imperial College London, in partnership with Public Health England (PHE). The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, PHE or the Department of Health and Care.

## Abstract

**Aims:** To describe the characteristics of medication administration (MA) incidents reported to have occurred in patients' own homes (reporters' profession, incident types, contributing factors, patient consequence, and most common medications involved) and to identify the connection terms related to the most common contributing factors based on free text descriptions.

**Design:** A retrospective study using descriptive statistical analysis and text mining.

**Methods:** Medication administration incidents ( $N = 19,725$ ) reported to have occurred in patients' homes between 2013–2018 in one district in Finland were analysed, describing the data by the reporters' occupation, incident type, contributing factors, and patient consequence. SAS® Text Miner was used to analyse free text descriptions of the MA incidents to understand contributing factors, using concept linking.

**Results:** Most MA incidents were reported by practical (lower level) nurses (77.8%,  $N = 15,349$ ). The most common category of harm was 'mild harm' (40.1%,  $N = 7,915$ ) and the most common error type was omissions of drug doses (47.4%,  $N = 9,343$ ). The medications most commonly described were Marevan [warfarin] ( $N = 2,668$ ), insulin ( $N = 811$ ), Furesis [furosemide] ( $N = 590$ ), antibiotic ( $N = 446$ ), and Panadol [paracetamol] ( $N = 416$ ). The contributing factors most commonly reported were 'communication and flow of information' (25.5%,  $N = 5,038$ ), 'patient and relatives' (22.6%,  $N = 4,451$ ), 'practices' (9.9%,  $N = 1,959$ ), 'education and training' (4.8%,  $N = 949$ ), and 'work environment and resources' (3.0%,  $N = 598$ ).

**Conclusion:** There is need for effective communication and clear responsibilities between home care patients and their relatives and health providers, about MA and its challenges in home environments. Knowledge and skills relating to safe MA are also essential.

**Impact:** These findings about MA incidents that have occurred in patients' homes and have been reported by home care professionals demonstrate the need for medication safety improvement in home care.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. Journal of Advanced Nursing published by John Wiley & Sons Ltd

## KEYWORDS

home care, incident report, medication administration, nursing, text mining

## 1 | INTRODUCTION

Care provided by health professionals in patients' own homes (referred to here as 'home care') poses challenges in medication administration (MA), as patients mostly self-administer their own medicines (Olaniyan, Ghaleb, Dhillon, & Robinson, 2015). This is provided sometimes with the help of relatives or informal caregivers who do not have any formal education for MA and with the support of educated home care professionals.

Medication errors are common in the home context and can threaten patient safety resulting in morbidity and mortality (Berland & Bentsen, 2017). In this study, we use the term 'incident' to include near misses that are prevented before they reach the patient as well as errors that reach the patient, whether or not they cause harm.

### 1.1 | Background

Home care aims to enable patients to live at home for as long as possible (Turjamaa, Hartikainen, Kangasniemi, & Pietilä, 2014). This presents new challenges as a growing number of medically complex patients with multiple medications are receiving care in their own homes with the support of home care professionals together with informal caregivers. Many informal caregivers, such as family members, have limited skills for managing complex medications (ISMP, 2014). Based on a previous systematic review of carers' MA errors in the domiciliary setting (Parand, Garfield, Vincent, & Franklin, 2016), home MA errors made by carers are common and carers made similar errors to those made by professionals in other contexts. Home care professionals in Finland are usually trained practical (licensed, lower level) or Registered (higher level) Nurses. They practice independently in patients' homes where no other healthcare workers are accessible to consult with (Absulem & Hardin, 2011; Olaniyan et al., 2015).

The medication management process in home care can be challenging and complex with unclear boundaries of responsibilities and variable work conditions (Lindblad, Flink, & Ekstedt, 2017; Norri-Sederholm, Saranto, & Paakkonen, 2016). Communication failures are common (Absulem & Hardin, 2011; Berland & Bentsen, 2017; ISMP, 2014), as are medication reconciliation problems at hospital discharge (Kuusisto, Asikainen, & Saranto, 2014). Based on the findings of Hale et al. (2015), almost all home care patients (94%) have at least one medication discrepancy following hospital discharge based on comparison of the hospital discharge medication list and what the patient was taking at the first home healthcare visit. Almost half of these discrepancies are omissions (46%), where the patient is not taking medication as indicated on

their discharge medication list. In addition, lack of competence in home care professionals (Berland & Bentsen, 2017) can contribute to these incidents.

Home settings can be disorganized environments, with noise and other distractions. Older patients can be physiologically and psychologically frail, such that medication incidents can have a greater impact on their health (Absulem & Hardin, 2011). In addition, home care patients have risk factors for medication-related problems, including multiple co-morbidities, cognitive impairment, polypharmacy, and use of high-risk medicines (Elliot, Lee, Beanland, Vakil, & Goeman, 2016). Based on previous findings (Sears et al., 2018), patients' knowledge about their medications decreases as the number of medications increases. Therefore, these patients may be at high risk of medication errors, although it is often difficult to determine these contributing factors as most of the care provided in patients' homes is unobserved (Doran et al., 2013). To avoid home care MA incidents, there should be transparency, openness, and accurate reporting of incidents to identify risk factors associated with the process (Berland & Bentsen, 2017).

Nowadays, many healthcare organizations worldwide gather information on such incidents using incident reporting schemes (Härkänen, Vehviläinen-Julkunen, Franklin, Murrells, & Rafferty, 2020; Härkänen, Vehviläinen-Julkunen, Murrells, et al., 2020). However, only limited data are available about MA incidents that occur in patients' homes (Sears, Baker, Barnsley, & Short, 2013) and are reported by the home care staff responsible for those patients. The information in incident reports is typically both structured (e.g. categories for type of incident) and unstructured (e.g. free text descriptions). Free text information can include valuable information about contributing factors that may remain hidden if solely relying on structured information (Verma & Maiti, 2018). However, manual analysis of free text found in the incident reports is challenging using traditional qualitative text-based analysis methods, as the text can include considerable amounts of extraneous information and incident report datasets can also be much larger than those normally analysed manually or using qualitative software. Thus, this study aimed to use text mining for analysing home care MA incidents.

## 2 | THE STUDY

### 2.1 | Aim

The aim of this study was to describe the characteristics of MA incidents reported to have occurred in patients' own homes, including reporters' professions, incident types, contributing factors, patient consequences, and the most common medications involved. In addition, we aimed to identify the terms that are closely connected

to the most common contributing factors based on free text descriptions.

## 2.2 | Design and setting

We conducted a retrospective study of an existing dataset, using descriptive statistical analysis and text mining. The Home Care Services (domestic services and home nursing) unit of the Helsinki Health Centre organizes nursing, care, and the necessary support services to maintain the health, functionality, and offer care in cases of illness or disability of older people, convalescents, patients suffering from chronic illnesses, and disabled people over the age of 18. Home nursing entails nursing and rehabilitation services prescribed by a doctor, taking place at the patients' home. A doctor's referral is required to receive home nursing services (City of Helsinki, 2020). Some of the home care visits are conducted remotely using two-way video and audio via a tablet computer. At the beginning of 2019, there were approximately 790 clients receiving remote care in Helsinki, with approximately 24,500 remote visits per month. These typically include reminder and monitoring of medication intake, nutrition and health monitoring, and supporting physical activity.

Most of the home care nurses are licenced practical nurses (upper secondary level qualification at vocational school, 180 ECTS credits); few are Registered Nurses (higher education at the University of Applied Sciences, 210–270 ECTS credits). Both nurse groups take care of basic MA, but intravenous administration is usually the responsibility of Registered Nurses, who receive greater training in their use. In Finland, nurses need to pass medication tests at the beginning of their employment and then every third or fifth year (depending on their employer) to be allowed to administer medication. Nurses need written permission from their employer to administer more complex medications after their competence has been verified, for example practical nurses need to have permission for administration of injections and Registered Nurses for administration of intravenous medications (Valvira, 2020).

## 2.3 | Sample

The HaiPro incident reporting system is used for reporting patient safety incidents in Finland; it is anonymous, voluntary, computerized, and is used in over 200 social service and healthcare organizations (HaiPro, 2020). The data in this study comprised MA incidents ( $N = 19,725$ ) reported to HaiPro as having occurred in patients' own homes between 2013–2018 in Helsinki, including where patients lived in sheltered housing.

## 2.4 | Data collection

The license for use of the HaiPro data was obtained from the City of Helsinki. IT company Awanic Company (Ltd) provided the data for analysis.

## 2.5 | Data analysis

The incident report data included both structured information and free text. For this study, pre-specified structured categories analysed included: Reporters of incidents (Practical nurses, Registered nurses, Supervisors/directors, Public health nurses, Students, Social workers, Physicians, Other, and Missing data. Reporters of incidents may or may not be the same as the person involved with an incident); Consequences of incidents (No Harm, Mild harm, Moderate harm, Serious harm, and Not known. Consequences are described by the reporter in free text and later classified by the handler of the incident; this describes the perceived severity of the consequences that actually occurred to the patient); Type of incidents (Omission, Patient did not take the drug, Wrong timing, Wrong dose, Wrong patient, Wrong drug, Expired drug administered, Wrong administration technique, Other, Not known, or missing. The type of incident is classified by the reporter and later verified by the handler); and Contributing factors (Communication and flow of information, Patient and relatives, Practices, Education and training, Work environment and resources, Team, Medications, Equipment and supplies, Organization and leadership, Not known, or no identified contributing factors. Contributing factors are described by the reporter in free text and later classified by the handler, multiple categories are possible). Examples of the contributing factors can be found in online file 1. First, this structured information was described using IBM SPSS statistics 25 (frequencies & percentages).

Second, we divided incident reports into samples based on pre-defined contributing factors and chose incident reports that had the most common contributing factors (Communication and flow of information, Patient and relatives, Practices, Education and training, Work environment, and resources). These samples of incident reports' free text descriptions (what happened) were read by the first author of this paper and the common terms (keywords) for describing these incidents were identified and manually recorded until new terms were no longer required. Finally, the research group agreed these selected terms.

Then, SAS® Enterprise Miner 13.2 and its Text Miner tool were used to analyse the free text descriptions across the whole dataset. 'Text parsing' (processing unstructured text to a structured form, including tokenization, stemming, and part-of-text tagging) and 'text filtering' (reducing the total number of parsed terms and check the spellings) were conducted automatically using SAS Text Miner. The method is described more detail in our previous studies (Härkänen, Paananen, Murrells, Rafferty, & Franklin, 2019; Härkänen, Vehviläinen-Julkunen, Murrells, et al., 2020; Härkänen, Vehviläinen-Julkunen, Murrells, Rafferty, & Paananen, 2019). The Finnish language was chosen for parsing and filtering the text. A SAS Text Miner stop list was used to ignore some parts of the text such as auxiliary verbs, conjunctions, possessive pronoun, interjections, numbers, participles, and prepositions. Using an interactive filter viewer, synonyms were combined manually. Unwanted terms (such as most abbreviations) were excluded, as well as terms occurring in

fewer than in 10 reports. The medications most commonly mentioned in the free text descriptions were identified using the interactive filter viewer. A screenshot of the interactive filter viewer page can be found in online material 2.

A specific focus of this study was to explore connection terms related to the most common contributing factors; thus, the number of each of previously (manually) identified keywords describing contributing factors was determined using SAS Text Miner and its interactive filter viewer. The concept linking method was then used for identifying terms that are highly connected (connection terms) with these selected terms. In concept linking, the selected term is shown at the centre of a link diagram and the terms that circle this are those that occur together most often with that central term (Härkänen, Paananen, et al., 2019; SAS, 2012). The strength of association between terms in a corpus of documents is calculated using the binomial distribution (SAS, 2020). An example of concept linking with expanded links for the term 'Marevan' can be found in online material 3 (Finnish language). Finally, verbatim quotes were used to illustrate these findings (Figure 1).

## 2.6 | Validity and reliability/rigour


The text mining approach allowed for analysis of a large dataset including about 19,000 free text descriptions that would be difficult to analyse manually. Text mining applications' algorithms were

effective for identifying the concept links between terms. The credibility of text mining has already been recognized and tested (Verma & Maiti, 2018). Its accuracy, sensitivity, and specificity have been shown to be high when compared with manual analysis (Ruud, Johnson, Liesinger, Grafft, & Naessens, 2010), which was also confirmed in our previous study comparing automated concept linking and manual analysis (Härkänen, Vehviläinen-Julkunen, Murrells, et al., 2020).

Data analyses were discussed within the research group to ensure methodological coherence, adequate sampling, and responsiveness. The first author of this paper conducted the analyses independently, but the other authors of this paper critically reviewed the findings and consensus was reached on the themes.

## 2.7 | Ethical considerations

According to the guidelines of the Finnish National Advisory Board on Research Ethic (TENK, 2019, 62), the use of existing anonymous (register) data does not require an approval from the National Committee of Research Ethics. Permission to access the register of incident reports was granted from the relevant hospital district in 2019. Incident reports were anonymous; thus, anonymity of the reporters, patients, and other involved persons could be guaranteed. All data handling was conducted following the 'responsible conduct of research' (TENK, 2012).



**FIGURE 1** Analysis process of home care medication administration incident reports' free text descriptions [Colour figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

### 3 | RESULTS

#### 3.1 | Characteristics of MA incidents in home care

The reporters of the home care MA incidents were mostly (77.8%,  $N = 15,349$ ) practical nurses, with 14.9% ( $N = 2,935$ ) Registered Nurses and only three incidents reported by physicians. Of these incidents, 40.1% ( $N = 7,915$ ) caused mild harm to the patient based on the handler's evaluation. Moderate harm was caused in 5.3% ( $N = 1,047$ ) and serious harm in 0.1% ( $N = 22$ ). For 27.3% ( $N = 5,387$ ) of incidents, it was evaluated that 'no harm' resulted.

The most common error types based on the pre-specified categories were omissions of drug doses (47.4%,  $N = 9,343$ ) and 'patient did not take the drug' (29.6%,  $N = 5,843$ ), followed by wrong time errors (6.1%,  $N = 1,202$ ). Wrong patient incidents (drug administered to wrong patient) were the rarest, occurring in only 0.6% ( $N = 128$ ) of incidents (Table 1).

The most commonly reported contributing factor based on the pre-specified categories was 'communication and flow of information', reported in 25.5% ( $N = 5,038$ ) of incidents. Other common reported contributing factors were 'patient and relatives' (22.6% of incidents,  $N = 4,451$ ), followed by 'practices' (9.9%,  $N = 1,959$ ), 'education and training' (4.8%,  $N = 949$ ), and 'work environment and resources' (3.0%,  $N = 598$ ) (Table 1). The most common medications described in the free text descriptions were Marevan [warfarin] ( $N = 2,668$ ), insulin ( $N = 811$ ), Furesis [furosemide] ( $N = 590$ ), 'antibiotic' ( $N = 446$ ), and Panadol [paracetamol] ( $N = 416$ ) (Table 2).

#### 3.2 | Connection terms related to the most common contributing factors

##### 3.2.1 | Communication and flow of information

The most common terms (identified manually and counted by text mining) as being related to the contributing factor 'communication and flow of information' were: information ( $N = 2054$ ), tell ( $N = 1859$ ), call ( $N = 1,201$ ), and say ( $N = 1,080$ ) (Table 3). Other terms and connection terms can be found in online material 4.

Connection terms identified using text mining for the term 'information' were: 'anomaly, find, occurrence, information about when, precise information, factor, contributing factors':

"Antibiotic eyedrops have been prescribed to the resident starting on 21 May 2018. Due to an information breakdown, the nurses did not receive the information in time and the drops were not given. The error was detected on 28 May 2018 and the eyedrops were ordered from a pharmacy... Information breakdown among nursing staff."

[Incident report (IR) 123635].

**TABLE 1** Home care medication administration incidents' ( $N = 19,725$ ) pre-specified categories about reporters, severity, type, and contributing factors

	Home care medication administration incidents (No., % of all incidents)
<b>Reporters of incidents</b>	
Practical nurses	15,349 (77.8)
Registered Nurses	2,935 (14.9)
Supervisors	170 (0.9)
Public health nurses	359 (1.8)
Students	227 (1.2)
Social workers	62 (0.3)
Physicians	3 (0)
Other	185 (0.9)
Missing data	435 (2.2)
Total	19,725 (100)
<b>Consequences of incidents</b>	
No Harm	5,387 (27.3)
Mild harm	7,915 (40.1)
Moderate harm	1,047 (5.3)
Serious harm	22 (0.1)
Not known	5,354 (27.1)
Total	19 725 (100)
<b>Type of incidents</b>	
Omission	9,343 (47.4)
Patient did not take the drug	5,843 (29.6)
Other	1,685 (8.5)
Wrong timing	1,202 (6.1)
Wrong dose	804 (4.1)
Wrong patient	128 (0.6)
Wrong drug	182 (0.9)
expired drug administered	152 (0.8)
Wrong administration technique	189 (1.0)
Not known/missing	197 (1.0)
Total	19 725 (100)
<b>Contributing factors<sup>a</sup></b>	
Not known/no identified contributing factors	6,508 (33.0)
Communication and flow of information	5,038 (25.5)
Patient and relatives	4,451 (22.6)
Practices	1,959 (9.9)
Education and training	949 (4.8)
Work environment and resources	598 (3.0)
Team	234 (1.1)
Medications	56 (0.2)
Equipment and supplies	15 (0)
Organization and leadership	4 (0)
Total	19 725 (100)

<sup>a</sup>classified based on the first contributing factor, if many factors were mentioned.

**TABLE 2** Twenty of the most commonly mentioned medications in the free text descriptions of incidents ( $N = 19,725$ )

	freq	docs <sup>a</sup>
Marevan [warfarin]	2,668	1,334
insulin	811	341
Furesis [furosemide]	590	435
'Antibiotic'	446	227
Panadol [paracetamol]	416	315
Norspan [buprenorphine]	272	195
Thyroxin [levothyroxine]	229	166
Exelon [rivastigmine]	228	153
Mirtazapin [mirtazapine]	222	182
Para-tabs [paracetamol]	219	185
Novorapid [aspart insulin]	218	135
Klexane [enoxaparin]	198	90
Calcichew [calcium carbonate]	183	165
Bisoprolol [bisoprolol]	174	157
Lantus [glargine insulin]	167	97
Risperidon [risperidone]	161	131
Madopar [levodopa/benserazide hydrochloride]	152	125
Lyrica [pregabalin]	126	108
Kaleorid [potassium chloride]	126	102
Ketipinor [quetiapine]	119	96

<sup>a</sup>a drug might get mentioned more than once in the same document

Connection terms for the term 'tell' were: 'visit, home care, evening, contributing factors, take, anyway, issue, client, also':

"The resident was not given his/her daytime medications as there was an interruption in the information flow. The person working the morning shift failed to **tell** that he/she had not given the client his/her daytime medications. The person presumed that the evening shift worker was responsible for giving the client daytime medications. The worker was unsure that the team had agreed that the morning shift worker was responsible for giving the client daytime medications..."

[IR 132278].

Connection terms for the term 'call' were: 'morning, day, blood glucose, nurse, visit, home care, family member, issue':

"The virtual nurse calls clients in the evenings and makes sure that the clients take their medications. The entries say that the client has taken his/her evening medications, but the medications were actually still in a sachet on a table."

[IR 124122].

**TABLE 3** Most common keywords related to contributing factors

Keywords in English	Original keywords in Finnish	Freq	Docs
'Communication and flow of information'			
Information	tieto	2,054	1,885
Tell	kertoa	1,859	1,587
Call	soittaa	1,201	1,020
Say	sanoa	1,080	970
Read	lukea	965	843
Consult	konsultoida	505	460
'Patient and relatives'			
Refuse	kieltäytyä	420	370
Forgetful	muistamaton	232	206
Memory-disordered	muistisairas	272	218
Alcohol	alkoholi	139	119
Tired	väsynyt	213	206
Spouse	puoliso	357	215
Daughter	tytär	310	178
Practices			
Guide, instructions	ohje	980	820
Treatment plan	hoitosuunnitelma	793	666
Under supervision	valvotusti	764	706
Virtual care	virtuaalihoito	248	190
Remote care	etähoito	154	92
Education and training			
Temporary employee (fill-in)	sijainen	613	482
Know	tietää	381	360
Know-how/Can	osata	220	217
Deputy	keikkalainen	171	147
Student	opiskelija	131	82
Work environment			
Hurry, rush	kiire	523	543
Negligence	huolimattomuus	327	324
Excessive	liian	236	213
Much	paljon	247	232

Connection terms for the term 'say' were: 'tell, know, no, part, ask, call, take, issue':

"The client had not taken the previous nights' medication at 22 (Ketipinor [quetiapine] 25 mg). The medication was in a cup. The client was unable to say why he/she had not taken the medication and regretted always forgetting it. The client uses virtual remote care."

[IR 115676].



### 3.2.2 | Patient and relatives

Most common terms identified as being related to the contributing factor 'Patient' were; refuse ( $N = 420$ ), memory-disordered ( $N = 272$ ), and forgetful ( $N = 232$ ), and terms related to substance use such as alcohol ( $N = 139$ ). Most common terms related to 'relatives' were relative ( $N = 494$ ), spouse ( $N = 357$ ), and daughter ( $N = 310$ ) (Table 3). Other terms and connection terms can be found in online material 5.

Connection terms by concept linking for the term 'refuse' were 'negative towards care, persuade, many, agree, offer, try to take'. Identified similar terms were 'negative towards care' and 'negative towards medications':

"The client often neglects taking medications. The client is often aggressive and negative towards treatment and gets angry if you mention that he/she should take his/her medications..."

[IR 131601].

Connection terms for the term 'memory-disordered' were 'live, remember, spouse, client, well, home care, take', and for 'forgetful' were 'negative towards medications, visit, client, still, home care, well, take, leaving':

"The client has failed to take his/her medications [negative towards medications] put in a cup for today and yesterday. The client is forgetful and unable to take care of his/her medication. The client's closest family member also has memory problems."

[IR 112567].

Terms explaining substance use of patients were 'alcohol', 'intoxicated', and 'drunkenness'. Connection terms for the term 'alcohol' were 'drunkenness, use, effect, enjoy, drink, under influence, use':

"The client is considerably intoxicated and has failed to take his/her medications... The client passed out and, on the afternoon, after sobering up a bit, took his/her medications again."

[IR 132064].

Most common terms explaining relatives were 'relative', 'spouse', and 'daughter'. Connected terms for these explained their roles as caregivers, such as connection terms for the term 'relative' were 'told, agree, client, relative, place, inform, call, home care' (online material 2):

"The wife had put on one of her own Norspan [buprenorphine] 10 mikrog/h patches on her husband even though the husband is not under that medication. The wife had informed the nurse of this issue after the nurse had noticed the patch on the husband's arm. The wife said that the patch had reduced

the client's use of Opamox [benzodiazepine]. The patch caused no obvious harm to the client. The client has Alzheimer's disease, while the wife has no memory disorder."

[IR 110642].

### 3.2.3 | Practices

Most common terms identified as being related to the contributing factor 'Practices' were guide, instructions ( $N = 980$ ), treatment plan ( $N = 793$ ), under supervision ( $N = 764$ ), virtual care ( $N = 248$ ), and remote care ( $N = 154$ ) (Table 3). Other terms and connection terms can be found in online material 6. Connection terms by concept linking for 'guide, instructions' were 'next, day, blood glucose, keep, nurse, thing, also, Marevan [warfarin]':

"During a morning visit, the client's blood glucose levels revealed that the client had been given an extra medicine (Novorapid [insulin] 2 IU [international units]) even though his/her blood glucose levels did not indicate that this was necessary. The list of medications and treatment plan included clear instructions by a physician for mealtime insulin..."

[IR 125006].

Connection terms for the term 'treatment plan' were 'medication list, read, controlled, keep, nurse, visit, contributing factors, also':

"The client should have been given eyedrops in both eyes in the morning. Pred Forte [prednisolone acetate] and Oftagel [carbomerum]. The treatment plan had not been brought up to date."

[IR 127175].

Connection terms for 'under supervision' were 'table, visit, cup, treatment plan, read, morning medicine, give':

"A nurse gives the client medications each morning from dose sachets and a pill dispenser under supervision. The medications are kept in a locked medication kit at the client's home. During my visit this morning, I noticed that the client's Marevan [warfarin] for the previous evening was still in the pill dispenser, which means that the nurse had failed to give it when visiting the client in the morning...It seems that the nurse had not read the list of medications or treatment plan."

[IR 124173].

Connection terms for the term 'virtual care' were 'call, sample, evening, client, remind, call, take'. Connection terms for the term 'remote care' were quite similar to those for virtual care:

"The remote care service was not used. The client has diabetes and the purpose of the remote visit is to check how the client is doing. However, the remote care services did not contact the client during the entire evening. The remote care services also failed to inform the home care provider of the failure to contact the client. The client also noted that this was not the first time when the remote care services had failed to contact him/her..."

[IR 124599].

### 3.2.4 | Education and training

The most common terms identified as being related to the contributing factor 'Education and training' were: temporary employee ( $N = 613$ ), know ( $N = 381$ ), know-how/can ( $N = 220$ ), deputy ( $N = 171$ ), and student ( $N = 131$ ) (Table 3). Other terms and connection terms can be found in online material 7.

Connection terms by concept linking for the term 'temporary employee' were 'do, evening shift, work shift, treatment plan, permanent (staff), visit'. Some of the new employees and temporary staff members were not permitted to administer medications causing medication to be omitted:

"Forgot to put in a vaginal suppository, no knowledge of the consequences to the client. A temporary employee had been working the shift for a short time, was unaware of the issue."

[IR 133978].

"On Friday morning, it was observed that the client had not been given an Innohep [tinzaparin] injection. A public health nurse was called and consulted to decide what to do. The nurse visiting the client on Thursday had no medication permit and did not give the client the injection because of this."

[IR 105928].

Some of the terms described lack of knowledge, skills, and lack of experience ('know-how', 'can', and 'inexperience') by health professionals:

"Home care services inject the client with long-acting insulin every morning. The client uses safety needles in his/her insulin pen. The nurse did not know how to use the needle/ensure that the needle is used correctly, as a result of which the client was probably not given the day's insulin dose."

[IR 102118].

Connected terms to keyword 'student' were 'injury, insulin, "put in a cup", instantly, instructor, screen, inject, nurse':

"A student gave the client eyedrops that are meant to be given in the evenings. The nurse noticed that the student had done this. A registered nurse was informed about this and it was agreed that Xalatan [latanoprost] drops would not be given in the evening. A situation involving student supervision."

[IR 108116].

### 3.2.5 | Work environment and resources

The most common terms identified as being related to the contributing factor 'Work environment and resources' were hurry, rush ( $N = 523$ ), negligence ( $N = 327$ ), much ( $N = 247$ ), and excessive ( $N = 236$ ) (Table 3). Other terms and connection terms can be found in online material 8.

Connection terms by concept linking for 'hurry, rush' were 'tiredness, maker, a lot, error, nurse, busy nurse, negligence, contributing'. Connection terms for the term 'busy' were 'day, new, go, contribute, other, client, situation, morning shift, evening shift':

"Simvastatin 10 mg, in the client's pill dispenser, not given to patient. Shift with excessive workload. The nurse who made the mistake is typically careful and diligent. Things happen when you are busy."

[IR 63560].

Connection terms for 'negligence' were 'fuss, error, maker, nurse negligence, nurse, rush, contributing':

"During an evening visit, I forgot to give the client Klexane [dalteparin] 40 mg. Negligence, stressful evening, high stress level, prone to errors"

[IR 114609].

Connection terms for 'excessive' were 'reach, a lot, inject, get, short, too little, late' and for 'much' were 'work, also, how, goods, visit, too, time, rush':

"The nurse working the morning shift had so many visits between 8 and 10 a.m. that this client could only be visited so late as to get his/her morning medications at 10:30 a.m. Excessive workload in the morning"

[IR 130510].

## 4 | DISCUSSION

The aim of this study was to describe the characteristics of home care MA incidents and analyse the factors most commonly reported as contributing factors, based on the free text descriptions. The number of 'no harm' situations was relatively low in this dataset, representing only a third of all reported incidents, with other incidents



causing at least mild harm. The corresponding number of 'no harm' MA incidents was over 80% in over 500,000 acute care MA incidents reported within 10 years in England and Wales (Härkänen, Vehviläinen-Julkunen, Franklin, et al., 2020). Based on previous analysis of medication incidents that occurred at home in Canada, 37% resulted in harm to the patient (ISMP, 2014). Our finding, therefore, raises the questions as to whether home care incidents are more likely to cause patient harm or whether this difference can be explained by differences in reporting cultures. Based on previous findings, it has been estimated that self-reporting systems detect only 7–15% of all medication incidents (Elliott et al., 2018), but the actual percentage may be even lower.

Reporting of home care incidents is still a relatively new practice in Finland, which may result in near miss situations being even less likely to be reported. It may also be more difficult to identify incidents in the home care setting. Additionally, there may be differences in evaluating the reported severity, which should relate to the actual harm resulting directly from the incident rather than perceived potential harm (Härkänen, Vehviläinen-Julkunen, Franklin, et al., 2020).

The most common MA error types in this study were 'omissions of drugs' representing almost half of the incidents and 'patient did not take the drug' representing a third of incidents. Interestingly, omission cases were divided into these two groups, probably for the purposes of highlighting the patient's role in the case of the latter. Findings are similar to Hale et al. (2015), who found that almost half of medication discrepancies in home care were omissions. Patients in home care mostly self-administer their own medicines; thus, patients' coping with medication management should be encouraged in accordance with individual patient preferences (Bucknall et al., 2019). One of the key challenges in home care is how to bridge unclear boundaries of responsibility in the patient's home and maintain a safe medication process, while at the same time preserving the patient's autonomy and integrity (Lindblad et al., 2017). This means clear distribution of MA responsibilities among health professionals, patients, and families and understanding that not all patients are interested in taking more control (Bucknall et al., 2019).

'Communication and flow of information' was the most common contributing factor for MA home care incidents, as is also the case in acute care incidents (Syyrilä, Vehviläinen-Julkunen, & Härkänen, 2020). We found communication challenges among health professionals, and between health professionals, patients, and family members. Other studies have highlighted insufficient exchange of information and poor communication between home care health services and professionals (Absulem & Hardin, 2011; Berland & Bentsen, 2017). Suitable means of communication among home care professionals is required to ensure safe care for home care patients.

Our study demonstrated that patients who decline to take medications, people with memory disorders, and substance abuse were associated with MA incidents, such as omission of drugs. Other research found that patients who are visually impaired or who live alone with cognitive impairment, the presence of polypharmacy, and

the lack of an individual caregiver who can assume the professional role for MA are also risk areas (HPNA, 2011). An obvious contributing factor is the lack of around-the-clock supervision by a professional at home. Home care professionals have previously reported that patients may not have taken their medication because it was felt to be burdensome or not needed based on the patient's or his/her caregiver's opinion (HPNA, 2011). Based on our findings, even caregivers and family members have very important roles in home care patients' everyday coping, they were sometimes sources of MA incidents. Thus, patients and family caregivers need to have competence for safe MA and they should be active partners in the medication process for resilient home care (Lindblad et al., 2017). It is also important to understand that informal caregivers might experience a considerable burden associated with managing medications, a task which they may not be equipped to do.

Lack of health professional competence (Berland & Bentsen, 2017) can also contribute to home care MA incidents. Our data suggested that temporary staff were not always well inducted into the workplace. Some new employees did not have the required qualification for MA, contributing to medication errors. Findings related to work environmental issues, such as rushing because of excessive workload or lack of (competent) employees, were found in our free text descriptions. Similar issues have been found in other studies concerning MA incidents demonstrating that inadequate staffing levels, workload, and working in haste (Härkänen, Vehviläinen-Julkunen, Murrells, et al., 2020), as well as nurse workload or distractions during medication tasks (Sessions, Nemeth, Catchpole, & Kelechi, 2019) may increase the risk of omissions and other types of errors.

## 4.1 | Limitations

Our analysis required the researchers to make some subjective decisions, such as identifying and selecting the keywords for analysis. It is possible that some have been missed as there is no previous guidance to follow. Verbatim quotes were translated from Finnish to English in this paper, thus, because of the difference between linguistic expressions, some connection terms might be expressed slightly differently in these examples than in original expression in Finnish. In addition, incident report data were not originally meant for research purposes and thus poses several limitations, as incidents are likely to be under-reported which may introduce bias (Härkänen, Vehviläinen-Julkunen, Murrells, et al., 2020). The quality of the reports may also vary in terms of detail and accuracy (NHS, 2014).

## 5 | CONCLUSIONS

Challenges in home care MA process are manifold and reported MA incidents seem to be more serious compared with acute care settings, although this may reflect lack of reporting of near misses.

There is need for effective communication between patients and home care professionals to guarantee timely information of the current state of MA fulfilment and its challenges. Complex roles and responsibilities among health professionals, patients, and their family members exist in the MA process in home care. However, it might not be possible to control all patients changing preferences around taking their medications to ensure practice is safe; we may need to accept some inherent risks in medication management in patients' homes to preserve their autonomy and integrity.

## Peer Review

The peer review history for this article is available at <https://publons.com/publon/10.1111/jan.14532>.

## ORCID

Marja Härkänen  <https://orcid.org/0000-0001-8542-8193>  
 Bryony Dean Franklin  <https://orcid.org/0000-0002-2892-1245>  
 Trevor Murrells  <https://orcid.org/0000-0001-6716-0170>  
 Anne Marie Rafferty  <https://orcid.org/0000-0003-1010-6581>  
 Katri Vehviläinen-Julkunen  <https://orcid.org/0000-0002-6828-8261>

## TWITTER

Marja Härkänen  @MHarkanen  
 Bryony Dean Franklin  @BryonyDF  
 Trevor Murrells  @TJMurrells  
 Anne Marie Rafferty  @annemarieraffer  
 Katri Vehviläinen-Julkunen  @KVehvilainen

## REFERENCES

- Absulem, S., & Hardin, H. (2011). Home health nurses' perceived care errors. *Rehabilitation Nursing, 36*(3), 98–105. <https://doi.org/10.1002/j.2048-7940.2011.tb00073.x>
- Berland, A., & Bentsen, S. B. (2017). Medication errors in home care: A qualitative focus group study. *Journal of Clinical Nursing, 26*(21–22), 3734–3741. <https://doi.org/10.1111/jocn.13745>
- Bucknall, T., Digby, R., Fossum, M., Hutchinson, A. M., Considine, J., Dunning, T., ... Manias, E. (2019). Exploring patient preferences for involvement in medication management in hospitals. *Journal of Advanced Nursing, 75*(10), 2189–2199. <https://doi.org/10.1111/jan.14087>
- City of Helsinki (2020). Home Care. Retrieved from <https://www.hel.fi/helsinki/en/social-health/elderly/home/>
- Doran, D. M., Hirdes, J. P., Blais, R., Baker, G. R., Poss, J. W., Li, X., ... Jantzi, M. (2013). Adverse events among Ontario home care clients associated with emergency room visit or hospitalization: A retrospective cohort study. *BMC Health Services Research, 22*(13), 227. <https://doi.org/10.1186/1472-6963-13-227>
- Elliott, R. A., Camacho, E., Campbell, F., Jankovic, D., St James, M. M., Kaltenthaler, E., ... Faria, R. (2018). Prevalence and economic burden of medication errors in the NHS in England. Rapid evidence synthesis and economic analysis of the prevalence and burden of medication error in the UK. Policy Research Unit in Economic Evaluation of Health & Care Interventions (EPRU). Retrieved from <http://www.epru.org.uk/wp-content/uploads/2020/03/medication-error-report-edited-27032020.pdf>
- Elliott, R. A., Lee, C. Y., Beanland, C., Vakil, K., & Goeman, D. (2016). Medicines management, medication errors and adverse medication events in older people referred to a community nursing service: A retrospective observation study. *Drugs Real World Outcomes, 3*, 13–24.
- NHS England. (2014). Patient Safety Alert. Improving medication error incident reporting and learning. Retrieved from <https://www.england.nhs.uk/wp-content/uploads/2014/03/psa-sup-info-med-error.pdf>
- HaiPro. (2020). Reporting System for Safety Incidents in Health Care Organizations. Retrieved from <http://awanic.com/haipro/eng/>
- Hale, J., Neal, E. B., Myers, A., Wright, K. H. S., Triplett, J., Brown, L. B., ... Mixon, A. S. (2015). Medication discrepancies and associated risk factors identified in home health patients. *Home Healthcare now, 33*(9), 493–499. <https://doi.org/10.1097/NHH.0000000000000290>
- Härkänen, M., Paananen, J., Murrells, T., Rafferty, A. M., & Franklin, B. D. (2019). Identifying risks areas related to medication administrations - Text mining analysis using free-text descriptions of incident reports. *BMC Health Services Research, 19*, 791. <https://doi.org/10.1186/s12913-019-4597-9>
- Härkänen, M., Vehviläinen-Julkunen, K., Franklin, B. D., Murrells, T., & Rafferty, A. M. (2020). Factors related to medication administration incidents in England and Wales between 2007 and 2016: A retrospective trend analysis. *Journal of Patient Safety*. March 11, 2020 Publish Ahead of Print. <https://doi.org/10.1097/PTS.0000000000000639>
- Härkänen, M., Vehviläinen-Julkunen, K., Murrells, T., Paananen, J., Franklin, B. D., & Rafferty, A. M. (2020). The Contribution of Staffing to Medication Administration Errors: A Text Mining Analysis of Incident Report Data. *Journal of Nursing Scholarship, 52*(1), 113–123. <https://doi.org/10.1111/jnu.12531>
- Härkänen, M., Vehviläinen-Julkunen, K., Murrells, T., Rafferty, A. M., & Paananen, J. (2019). Text mining method for studying medication administration incidents and nurse-staffing contributing factors – a pilot study. *Computers, Informatics, Nursing, 37*(7), 357–365. <https://doi.org/10.1097/CIN.0000000000000518>
- HPNA Position Statement (2011). Medication Error Reporting in the Home Setting. *Journal of Hospice & Palliative Nursing, 13*(5), 268–269. <https://doi.org/10.1097/NJH.0b013e31822a5b6e>
- ISMP Canada Safety Bulletin. (2014). Aggregate Analysis of Medication Incidents in Home Care. Retrieved from [https://www.ismp-canada.org/download/safetyBulletins/2014/ISMPCSB2014-8\\_MedicationIncidentsHomeCare.pdf](https://www.ismp-canada.org/download/safetyBulletins/2014/ISMPCSB2014-8_MedicationIncidentsHomeCare.pdf)
- Kuusisto, A., Asikainen, P., & Saranto, K. (2014). Medication documentation in nursing discharge summaries at patient discharge from special care to primary care. *Nursing & Care, 3*, 147. <https://doi.org/10.4172/2167-1168.1000147>
- Lindblad, M., Flink, M., & Ekstedt, M. (2017). Safe medication management in specialized home healthcare - an observational study. *BMC Health Services Research, 24*, 17(1), 598. <https://doi.org/10.1186/s12913-017-2556-x>
- Norri-Sederholm, T., Saranto, K., & Paakkonen, H. (2016). Medication information flow in home care. *Studies in Health Technology and Informatics, 225*, 359–363.
- Olanian, J. O., Ghaleb, M., Dhillon, S., & Robinson, P. (2015). Safety of medication use in primary care. *International Journal of Pharmacy Practice, 23*, 3–20. <https://doi.org/10.1111/ijpp.12120>
- Parand, A., Garfield, S., Vincent, C., & Franklin, B. D. (2016). Carers' Medication Administration Errors in the Domiciliary Setting: A Systematic Review. *PLoS One, 11*(12), e0167204. <https://doi.org/10.1371/journal.pone.0167204>
- Ruud, K. L., Johnson, M. G., Liesinger, J. T., Grafft, C. A., & Naessens, J. M. (2010). Automated detection of follow-up appointments using text mining of discharge records. *International Journal of Quality in Health Care, 22*(3), 229–235. <https://doi.org/10.1093/intqhc/mzq012>
- SAS. (2020). SAS® Text Miner 15.1: Strength of Association for Concept Linking. Retrieved from <http://documentation.sas.com/?docse>

- tld=tmref&docsetTarget=n0chdwsd64uafcn164shwz4793a2.htm&docsetVersion=15.1&locale=en
- SAS Institute Inc. (2012). *Getting Started with SAS® Text Miner 12.1*. Cary, NCSAS Institute Inc.
- Sears, K., Woo, K. Y., Almost, J., Wilson, R., Frymire, E., Whitehead, M., & Van Den Kerkhof, E. (2018). Medication knowledge among older adults admitted to home care in Ontario during 2012–2013. *Journal for Healthcare Quality*, 40(3), e33–e45. <https://doi.org/10.1097/JHQ.0000000000000055>
- Sears, N. A., Baker, G. R., Barnsley, J., & Short, S. (2013). The incidence of adverse events among home care patients. *International Journal of Quality in Health Care*, 1, 1–13. <https://doi.org/10.1093/intqhc/mzs075>
- Sessions, L. C., Nemeth, L. S., Catchpole, K., & Kelechi, T. J. (2019). Nurses' perceptions of high-alert medication administration safety: A qualitative descriptive study. *Journal of Advanced Nursing*, 75(12), 3654–3667. <https://doi.org/10.1111/jan.14173>
- Syrrilä, T., Vehviläinen-Julkunen, K., & Härkänen, M. (2020). Communication issues contributing to medication incidents: Mixed method analysis of hospitals' incident reports using indicator phrases based on literature. *Journal of Clinical Nursing*, 29(13–14), 2466–2481. <https://doi.org/10.1111/jocn.15263>
- TENK. (National Advisory Board on Research Ethics). (2012). Responsible conduct of research and procedures for handling allegations of misconduct in Finland. Retrieved from [https://www.tenk.fi/sites/tenk.fi/files/HTK\\_ohje\\_2012.pdf](https://www.tenk.fi/sites/tenk.fi/files/HTK_ohje_2012.pdf)
- TENK. (National Advisory Board on Research Ethics). (2019). The ethical principles of research with human participants and ethical review in the human sciences in Finland. Retrieved from [https://www.tenk.fi/sites/tenk.fi/files/lhmistieteiden\\_eettisen\\_ennakkoarviointi\\_in\\_ohje\\_2019.pdf](https://www.tenk.fi/sites/tenk.fi/files/lhmistieteiden_eettisen_ennakkoarviointi_in_ohje_2019.pdf)
- Turjamaa, R., Hartikainen, S., Kangasniemi, M., & Pietilä, A. M. (2014). Living longer at home: A qualitative study of older clients' and practical nurses' perceptions of home care. *Journal of Clinical Nursing*, 23, 3206–3217. <https://doi.org/10.1111/jocn.12569>
- Valvira. (National Supervisory Authority for Welfare and Health). (2020). Lääkehoidon toteuttaminen (in Finnish). Retrieved from [https://www.valvira.fi/terveydenhuolto/hyva-ammattinharjoittaminen/laakehoito/laakehoidon\\_toteuttaminen](https://www.valvira.fi/terveydenhuolto/hyva-ammattinharjoittaminen/laakehoito/laakehoidon_toteuttaminen)
- Verma, A., & Maiti, J. (2018). Text-document clustering-based cause and effect analysis methodology for steel plant incident data. *International Journal of Injury Control and Safety Promotion*, 7, 1–11. <https://doi.org/10.1080/17457300.2018.1456468>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

**How to cite this article:** Härkänen M, Franklin BD, Murrells T, Rafferty AM, Vehviläinen-Julkunen K. Factors contributing to reported medication administration incidents in patients' homes – A text mining analysis. *J Adv Nurs*. 2020;76:3573–3583. <https://doi.org/10.1111/jan.14532>

The *Journal of Advanced Nursing (JAN)* is an international, peer-reviewed, scientific journal. *JAN* contributes to the advancement of evidence-based nursing, midwifery and health care by disseminating high quality research and scholarship of contemporary relevance and with potential to advance knowledge for practice, education, management or policy. *JAN* publishes research reviews, original research reports and methodological and theoretical papers.

For further information, please visit *JAN* on the Wiley Online Library website: [www.wileyonlinelibrary.com/journal/jan](http://www.wileyonlinelibrary.com/journal/jan)

### Reasons to publish your work in *JAN*:

- **High-impact forum:** the world's most cited nursing journal, with an Impact Factor of 1.998 – ranked 12/114 in the 2016 ISI Journal Citation Reports © (Nursing (Social Science)).
- **Most read nursing journal in the world:** over 3 million articles downloaded online per year and accessible in over 10,000 libraries worldwide (including over 3,500 in developing countries with free or low cost access).
- **Fast and easy online submission:** online submission at <http://mc.manuscriptcentral.com/jan>.
- **Positive publishing experience:** rapid double-blind peer review with constructive feedback.
- **Rapid online publication in five weeks:** average time from final manuscript arriving in production to online publication.
- **Online Open:** the option to pay to make your article freely and openly accessible to non-subscribers upon publication on Wiley Online Library, as well as the option to deposit the article in your own or your funding agency's preferred archive (e.g. PubMed).