

Research

Open Access

Effect of long-term treatment with steroid hormones or tamoxifen on the progesterone receptor and androgen receptor in the endometrium of ovariectomized cynomolgus macaques

Hong Wang*¹, Erika Isaksson², Bo von Schoultz³, J Mark Cline⁴ and Lena Sahlin¹

Address: ¹Division for Reproductive Endocrinology, Karolinska Hospital, Stockholm, Sweden, ²Department of Oncology, Radiumhemmet, Karolinska Hospital, 17176 Stockholm, Sweden, ³Division for Obstetrics and Gynecology, Department of Woman and Child Health, Karolinska Hospital, Stockholm, Sweden and ⁴Department of Pathology, Section on Comparative Medicine, Wake Forest University School of Medicine, Winston-Salem, NC, USA

Email: Hong Wang* - Hong.Wang@cgb.ki.se; Erika Isaksson - erika.isaksson@ks.se; Bo von Schoultz - bo.vonschoultz@ks.se; J Mark Cline - Jmcline@wfubmc.edu; Lena Sahlin - lena.sahlin@kbh.ki.se

* Corresponding author

Published: 5 February 2003

Received: 27 January 2003

Reproductive Biology and Endocrinology 2003, 1:7

Accepted: 5 February 2003

This article is available from: <http://www.RBEj.com/content/1/1/7>

© 2003 Wang et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

The progesterone receptor (PR) and androgen receptor (AR) belong to the nuclear receptor superfamily. Two isoforms of PR (A and B) have been identified with different functions. The expression of AR, each isoform of PR and their involvement in long-term effects on the endometrium after hormonal replacement therapy (HRT) or tamoxifen (TAM) treatment is not known. The aims of this study were to determine PR(A+B), PRB and AR distribution by immunohistochemistry in the macaque (*Macaca fascicularis*) endometrium. Ovariectomized (OVX) animals were orally treated continuously for 35 months with either conjugated equine estrogens (CEE); medroxyprogesterone acetate (MPA); the combination of CEE/MPA; or TAM. Treatment with CEE/MPA tended to down-regulate PR in the superficial glands, but increased it in the stroma. TAM treatment increased both the PR and PRB levels in the stroma. Overall, less than 20% of the cells were positive for the PRB isoform and less variation was observed after steroid treatment. AR was found in the stroma, mainly distributed in the basal layer of the endometrium in the OVX and steroid treated groups, but was absent in the TAM treated group. No AR was found in the glandular epithelium. The present data show that long-term hormone treatment affects the PR level, and also the ratio between PRA and PRB in the endometrium.

Background

Endometrial carcinomas are one of the most common cancers in women. Hormone replacement therapy (HRT) is commonly used for reducing menopausal symptoms, but is associated with an increased risk of endometrial carcinoma [1,2]. Tamoxifen (TAM, estrogen antagonist in breast, agonist in uterus) is the most commonly used treatment for women with estrogen receptor (ER) positive

breast cancers but the risk of developing endometrial cancer is also increased in those patients [3]. Progesterone provides a protective effect in HRT by interrupting continued estrogenic stimulation of the endometrium [4]. The risk of endometrial cancer has been shown to be decreased both after contraceptive treatment with progestin alone, and after continuous-combined HRT [5,6].

However, the association between HRT or TAM treatment and endometrial cancer is still not clearly understood.

The effects of progesterone in humans are mediated by two specific intracellular progesterone receptor isoforms, PRA and PRB [7,8]. PRA is a truncated form of PRB, lacking 164 amino acids at the N-terminal region [9]. The PR isoforms have different biological activities at the transcriptional level [9–11], and the ratio of the receptor isoforms can influence the responsiveness of the tissue to progesterone signals [12,13]. Studies have shown that PRB is an effective activator of gene transcription [11,14], and PRA has the ability to suppress the activity of PRB as well as other steroid hormone receptor family members, such as AR, glucocorticoid receptor and mineralocorticoid receptor [11,14,15]. In addition, liganded PRA and PRB can each suppress E₂-stimulated ER activity, with the magnitude of repression dependent on the PR isoform, progesterin ligand, promoter, and cell type [16].

Androgens are generally believed to be hormones of the male, but androgens also have effects on the female reproductive system [17,18]. Studies have shown that the androgen receptor (AR) is present in the human and nonhuman primate endometrium, but the results of AR distribution and regulation are contradictory [19–23]. A recent study indicated that androgens can antagonize estradiol (E₂) action in the macaque and human endometrium, and that AR can play a role in the antiproliferative, "antiestrogenic" effects of antiprogestin in primates [24]. However, the involvement of PR isoforms and AR in the pathogenesis of endometrial cancer is still not clear. Female macaques are similar to human beings in many aspects of reproductive biology, including the menstrual cycle [25,26]. The aim of this study was to use the macaques as a model to compare the protein expression of PR, PRB and AR in the endometrium after long-term exposure to CEE (conjugated equine estrogens), MPA (medroxyprogesterone acetate), CEE+MPA or TAM, in order to clarify the possible roles of these receptors on the endometrium after long-term treatment with these compounds.

Material and Methods

Animals

The twenty-five female adult cynomolgus macaques (*Macaca fascicularis*) used in this study were imported from Indonesia to the United States. All the macaques were 4–6 years old at the beginning of the study. They were housed in social groups of 4–8 monkeys each in a facility. Bilateral ovariectomy (OVX) was performed three months before initiation of hormone treatment. Twenty monkeys were treated continuously with either 0.625 mg/day CEE (n = 4); 2.5 mg/day MPA (n = 5); the combination of CEE + MPA (n = 5); or 20 mg/day TAM (n = 6). Five

animals were without hormone treatment and used as OVX controls. The compounds were administered in the diet for 35 months and are expressed as doses equivalent, on a caloric basis, to those given to women [27]. The animal studies were approved by the association for the Advancement and Accreditation of Laboratory Animal Care and use committee in USA.

Tissue collection

Uterine tissues were collected at the end of the treatment period, when all the animals were euthanized. Tissues were trimmed from fat and connective tissues, and fixed in 4% paraformaldehyde for 24 hours and stored at 4 °C in 70% ethanol. Thereafter tissues were trimmed to 3 mm in thickness and embedded in paraffin.

Immunohistochemistry

Paraffin sections (5 µm) were used and a standard immunohistochemical technique (avidin-biotin-peroxidase) was carried out as described before [28] for immunolocalization of PR (A+B), PRB and AR. The different immunostaining procedures for each specific primary antibody are given in table 1. The samples were incubated with the primary monoclonal antibody, followed by incubation with the secondary biotinylated horse anti-mouse antibody (BA-2000, Vector Laboratory, Burlingame, CA, USA) and an avidin-biotin-peroxidase complex at room temperature. Replacement of the primary antibody with an equivalent concentration of non-immune mouse IgG was used for negative controls of PR, PRB and AR, respectively.

The peroxidase substrate diaminobenzidine (DAB) was used to visualize the immunostaining reaction (SK 4100; Vector Laboratory, Burlingame, CA, USA). Thereafter the procedure was as described before [28].

Image analysis and scoring

A Leica microscope and Sony video camera (Park Ridge, NJ, USA) connected to a computer with an image analysis software (Leica Imaging System Ltd. Cambridge, UK) were used to quantitate PR and PRB immunohistochemistry results, as described previously [28]. In short, by using colour discrimination software the total area of positively stained nuclei (brown) was measured, and expressed as a percentage of the total area of cell nuclei (brown + blue). Since AR immunostaining was weak it was not possible to measure by the image analysis software and therefore the immunostaining intensity and distribution of AR were assessed semiquantitatively by manual scoring on a two point scale: - = no staining, + = faint staining, by one observer blinded to treatment.

Statistics

Statistical calculations for the immunohistochemistry results were performed by ANOVA on ranks (Kruskal-Wallis

Table 1: Primary antibodies used in the study. Since a specific PRA antibody is not commercially available, the PRA level is difficult to measure directly by immunohistochemistry.

Antibody	[Specificity]	[Working condition]	[Sources]
PR (18-0172)	N-terminal of Human PR(A+B)	Diluted to 1:100 Incubated at RT, 60 min	Zymed Lab. Inc., San Francisco, CA
PRB (MA1-411)	The 120 kDa of Human PRB	Diluted to 1:50 Incubated at RT, 60 min	Affinity Bioreagents, Inc., Golden, CO
AR (NCL-AR-2F12)	N-terminal of Human AR	Diluted to 1:25 Incubated at 4°C, 18 h	Novacastra, England UK

test) and significances were evaluated by Dunn's test (all compared to all) or Dunnett's test (all compared to control). Values with different letter designations are significantly different ($p < 0.05$). Values with * are significantly different ($p < 0.05$) to OVX control.

Results

Immunohistochemistry showed that PR, PRB and AR were present in the nuclei of cells in the compartments of the endometrium (Fig. 1 and 2). The treatments induced changes of the histological features of the endometrium, e.g. TAM induced cystic endometrial hyperplasia (Fig. 1i,1j)

PR

Variations were found in PR immunostaining in the superficial endometrium, whereas the immunostaining pattern was less varying in the basal layer. The statistical analysis was therefore performed on image analyses of the superficial layer.

In the stromal cells, PR immunoreactivity was low in the OVX control group (Figure 1a). The immunostaining intensity and number of positive cells increased after treatments (Figure 1c,1e,1g and 1i). In addition, the stromal cells in the CEE/MPA group showed decidualization, and PR staining was intense (Fig. 1g). The image analysis showed an increased level in the CEE/MPA and TAM groups as compared to the OVX control (Figure 3a).

In the glandular and luminal epithelium, around 30–40% of the cells were positive in the OVX group (Figure 3b,3c). In the glands of the superficial layer, a faint PR immunoreactivity was observed in the MPA group (Figure 1e, and figure 3b). Only weak or no immunostaining was observed in glands of the CEE/MPA group (Figure 1g, black arrow and Figure 3b).

Intense PR immunostaining was observed after TAM treatment in both glandular and luminal epithelium (Fig. 1i and Fig. 3b,3c). No positive immunostaining was found in the negative controls (Fig. 1k).

PRB

PRB immunostaining was faint or absent in all compartments of the endometrium and in all studied groups (Fig. 1b,1d,1f,1h,1j). Intensive immunoreactivity of PRB was observed in the glands of human endometrium from the proliferative stage of the menstrual cycle (Fig. 1l) and also in the breast tissue of the CEE treated group of monkeys (Fig. 1m), which were used as positive controls.

No PRB immunostaining was found in the majority of stromal cells of the OVX and CEE groups (Figure 1b,1d and Figure 3d). Around 1–5% of the stroma was weakly stained for PRB in the MPA, CEE/MPA and TAM groups. Image analysis results showed an increase after TAM treatment as compared to the OVX group (Fig. 1j and Fig. 3d).

In the luminal epithelium, the PRB immunostaining was almost absent in the OVX control group. Around 10% of glandular and luminal epithelia were faintly stained after TAM treatment (Figure 1j and Figure 3e,3f), as well as in the glands of the CEE group (Figure 1d and Figure 3e). No significant differences were found between groups in the luminal epithelium and in the glands.

PRB/PR

In order to evaluate the relative level of PRA and PRB in the endometrium, the ratio of PRB/PR are shown in table 2. PRA is by far the most common isoform in macaque endometrium. The highest ratios of PRB/PR were found in the glands of the CEE group and in all compartments of the TAM group, suggesting an increased influence of PRB in these cell types as compared to the other compartments and treatment groups.

AR

No AR immunostaining was found in the glands in any of the studied groups. Positive immunostaining was observed in some of the stromal cells, mainly distributed in the basal layer of the endometrium in the OVX (Fig. 2a), MPA (Fig. 2b) and CEE/MPA (not shown) treated animals (Table 3). No staining was present in the endothelial and smooth muscle cells of the vessels (Figure 2b, arrow). The TAM treated animals showed no positive AR immunostaining (Figure 2c and Table 3). Sweat glands from the





Figure 1

Immunohistochemical localization of PR and PRB in the superficial endometrial stroma (Str), glands (G) and luminal epithelium (Le). Positive nuclear immunostaining for PR is present in the study groups as shown: OVX (a); CEE (c); MPA (e); CEE /MPA (g), and TAM (i). Nuclear PRB immunostaining is shown for: OVX (b); CEE (d); MPA (f); CEE /MPA (h) and TAM (j) groups, (bar = 30 μm). A negative control is shown for PR where the first antibody is replaced by mouse IgG (k). Positive controls for PRB immunostaining in human endometrium (l) and breast tissue from a CEE treated monkey (m), (bar = 30 μm).

**Figure 2**

AR immunostaining from the basal layer of the endometrium is shown in: OVX (a); MPA (b) and TAM (c) treated animals. As positive control for AR immunostaining sweat glands from monkey breast tissue is shown (d), (bar = 20 µm).

monkey breast tissue, which was used as positive control, showed an intense immunostaining for AR (Figure 2d).

Discussion

In vitro studies have shown that PRA and PRB isoforms can activate different target genes [9]. In vivo studies have demonstrated that the relative levels of PRA and PRB within endometrial cells may determine the nature and magnitude of their functional responses to progesterone and estrogen during the menstrual cycle [13]. The results from the present study show that long-term treatment with the combination of CEE and MPA to OVX macaques, increased the PR level in the stroma, but reduced PR immunoreactivity in the superficial glands. TAM, a partial estrogen agonist in the uterus, increased the PR levels in the stroma, with a similar tendency in glandular epithelium. CEE tended to increase the PR level in the stroma and luminal epithelium, but no significant difference was observed. PRB immunoreactivity was weak in all groups. Less than 20% of the endometrial cells were positively stained for PRB in almost all studied samples, however, TAM increased the stromal PRB level.

It has been demonstrated that estrogen increases, but progesterone reduces cell proliferation in the endometrium during the normal menstrual cycle [29], and also in long term treated OVX macaques [30]. Both PRA and PRB can be regulated by estrogen and progesterone in the endometrium, and the distinct differences in the relative expression of PRA and PRB may be associated with different roles for the isoforms in the different compartments of the uterus [13]. Pike et al. [6] have reported that the continuous-combined estrogen and progesterone replacement therapy is not associated with an increased risk of endometrial cancer. The antiproliferative activity of progesterone in the endometrium may be the cause of reduced

cancer risk. Although the mechanism of PRA mediated transdominant inhibition is still unknown, PRA can antagonize ER action [15] and has an ability to suppress the activity of PRB [11,14]. The ER has important interrelationships with the PR system in modulation of biological responses. In many estrogen target tissues, e.g. uterus, estrogen increases *c-fos* mRNA levels, cell proliferation, PR mRNA and protein levels, gap junction formation and myometrial contractility. These effects are largely antagonized by progesterone [31]. These findings give evidence of interactions and cross talk between members of the steroid hormone receptor family, which make the mechanism of the receptor activity even more complex.

It has been shown in humans that a relatively high level of PRA continued to be expressed in the stroma, but not in epithelial cells of the progestin dependent luteal phase [32]. It was suggested that the cell-specific effect of progestin indicated a different pathway in the regulation of PR isoforms in endometrial stromal cells [33]. In the present study, the up- and down-regulation of PR (mainly PRA) by MPA/CEE found in the stroma and epithelium, respectively, is in agreement with a previous study on the PR mRNA expression [33]. Taken together, the functional differences of PRA and PRB together with their different expression pattern in the different compartments of the endometrium suggest an important role for both isoforms in the cellular response to steroids. The relative level of PRA and PRB may determine the expression of specific genes upon hormonal stimulation. As for ER, PR function in the epithelium could be regulated by paracrine signals in the stroma, such as IGF-I and TGF- β 4 [34,35].

Studies have shown that endometrial cancers exhibit lower PR expression and a predominance of one PR isoform as compared with the epithelial cells of the normal




Figure 3

Image analysis scores of positive PR immunoreactivity in superficial stroma (a), superficial glands (b) and luminal epithelium (c). PRB immunostaining is shown in superficial stroma (d), superficial glands (e) and luminal epithelium (f). Box and whisker plots representing the median value with 50% of all data falling within the box. The whiskers extend to the 5th and 95th percentiles. In OVX, MPA and CEE/MPA groups: n = 5, in CEE group n = 4 and in the TAM group n = 6. Values with different letter designations are significantly different (p < 0.05). Values with * is significantly different (p < 0.05) to OVX control.

Table 2: Ratios of PRB/PR from the median values shown in Fig 2. No significant difference was observed. GE = glandular epithelium. LE = luminal epithelium.

Cell type	OVX	CEE	MPA	CEE/MPA	TAM
Superficial stroma	0.040	0.005	0.060	0.020	0.110
Superficial GE	0.030	0.195	0.050	0.000	0.130
LE	0.020	0.015	0.010	0.055	0.160

Table 3: Manual scoring data of AR immunostaining activity found in basal layer of endometrium.

Group	AR immunostaining		
	Stroma	Gland	Luminal epithelium
OVX (n = 5)	+, -, -, +, +.	-, -, -, -, -.	-, -, -, -, -.
CEE (n = 4)	+, -, -, -.	-, -, -, -.	-, -, -, -.
MPA (n = 5)	+, +, +, +, +.	-, -, -, -, -.	-, -, no, -, -.
CEE+MPA (n = 5)	+, +, +, +, -.	-, -, -, -, +.	-, no, -, no, -.
TAM (n = 6)	-, -, -, -, -, -.	+, -, -, -, -, -.	no, -, -, -, -, -.

(-) = no staining, (+) faint staining, no = no luminal epithelium in the sample.

endometrium, where the PR level was high and both PRA and PRB usually coexpressed [4]. These authors suggest that loss of coordinated expression of PR isoforms is an early event in the development of endometrial cancer, since loss of expression of one PR isoform (A or B) was observed in hyperplasia [4]. In the present study, TAM treatment induced cystic hyperplasia, but the PR level was increased in the glands and there was a higher ratio of PRB/ PR in this group. This suggests that the two PR isoforms coexist or that the ratio of PRB/PRA is relatively higher than in the other studied groups. However, PRA is still the predominant isoform in the epithelium as well as stroma in this group, since the PRB level is very low, which is partly consistent with the conclusion made by Arnett-Mansfield et al. [4].

There is a tendency to increase in stromal PR levels after CEE treatment in the present study, but not even a tendency can be seen for PRB. Both PR and PRB are increased by TAM treatment. The conclusion could be that CEE regulates only PRA, while TAM regulates both PR subtypes.

The mechanism of TAM-induced carcinogenesis is thought to be due to its estrogenic actions on the reproductive tract [36]. However, the mechanism that controls the relative level of PRA and PRB in endometrial cancer and how TAM induces cancer needs to be further clarified. Studies have shown that exogenous androgen can have inhibitory effects on the female reproductive system [17,18]. During the menstrual cycle, AR was mainly distributed in the endometrial stroma of human and nonhuman primates [23,24,37], and in occasional cells of the glandular epithelium, but was absent in the vascular wall [24]. Though we found faint immunostaining of AR in the macaque endometrium, the stromal distribution was similar to the previous report [24]. Slayden et al. [24] have implied that under normal cyclic conditions, stromal AR levels seem to be up-regulated by E₂ and suppressed by progesterone, which is contradictory to the impression of the present results on long term treatment. The AR-in-

duced effects of endogenous androgens in the endometrium may be mediated mainly via the AR receptor in the stroma, but not the epithelium. A most interesting finding by these authors was that AR expression is enhanced in the endometrial stroma and induced in the glandular epithelium by antiprogestin (RU486) treatment. They suggest that the increased AR could mediate androgen dependent antiestrogenic effects induced by antiprogestin [24]. In addition, we found no positive AR immunostaining in the TAM treated group, which suggests that TAM may down-regulate AR expression in the stroma.

We have shown that in the superficial stroma the ER α level was lower in the CEE/MPA group than in the CEE and MPA groups [38]. ER β immunostaining was faint with minor variations in response to treatments, but increased in the superficial stroma after MPA treatment. The ratio of ER β /ER α increased in superficial stroma and glands after CEE/MPA treatment, and also in stroma after MPA and TAM group. Cystic endometrial hyperplasia was observed in TAM treated animals, in combination with a high level of ER α protein expression. These data show that long-term hormone treatment also affects the ER α and ER β protein levels in the endometrium, where the balance between ER α and ER β seems to be important for the proliferative response to estrogen [38].

The results on ER α [38] showed that there are more cells positive for ER α than for PR, in all tissues we examined. Similarities in expression pattern between ER α [38] and PR are that in the CEE/MPA group both proteins are decreased in the epithelium of the superficial glands, whereas TAM treatment exhibits the highest levels of both ER α [38] and PR in the luminal epithelium. In the stroma the combined CEE/MPA treatment reduced the ER α level, while the PR level was increased [38].

The results of the present study indicate that PR is an important indicator of hormone responses in endometrium. The different expression of PRA and PRB might be related to the responses to hormone therapy and patient out-

come. The endometrial hyperplasia found after TAM treatment could be due to the different ratio of PRA to PRB in the endometrium, possibly in combination with the increased ER α dominance after TAM treatment [38]. Long-term treatment with steroids did not show a different distribution pattern of AR as compared to the OVX control, but AR was almost absent in the TAM group as well as in the CEE group. Further studies are needed to determine the importance of the interactions among the steroid hormone receptor members (e.g. AR, PRA and PRB, ER α and ER β) for proliferative activity in the endometrium and possible side effects of HRT and TAM therapy.

Acknowledgements

The present study received financial support from The Cancer Society in Stockholm, the Swedish Medical Research Council (grants 3972 and 5982), the Swedish Society for Medical Research and Karolinska Institutet.

References

- Grady D, Rubin SM, Petitti DB, Black D, Ettinger B, Ernster VL and Cummings SR **Hormone therapy to prevent disease and prolong life in postmenopausal women.** *Ann Intern Med* 1992, **117**:1016-1037
- Grady D, Gebretsadik T, Kerlikowske K, Ernster V and Petitti D **Hormone replacement therapy and endometrial cancer risk: a meta-analysis.** *Obstet Gynecol* 1995, **85**:304-313
- Malfetano JH **Tamoxifen-associated endometrial carcinoma in postmenopausal breast cancer patients.** *Gynecol Oncol* 1990, **39**:82-84
- Arnett-Mansfield RL, deFazio A, Wain GV, Jaworski RC, Byth K, Mote PA and Clarke CL **Relative expression of progesterone receptors A and B in endometrioid cancers of the endometrium.** *Cancer Res* 2001, **61**:4576-4582
- Vihko R and Isomaa V **Endocrine aspects of endometrial cancer.** In: *Endocrine Dependent Tumor* (Edited by: Voight K, Knabbe C) New York: Raven Press 1982, 197-212
- Pike MC and Ross RK **Progestins and menopause: epidemiological studies of risks of endometrial and breast cancer.** *Steroids* 2000, **65**:659-664
- Horwitz KB and Alexander PS **In situ photolinked nuclear progesterone receptors of human breast cancer cells: subunit molecular weights after transformation and translocation.** *Endocrinology* 1983, **113**:2195-2201
- McDonnell DP **Unraveling the human progesterone receptor signal transduction pathway.** *Trends Endocrinol Metab* 1995, **6**:133-138
- Kastner P, Krust A, Turcotte B, Stropp U, Tora L, Gronemeyer H and Chambon P **Two distinct estrogen-regulated promoters generate transcripts encoding the two functionally different human progesterone receptor isoforms A and B.** *EMBO J* 1990, **9**:1603-1614
- Tora L, Gronemeyer H, Turcotte B, Gaub MP and Chambon P **The N-terminal region of the chicken progesterone receptor specifies target gene activation.** *Nature* 1988, **333**:185-188
- Vegeto E, Shahbaz MM, Wen DX, Goldman ME, O'Malley BW and McDonnell DP **Human progesterone receptor A form is a cell- and promoter-specific repressor of human progesterone receptor B function.** *Mol Endocrinol* 1993, **7**:1244-1255
- Graham JD, Yeates C, Balleine RL, Harvey SS, Milliken JS, Bilous AM and Clarke CL **Characterization of progesterone receptor A and B expression in human breast cancer.** *Cancer Res* 1995, **55**:5063-5068
- Mote PA, Balleine RL, McGowan EM and Clarke CL **Colocalization of progesterone receptors A and B by dual immunofluorescent histochemistry in human endometrium during the menstrual cycle.** *J Clin Endocrinol Metab* 1999, **84**:2963-2971
- Tung LT, Mohamed MK, Hoeffler JP, Takimoto GS and Horwitz KB **Antagonist-occupied human progesterone B-receptors activate transcription without binding to progesterone response elements and are dominantly inhibited by A-receptors.** *Mol Endocrinol* 1993, **7**:1256-1265
- McDonnell DP, Shahbaz MM, Vegeto E and Goldman ME **The human progesterone receptor A-form functions as a transcriptional modulator of mineralocorticoid receptor transcriptional activity.** *J Steroid Biochem Mol Biol* 1994, **48**:425-432
- Katzenellenbogen BS **Estrogen receptors: bioactivities and interactions with cell signalling pathways.** *Biol Reprod* 1996, **54**:287-293
- Miller N, Bedard YC, Cooder NB and Shaul DL **Histological changes in the genital tract in transsexual women following androgen therapy.** *Histopathology* 1986, **10**:661-669
- Futterweit W and Deligdisch L **Histopathological effects of exogenously administered testosterone in 19 female to male transsexuals.** *J Clin Endocrinol Metab* 1986, **62**:16-21
- Ruizeveld de Winter JA, Trapman J, Vermey M, Mulder E, Zegers ND and van der Kwast TH **Androgen receptor expression in human tissues: an immunohistochemical study.** *J Histochem Cytochem* 1991, **39**:927-936
- Horie K, Takakura K, Imai K, Liao S and Mori T **Immunohistochemical localization of androgen receptor in the human endometrium, decidua, placenta and pathological conditions of the endometrium.** *Hum Reprod* 1992, **7**:1461-1466
- Kimura N and Mizokami A **Immunocytochemical localization of androgen receptor with polyclonal antibody in paraffin-embedded human tissues: authors' response.** *J Histochem Cytochem* 1994, **42**:125-126
- Chadha S, Pache TD, Huikeshoven JM, Brinkmann AO and van der Kwast TH **Androgen receptor expression in human ovarian and uterine tissue of long term androgen-treated transsexual women.** *Hum Pathol* 1994, **25**:1198-1204
- Mertens HJ, Heineman MJ, Koudstaal J, Theunissen P and Evers JL **Androgen receptor content in human endometrium.** *Eur J Obstet Gynecol Reprod Biol* 1996, **70**:11-13
- Slayden OD, Nayak NR, Burton KA, Chwalisz K, Cameron ST, Critchley HOD, Baird DT and Brenner RM **Progesterone antagonists increase androgen receptor expression in the rhesus macaque and human endometrium.** *J Clin Endocrinol Metab* 2001, **86**:2668-2679
- Brenner RM, West NB and McClellan MC **Estrogen and progestin receptors in the reproductive tract of male and female primates.** *Biol Reprod* 1990, **42**:11-19
- Ace CI and Okulicz WC **Differential gene regulation by estrogen and progesterone in the primate endometrium.** *Mol Cell Endocrinol* 1995, **115**:95-103
- Adams MR, Register TC, Golden DL, Wagner JD and Williams JK **Medroxyprogesterone acetate antagonizes inhibitory effects of conjugated equine estrogens on coronary artery atherosclerosis.** *Arterioscler Thromb Vasc Biol* 1997, **17**:217-221
- Wang H, Masironi B, Eriksson H and Sahlin L **A comparative study of estrogen receptors α and β in the rat uterus.** *Biol Reprod* 1999, **61**:955-964
- Persson I **Estrogens in the causation of breast, endometrial and ovarian cancers – evidence and hypotheses form epidemiological findings.** *J Steroid Biochem Mol Biol* 2000, **74**:357-364
- Cline JM, Söderqvist G, Register TC, Williams JK, Adams MR and Schoultz BV **Assessment of hormonally active agents in the reproductive tract of female nonhuman primates.** *Toxicol pathol* 2001, **29**:84-90
- Katzenellenbogen BS **Mechanisms of action and cross-talk between estrogen receptor and progesterone receptor pathways.** *J Soc Gynecol Invest* 2000, **S33**-7
- Wang H, Critchley HO, Kelly RW, Shen D and Baird DT **Progesterone receptor subtype B is differentially regulated in human endometrial stroma.** *Mol Hum Reprod* 1998, **4**:407-412
- Tseng L and Zhu HH **Regulation of progesterone receptor messenger ribonucleic acid by progestin in human endometrial stromal cells.** *Biol Reprod* 1997, **57**:1360-1366
- Smith CL **Cross-talk between peptide growth factor and estrogen receptor signaling pathway.** *Biol Reprod* 1998, **58**:627-732
- Tabibzadeh S, Lessey B and Satyaswaroop PG **Temporal and site-specific expression of transforming growth factor-beta4 in human endometrium.** *Mol Hum Reprod* 1998, **4**:595-602
- Schwartz LB, Krey L, Demopoulos R, Goldstein SR, Nachtigall LE and Mittal K **Alterations in steroid hormone receptors in the**

- tamoxifen-treated endometrium.** *Am J Obstet Gynecol* 1997, **176**:129-137
37. Adesanya-Famuyiwa OO, Zhou J, Wu G and Bondy C **Localization and sex steroid regulation of androgen receptor gene expression in rhesus monkey uterus.** *Obstet Gynecol* 1999, **93**:265-70
38. Wang H, Isaksson I, von Schoultz B, Cline JM and Sahlin L **The effect of long-term treatment with steroid hormones or tamoxifen on oestrogen receptors (α and β) in the endometrium of ovariectomized cynomolgus macaques.** *J Endocrinol* 2002, **175**:673-681

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

