

Review Article

Polyphenols in Colorectal Cancer: Current State of Knowledge including Clinical Trials and Molecular Mechanism of Action

Md Nur Alam, Muhammad Almoyad, and Fazlul Huq 

Discipline of Biomedical Sciences, Sydney Medical School, The University of Sydney, Cumberland Campus C42, East Street, Lidcombe, NSW 1825, Australia

Correspondence should be addressed to Fazlul Huq; fazlul.huq@sydney.edu.au

Received 1 August 2017; Revised 8 November 2017; Accepted 17 December 2017; Published 15 January 2018

Academic Editor: Michael Linnebacher

Copyright © 2018 Md Nur Alam et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Polyphenols have been reported to have wide spectrum of biological activities including major impact on initiation, promotion, and progression of cancer by modulating different signalling pathways. Colorectal cancer is the second most major cause of mortality and morbidity among females and the third among males. The objective of this review is to describe the activity of a variety of polyphenols in colorectal cancer in clinical trials, preclinical studies, and primary research. The molecular mechanisms of major polyphenols related to their beneficial effects on colorectal cancer are also addressed. Synthetic modifications and other future directions towards exploiting of natural polyphenols against colorectal cancer are discussed in the last section.

1. Introduction

Epidemiological studies exhibiting protective effect of diets rich in fruits and vegetables against different types of cancer have drawn increased attention to the possibility of exploiting biologically active secondary metabolites of plants to fight against cancer. Among the vast array of phytochemicals, compounds called “polyphenols” constitute one of the most numerous and widely distributed groups, covering more than 10,000 different chemical structures [1]. Polyphenols (PP) are reported to have antioxidant, anticarcinogenic, antiatherosclerotic, anti-inflammatory, spasmolytic, hepatoprotective, antiviral, antiallergic, antidiarrheal, antimicrobial, and oestrogenic activity [2].

Colorectal cancer (CRC) is the third most common diagnosed cancer in men after lung and prostate cancer throughout the world. While in women CRC occupies the second position after breast cancer worldwide. Prevalence of CRC is 18% higher in developed countries than developing and undeveloped nations. People of more than 50 years old are more prone to be affected by CRC, and incidence in males is greater than in females. Although diet and Western lifestyle are still considered as being the main factors responsible for CRC, no specific food or other environmental agent has been identified as an exact causative factor [3]. Thus

far, clearly identified types or causes of CRC are hereditary nonpolyposis colorectal cancer, familial adenomatous polyposis, inflammatory bowel diseases, human papillomavirus, and acquired immunodeficiency syndrome [4]. Although surgical resection remains the only curative treatment for CRC, an alternative approach to reduce the mortality rate is chemoprevention, use of synthetic or natural compounds in pharmacologic doses [5].

Colon cancers result from a series of pathologic changes that transform normal colonic epithelium into invasive carcinoma. Dietary PP affect these different cellular processes by acting as chemopreventive blockers. So far, only one review article that has been published concentrated on the effect of polyphenols on colorectal cell lines [6], and only a limited number of polyphenols have been considered. This review focuses on the updated research on a wider variety of polyphenols as applied to colorectal cancer.

2. Chemistry of PP and Their Dietary Sources

PP are also known as polyhydroxyphenols and characterized by the presence of large number of phenol units in their structures, usually existing in plants as glycosides. Polyphenols can be classified according their sources, chemical structures, therapeutic actions, and so on. A classification system of




FIGURE 1: Classification of Polyphenols.

PP has been given in Figure 1 on the basis of the chemical structures of the aglycone portions and Figure 2 gives the basic structures of major groups [7].

A list of the 100 richest dietary sources of PP has been produced using comprehensive Phenol-Explorer data [8]. The richest sources are various spices and dried herbs, cocoa products, some dark coloured berries, some seeds (flaxseed) and nuts (chestnut, hazelnut), and some vegetables, including olive and globe artichoke heads. Top ten of the list containing the highest amount of PP is in the following order: cloves > peppermint (dried) > star anise > cocoa powder > Mexican oregano (dried) > celery seed > black chokeberry > dark chocolate > flaxseed meal > black elderberry.

3. Pathogenesis of CRC and Its Signalling Pathways

Acquired functional capabilities of cancer cells that would allow them to survive, proliferate, and disseminate are known as the hallmarks of cancer, that is, sustaining proliferative signalling, evading growth suppressors, resisting cell death, enabling replicative immortality, inducing angiogenesis, activating invasion and metastasis, reprogramming of energy metabolism, and evading immune destruction [9]. Underpinning these hallmarks are genomic instability and inflammation. While genomic instability confers random mutations including chromosomal rearrangements, causing genetic diversity that expedites the acquisition of hallmarks of cancer, the inflammatory state of premalignant and frankly malignant lesions that is driven by cells of the immune system also fosters multiple hallmark functions.

Based on investigation of different stages of tumour initiation and progression, Fearon and Vogelstein proposed a model of colorectal carcinogenesis that correlated specific genetic events with evolving tissue morphology [10]. The Wnt/ β -catenin pathway plays a dominant role in an initial stage of CRC development. Inactivation of the adenomatous polyposis coli gene is a key starting event in carcinogenesis of more than 60% of colorectal adenomas and carcinomas leading to stimulation of the Wnt pathway via free β -catenin [10].

Stimulation of the epidermal growth factor receptor (EGFR) leads to the activation of KRAS or phosphatidylinositol-3-kinase pathways, which is important in CRC development from early adenoma to intermediate adenoma. Subsequently, numerous signal transduction molecules initiate a cascade of downstream effectors that trigger tumour growth, angiogenesis, and metastasis [11].

Transforming growth factor- β (TGF- β) is a multifunctional polypeptide that binds to specific TGF- β receptors for paracrine and autocrine signalling. This ligand and receptor complex triggers intracellular signalling cascades that include the canonical Smad2 signalling pathway, which complexes with Smad4 and accumulates and translocates into the nucleus. In the nucleus, activated Smad complexes regulate the transcription of specific genes and ultimately regulate cell cycle and tissue repair [12]. TGF β pathway contributes to a favourable microenvironment for tumour growth and metastasis throughout all the steps of carcinogenesis [13]. TGF- β also induces apoptosis, with the association of death-associated protein 6 (DAXX) from the death receptor Fas. After binding, DAXX is then phosphorylated by




FIGURE 2: Basic structures of major groups of polyphenols.

homeodomain-interacting protein kinase 2 (HIPK2), which then activates apoptosis signal-inducing kinase 1 (ASK1). ASK1 activates the Jun amino-terminal kinase (JNK) pathway that causes apoptosis [14–16]. Inactivation of TGF- β pathway components is first detected in advanced adenomas and affects 40–50% of all CRCs [17].

Almost 50% of all CRCs show p53 gene mutations, with higher frequencies observed in distal colon and rectal tumours and lower frequencies in proximal tumours and those with the microsatellite instability or methylator phenotypes [18]. The mutations in p53 or the loss of its functionality occurs mainly at the transition from adenoma to cancer, and the frequency of alterations in the gene increases with the corresponding progression of the lesion [19].

CRC cells share many properties in common with stem cells which are conserved in both dormant and actively proliferating cancer cells [20]. On top of maintaining “stemness” characteristics, CRC cells with metastatic potential dissociate from the tumour mass and spread to other organs in the body [21]. This is achieved through a dedifferentiation program called epithelial-mesenchymal transition (EMT). This key developmental program allows stationary and polarized epithelial cells to undergo multiple biochemical changes that enable them to disrupt cell-cell adherence, lose apical-basal polarity, dramatically remodel the cytoskeleton, and acquire mesenchymal characteristics such as enhanced migratory capacity, invasiveness, and elevated resistance to apoptosis [22]. Adhesion molecules that maintain cell-cell contact in the differentiated tumour cells, such as E-cadherin, are downregulated in the undifferentiated cells, while molecules that impart invasive and migratory behaviour would be upregulated. To accommodate both the “stemness” and mesenchymal properties of invasive CRC cells, it has been proposed that CRC cells with metastatic potential are like “migratory stem cells” [23]. The EMT process is initially driven by three core groups of transcriptional regulators described as follows. The first is a group of transcription factors (TFs) of the Snail zinc-finger family, including SNAI1 and SNAI2 (SLUG) [24]. The second group is the distantly related zinc-finger E-box-binding homeobox family of proteins ZEB1 and ZEB2 (SIP1) [25]. The third group is the basic helix-loop-helix (bHLH) family of transcription factors, including TWIST1, TWIST2, and E12/E47 [26]. In CRC, 85% of resected specimens have moderate to strong TWIST1 expression [27]. The earlier steps of the metastatic cascade EMT program include local invasion, intravasation, survival while transiting through the circulation, and extravasation. EMT programs are dynamically regulated, and during the last step of the metastatic cascade, colonization, carcinoma cells are thought to switch back to an epithelial state through the reverse process, mesenchymal-epithelial transition (MET) [28]. The final stage of the invasion-metastasis cascade, colonization, is likely to require adaptation of propagated CRC cells to the microenvironment of a distant tissue [29].

Increased matrix metalloproteinases (MMPs) expression and their activation generally promote hallmarks of CRC progression including angiogenesis, invasion, and metastasis and correlate with shortened survival. MMPs comprise a large family of at least 25 zinc-dependent endopeptidases

capable of degrading all components of the extracellular matrix (ECM) and are categorized primarily by their structural features as gelatinases, collagenases, membrane-type, stromelysins, and matrilysins [30]. Intercellular adhesion molecule-1 (ICAM-1) is a 90-kDa cell surface glycoprotein that is known to be a member of the immunoglobulin gene superfamily of adhesion molecules. ICAM-1 expression is closely associated with metastasis and may be a useful indicator of prognosis in patients with colorectal cancer [31].

It is evident from the above discussion that the pathogenesis of CRC is characterized by regulatory pathways that are complex involving several layers of communication, cascades, crosstalk, and extensive networking. CRC usually develops through interaction of cytokines, the chemical mediators of inflammation; cytokine receptors, present on the surface of a variety of cell types; secondary messengers which convey signals from cell surface to the interior; transcription factors, which regulate the expression of several genes that affect CRC. Figure 3 depicts the signalling pathways involved in CRC.

4. Roles of PP in CRC Related to Chemoprevention and Apoptosis

Consumption of PP rich food proved to be beneficial in occurrence of CRC in a national prospective cohort study [32]. Numerous studies have evaluated the efficacy of dietary polyphenols against CRC *in vivo*, *in vitro* model and in clinical trials [33–35]. Polyphenols can affect the overall process of carcinogenesis by several mechanisms and cause tumour cell death through apoptotic pathway.

PP have been shown to be highly effective in scavenging singlet oxygen and various free radicals, which leads to DNA damage and tumour promotion [36]. PP also displayed chemopreventive effect through their impact on the bioactivation of carcinogens. Most carcinogens of chemical origin undergo biotransformation by Phase I metabolizing enzymes to be converted into more reactive form suitable for binding with DNA and proceed towards carcinogenesis process. PP were found to inhibit cytochrome P450 enzymes of the CYP1A family and thus act as chemopreventive agents [37]. On the other hand, by increasing the activity of Phase II metabolizing enzymes (glutathione reductase, glutathione peroxidase, glutathione S-reductase, catalase, and quinone reductase), PP are able to provide beneficial effects against CRC [38, 39]. For example, PP obtained from apple inhibited growth of HT-29 human colon cancer cells by modulating expression of genes (*GSTP1*, *GSST2*, *MGST2*, *CYCP4F3*, *CHST5*, *CHST6*, and *CHST7*) involved in the biotransformation of xenobiotics [40].

Orner et al. demonstrated that epigallocatechin-3-gallate (EGCG) attenuated the expression of β -catenin and inhibited intermediate and late stages of colon cancer, via effects on the Wnt/ β -catenin/TCF signalling pathway [41]. EGFR signalling mechanism of CRC progression has been reported to be inhibited by apple procyanidins [42]. Expression of p53 gene has been increased by EGCG that can impede the conversion of colorectal adenoma to colorectal carcinoma during carcinogenesis [43].




FIGURE 3: Signalling pathways in colorectal cancer pathogenesis (adapted from [168]). (EGFR: epidermal growth factor receptor, TGFβ R1/2: transforming growth factor, beta receptor 1/2, EMT: epithelial-mesenchymal transition, ICAMs: intercellular adhesive molecules, MMPs: matrix metalloproteinases).

Apoptosis is a vital physiological process in the normal development, and induction of apoptosis is highly anticipated mode as a therapeutic strategy for cancer control [44–46]. Bcl family of protein, caspase signalling proteins, and p53 genes are the key factors that regulate apoptosis [47]. PP are effective general inhibitor of cancer cell growth and inducers of apoptosis in different cancer cell lines, including leukaemia, skin, lung, stomach, colorectal, and prostate cancer cells [34, 48–52]. Anthocyanin, ellagic acid, curcumin, flavone induced apoptosis in various colon cancer cell lines by different mechanisms in miscellaneous observations [34, 53–55].

PP can prevent the DNA damage caused by free radicals or carcinogenic agents through diverse mechanisms: (a) direct radical scavenging [56, 57], (b) chelating divalent cations involved in Fenton reaction [58], and (c) modulation of enzymes related to oxidative stress (glutathione peroxidase, glutathione reductase, superoxide dismutase, nitric oxide synthase, lipooxygenase, xanthine oxidase, etc.) [59]. Dietary

PP can also act as prooxidants depending on the cell type, dose, and/or time of treatment, as they can enhance reactive oxygen species production and therefore induce apoptosis [58, 60, 61]. In colon cancer HT-29 cells, flavone enriched the mitochondrial pyruvate or lactate uptake, which augmented the superoxide radical production and led to apoptosis [62].

5. Recent Update of Key PP as Applied to CRC

Reported antitumour activity of PP against CRC is largely based on *in vitro* studies, rodent model studies, and even human clinical trials. During *in vitro* studies on antitumour activity of PP, different colorectal cancer cells (HT-29, SW480, Caco-2, Colo-205, Colo-115, HCT-115, HCT-116, DLD-1, LoVo etc.) were cultured, and cell viability was determined via MTT [3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide] reduction assay [154], SRB (Sulforhodamine B) colorimetric assay [155], and crystal violet method [156]. *In vivo* animal model models were produced

by inducing tumour chemically, resulting in APC^{Min/+} mouse and rodent xenograft models. Carcinogen [azoxymethane (AOM), dimethylhydrazine (DMH), dextran sodium sulphate (DSS)] induced colon cancer in rodents can recapitulate in a highly reliable way and frequently used to assess activity of PP. Mutations in the adenomatous polyposis coli (APC) gene are required to initiate familial adenomatous polyposis (FAP) and are also important in CRC tumorigenesis. Several studies have been conducted with PP in APC^{Min/+} mouse that contains (multiple intestinal neoplasia *Min*) a point mutation in the APC gene and develops numerous adenomas. The role of PP has also been investigated in xenograft model where human tumours are injected and established in immunodeficient mouse strains (nude or SCID mice). This section contains the outcome from the studies conducted with PP against CRC.

5.1. Phenolic Acids

5.1.1. Benzoic Acid Derivatives. From literature, only gallic acid among benzoic acid derivatives showed anticancer activity against CRC *in vitro* and *in vivo* model [157, 158], but no study has been conducted to identify the anticancer mechanism of gallic acid in CRC. However, gallic acid is believed to exhibit its anticancer effect by upregulating Bax and downregulating Bcl-2 in other tumour models [157]. Vanillic acid showed significant activity with IC₅₀ values less than 30 μ M in three different CRC cell lines but mechanism has not been studied [159] although vanillic acid and protocatechuic acid did not show significant anticancer activity against CRC [160, 161].

5.1.2. Cinnamic Acid Derivatives. Caffeic acid showed apoptotic cell death against HCT 15 cell lines although IC₅₀ value was very high (800 μ M). Similar findings were made by other researchers [162, 163]. In a recent study caffeic acid did not show any significant activity against HT-29 cell lines up to 200 μ M concentration nor did chlorogenic acid [164] that did not show any significant activity against different human colorectal carcinomas [161]. IC₅₀ values of *p*-coumaric acid against some other CRC cell lines were around 1 mM and apoptosis was the mechanism of cell death [163, 165, 166]. Ferulic acid inhibited CRC progression at adhesion and migration steps but no IC₅₀ value was greater than 1 mM concentration [163].

Carnosic acid showed IC₅₀ values in the range of 24–96 μ M against Caco-2, HT29, and LoVo cell lines. It inhibited cell adhesion and migration, possibly by reducing the activity of secreted proteases such as urokinase plasminogen activator and metalloproteinases. These effects may be mediated through a mechanism involving the inhibition of the COX-2 pathway [167]. Sinapic acid showed IC₅₀ values of less than 25 μ M in three different CRC cell lines but mechanism has not been studied [159].

5.2. Flavonoids

5.2.1. Isoflavones, Neoflavonoids, and Chalcones. Among isoflavones, biochanin A showed ID₅₀ values below 15 μ g/mL

against two CRC cell lines and was found to enhance radiotoxicity *in vitro* [170, 171]. Formononetin that showed dose dependent cell killing, both *in vitro* and *in vivo* in RKO cell line, induces apoptosis by modulating Bax/Bcl-2 activities, inactivating ERK pathway and TNF- α /NF- κ B pathway [172]. Formononetin also showed anticarcinogenic activity in HCT-116 cells via promotion of caspase-dependent apoptosis and inhibition of cell growth, with contribution by downregulation of the antiapoptotic proteins Bcl-2 and Bcl-xL [173]. Daidzein killed 50% of HCT cells, LoVo cells, and DLD-1 cells at concentration of 40 μ M, 68.8 μ M, and 46.3 μ M, respectively, but against LoVo cells it exhibited biphasic effects by killing cells in dose dependent manner at higher concentrations (≥ 5 μ M) and vice versa at lower concentrations (≤ 1 μ M) [75, 174, 175]. Most commonly studied isoflavone, genistein, showed cytotoxicity against HCT, LoVo, and DLD-1 cell lines with IC₅₀ values of 15 μ M, 57.3 μ M, and 56.1 μ M, respectively, whereas in HCC-44B2 cells and HCC50-D3 the value was 11.5 μ g/mL and 9.5 μ g/mL [75, 170, 174]. Genistein reduces the density of cell surface charge and increases the order in membrane protein conformation which might be one of the mechanisms of its anticancer effect [174].

No literature reporting neoflavonoids activity against CRC was found. Among chalcones, phloretin caused apoptotic cell death to HT-29 cells with an IC₅₀ value close to 100 μ M. The mechanism involved changes in mitochondrial membrane permeability and activation of the caspase pathways [176]. Phloretin also has the potential to increase adoptive cellular immunotherapy against SW-1116 CRC cells [177]. Xanthohumol, another important chalcone, is found to show cytotoxicity in different CRC cells *in vivo* and *in vitro* with IC₅₀ values less than 5 μ M [178–180]. The apoptosis involved downregulation of Bcl-2, activation of the caspase cascade, and inhibition of topo I activity. In combination with chemotherapy, it is recommended for use in HCT-15 cell lines, being aimed to reduce drug resistance by inhibition of efflux transporters [180]. Xanthohumol inhibits metastasis by inhibiting expression of CXCR4 chemokine receptor [181].

5.2.2. Flavones, Flavonols, Flavanones, and Flavanonols. Among all different types of flavones, apigenin and luteolin were most commonly investigated phytochemicals for their anticancer activity against CRC. Important flavones that have been studied against CRC are given in Table 1.

Among all different types of flavonols quercetin, chrysin and rutin were studied most for their anticancer activity against colorectal cancer models. Important flavonols that have been investigated against CRC are given in Table 2.

Naringenin appears to be the most commonly studied phytochemicals among the flavanones that can act against colorectal cancer. It suppressed colon carcinogenesis through the aberrant crypt stage in azoxymethane-treated rats [74]. Another study showed that antiproliferative activity of naringenin was estrogen receptor dependent [182], while other *in vitro* studies gave mixed results in different CRC cell lines [65, 80, 152, 183]. Another flavanone, hesperetin, significantly reduced the formation of preneoplastic lesions and effectively

TABLE 1: Important flavones studied against CRC.

Name	Cell line/animal	Comments	Ref.
Apigenin	SW480, HT-29, and Caco-2	Inhibited colon carcinoma cell growth by inducing a reversible G2/M arrest, associated with inhibited activity of p34cdc2 kinase, reduced accumulation of p34cdc2 and cyclin B1 proteins.	[63]
Apigenin	HCT-8	Suppressed tumour angiogenesis via HIF-1 and VEGF expression.	[64]
Apigenin	HCT-116, SW480, HT-29 and LoVo; APC ^{Min/+} mice	Cell death due to apoptosis is mediated by induction of proapoptotic proteins (NAG-1 and p53), cell cycle inhibitor (p21), and kinase pathways. <i>In vivo</i> data also supported <i>in vitro</i> results.	[65]
Apigenin	HT-29	Cytotoxic activity is related to cell cycle arrest through activation of caspase cascade and stimulation of apoptosis. Synergistic activity observed with 5-FU.	[66]
Apigenin	HT-29 and HRT-18	Inhibited metastasis by upregulating CD26 and degrades CXCL12 by increasing DPPIV activity.	[67]
Apigenin	Xenograft of SW480 cells in nude mice	Suppressed growth of colorectal cancer xenografts via phosphorylation and upregulated FADD expression.	[68]
Apigenin	SW480, DLD-1, and LS174T	Inhibited tumour growth and metastasis both <i>in vitro</i> and <i>in vivo</i> by upregulating TAGLN, downregulating MMP-9 expression, decreasing phosphorylation of Akt at Ser473 and in particular Thr308.	[69]
Apigenin	Xenograft study using DLD1, HCT-116, HT-29, HCT-8, and SW480	Synergistic effect was observed with ABT-263 and cell death is mediated via inhibition of Mcl-1, AKT, and ERK pathways.	[70]
Apigenin	HCT116	Induced cell death due to apoptosis and autophagy where apoptosis is via decreased expression of cyclin B1, Cdc2, and Cdc25c; increased expression of p53 and p21 ^{CIP1/WAF1} ; decreased levels of procaspase-8, -9, and -3.	[71]
Apigenin	HT-29 and HCT-15	Oxidative stress resulted in senescence and chemotherapeutic effect.	[72]
	SW480 and HCT-15	Suppressed cell proliferation, migration, and invasion via inhibition of the Wnt/ β -catenin signalling pathway.	[73]
	Sprague Dawley rats	Lowered the number of aberrant crypt foci (ACF) significantly.	[74]
Apigenin, luteolin, baicalein	LoVo and DLD-1	Apigenin had IC ₅₀ values in LoVo and DLD-1 cells lines at 44.7 μ M and 29.6 μ M, luteolin at 57.6 and 40.1 respectively. Baicalein has IC ₅₀ value 51.4 μ M in DLD-1 cell line but no significant activity in LoVo cell lines.	[75]
Apigenin, luteolin, tangeretin, nobiletin	Colo 205	After 24-hour exposure, IC ₅₀ value for apigenin was greater than 100 μ M. For luteolin, tangeretin, and nobiletin the values were 47.6 μ M, 37.5 μ M, and 66.2 μ M, respectively.	[76]
Apigenin, baicalein, luteolin, tangeretin, diosmetin	HT-29 and Caco-2	IC ₅₀ values ranged from 49.4 μ M to 203.6 μ M in HT-29 cell lines and the trend was baicalein < tangeretin < luteolin < apigenin < diosmetin. For Caco-2 cell lines the trend was baicalein < tangeretin < luteolin < diosmetin < apigenin with values ranging from 56.4 μ M to 115.4 μ M.	[77]

TABLE 1: Continued.

Name	Cell line/animal	Comments	Ref.
Luteolin	HT-29	Downregulated the activation of the PI3K/Akt and ERK1/2 pathways via reduction in IGF-IR signalling which may be one of the mechanisms responsible for the observed apoptosis and cell cycle arrest.	[78]
Luteolin	HT-29, SW480	In HT-29 cells, IC ₅₀ value was greater than 200 μ M but in SW480 cells it is 90 μ M.	[79, 80]
	Male Balb/c mice	Inhibited azoxymethane-induced colorectal cancer growth through activation of Nrf2 signalling; altered carbohydrate metabolizing enzymes; decreased expressions of iNOS and COX-2; restored reduced glutathione and protein thiols; decreased lysosomal enzymes, induced apoptosis by modulating Bcl2, Bax, and caspase-3; decreased mucin depleted foci, levels of glycoconjugates; controlled cell proliferation by inhibiting wnt/ β -catenin/GSK-3 β pathway. Luteolin also acts as antimetastatic agent by decreasing MMP-9 and MMP-2.	[81-88]
	HCT-15	Induced growth arrest by inhibiting wnt/ β -catenin/GSK-3 β signalling pathway, induces apoptosis by caspase-3 mediated manner.	[89]
	HT-29	Induced cell cycle arrest by inhibiting CDK2 and cyclin D1, induces apoptosis by activating caspase-3, -7, and -9.	[90]
	Wistar rats	Decreased the number and volume of 1,2-dimethyl hydrazine induced colon cancer and increased activities of enzymic and nonenzymic antioxidants.	[91, 92]
Pinocembrin	HCT-116, SW480	IC ₅₀ value in SW480 cell line was 50 μ M and <100 μ M in HCT-116 cell line. Pinocembrin triggers Bax-dependent mitochondrial apoptosis.	[93]
Tangeretin	HCT-116, HT-29	IC ₅₀ values were 22 μ M and 26 μ M, respectively.	[94]
	Colo 205	Induced cell-cycle G1 arrest through inhibiting cyclin-dependent kinases 2 and 4 activities as well as elevating CDK inhibitors p21 and p27.	[76]
	LoVo and multidrug resistant LoVo/Dx	Greater activity was observed against resistant cells more than LoVo cells and gave synergistic effects with doxorubicin by increasing accumulation and sensitizing doxorubicin. It also induced caspase-3 activation and elevated surface phosphatidylserine exposure.	[95]
	HCT-116 and HT-29	<i>In vitro</i> and <i>in vivo</i> anticancer activity of tangeretin against colorectal cancer was enhanced by emulsion-based delivery system.	[96]
Vitexin-2-O-xyloside	LoVo and Caco-2	Showed IC ₅₀ values greater than 100 μ g/mL in both cell lines but synergistically affected cell growth and apoptosis with raphasatin and (-)-epigallocatechin-3-gallate.	[97]
Nobiletin	F344 rats Sprague Dawley rats	Study on PhIp-induced cancer in F344 rats indicated that nobiletin did significantly reduce the total number of colonic aberrant crypt foci (ACF) compared to the control value.	[74, 98]
Baicalein, wogonin	HT-29 Xenograft assay in nude mouse	IC ₅₀ values for baicalein and wogonin after 48 h exposure were 100 μ M and 150 μ M, respectively. <i>In vivo</i> data supported the activity of baicalein but wogonin proved to be ineffective. Baicalein induced apoptosis in HT-29 cells via Akt inactivation and in a p53-dependent manner.	[99]
Baicalein	DLD-1 (mutant p53), SW48 (p53 wild-type), and HaCaT	Proteomic study proved that baicalein upregulated the expression of PRDX6, which attenuates the generation of ROS and inhibits the growth of CRC cells.	[100]

TABLE 2: Important flavonols studied against CRC.

Name	Cell line/animal	Comments	Ref.
Quercetin	SW480 and HT-29	Inhibited cell growth and induced apoptosis via downregulation of ErbB2/ErbB3 signalling and the Akt pathway.	[101]
	Wistar rats	During DMH induced colon cancer assay, quercetin inhibited intestinal crypt cell proliferation <i>in vivo</i> , but the effect diminished as the level of dietary exposure increased.	[33]
	SW480	Inhibited β -catenin/TCF signalling.	[102]
	CACO-2 and HT-29	Had IC ₅₀ values in the range 30–40 μ M.	[103]
	CO115 and HCT15	Produced synergistic effect in combination with 5-FU by increasing apoptosis via modulating p53.	[104]
	HT-29 xenografts in male nude mice	Induced apoptosis via AMPK activation and p53-dependent apoptotic cell death. Another study using HT29 cell line indicated that quercetin inhibited phosphorylation of EGFR and the ErbB2 receptor.	[105, 106]
	SW480	Antitumour action in SW480 colon cancer cells is related to the inhibition of expression of cyclin D1 and survivin through Wnt/ β -catenin signalling pathway.	[107]
	HT-29	Resveratrol and quercetin in combination showed anticancer activity in colon cancer cells and repressed oncogenic microRNA-27a.	[108]
Quercetin, myricetin, fisetin, galangin, chrysin, morin	HT-29 xenografts in female nude mice	Quercetin and trans-pterostilbene in combination facilitated elimination of colorectal cancer by chemoradiotherapy through a Bcl-2- and superoxide dismutase 2-dependent mechanism.	[109]
	CF1 mice, F344 rats, Wistar rats	Azoxymethane and dimethylhydrazine induced colon cancer study showed reduction of aberrant crypt foci and focal areas of dysplasia.	[110–115]
	APC ^{Min/+} mouse	Quercetin reduced polyp number and size distribution, which might be due to a reduction in macrophage infiltration.	[116]
Quercetin, chrysin, kaempferol	LoVo and DLD-1	In LoVo cell lines the trend of IC ₅₀ values was fisetin < myricetin < quercetin < galangin < chrysin, whereas in DLD-1 cell line it was fisetin < myricetin < galangin < quercetin < chrysin. No significant antitumour effect was observed for Morin.	[75]
	SW480	Quercetin, chrysin, and kaempferol gave IC ₅₀ values of 85, 165, and 100 μ M, respectively.	[80]
Myricetin	HCT-115, Colo-205	Myricetin induced cell death of human HCT-115 cells via Bax/Bcl2-dependent pathway. It inhibited matrix metalloproteinase 2 protein expression and enzyme activity in Colo-205 cells.	[117, 118]
Rutin	SW480, Nude mice	Rutin gave IC ₅₀ value of 125 μ M and exerted <i>in vivo</i> antitumor and antiangiogenic activities.	[119]
	HT-29	Induced mitochondrial apoptosis through a caspase-dependent mechanism.	[120]
	CF1 –female mice	Inhibited azoxymethane-induced colonic neoplasia.	[113]

TABLE 2: Continued.

Name	Cell line/animal	Comments	Ref.
Chrysin	HT-29	Had IC ₅₀ value of 3.1 μ M.	[121]
	HCT-116	Chrysin sensitized tumour necrosis factor- α -induced apoptosis in human tumor cells via suppression of nuclear factor-kappaB.	[122]
	HCT-116	Promoted tumour necrosis factor- (TNF-) related apoptosis-inducing ligand (TRAIL) induced apoptosis.	[123]
	SW480	Chrysin caused cell-cycle arrest at the G2/M phase in a dose-dependent manner.	[80]
	HCT116, DLD1 and SW837	Aryl hydrocarbon receptor was required for the chrysin induced apoptosis and the upregulation of <i>TNF-α</i> and - β gene expression and consequent activation of the TNF-mediated transcriptional pathway.	[124]
	Caco-2	Blocked topotecan-induced apoptosis in spite of inhibition of ABC-transporters.	[125]
Kaempferol	SW480	Sensitized TRAIL-induced apoptosis.	[126]
	HCT-116	The IC ₅₀ of kaempferol was 53.6 μ M in HCT116 (p53+/+) cells and 112.7 μ M in HCT116 (p53-/-) cells. It induced via ataxia-telangiectasia mutated-p53 pathway with the involvement of p53 up-regulated modulator of apoptosis. Kaempferol increased chromatin condensation, DNA fragmentation, and the number of early apoptotic cells in a dose-dependent manner. Kaempferol increased the levels of cleaved caspase-9, caspase-3, and caspase-7 as well as those of cleaved poly (ADP-ribose) polymerase. Moreover, it increased mitochondrial membrane permeability and cytosolic cytochrome c concentrations.	[127]
	HT-29		[128]
Isorhamnetin	HT-29, FVB/N mice	Chemoprotective effects of isorhamnetin were linked to its inhibition of oncogenic Src activity and consequential loss of nuclear β -catenin, activities that were dependent on CSK expression.	[129]
	HCT-116, SW480 and HT-29	IC ₅₀ values for isorhamnetin in HCT-116, SW480, and HT-29 cell lines were 54.87, 56.24, and 43.85 μ M, respectively. The mechanism of cell death was linked with PI3K/Akt/mTOR pathway.	[130]
Fisetin	HT-29	Fisetin inhibited cyclin-dependent kinases leading to cell cycle arrest.	[131]
	HT-29	Enhanced radiosensitivity of p53-mutant HT-29 human colorectal cancer cells.	[132]
	HCT-116, HT-29	IC ₅₀ values for fisetin in HCT-116 and HT-29 cell lines were 132.2 and 57.7 μ M after 72 h, respectively. The mechanism was induction of apoptosis by inhibition of COX2 and Wnt/EGFR/NF- κ B-signalling pathways.	[133]
	HCT-116	Securin depletion sensitizes human colon cancer cells to fisetin-induced apoptosis.	[134]
Galangin	HCT-15, HT-29	Induced cell death via mitochondrial dysfunction and caspase-dependent pathway.	[135]
Morin	HCT-116	Had IC ₅₀ value less than 350 μ g/mL after 48 h and induced apoptosis by modulation of Bcl-2 family members and Fas receptor.	[136]

modulated the xenobiotic-metabolizing enzymes in rats during DMH-induced colon cancer study [184, 185].

Among flavanonols, only taxifolin acts as an effective chemopreventive agent against colon carcinogenesis due to

its antioxidant mediated apoptosis and antiproliferative activities [186, 187]. Taxifolin is found to control NF- κ B-mediated Wnt/ β -catenin signalling via upregulating Nrf2 pathway [188].




FIGURE 4: Molecular mechanism for anticancer action of EGCG and quercetin in CRC.

5.2.3. *Flavanols and Proanthocyanidins.* Epigallocatechin gallate (EGCG) is the most studied flavanols against CRC. EGCG showed IC₅₀ values of 42.2 μM, 47.7 μM, 50.2 μM, 80.1 μM, and 43.1 μM against HCT116, HT29, SW480, and SW837 cell lines, respectively. Mechanism of action has been linked to the inhibition of growth and activation of the epidermal growth factor receptor and human epidermal growth factor receptor-2 signalling pathways [189]. Among eleven different types of flavanol, EGCG showed the highest antiproliferative activity against HCT-116 cells at 50 μM [190]. In APC^{Min/+} mice, EGCG significantly inhibited intestinal tumorigenesis. Oral administration of EGCG showed increased levels of E-cadherin and decreased levels of nuclear β-catenin, c-Myc, phospho-AKT, and phospho-extracellular signal-regulated kinase 1/2 (ERK1/2) in small intestinal tumours [191]. In another mice model, EGCG reduced inflammation-related colon carcinogenesis induced by azoxymethane and dextran sodium sulphate. Antitumour activity has been ascribed to decrease in mRNA expression

levels of COX-2 and inflammatory cytokines (*TNF-α*, *IFN-γ*, *IL-6*, *IL-12*, and *IL-18*) in the colonic mucosa [192]. Other studies mentioned the proposed anticancer mechanisms of EGCG including cell cycle arrest and apoptosis through inhibition of cyclooxygenase-2 expression, activation of AMP-activated protein kinase, cyclin D1 degradation and p21 transcriptional activation, and inhibition of HES1 and Notch2 signalling in different colorectal cancer cell lines [43, 193–196]. EGCG also inhibited invasion and MMP expression, angiogenesis, through blocking the induction of VEGF [197, 198]. Molecular mechanism for antitumour activity of EGCG and quercetin is shown in Figure 4.

Chemopreventive effects of theaflavin have been reported in azoxymethane induced colon cancer study in male Sprague Dawley rats [199]. The mechanism behind the beneficial effects of proanthocyanidins against CRC has been related to inhibition of angiogenesis through suppressing the expression of VEGF and Ang1 [200], induction of apoptosis [201], and antioxidant activity [202].

5.2.4. Anthocyanidins. Malvidin and pelargonidin showed IC_{50} values of 71.7 $\mu\text{g/mL}$ and 154.3 $\mu\text{g/mL}$ against HCT-116 cell line, whereas cyanidin, delphinidin, and petunidin did not induce 50% cell killing even at a concentration of 200 $\mu\text{g/mL}$ [203]. Cyanidin, delphinidin, malvidin, and pelargonidin exhibited no cytotoxicity against Caco-2 cell line, but against metastatic LoVo and LoVo/ADR cell line cyanidin and delphinidin showed significant antitumour activity with low IC_{50} values [204]. Anthocyanin or anthocyanidin containing extracts obtained from a variety of sources were reported to have significant antiproliferative activity against CRC in several animal model ($APC^{\text{Min/+}}$ mouse model, chemically induced) and cell lines [205–213]. Anticancer activity of anthocyanidins is believed to be due to its antimetastatic property through modulation of claudins, matrix metalloproteinases (MMPs), nuclear factor κB (NF- κB) activation, and demethylation of tumour suppressor genes [214–216]. A pilot study involving 25 CRC patients showed 7% decrease in proliferation after consumption of anthocyanin rich diet [217].

5.3. Polyphenolic Amides. Capsaicin is the most studied polyphenolic amide against CRC. *In vitro* and *in vivo* studies in mice bearing Colo-205 tumour xenografts showed that capsaicin significantly reduced tumour progression by activating caspase-3, caspase-8, caspase-9, Bax, Fas and reducing Bcl-2 [218]. Capsaicin and 3,3'-Diindolylmethane worked synergistically against CRC via modulating transcriptional activity of NF- κB , p53, and target genes linked with apoptosis [219]. Other studies showed that anticancer action of capsaicin was related to nitric oxide production, reactive oxygen species generation, and suppression of transcriptional activity of β -catenin/TCF pathway [220–222]. In our study, we have found synergism in binary sequenced combination of capsaicin with oxaliplatin in all sequences (0,0 h; 0,4 h; 4,0 h) and more so in higher concentration in Lim 2405 cell line (unpublished data). Dihydrocapsaicin, a saturated structural analogue of capsaicin, was found to possess greater activity than capsaicin against HCT-116 cells and induced autophagy in a catalase-regulated manner [223]. Avenanthramides significantly inhibited proliferation of HT29, Caco-2, LS174T, and HCT116 human colon cancer cells [224].

5.4. Other Polyphenols

5.4.1. Resveratrol. 3,5,4-Trihydroxystilbene, known as resveratrol, is one of the most studied polyphenols against CRC. It entered into clinical trial after a number of preclinical studies for its encouraging activity and nontoxicity. All studies conducted in rodent model and clinical trial regarding the activity of resveratrol up to 2009 have been described by Bishayee in his review [225]. The authors mentioned 9 *in vivo* studies related to CRC, among which three were conducted in $APC^{\text{Min/+}}$ mice model and others related to chemically induced tumour [226–234]. All the studies discussed in the review (except one on $APC^{\text{Min/+}}$ mice study) provided support for therapeutic potential of resveratrol against CRC. Later on, in 2012, Juan et al. published another review article that focused on the effects of resveratrol on CRC

from conducted *in vivo* studies and clinical trials [235]. The authors provided details on the molecular mechanism of action of resveratrol against CRC. Two more research articles recently described the promising effect of resveratrol in mouse model against CRC, which were not mentioned in earlier reviews [236, 237]. The results from the investigations of the anticancer activity of resveratrol against CRC have not been detailed here because they have been considered well in the previously mentioned reviews. However, a pictorial representation of the molecular mechanism of action of resveratrol against CRC is given in Figure 5. The effect of 5-fluorouracil increased in combination with resveratrol due to chemosensitizing property [238]. Antimetastatic activity of resveratrol in CRC has also been reported [237, 239]. Since resveratrol is found to downregulate multidrug resistant protein 1 by preventing activation of NF- κB signalling and suppressing cAMP-responsive element transcriptional activity, it can be used to overcome drug resistance by combining with other chemotherapeutic drugs [240]. We have found synergism at higher concentration in binary sequenced combination (at 0,0 h; 0,4 h; 4,0 h) of resveratrol with cisplatin but additive to antagonism at lower concentration in HT-29, Caco-2, and Lim 2405 cell lines (unpublished data).

5.4.2. Curcumin. Curcumin is the main active compound in turmeric (dried rhizome of *Curcuma longa*). We have found 82 research articles (*in vitro* and *in vivo* preclinical studies, clinical trial) describing effect of curcumin including mechanism of action against CRC. Here, we have not considered the anticancer activity of curcumin in detail because three review articles discussed well the therapeutic potential of curcumin for CRC along with its mechanism of action [241–243]. Curcumin can inhibit the initiation of carcinogenesis by increasing glutathione S-transferase, induce cell cycle arrest in S and G2/M phase, induce apoptosis, and inhibit metastasis by decreasing CD31, VEGF, IL-8, and mir-21 expression [244–247]. Mechanism of curcumin in CRC as applied to apoptosis is shown in Figure 6. Curcumin has proved to be beneficial in combination with chemotherapy and radiotherapy as well [248, 249]. Synergistic activity of curcumin has also been observed in our study in different sequences and doses with oxaliplatin and cisplatin in four colorectal cell lines (unpublished data).

5.4.3. Rosmarinic Acid and Gingerol. Rosmarinic acid is the main component of Rosemary which at high dose showed antitumour activity applying to *in vitro* and DMH-induced *in vivo* study against CRC [250, 251]. It is thought that MAPK/ERK pathway is linked with the apoptotic mechanism of rosmarinic acid in CRC [252]. Rosmarinic acid has also been reported to possess antimetastatic activity [253].

6-Gingerol is the most important ingredient of ginger showing antiproliferative activity in a dose dependent manner against LoVo cell lines via G2/M cell cycle arrest. Exposure to 6-gingerol induced intracellular ROS and upregulate p53, p27Kip1, and p21Cip1 levels leading to consequent decrease of CDK1, cyclin A, and cyclin B1 [254]. In HCT-115 cell line, its anticancer action was found to be mediated by inhibition of Leukotriene A4 hydrolase [255]. Another




FIGURE 5: Molecular mechanism for anticancer action of resveratrol in CRC.




FIGURE 6: Different pathways involved in apoptosis by curcumin in CRC.

study showed that 6-gingerol stimulated apoptosis through upregulation of NAG-1 and G1 cell cycle arrest through downregulation of cyclin D1 that involved protein degradation as well as β -catenin, PKC ϵ , and GSK-3 β pathways [256]. However, 6-gingerol did not show anticancer activity in Colo-205 cell line [257].

5.4.4. *Ellagic Acid, Secoisolaricresinol, and Matairesinol.* Ellagic acid is a dilactone of hexahydroxydiphenic acid that occurs naturally in berries and nuts such as the raspberry, strawberry, walnut, and pecan. It inhibited growth of Caco-2 cell line, possibly mediated by regulation of matrix metalloproteinases, vascular endothelial growth factor expression,

and induction of apoptosis [258]. According to others, the anticancer action is mediated through downregulation of cyclins A and B1 and upregulation of cyclin E, cell cycle arrest in S phase, induction of apoptosis via intrinsic pathway (Fas-independent, caspase 8-independent) via Bcl-XL downregulation with mitochondrial release of cytochrome c into the cytosol, and activation of initiator caspase-9 and effector caspase-3 [54]. Studies on DMH-induced colon carcinogenesis also proved the beneficial effect of ellagic acid and showed the mechanism to be linked with reduced expressions of NF- κ B, COX-2, iNOS, TNF- α , and IL-6 as well as inhibition of AKT-phosphoinositide-3 kinase pathway [259, 260]. However, metabolic products urolithins were found to be more potent against CRC compared to ellagic acid itself [261]. Mixed urolithins and ellagic acid inhibited phenotypic and molecular colon cancer stem cell features as well [262].

Secoisolariciresinol and matairesinol are lignans. They constitute one of the major groups of phytoestrogens that have been investigated against CRC but the activity remains controversial. Earlier study with flax seed diet containing secoisolariciresinol and matairesinol showed significant antitumour activity [263]. However, *in vivo* and *in vitro* studies done later did not provide insights into the anticancer potential of secoisolariciresinol and matairesinol [264, 265].

6. Chemical Modifications of PP in Nature and Synthetic Analogues

The most important impediment to the successful development of natural PP as clinical therapy against CRC is their low bioavailability. To overcome the problem towards reaching therapeutic concentrations of PP and increasing efficacy, many researchers tried to produce a number of synthetic analogues through structural modifications. Increase in potency *in vitro* and bioavailability *in vivo* has been observed in many studies. Some selected reports concerning chemical modifications of PP applied against CRC are represented in Table 3.

7. Combination of Polyphenol with Chemotherapy/Radiotherapy and Other PP

Emerging evidence suggests that a single-agent approach is probably less likely to be very effective in the management cancer. The rationale for recommending a multidrug regimen is to attack cancer cells through multiple targets and diverse mechanisms of actions with reduced toxicity, ultimately leading towards improved clinical outcomes. With that aim, PP have been investigated with chemotherapy and radiotherapy in various cancer models. For example, curcumin given in combination potentiated the cytotoxic effects of doxorubicin, 5-FU, and paclitaxel in prostate cancer cells [266]; enhanced the antitumour activities of cisplatin, doxorubicin, and Taxol in HA22T/VGH hepatic cancer cells, HeLa cells, or CAO3 and SKOV-3 ovarian

cancer cells [267, 268]; sensitized multiple myeloma cells to vincristine and melphalan [269]. Similar evidence is available in literature for resveratrol, EGCG, quercetin, genistein, proanthocyanidin, and daidzein in various types of cancer with different classes of chemotherapeutic drugs [169, 270–273]. A few studies also have been conducted against CRC where it has been observed that PP in combination with other chemotherapeutic drugs produced synergism, for example, curcumin with 5-fluorouracil against HCT-116 and HT-29 cell line, resveratrol metabolites and oxaliplatin in SW-480 and SW-620 cell lines, genistein with 5-fluorouracil in HT-29 cell line [274–277]. Outcomes of some other combination studies already have been mentioned in describing the effect of individual PP. Like chemosensitization, PP also have shown the potential of radiosensitization in various cancers, but investigations of the same effects against CRC is very scarce [278–281]. In one study quercetin has been shown to increase chemoradiosensitivity against colorectal cancer in xenograft mouse model [109].

Plenty of evidence exists in literature regarding the benefits of the effect of one polyphenol combined with another polyphenol against different types of cancer. Combination of resveratrol with black tea polyphenols resulted in a synergistic tumour suppressive response in mouse skin tumour [282]. Resveratrol showed better chemopreventive response when combined with curcumin by maintaining adequate zinc and modulating Cox-2 and p21 level in mouse model of lung cancer [283]. Combination of genistein with resveratrol reduced the most severe grade of prostate cancer in SV40 Tag-targeted probasin promoter rat model [284]. EGCG in combination with curcumin synergistically inhibited oral premalignant epithelial cells [285]. Quercetin and resveratrol in combination with ellagic acid showed synergism against leukaemia [286]. However, studies on the effect of combination of pure polyphenols against CRC are not numerous. Curcumin showed synergistic antitumour effects in combination with resveratrol in one report of colon cancer model in SCMD mice [287]. Combination of epicatechin and EGCG also exhibited synergistic outcome against HT-29 cancer cells. There are few reports in literature related to the beneficial effects of plant extracts or juices that possess mixture of polyphenols against CRC [288, 289].

Bioprospecting and molecular pharmacology studies have shown that PP can modulate the survival pathways induced by cancer cells, carcinogens, and chemotherapeutics. The possible mechanisms of chemoresistance are shown in Figure 7.

In Section 5, we have considered the molecular mechanisms by which PP would produce anticancer action in CRC. It is thought that PP have the ability to effectively modulate the various mechanisms of chemoresistance. For example, EGCG and quercetin can directly inhibit PI3/AKT pathway, NF κ B pathway, EGFR family pathway, and IAP family pathway and increase p53 (Figure 4). Similarly curcumin can inhibit Bcl-2 family pathway, EGFR family pathway, and NF κ B pathway (Figure 6).

TABLE 3: Important synthetic polyphenols studied against CRC.

Parent PP	Synthetic analogue	Activity	Ref.
Pterostilbene	3'-Hydroxy-pterostilbene	In terms of IC ₅₀ values, synthetic analogue found to be more sensitive against 3 CRC cell lines. <i>In vivo</i> study also proved its greater activity.	[137]
Resveratrol	3, 5, 4'-Trimethoxystilbene, 3, 3', 4, 5'-tetramethoxystilbene	Inhibited HT-29 cell growth.	[138]
	Digalloyl resveratrol	Inhibited HT-29 cell growth more effectively than gallic acid and resveratrol.	[139]
Flavone	3', 4', 5', 5, 7-Pentamethoxyflavone	More active compared to tricrin and apigenin in APC ^{Min/+} mice model.	[140]
Curcumin	Dimethoxycurcumin	In HCT-116 cell lines, dimethoxycurcumin is more potent in terms of ability to kill cancer cells by apoptosis, less extensively metabolized in microsomal systems, and more stable <i>in vivo</i> compared to curcumin.	[141]
	Curcumin difluorinated	At higher concentration synthetic analogue showed greater potency than curcumin in HCT-116 cells.	[142]
	EF31 and UBS109	Both analogues showed significant antitumour activity in colorectal xenograft model possibly via inhibition of NF- κ B and cell cycle progression at G0/G1 phase.	[143]
	GO-Y030, FLLL-11, and FLLL-12	All of the analogues exhibited 4 to 20 times greater activity than curcumin against SW480, HT-29, and HCT116 cell lines but with minimal toxicity against normal cell line.	[144]
EGCG	Peracetylated EGCG	Administration of the synthetic analogue was more effective than EGCG in preventing the shortening of colon length and the formation of aberrant crypt foci and lymphoid nodules in mouse.	[145]
Procyanidin dimer	[3-O-Galloyl]-(-)-epicatechin-(4 β ,8)-(+)-catechin-3-O-gallate	Compared to parent compound synthetic analogue showed increased cytotoxicity against twelve different cell lines including two colorectal cell lines.	[146]
Catechin and/or epicatechin	(2R, 3S)-3', 4', 5,7-tetrahydroxyflavone-3-yl decanoate, (2R, 3S)-3', 4', 5,7-tetrahydroxyflavone-3-yl octadecanoate	Both of them exerted greater cytotoxicity in HCT-116 cells than catechin.	[147, 148]
Genistein	4'-O-(3,4-Di-O-acetyl- α -L-arabino-hexopyranosyl)genistein, 7-O-(2,3,4,6-tetra-O-acetyl- β -D-galactopyranosyl)-(1 \rightarrow 4)-(6-O-acetyl-hex-2-ene- α -D-erythro-pyranosyl)genistein, 7-O-(2,3,5-tri-O-benzyl- β -D-arabinofuranosyl)genistein and 7-O-(4,6-di-O-acetyl-hex-2-ene- α -D-erythro-pyranosyl)genistein	The derivatives showed greater cytostatic and cytotoxic effect than genistein in Colo-205 cell lines.	[149]
Epicatechin	3-O-(3,4,5-trimethoxybenzoyl)-(-)-epicatechin	Synthetic analogue showed IC ₅₀ values at 33 μ M against Caco-2 cell lines and greater activity compared to epicatechin.	[150]

TABLE 3: Continued.

Parent PP	Synthetic analogue	Activity	Ref.
Naringenin	6-C-(E-phenylethenyl)-naringenin	6-C-(E-phenylethenyl)-naringenin suppressed CRC without any toxicity by inhibiting cyclooxygenase-1.	[151]
	5-Hydroxy-2-(4-hydroxyphenyl)-4-oxochroman-7-yl thiophene-2-carboxylate, 5-hydroxy-2-(4-hydroxyphenyl)-4-oxochroman-7-yl 2-methylbenzoate, 5-hydroxy-2-(4-hydroxyphenyl)-4-oxochroman-7-yl isobutyrate, 7-(allyloxy)-5-hydroxy-2-(4-hydroxyphenyl) chroman-4-one and 5-hydroxy-2-(4-hydroxyphenyl)-4-oxochroman-7-yl phenyl carbonate	All of the derivatives gave lower IC ₅₀ values compared to naringenin in HCT-116 cell lines.	[152]
Chrysin	5,7-dimethoxy-8-iodochrysin, 8-bromo-5-hydroxy-7-methoxychrysin and 5,7-Dihydroxy-8-nitrochrysin	These three derivatives among twelve prepared analogues showed prominent activity against CRC compared to chrysin.	[121]
Nobiletin or/and tangeretin	5-hydroxy-6,7,8,3',4'-pentamethoxyflavone, 5-hydroxy-3,6,7,8,3',4'-hexamethoxyflavone, and 5-hydroxy-6,7,8,4'-tetramethoxyflavone.	All synthetic analogues showed lower IC ₅₀ values than nobiletin and tangeretin.	[94]

8. Current Status of PP in Clinical Trials Related to CRC

Following the discovery of significant anticancer potential of curcumin, resveratrol, EGCG, and genistein seen in studies related to *in vitro* and *in vivo* rodent model, the compounds entered into clinical trials for efficacy and toxicity study in human model. Many of the reported Phase I, Phase II, and Phase III studies further validated the potential of using PP against CRC. Benamouzig and Uzzan provided a summary of 21 clinical trials conducted by 2016 related to the use of curcumin against CRC in their review [243]. Similarly 17 clinical trials on resveratrol have been reviewed elsewhere [290]. Few other chemotherapeutics and chemoprevention clinical trials conducted on PP have been summarized by Vinod et al. [169]. Table 4 represents the recently completed or ongoing clinical trials on PP against CRC, which have not been covered by others.

9. Future Perspective and Directions

From our study we have found that curcumin, resveratrol, quercetin, luteolin, apigenin, EGCG are the most investigated

polyphenols against CRC. In terms of *in vitro* cytotoxicity these polyphenols gave average IC₅₀ value around 15–60 μ M against different colorectal cancer cell lines, which is comparatively larger than clinically used anticancer drugs. Moreover some results from single administration of curcumin or resveratrol produced contradictory evidence against *in vitro* and *in vivo* model data. It would be unwise to be overoptimistic and battle against CRC with a single polyphenol only. Rather the strength of polyphenols can be exploited by combining them with clinically used chemotherapeutic drugs to reduce dose related side effects of chemotherapy and overcoming drug resistance. Therefore, we can search among polyphenols for the ones that give synergistic effect with chemotherapeutic drugs or with other phytochemicals. In our laboratory we have been working with curcumin, resveratrol, EGCG, quercetin, capsaicin, 6-gingerol, genistein in combination with cisplatin and oxaliplatin against different CRC cell lines. In many cases we found synergism (unpublished data) like others [291]. Chemical modifications of natural PP can be continued to improve their activity and bioavailability.

Another area of research that could be explored is related to exploiting ability of PP to interact with stem cells in




FIGURE 7: Mechanisms of chemoresistance (adapted from [169]).

CRC. Cancer stem cells are multipotent cells that possess self-renewal capacity and high proliferative capacity and lead to metastasis through migration. Although cancer stem cells represent less than 2.5% of the tumour mass, they may be responsible for the resistance to cancer therapies and relapse in CRC [292]. Wnt/ β -catenin, Hedgehog, and Notch have been identified to play pivotal roles in cancer stem cells self-renewal. Presently researchers are targeting the hallmark stem cell-like properties of tumour cells to overcome cancer. A number of phytochemicals have also been investigated against cancer stem cells in several studies. Curcumin suppressed mammosphere formation along serial passage in breast cancer, and the effect of curcumin on breast cancer stem/progenitor cells was seen to be mediated through its potent inhibitory effect on Wnt/ β -catenin signalling [293]. Genistein also reduced breast cancer stem cells by inhibiting AKT and increasing PTEN [294]. Likewise, resveratrol inhibited pancreatic cancer stem cell characteristics in human and mouse model by inhibiting EMT [295]. In pancreatic cancer model quercetin decreased ALDH1 activity, induced apoptosis, and reduced the expression of proteins implicated in EMT *in vitro*, while it inhibited stem cell-derived xenografts *in vivo*, reducing the expression of proliferation, stemness, and angiogenesis related genes [296]. However, very little has been studied to modulate the stem cells by PP in CRC.

10. Conclusion

As oxidative stress is an inescapable part of aerobic life, it can be said that cancer with its origin in mutations is a disease of living even though it evokes death sentence in many minds. However, as nature creates problems, it also provides solutions. As tumour active polyphenols have been a part of human diet for thousands of years, but without any adverse side effect, it is thought that selected tumour active polyphenols or their derivatives in combination with targeted therapy may provide an affordable means of overcoming drug resistance and reducing side effects in colorectal cancer and indeed in many other cancers.

Abbreviations

- ABCBI: ATP-binding cassette subfamily B member 1
- AKT: Protein kinase B
- AMPK: 5' AMP-activated kinase
- AP-1: Activator protein-1
- Apaf-1: Apoptotic protease activating factor 1
- APC: Activated protein C
- Bax: Bcl-2-associated X protein
- Bcl-2: B-cell lymphoma 2

TABLE 4: Recent clinical trials on PP against CRC.

Polyphenol	Study description (patients)	Institution and status
Curcumin	A Phase II, randomized, double blind, placebo controlled trial for the effectiveness of holistic turmeric supplementation on polyp burden among patients with FAP (40)	Tel Aviv Sourasky Medical Center, Israel; started February 2017
Curcumin	Early Phase I, curcumin in combination with 5-FU in chemoresistant metastatic colorectal cancer (14)	Baylor Research Institute, USA; started March 2016
Curcumin	Randomized Phase II trial studies in treating patients with FAP (44)	Johns Hopkins University USA; completed in 2017 but results have not been published yet
Curcumin	Phase I, pharmacokinetic trial of curcuminoids (24)	University of Michigan Cancer Center, USA; completed but no publication
Curcumin	Phase I, microarray analysis to identify genes that are modified by curcumin that could be used as biomarkers (40)	University of North Carolina, USA; completed but no publication
EGCG	Phase I, chemopreventive effects in patients with curative resections (50)	The University of Texas Health Science Center at San Antonio, USA; started January 2017 and recruiting
EGCG	Green tea extracts for the prevention of colorectal adenomas and colorectal cancer (176)	Seoul National University Hospital, South Korea; completed and found favourable outcome for the chemoprevention [153]
Genistein	Phase I/II, incorporation of genistein in FOLFOX treatment regimen against metastatic CRC (13)	Sofya Pintova, Icahn School of Medicine at Mount Sinai in collaboration with DSM Nutritional Products, Inc., USA; completed January 2017 but result has not been published yet

BCRP:	Breast cancer resistance protein	ICAMs:	Intercellular cell-adhesion molecule-1
CASP3:	Caspase-3	IGF-1R:	Insulin-like growth factor 1 (IGF-1) receptor
Cdc2:	Cell division cycle protein 2	IKK:	I kappa B kinase
CDK4:	Cyclin-dependent kinase 4	IL-8:	Interleukin 8
c-Fos:	Protooncogene	ILs:	Interleukins
CHOP/GADD 153:	Homologous protein/growth arrest- and DNA damage-inducible gene 153	INOS:	Inducible nitric oxide synthase
cIAP:	Cellular inhibitor of apoptosis protein	IRS-1:	Insulin receptor substrate 1
Cox-2:	Cyclooxygenase-2	IRS-2:	Insulin receptor substrate 2
CRC:	Colorectal cancer	JAK:	Janus kinase
CTGF:	Connective tissue growth factor	JNK:	C-Jun N-terminal kinases
Cyt-c:	Cytochrome complex	PGE2:	Prostaglandin E2
DcR3:	Decoy receptor 3	KRAS:	Kirsten rat sarcoma
DR5:	Death receptor 5	LOX:	Liquid oxygen
ECM:	Extracellular matrix	LRP:	Lipoprotein receptor-related protein
EGFR:	Epidermal growth factor receptor	MALAT1:	Metastasis associated lung adenocarcinoma transcript 1
EMT:	Epithelial-mesenchymal transition	MAPK:	Mitogen-activated protein kinase
ERK:	Extracellular signal-regulated kinases	MEK:	Mitogen-activated protein kinase
FAP:	Familial adenomatous polyposis	miR-34a:	MicroRNA 34a
GADD153:	Growth arrest- and DNA damage-inducible gene 153	MMPs:	Matrix metalloproteinases
HIF-1:	Hypoxia-inducible factor-1	MRP1:	Multidrug resistance-associated protein 1
Hsp70:	Heat shock protein 70	mTOR:	Mammalian target of rapamycin
IAP:	Inhibitor of apoptosis	Myc:	Myelocytomatosis

NADPH:	Nicotinamide adenine dinucleotide phosphate
NF- κ B:	Nuclear factor kappa-light-chain-enhancer of activated B cells
P-450:	Cytochromes P450
P53:	Phosphoprotein p53
PARP:	Poly (ADP-ribose) polymerase
PDCD4:	Programmed cell death protein 4
P-gp:	P-glycoprotein 1
PI3K:	Phosphoinositide 3-kinase
PP:	Polyphenols
PTEN:	Phosphatase and tensin homolog
RECK:	Reversion-inducing-cysteine-rich protein with kazal motifs
ROS:	Reactive Oxygen Species
SIRT1:	Sirtuin 1
TCF:	Transcription factor
TGF β :	Transforming growth factor beta
TGF β R1/2:	Transforming growth factor beta receptor 1
TIMP3:	Metalloproteinase inhibitor 3
TNF- α :	Tumour necrosis factor alpha
TRAIL-R1:	Tumour necrosis factor-related apoptosis-inducing ligand-receptor 1
TRAIL-R2:	Tumour necrosis factor-related apoptosis-inducing ligand-receptor 2
TSPI:	Thrombospondin 1
uPAR:	Urokinase receptor
xIAP:	X-linked inhibitor of apoptosis protein
ZEB1/2:	Zinc-finger E-box-binding homeobox 1 and 2.

Conflicts of Interest

Md Nur Alam, Muhammad Almoayad, and Fazlul Huq declare that they have no conflicts of interest.

Authors' Contributions

Muhammad Almoayad conducted the literature review, Md Nur Alam has drafted the entire review, and Fazlul Huq supervised the whole process and critically reviewed the manuscript.

Acknowledgments

Md Nur Alam is grateful to the Australian Government for the Endeavour Postgraduate Award and to his employer Jahangirnagar University, Bangladesh, for providing study leave in support of his Ph.D. program. Muhammad Almoayad expresses his gratitude towards the Government of Saudi Arabia for providing him scholarship and to his employer King Khalid University, Saudi Arabia.

References

- [1] A.-N. Li, S. Li, Y.-J. Zhang, X.-R. Xu, Y.-M. Chen, and H.-B. Li, "Resources and biological activities of natural polyphenols," *Nutrients*, vol. 6, no. 12, pp. 6020–6047, 2014.
- [2] P. Pietta, M. Minoggio, and L. Bramati, "Plant polyphenols: Structure, occurrence and bioactivity," *Studies in Natural Products Chemistry*, vol. 28, pp. 257–312, 2003.
- [3] J. D. Potter, "Colorectal cancer: molecules and populations," *Journal of the National Cancer Institute*, vol. 91, no. 11, pp. 916–932, 1999.
- [4] M. Ponz De Leon, P. Benatti, F. Borghi et al., "Aetiology of colorectal cancer and relevance of monogenic inheritance," *Gut*, vol. 53, no. 1, pp. 115–122, 2004.
- [5] P. A. Jänne and R. J. Mayer, "Chemoprevention of colorectal cancer," *The New England Journal of Medicine*, vol. 342, no. 26, pp. 1960–1968, 2000.
- [6] J. R. Araújo, P. Gonçalves, and F. Martel, "Chemopreventive effect of dietary polyphenols in colorectal cancer cell lines," *Nutrition Research*, vol. 31, no. 2, pp. 77–87, 2011.
- [7] R. Tsao, "Chemistry and biochemistry of dietary polyphenols," *Nutrients*, vol. 2, no. 12, pp. 1231–1246, 2010.
- [8] J. Pérez-Jiménez, V. Neveu, F. Vos, and A. Scalbert, "Identification of the 100 richest dietary sources of polyphenols: an application of the Phenol-Explorer database," *European Journal of Clinical Nutrition*, vol. 64, supplement 3, pp. S112–S120, 2010.
- [9] D. Hanahan and R. A. Weinberg, "Hallmarks of cancer: the next generation," *Cell*, vol. 144, no. 5, pp. 646–674, 2011.
- [10] E. R. Fearon and B. Vogelstein, "A genetic model for colorectal tumorigenesis," *Cell*, vol. 61, no. 5, pp. 759–767, 1990.
- [11] F. Ciardiello and G. Tortora, "EGFR antagonists in cancer treatment," *The New England Journal of Medicine*, vol. 358, no. 11, pp. 1096–1174, 2008.
- [12] J. Massague, *How cells read TGF-beta signals*. *Nat Rev Mol Cell Biol*, 1, 69-78, 2000.
- [13] C. Neuzillet, A. Tijeras-Raballand, R. Cohen et al., "Targeting the TGF β pathway for cancer therapy," *Pharmacology & Therapeutics*, vol. 147, pp. 22–31, 2015.
- [14] X. Yang, R. Khosravi-Far, H. Y. Chang, and D. Baltimore, "Daxx, a novel fas-binding protein that activates JNK and apoptosis," *Cell*, vol. 89, no. 7, pp. 1067–1076, 1997.
- [15] R. Perlman, W. P. Schiemann, M. W. Brooks, H. F. Lodish, and R. A. Weinberg, "TGF- β -induced apoptosis is mediated by the adapter protein Daxx that facilitates JNK activation," *Nature Cell Biology*, vol. 3, no. 8, pp. 708–714, 2001.
- [16] T. G. Hofmann, N. Stollberg, M. L. Schmitz, and H. Will, "HIPK2 Regulates Transforming Growth Factor- β -Induced c-Jun NH 2-Terminal Kinase Activation and Apoptosis in Human Hepatoma Cells," *Cancer Research*, vol. 63, no. 23, pp. 8271–8277, 2003.
- [17] S. D. Markowitz and M. M. Bertagnolli, "Molecular basis of colorectal cancer," *The New England Journal of Medicine*, vol. 361, no. 25, pp. 2404–2460, 2009.
- [18] B. Iacopetta, "TP53 mutation in colorectal cancer," *Human Mutation*, vol. 21, no. 3, pp. 271–276, 2003.
- [19] D. L. Worthley, V. L. Whitehall, K. J. Spring, and B. A. Leggett, "Colorectal carcinogenesis: road maps to cancer," *World Journal of Gastroenterology*, vol. 13, no. 28, pp. 3784–3791, 2007.
- [20] E. Vincan and N. Barker, "The upstream components of the Wnt signalling pathway in the dynamic EMT and MET associated with colorectal cancer progression," *Clinical & Experimental Metastasis*, vol. 25, no. 6, pp. 657–663, 2008.
- [21] J. P. Thiery and J. P. Sleeman, "Complex networks orchestrate epithelial-mesenchymal transitions," *Nature Reviews Molecular Cell Biology*, vol. 7, no. 2, pp. 131–142, 2006.

- [22] B. Boyer and J. P. Thiery, "Epithelium-mesenchyme interconversion as example of epithelial plasticity," *APMIS-Acta Pathologica, Microbiologica et Immunologica Scandinavica*, vol. 101, no. 1-6, pp. 257-268, 1993.
- [23] T. Brabletz, A. Jung, S. Spaderna, F. Hlubek, and T. Kirchner, "Migrating cancer stem cells—an integrated concept of malignant tumour progression," *Nature Reviews Cancer*, vol. 5, no. 9, pp. 744-749, 2005.
- [24] E. Batlle, E. Sancho, C. Franci et al., "The transcription factor Snail is a repressor of E-cadherin gene expression in epithelial tumour cells," *Nature Cell Biology*, vol. 2, no. 2, pp. 84-89, 2000.
- [25] J. Comijn, G. Berx, P. Vermassen et al., "The two-handed E box binding zinc finger protein SIP1 downregulates E-cadherin and induces invasion," *Molecular Cell*, vol. 7, no. 6, pp. 1267-1278, 2001.
- [26] J. Yang, S. A. Mani, J. L. Donaher et al., "Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis," *Cell*, vol. 117, no. 7, pp. 927-939, 2004.
- [27] F. Kroepil, G. Fluegen, D. Vallböhmer et al., "Snail expression in colorectal cancer and its correlation with clinical and pathological parameters," *BMC Cancer*, vol. 13, article no. 145, 2013.
- [28] C. Scheel and R. A. Weinberg, "Cancer stem cells and epithelial-mesenchymal transition: concepts and molecular links," *Seminars in Cancer Biology*, vol. 22, no. 5-6, pp. 396-403, 2012.
- [29] T. Shibue and R. A. Weinberg, "Metastatic colonization: settlement, adaptation and propagation of tumor cells in a foreign tissue environment," *Seminars in Cancer Biology*, vol. 21, no. 2, pp. 99-106, 2011.
- [30] A. H. Said, J.-P. Raufman, and G. Xie, "The role of matrix metalloproteinases in colorectal cancer," *Cancers*, vol. 6, no. 1, pp. 366-375, 2014.
- [31] K. Maeda, S.-M. Kang, T. Sawada et al., "Expression of intercellular adhesion molecule-1 and prognosis in colorectal cancer," *Oncology Reports*, vol. 9, no. 3, pp. 511-514, 2002.
- [32] L. J. Su and L. Arab, "Tea consumption and the reduced risk of colon cancer - Results from a national prospective cohort study," *Public Health Nutrition*, vol. 5, no. 3, pp. 419-425, 2002.
- [33] J. M. Gee, H. Hara, and I. T. Johnson, "Suppression of intestinal crypt cell proliferation and aberrant crypt foci by dietary quercetin in rats," *Nutrition and Cancer*, vol. 43, no. 2, pp. 193-201, 2002.
- [34] U. Wenzel, S. Kuntz, M. D. Brendel, and H. Daniel, "Dietary flavone is a potent apoptosis inducer in human colon carcinoma cells," *Cancer Research*, vol. 60, no. 14, pp. 3823-3831, 2000.
- [35] R. A. Sharma, H. R. McLelland, K. A. Hill et al., "Pharmacodynamic and pharmacokinetic study of oral Curcuma extract in patients with colorectal cancer," *Clinical Cancer Research*, vol. 7, no. 7, pp. 1894-1900, 2001.
- [36] L. Bravo, "Polyphenols: chemistry, dietary sources, metabolism, and nutritional significance," *Nutrition Reviews*, vol. 56, no. 11, pp. 317-333, 1998.
- [37] S. Muto, K.-I. Fujita, Y. Yamazaki, and T. Kamataki, "Inhibition by green tea catechins of metabolic activation of procarcinogens by human cytochrome P450," *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*, vol. 479, no. 1-2, pp. 197-206, 2001.
- [38] S. G. Khan, S. K. Katiyar, R. Agarwal, and H. Mukhtar, "Enhancement of antioxidant and phase II enzymes by oral feeding of green tea polyphenols in drinking water to SKH-1 hairless mice: possible role in cancer chemoprevention," *Cancer Research*, vol. 52, no. 14, pp. 4050-4052, 1992.
- [39] C. Luceri, G. Caderni, A. Sanna, and P. Dolara, "Red wine and black tea polyphenols modulate the expression of cyclooxygenase-2, inducible nitric oxide synthase and glutathione-related enzymes in azoxymethane-induced F344 rat colon tumors," *Journal of Nutrition*, vol. 132, no. 6, pp. 1376-1379, 2002.
- [40] S. Veeriah, T. Kautenburger, N. Habermann et al., "Apple flavonoids inhibit growth of HT29 human colon cancer cells and modulate expression of genes involved in the biotransformation of xenobiotics," *Molecular Carcinogenesis*, vol. 45, no. 3, pp. 164-174, 2006.
- [41] G. A. Orner, W.-M. Dashwood, C. A. Blum et al., "Response of Apcmin and A33ΔNβ-cat mutant mice to treatment with tea, sulindac, and 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP)," *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*, vol. 506-507, pp. 121-127, 2002.
- [42] F. Gossé, S. Guyot, S. Roussi et al., "Chemopreventive properties of apple procyanidins on human colon cancer-derived metastatic SW620 cells and in a rat model of colon carcinogenesis," *Carcinogenesis*, vol. 26, no. 7, pp. 1291-1295, 2005.
- [43] J.-T. Hwang, J. Ha, I.-J. Park et al., "Apoptotic effect of EGCG in HT-29 colon cancer cells via AMPK signal pathway," *Cancer Letters*, vol. 247, no. 1-2, pp. 115-121, 2007.
- [44] M. Y. Hong, R. S. Chapkin, L. A. Davidson et al., "Fish oil enhances targeted apoptosis during colon tumor initiation in part by downregulating Bcl-2," *Nutrition and Cancer*, vol. 46, no. 1, pp. 44-51, 2003.
- [45] H.-N. Koo, H.-J. Jeong, S.-H. Hong, J.-H. Choi, N.-H. An, and H.-M. Kim, "High molecular weight water-soluble chitosan protects against apoptosis induced by serum starvation in human astrocytes," *The Journal of Nutritional Biochemistry*, vol. 13, no. 4, pp. 245-249, 2002.
- [46] C. B. Thompson, "Apoptosis in the pathogenesis and treatment of disease," *Science*, vol. 267, no. 5203, pp. 1456-1462, 1995.
- [47] M. Hollstein, K. Rice, M. S. Greenblatt et al., "Database of p53 gene somatic mutations in human tumors and cell lines," *Nucleic Acids Research*, vol. 22, no. 17, pp. 3551-3555, 1994.
- [48] N. Ahmad, D. K. Feyes, A.-L. Nieminen, R. Agarwal, and H. Mukhtar, "Green tea constituent epigallocatechin-3-gallate and induction of apoptosis and cell cycle arrest in human carcinoma cells," *Journal of the National Cancer Institute*, vol. 89, no. 24, pp. 1881-1886, 1997.
- [49] H. Mukhtar and N. Ahmad, "Mechanism of cancer chemopreventive activity of green tea," *Proceedings of the Society for Experimental Biology and Medicine*, vol. 220, no. 4, pp. 234-238, 1999.
- [50] S. Okabe, Y. Ochiai, M. Aida et al., "Mechanistic aspects of green tea as a cancer preventive: Effect of components on human stomach cancer cell lines," *Japanese Journal of Cancer Research*, vol. 90, no. 7, pp. 733-739, 1999.
- [51] G.-Y. Yang, J. Liao, K. Kim, E. J. Yurkow, and C. S. Yang, "Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols," *Carcinogenesis*, vol. 19, no. 4, pp. 611-616, 1998.
- [52] I. Romero, A. Pérez, A. Ferruelo, M. Luján, and A. Berenguer, "Polyphenols in red wine inhibit the proliferation and induce apoptosis of LNCaP cells," *BJU International*, vol. 89, no. 9, pp. 950-954, 2002.
- [53] Q. K. Wu, J. M. Koponen, H. M. Mykkänen, and A. R. Törrönen, "Berry phenolic extracts modulate the expression of p21WAF1 and Bax but Not Bcl-2 in HT-29 colon cancer cells," *Journal of Agricultural and Food Chemistry*, vol. 55, no. 4, pp. 1156-1163, 2007.

- [54] M. Larrosa, F. A. Tomás-Barberán, and J. C. Espín, "The dietary hydrolysable tannin punicalagin releases ellagic acid that induces apoptosis in human colon adenocarcinoma Caco-2 cells by using the mitochondrial pathway," *The Journal of Nutritional Biochemistry*, vol. 17, no. 9, pp. 611–625, 2006.
- [55] R. Rashmi, T. R. Santhosh Kumar, and D. Karunakaran, "Human colon cancer cells differ in their sensitivity to curcumin-induced apoptosis and heat shock protects them by inhibiting the release of apoptosis-inducing factor and caspases," *FEBS Letters*, vol. 538, no. 1-3, pp. 19–24, 2003.
- [56] M. Alía, R. Mateos, S. Ramos, E. Lecumberri, L. Bravo, and L. Goya, "Influence of quercetin and rutin on growth and antioxidant defense system of a human hepatoma cell line (HepG2)," *European Journal of Nutrition*, vol. 45, no. 1, pp. 19–28, 2006.
- [57] M. Sonee, T. Sum, C. Wang, and S. K. Mukherjee, "The soy isoflavone, genistein, protects human cortical neuronal cells from oxidative stress," *NeuroToxicology*, vol. 25, no. 5, pp. 885–891, 2004.
- [58] H. Nakagawa, K. Hasumi, J.-T. Woo, K. Nagai, and M. Wachi, "Generation of hydrogen peroxide primarily contributes to the induction of Fe(II)-dependent apoptosis in Jurkat cells by (-)-epigallocatechin gallate," *Carcinogenesis*, vol. 25, no. 9, pp. 1567–1574, 2004.
- [59] M. Alía, S. Ramos, R. Mateos, A. B. Granado-Serrano, L. Bravo, and L. Goya, "Quercetin protects human hepatoma HepG2 against oxidative stress induced by *tert*-butyl hydroperoxide," *Toxicology and Applied Pharmacology*, vol. 212, no. 2, pp. 110–118, 2006.
- [60] S. C. Hyang, H.-J. Chang, H. C. Eun, J. K. Hyun, and H. K. Kyung, "Molecular and absorption properties of 12 soy isoflavones and their structure-activity relationship with selected biological activities," *Biotechnology Letters*, vol. 27, no. 15, pp. 1105–1111, 2005.
- [61] T. Nakazato, K. Ito, Y. Ikeda, and M. Kizaki, "Green tea component, catechin, induces apoptosis of human malignant B cells via production of reactive oxygen species," *Clinical Cancer Research*, vol. 11, no. 16, pp. 6040–6049, 2005.
- [62] U. Wenzel, K. Schoberl, K. Lohner, and H. Daniel, "Activation of mitochondrial lactate uptake by flavone induces apoptosis in human colon cancer cells," *Journal of Cellular Physiology*, vol. 202, no. 2, pp. 379–390, 2005.
- [63] W. Wang, L. Heideman, C. S. Chung, J. C. Pelling, K. J. Koehler, and D. F. Birt, "Cell-cycle arrest at G2/M and growth inhibition by apigenin in human colon carcinoma cell lines," *Molecular Carcinogenesis*, vol. 28, no. 2, pp. 102–110, 2000.
- [64] J. Fang, Q. Zhou, L.-Z. Liu et al., "Apigenin inhibits tumor angiogenesis through decreasing HIF-1 α and VEGF expression," *Carcinogenesis*, vol. 28, no. 4, pp. 858–864, 2007.
- [65] Y. Zhong, C. Krisanapun, S.-H. Lee et al., "Molecular targets of apigenin in colorectal cancer cells: Involvement of p21, NAG-1 and p53," *European Journal of Cancer*, vol. 46, no. 18, pp. 3365–3374, 2010.
- [66] M. Turktekin, E. Konac, H. I. Onen, E. Alp, A. Yilmaz, and S. Menevse, "Evaluation of the effects of the flavonoid apigenin on apoptotic pathway gene expression on the colon cancer cell line (HT29)," *Journal of Medicinal Food*, vol. 14, no. 10, pp. 1107–1117, 2011.
- [67] É. C. Lefort and J. Blay, "The dietary flavonoid apigenin enhances the activities of the anti-metastatic protein CD26 on human colon carcinoma cells," *Clinical & Experimental Metastasis*, vol. 28, no. 4, pp. 337–349, 2011.
- [68] Q. R. Wang, X. Q. Yao, G. Wen et al., "Apigenin suppresses the growth of colorectal cancer xenografts via phosphorylation and up-regulated FADD expression," *Oncology Letters*, vol. 2, no. 1, pp. 43–47, 2011.
- [69] L. Chunhua, L. Donglan, F. Xiuqiong et al., "Apigenin up-regulates transgelin and inhibits invasion and migration of colorectal cancer through decreased phosphorylation of AKT," *The Journal of Nutritional Biochemistry*, vol. 24, no. 10, pp. 1766–1775, 2013.
- [70] H. Shao, K. Jing, E. Mahmoud, H. Huang, X. Fang, and C. Yu, "Apigenin sensitizes colon cancer cells to antitumor activity of abt-263," *Molecular Cancer Therapeutics*, vol. 12, no. 12, pp. 2640–2650, 2013.
- [71] Y. Lee, B. Sung, Y. J. Kang et al., "Apigenin-induced apoptosis is enhanced by inhibition of autophagy formation in HCT116 human colon cancer cells," *International Journal of Oncology*, vol. 44, no. 5, pp. 1599–1606, 2014.
- [72] K. Banerjee and M. Mandal, "Oxidative stress triggered by naturally occurring flavone apigenin results in senescence and chemotherapeutic effect in human colorectal cancer cells," *Redox Biology*, vol. 5, pp. 153–162, 2015.
- [73] M. Xu, S. Wang, Y. Song, J. Yao, K. Huang, and X. Zhu, "Apigenin suppresses colorectal cancer cell proliferation, migration and invasion via inhibition of the Wnt/ β -catenin signaling pathway," *Oncology Letters*, vol. 11, no. 5, pp. 3075–3080, 2016.
- [74] T. Leonardi, J. Vanamala, S. S. Taddeo et al., "Apigenin and naringenin suppress colon carcinogenesis through the aberrant crypt stage in azoxymethane-treated rats," *Experimental Biology and Medicine*, vol. 235, no. 6, pp. 710–717, 2010.
- [75] H. Chang, M. Mi, W. Ling et al., "Structurally related cytotoxic effects of flavonoids on human cancer cells in vitro," *Archives of Pharmacological Research*, vol. 31, no. 9, pp. 1137–1144, 2008.
- [76] M.-H. Pan, W.-J. Chen, S.-Y. Lin-Shiau, C.-T. Ho, and J.-K. Lin, "Tangeretin induces cell-cycle G1 arrest through inhibiting cyclin-dependent kinases 2 and 4 activities as well as elevating Cdk inhibitors p21 and p27 in human colorectal carcinoma cells," *Carcinogenesis*, vol. 23, no. 10, pp. 1677–1684, 2002.
- [77] S. Kuntz, U. Wenzel, and H. Daniel, "Comparative analysis of the effects of flavonoids on proliferation, cytotoxicity, and apoptosis in human colon cancer cell lines," *European Journal of Nutrition*, vol. 38, no. 3, pp. 133–142, 1999.
- [78] D. Y. Lim, H. J. Cho, J. Kim, C. W. Nho, K. W. Lee, and J. H. Y. Park, "Luteolin decreases IGF-II production and downregulates insulin-like growth factor-I receptor signaling in HT-29 human colon cancer cells," *BMC Gastroenterology*, vol. 12, article no. 9, 2012.
- [79] S. Attoub, A. H. Hassan, B. Vanhoecke et al., "Inhibition of cell survival, invasion, tumor growth and histone deacetylase activity by the dietary flavonoid luteolin in human epithelioid cancer cells," *European Journal of Pharmacology*, vol. 651, no. 1-3, pp. 18–25, 2011.
- [80] W. Wang, P. C. VanAlstyne, K. A. Irons, S. Chen, J. W. Stewart, and D. F. Birt, "Individual and interactive effects of apigenin analogs on G2/M cell-cycle arrest in human colon carcinoma cell lines," *Nutrition and Cancer*, vol. 48, no. 1, pp. 106–114, 2004.
- [81] A. K. Pandurangan, S. K. Ananda Sadagopan, P. Dharmalingam, and S. Ganapasam, "Luteolin, a bioflavonoid inhibits Azoxymethane-induced colorectal cancer through activation of Nrf2 signaling," *Toxicology Mechanisms and Methods*, vol. 24, no. 1, pp. 13–20, 2014.
- [82] AK. Pandurangan.

- [83] S. Ganapasam, A. Pandurangan, S. Kumar, and P. Dharmalingam, "Luteolin, a bioflavonoid inhibits azoxymethane-induced colon carcinogenesis: Involvement of iNOS and COX-2," *Pharmacognosy Magazine*, vol. 10, no. 38, p. 306, 2014.
- [84] A.K. Pandurangan, *Ganapasam S: Luteolin modulates cellular thiols on azoxymethane-induced colon carcinogenesis*. *Asian J Exp Biol Sci*, 4, 25-250, 2013.
- [85] A. K. Pandurangan and S. Ganapsam, "Luteolin induces apoptosis in azoxymethane-induced colon carcinogenesis through the involvement of Bcl-2, Bax and Caspase-3," *Journal of Chemical and Pharmaceutical Research*, vol. 5, no. 4, pp. 143-148, 2013.
- [86] P. Ashok Kumar, P. Dharmalingam, A. S. Suresh Kumar, and S. Ganapasam, "Effect of luteolin on the levels of glycoproteins during azoxymethane-induced colon carcinogenesis in mice," *Asian Pacific Journal of Cancer Prevention*, vol. 13, no. 4, pp. 1569-1573, 2012.
- [87] P. Ashokkumar and G. Sudhandiran, "Luteolin inhibits cell proliferation during Azoxymethane-induced experimental colon carcinogenesis via Wnt/ β -catenin pathway," *Investigational New Drugs*, vol. 29, no. 2, pp. 273-284, 2011.
- [88] A. K. Pandurangan, P. Dharmalingam, S. K. A. Sadagopan, and S. Ganapasam, "Luteolin inhibits matrix metalloproteinase 9 and 2 in azoxymethane-induced colon carcinogenesis," *Human & Experimental Toxicology*, vol. 33, no. 11, pp. 1176-1185, 2014.
- [89] A. K. Pandurangan, P. Dharmalingam, S. K. A. Sadagopan, M. Ramar, A. Munusamy, and S. Ganapasam, "Luteolin induces growth arrest in colon cancer cells through involvement of Wnt/ β -catenin/gsk-3 β signaling," *Journal of Environmental Pathology, Toxicology and Oncology*, vol. 32, no. 2, pp. 131-139, 2013.
- [90] D. Y. Lim, Y. Jeong, A. L. Tyner, and J. H. Y. Park, "Induction of cell cycle arrest and apoptosis in HT-29 human colon cancer cells by the dietary compound luteolin," *American Journal of Physiology-Gastrointestinal and Liver Physiology*, vol. 292, no. 1, pp. G66-G75, 2007.
- [91] V. Manju and N. Nalini, "Chemopreventive potential of luteolin during colon carcinogenesis induced by 1,2-dimethylhydrazine," *The Italian journal of biochemistry*, vol. 54, no. 3-4, pp. 268-275, 2005.
- [92] V. Manju, V. Balasubramanian, and N. Nalini, "Rat colonic lipid peroxidation and antioxidant status: The effects of dietary luteolin on 1,2-dimethylhydrazine challenge," *Cellular & Molecular Biology Letters*, vol. 10, no. 3, pp. 535-551, 2005.
- [93] M. A. S. Kumar, M. Nair, P. S. Hema, J. Mohan, and T. R. Santhoshkumar, "Pinocembrin triggers Bax-dependent mitochondrial apoptosis in colon cancer cells," *Molecular Carcinogenesis*, vol. 46, no. 3, pp. 231-241, 2007.
- [94] P. Qiu, P. Dong, H. Guan et al., "Inhibitory effects of 5-hydroxy polymethoxyflavones on colon cancer cells," *Molecular Nutrition & Food Research*, vol. 54, no. 2, pp. S244-S252, 2010.
- [95] O. Wesołowska, J. Wiśniewski, K. Środa-Pomianek et al., "Multidrug resistance reversal and apoptosis induction in human colon cancer cells by some flavonoids present in citrus plants," *Journal of Natural Products*, vol. 75, no. 11, pp. 1896-1902, 2012.
- [96] Y. Ting, Y.-S. Chiou, M.-H. Pan, C.-T. Ho, and Q. Huang, "In vitro and in vivo anti-cancer activity of tangeretin against colorectal cancer was enhanced by emulsion-based delivery system," *Journal of Functional Foods*, vol. 15, pp. 264-273, 2015.
- [97] A. Papi, F. Farabegoli, R. Iori et al., "Vitexin-2-O-xyloside, raphasatin and (-)-epigallocatechin-3-gallate synergistically affect cell growth and apoptosis of colon cancer cells," *Food Chemistry*, vol. 138, no. 2-3, pp. 1521-1530, 2013.
- [98] M. X. Tang, K. Ogawa, M. Asamoto et al., "Effects of nobiletin on PhIP-induced prostate and colon carcinogenesis in F344 rats," *Nutrition and Cancer*, vol. 63, no. 2, pp. 227-233, 2011.
- [99] S.-J. Kim, H.-J. Kim, H.-R. Kim et al., "Antitumor actions of baicalein and wogonin in HT-29 human colorectal cancer cells," *Molecular Medicine Reports*, vol. 6, no. 6, pp. 1443-1449, 2012.
- [100] W.-S. Huang, Y.-H. Kuo, C.-C. Chin et al., "Proteomic analysis of the effects of baicalein on colorectal cancer cells," *Proteomics*, vol. 12, no. 6, pp. 810-819, 2012.
- [101] W. K. Kim, M. H. Bang, E. S. Kim et al., "Quercetin decreases the expression of ErbB2 and ErbB3 proteins in HT-29 human colon cancer cells," *The Journal of Nutritional Biochemistry*, vol. 16, no. 3, pp. 155-162, 2005.
- [102] C. H. Park, J. Y. Chang, E. R. Hahm, S. Park, H.-K. Kim, and C. H. Yang, "Quercetin, a potent inhibitor against β -catenin/Tcf signaling in SW480 colon cancer cells," *Biochemical and Biophysical Research Communications*, vol. 328, no. 1, pp. 227-234, 2005.
- [103] S.-M. Kuo, "Antiproliferative potency of structurally distinct dietary flavonoids on human colon cancer cells," *Cancer Letters*, vol. 110, no. 1-2, pp. 41-48, 1996.
- [104] C. P. R. Xavier, C. F. Lima, M. Rohde, and C. Pereira-Wilson, "Quercetin enhances 5-fluorouracil-induced apoptosis in MSI colorectal cancer cells through p53 modulation," *Cancer Chemotherapy and Pharmacology*, vol. 68, no. 6, pp. 1449-1457, 2011.
- [105] H.-J. Kim, S.-K. Kim, B.-S. Kim et al., "Apoptotic effect of quercetin on HT-29 colon cancer cells via the AMPK signaling pathway," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 15, pp. 8643-8650, 2010.
- [106] D. Fridrich, N. Teller, M. Esselen, G. Pahlke, and D. Marko, "Comparison of delphinidin, quercetin and (-)-epigallocatechin-3-gallate as inhibitors of the EGFR and the ErbB2 receptor phosphorylation," *Molecular Nutrition & Food Research*, vol. 52, no. 7, pp. 815-822, 2008.
- [107] B.-E. Shan, M.-X. Wang, and R.-Q. Li, "Quercetin inhibit human SW480 colon cancer growth in association with inhibition of cyclin D1 and survivin expression through Wnt/ β -catenin signaling pathway," *Cancer Investigation*, vol. 27, no. 6, pp. 604-612, 2009.
- [108] A. del Follo-Martinez, N. Banerjee, X. Li, S. Safe, and S. Mertens-Talcott, "Resveratrol and quercetin in combination have anticancer activity in colon cancer cells and repress oncogenic microRNA-27a," *Nutrition and Cancer*, vol. 65, no. 3, pp. 494-504, 2013.
- [109] S. Priego, F. Feddi, P. Ferrer et al., "Natural polyphenols facilitate elimination of HT-29 colorectal cancer xenografts by chemoradiotherapy: a Bcl-2- and superoxide dismutase 2-dependent mechanism," *Molecular Cancer Therapeutics*, vol. 7, no. 10, pp. 3330-3342, 2008.
- [110] A. A. Dihal, V. C. J. De Boer, H. Van Der Woude et al., "Quercetin, but not its glycosidated conjugate rutin, inhibits azoxymethane-induced colorectal carcinogenesis in F344 rats," *Journal of Nutrition*, vol. 136, no. 11, pp. 2862-2867, 2006.
- [111] S. R. Volate, D. M. Davenport, S. J. Muga, and M. J. Wargovich, "Modulation of aberrant crypt foci and apoptosis by dietary herbal supplements (quercetin, curcumin, silymarin, ginseng and rutin)," *Carcinogenesis*, vol. 26, no. 8, pp. 1450-1456, 2005.
- [112] A. P. Femia, G. Caderni, M. Ianni et al., "Effect of diets fortified with tomatoes or onions with variable quercetin-glycoside content on azoxymethane-induced aberrant crypt foci in the

- colon of rats," *European Journal of Nutrition*, vol. 42, no. 6, pp. 346–352, 2003.
- [113] E. E. Deschner, J. Ruperto, G. Wong, and H. L. Newmark, "Quercetin and rutin as inhibitors of azoxymethanol-induced colonic neoplasia," *Carcinogenesis*, vol. 12, no. 7, pp. 1193–1196, 1991.
- [114] M. A. Pereira, C. J. Grubbs, L. H. Barnes et al., "Effects of the phytochemicals, curcumin and quercetin, upon azoxymethane-induced colon cancer and 7, 12-dimethylbenz[a]anthracene-induced mammary cancer in rats," *Carcinogenesis*, vol. 17, no. 6, pp. 1305–1311, 1996.
- [115] G. Y. Wong and J. F. Ruperto, "The Effect Of Dietary Quercetin And Rutin On Aom-Induced Acute Colonic Epithelial Abnormalities In Mice Fed A High-Fat Diet," *Nutrition and Cancer*, vol. 20, no. 3, pp. 199–204, 1993.
- [116] E. A. Murphy, J. M. Davis, J. L. McClellan, and M. D. Carmichael, "Quercetin's effects on intestinal polyp multiplicity and macrophage number in the ApcMin/+ mouse," *Nutrition and Cancer*, vol. 63, no. 3, pp. 421–426, 2011.
- [117] M. E. Kim, T. K. Ha, J. H. Yoon, and J. S. Lee, "Myricetin induces cell death of human colon cancer cells via BAX/BCL2-dependent pathway," *Anticancer Reseach*, vol. 34, no. 2, pp. 701–706, 2014.
- [118] C.-H. Ko, S.-C. Shen, T. J. F. Lee, and Y.-C. Chen, "Myricetin inhibits matrix metalloproteinase 2 protein expression and enzyme activity in colorectal carcinoma cells," *Molecular Cancer Therapeutics*, vol. 4, no. 2, pp. 281–290, 2005.
- [119] A. J. Alonso-Castro, F. Domínguez, and A. García-Carrancá, "Rutin exerts antitumor effects on nude mice bearing SW480 tumor," *Archives of Medical Research*, vol. 44, no. 5, pp. 346–351, 2013.
- [120] T. E. Guon and H. S. Chung, "Hyperoside and rutin of nelumbo nucifera induce mitochondrial apoptosis through a caspase-dependent mechanism in HT-29 human colon cancer cells," *Oncology Letters*, vol. 11, no. 4, pp. 2463–2470, 2016.
- [121] X. Zheng, W.-D. Meng, Y.-Y. Xu, J.-G. Cao, and F.-L. Qing, "Synthesis and anticancer effect of chrysin derivatives," *Bioorganic & Medicinal Chemistry Letters*, vol. 13, no. 5, pp. 881–884, 2003.
- [122] X. Li, Q. Huang, C. Ong, X. Yang, and H. Shen, "Chrysin sensitizes tumor necrosis factor- α -induced apoptosis in human tumor cells via suppression of nuclear factor- κ B," *Cancer Letters*, vol. 293, no. 1, pp. 109–116, 2010.
- [123] X. Li, J. Wang, J. Huang et al., "Chrysin promotes tumor necrosis factor (TNF)-related apoptosis-inducing ligand (TRAIL) induced apoptosis in human cancer cell lines," *Toxicology in Vitro*, vol. 25, no. 3, pp. 630–635, 2011.
- [124] S. M. Ronnekleiv-Kelly, M. Nukaya, C. J. Díaz-Díaz et al., "Aryl hydrocarbon receptor-dependent apoptotic cell death induced by the flavonoid chrysin in human colorectal cancer cells," *Cancer Letters*, vol. 370, no. 1, pp. 91–99, 2016.
- [125] M. Schumacher, A. Hautzinger, A. Rossmann et al., "Chrysin blocks topotecan-induced apoptosis in Caco-2 cells in spite of inhibition of ABC-transporters," *Biochemical Pharmacology*, vol. 80, no. 4, pp. 471–479, 2010.
- [126] T. Yoshida, M. Konishi, M. Horinaka et al., "Kaempferol sensitizes colon cancer cells to TRAIL-induced apoptosis," *Biochemical and Biophysical Research Communications*, vol. 375, no. 1, pp. 129–133, 2008.
- [127] W. Li, B. Du, T. Wang, S. Wang, and J. Zhang, "Kaempferol induces apoptosis in human HCT116 colon cancer cells via the ataxia-telangiectasia mutated-p53 pathway with the involvement of p53 upregulated modulator of apoptosis," *Chemico-Biological Interactions*, vol. 177, no. 2, pp. 121–127, 2009.
- [128] H. S. Lee, H. J. Cho, R. Yu, K. W. Lee, H. S. Chun, and J. H. Y. Park, "Mechanisms underlying apoptosis-inducing effects of kaempferol in HT-29 human colon cancer cells," *International Journal of Molecular Sciences*, vol. 15, no. 2, pp. 2722–2737, 2014.
- [129] S. M. Saud, M. R. Young, Y. L. Jones-Hall et al., "Chemopreventive activity of plant flavonoid isorhamnetin in colorectal cancer is mediated by oncogenic Src and β -catenin," *Cancer Research*, vol. 73, no. 17, pp. 5473–5484, 2013.
- [130] C. Li, X. Yang, C. Chen, S. Cai, and J. Hu, "Isorhamnetin suppresses colon cancer cell growth through the PI3K-Akt-mTOR pathway," *Molecular Medicine Reports*, vol. 9, no. 3, pp. 935–940, 2014.
- [131] X. Lu, J. Jung, H. J. Cho et al., "Fisetin inhibits the activities of cyclin-dependent kinases leading to cell cycle arrest in HT-29 human colon cancer cells," *Journal of Nutrition*, vol. 135, no. 12, pp. 2884–2890, 2005.
- [132] W.-S. Chen, Y.-J. Lee, Y.-C. Yu et al., "Enhancement of P53-mutant human colorectal cancer cells radiosensitivity by flavonoid fisetin," *International Journal of Radiation Oncology • Biology • Physics*, vol. 77, no. 5, pp. 1527–1535, 2010.
- [133] Y. Suh, F. Afaq, J. J. Johnson, and H. Mukhtar, "A plant flavonoid fisetin induces apoptosis in colon cancer cells by inhibition of COX2 and Wnt/EGFR/NF- κ B-signaling pathways," *Carcinogenesis*, vol. 30, no. 2, pp. 300–307, 2009.
- [134] S.-H. Yu, P.-M. Yang, C.-W. Peng, Y.-C. Yu, and S.-J. Chiu, "Securin depletion sensitizes human colon cancer cells to fisetin-induced apoptosis," *Cancer Letters*, vol. 300, no. 1, pp. 96–104, 2011.
- [135] T. K. Ha, M. E. Kim, J. H. Yoon, S. J. Bae, J. Yeom, and J. S. Lee, "Galangin induces human colon cancer cell death via the mitochondrial dysfunction and caspase-dependent pathway," *Experimental Biology and Medicine*, vol. 238, no. 9, pp. 1047–1054, 2013.
- [136] H.-B. Hyun, W. S. Lee, S.-I. Go et al., "The flavonoid morin from Moraceae induces apoptosis by modulation of Bcl-2 family members and Fas receptor in HCT 116 cells," *International Journal of Oncology*, vol. 46, no. 6, pp. 2670–2678, 2015.
- [137] T.-C. Cheng, C.-S. Lai, M.-C. Chung et al., "Potent anti-cancer effect of 39-hydroxypterostilbene in human colon xenograft tumors," *PLoS ONE*, vol. 9, no. 11, Article ID e111814, 2014.
- [138] P. Saiko, M. Pemberger, Z. Horvath et al., "Novel resveratrol analogs induce apoptosis and cause cell cycle arrest in HT29 human colon cancer cells: inhibition of ribonucleotide reductase activity," *Oncology Reports*, vol. 19, no. 6, pp. 1621–1626, 2008.
- [139] A. Bernhaus, M. Fritzer-Szekeres, M. Grusch et al., "Digalloyl-resveratrol, a new phenolic acid derivative induces apoptosis and cell cycle arrest in human HT-29 colon cancer cells," *Cancer Letters*, vol. 274, no. 2, pp. 299–304, 2009.
- [140] H. Cai, S. Sale, R. Schmid et al., "Flavones as colorectal cancer chemopreventive agents - Phenol-O-methylation enhances efficacy," *Cancer Prevention Research*, vol. 2, no. 8, pp. 743–750, 2009.
- [141] C. Tamvakopoulos, K. Dimas, Z. D. Sofianos et al., "Metabolism and anticancer activity of the curcumin analogue, dimethoxy-curcumin," *Clinical Cancer Research*, vol. 13, no. 4, pp. 1269–1277, 2007.
- [142] S. Padhye, H. Yang, A. Jamadar et al., "New difluoro knoevenagel condensates of curcumin, their schiff bases and copper

- complexes as proteasome inhibitors and apoptosis inducers in cancer cells," *Pharmaceutical Research*, vol. 26, no. 8, pp. 1874–1880, 2009.
- [143] B. Rajitha, A. Belalcazar, G. P. Nagaraju et al., "Inhibition of NF- κ B translocation by curcumin analogs induces G0/G1 arrest and downregulates thymidylate synthase in colorectal cancer," *Cancer Letters*, vol. 373, no. 2, pp. 227–233, 2016.
- [144] L. Cen, B. Hutzen, S. Ball et al., "New structural analogues of curcumin exhibit potent growth suppressive activity in human colorectal carcinoma cells," *BMC Cancer*, vol. 9, article 99, 2009.
- [145] Y.-S. Chiou, N. J.-L. Ma, S. Sang, C.-T. Ho, Y.-J. Wang, and M.-H. Pan, "Peracetylated (-)-epigallocatechin-3-gallate (AcEGCG) potently suppresses dextran sulfate sodium-induced colitis and colon tumorigenesis in mice," *Journal of Agricultural and Food Chemistry*, vol. 60, no. 13, pp. 3441–3451, 2012.
- [146] L. Actis-Goretta, L. J. Romanczyk, C. A. Rodriguez, C. Kwik-Urbe, and C. L. Keen, "Cytotoxic effects of digalloyl dimer pro-cyanidins in human cancer cell lines," *The Journal of Nutritional Biochemistry*, vol. 19, no. 12, pp. 797–e2, 2008.
- [147] Y. Mizushima, A. Saito, K. Horikawa et al., "Acylated catechin derivatives: Inhibitors of DNA polymerase and angiogenesis," *Frontiers in Bioscience - Elite*, vol. 3, no. 4, pp. 1337–1348, 2011.
- [148] K. Ahmed, Z.-L. Wei, Q.-L. Zhao et al., "Role of fatty acid chain length on the induction of apoptosis by newly synthesized catechin derivatives," *Chemico-Biological Interactions*, vol. 185, no. 3, pp. 182–188, 2010.
- [149] K. Polkowski, J. Popiołkiewicz, P. Krzeczynski et al., "Cytostatic and cytotoxic activity of synthetic genistein glycosides against human cancer cell lines," *Cancer Letters*, vol. 203, no. 1, pp. 59–69, 2004.
- [150] L. Sánchez-del-Campo, F. Otón, A. Tárraga, J. Cabezas-Herrera, S. Chazarra, and J. N. Rodríguez-López, "Synthesis and biological activity of a 3,4,5-trimethoxybenzoyl ester analogue of epicatechin-3-gallate," *Journal of Medicinal Chemistry*, vol. 51, no. 7, pp. 2018–2026, 2008.
- [151] H. Li, F. Zhu, H. Chen et al., "6-C-(E-phenylethenyl)-naringenin suppresses colorectal cancer growth by inhibiting cyclooxygenase-1," *Cancer Research*, vol. 74, no. 1, pp. 243–252, 2014.
- [152] H. Yoon, T. W. Kim, S. Y. Shin et al., "Design, synthesis and inhibitory activities of naringenin derivatives on human colon cancer cells," *Bioorganic & Medicinal Chemistry Letters*, vol. 23, no. 1, pp. 232–238, 2013.
- [153] C. M. Shin, D. H. Lee, A. Y. Seo et al., "Green tea extracts for the prevention of metachronous colorectal polyps among patients who underwent endoscopic removal of colorectal adenomas: A randomized clinical trial," *Clinical Nutrition*, 2016.
- [154] M. Nikš and M. Otto, "Towards an optimized MTT assay," *Journal of Immunological Methods*, vol. 130, no. 1, pp. 149–151, 1990.
- [155] R. D. Blumenthal, *Chemosensitivity*, vol. 110, Humana Press, New Jersey, 2005.
- [156] M. Feoktistova, P. Geserick, and M. Leverkus, "Crystal violet assay for determining viability of cultured cells," *Cold Spring Harbor Protocols*, vol. 2016, no. 4, pp. 343–346, 2016.
- [157] A. Faried, D. Kurnia, L. S. Faried et al., "Anticancer effects of gallic acid isolated from Indonesian herbal medicine, Phaleria macrocarpa (Scheff.) Boerl, on human cancer cell lines," *International Journal of Oncology*, vol. 30, no. 3, pp. 605–613, 2007.
- [158] J. S. Giftson, S. Jayanthi, and N. Nalini, "Chemopreventive efficacy of gallic acid, an antioxidant and anticarcinogenic polyphenol, against 1,2-dimethyl hydrazine induced rat colon carcinogenesis," *Investigational New Drugs*, vol. 28, no. 3, pp. 251–259, 2010.
- [159] S. Gunasekaran, K. Venkatachalam, and N. Namasivayam, "p-Methoxycinnamic acid, an active phenylpropanoid induces mitochondrial mediated apoptosis in HCT-116 human colon adenocarcinoma cell line," *Environmental Toxicology and Pharmacology*, vol. 40, no. 3, pp. 966–974, 2015.
- [160] K. Ho, L. S. Yazan, N. Ismail, and M. Ismail, "Apoptosis and cell cycle arrest of human colorectal cancer cell line HT-29 induced by vanillin," *Cancer Epidemiology*, vol. 33, no. 2, pp. 155–160, 2009.
- [161] Q. Zheng, Y. Hirose, N. Yoshimi et al., "Further investigation of the modifying effect of various chemopreventive agents on apoptosis and cell proliferation in human colon cancer cells," *Journal of Cancer Research and Clinical Oncology*, vol. 128, no. 10, pp. 539–546, 2002.
- [162] C. V. Rao, D. Desai, B. Kaul, S. Amin, and B. S. Reddy, "Effect of caffeic acid esters on carcinogen-induced mutagenicity and human colon adenocarcinoma cell growth," *Chemico-Biological Interactions*, vol. 84, no. 3, pp. 277–290, 1992.
- [163] N. Nasr Bouzaïene, S. Kilani Jaziri, H. Kovacic, L. Chekir-Ghedira, K. Ghedira, and J. Luis, "The effects of caffeic, coumaric and ferulic acids on proliferation, superoxide production, adhesion and migration of human tumor cells in vitro," *European Journal of Pharmacology*, vol. 766, pp. 99–105, 2015.
- [164] D. W. Choi, M. S. Lim, J. W. Lee et al., "The cytotoxicity of kahweol in HT-29 human colorectal cancer cells is mediated by apoptosis and suppression of heat shock protein 70 expression," *Biomolecules & Therapeutics*, vol. 23, no. 2, pp. 128–133, 2015.
- [165] L. R. Ferguson, S.-T. Zhu, and P. J. Harris, "Antioxidant and antigenotoxic effects of plant cell wall hydroxycinnamic acids in cultured HT-29 cells," *Molecular Nutrition & Food Research*, vol. 49, no. 6, pp. 585–593, 2005.
- [166] S. K. Jaganathan, E. Supriyanto, and M. Mandal, "Events associated with apoptotic effect of p-Coumaric acid in HCT-15 colon cancer cells," *World Journal of Gastroenterology*, vol. 19, no. 43, pp. 7726–7734, 2013.
- [167] M. V. Barni, M. J. Carlini, E. G. Cafferata, L. Puricelli, and S. Moreno, "Carnosic acid inhibits the proliferation and migration capacity of human colorectal cancer cells," *Oncology Reports*, vol. 27, no. 4, pp. 1041–1048, 2012.
- [168] O. Slaby, M. Svoboda, J. Michalek, and R. Vyzula, "MicroRNAs in colorectal cancer: translation of molecular biology into clinical application," *Molecular Cancer*, vol. 8, article 102, 2009.
- [169] B. S. Vinod, T. T. Maliekal, and R. J. Anto, "Phytochemicals as chemosensitizers: from molecular mechanism to clinical significance," *Antioxidants & Redox Signaling*, vol. 18, no. 11, pp. 1307–1348, 2013.
- [170] K. Yanagihara, A. Ito, T. Toge, and M. Numoto, "Antiproliferative effects of isoflavones on human cancer cell lines established from the gastrointestinal tract," *Cancer Research*, vol. 53, no. 23, pp. 5815–5821, 1993.
- [171] A. Puthli, R. Tiwari, and K. P. Mishra, "Biochanin A enhances the radiotoxicity in colon tumor cells in vitro," *Journal of Environmental Pathology, Toxicology and Oncology*, vol. 32, no. 3, pp. 189–203, 2013.
- [172] J. Huang, M. Xie, P. Gao et al., "Antiproliferative effects of formononetin on human colorectal cancer via suppressing cell growth in vitro and in vivo," *Process Biochemistry*, vol. 50, no. 6, Article ID 10369, pp. 912–917, 2015.
- [173] K. K.-W. Auyeung and J. K.-S. Ko, "Novel herbal flavonoids promote apoptosis but differentially induce cell cycle arrest in

- human colon cancer cell," *Investigational New Drugs*, vol. 28, no. 1, pp. 1–13, 2010.
- [174] J. Yu, Y. Cheng, L. Xie, and R. Zhang, "Effects of genistein and daidzein on membrane characteristics of HCT cells," *Nutrition and Cancer*, vol. 33, no. 1, pp. 100–104, 1999.
- [175] J. M. Guo, B. X. Xiao, D. H. Liu et al., "Biphasic effect of daidzein on cell growth of human colon cancer cells," *Food and Chemical Toxicology*, vol. 42, no. 10, pp. 1641–1646, 2004.
- [176] S. Y. Park, E. J. Kim, H.-K. Shin et al., "Induction of apoptosis in HT-29 colon cancer cells by phloretin," *Journal of Medicinal Food*, vol. 10, no. 4, pp. 581–586, 2007.
- [177] S.-P. Zhu, G. Liu, X.-T. Wu et al., "The effect of Phloretin on human $\gamma\delta$ T cells killing colon cancer SW-1116 cells," *International Immunopharmacology*, vol. 15, no. 1, pp. 6–14, 2013.
- [178] F. Ferkl, W. W. Huber, M. Filipič et al., "Xanthohumol, a prenylated flavonoid contained in beer, prevents the induction of preneoplastic lesions and DNA damage in liver and colon induced by the heterocyclic aromatic amine amino-3-methylimidazo[4,5-f]quinoline (IQ)," *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis*, vol. 691, no. 1-2, pp. 17–22, 2010.
- [179] L. Pan, H. Becker, and C. Gerhäuser, "Xanthohumol induces apoptosis in cultured 40-16 human colon cancer cells by activation of the death receptor- and mitochondrial pathway," *Molecular Nutrition & Food Research*, vol. 49, no. 9, pp. 837–843, 2005.
- [180] S. H. Lee, H. J. Kim, J. S. Lee, I.-S. Lee, and B. Y. Kang, "Inhibition of topoisomerase I activity and efflux drug transporters' expression by xanthohumol from hops," *Archives of Pharmacol Research*, vol. 30, no. 11, pp. 1435–1439, 2007.
- [181] Y. Wang, Y. Chen, J. Wang et al., "Xanthohumol, a prenylated chalcone derived from hops, suppresses cancer cell invasion through inhibiting the expression of CXCR4 chemokine receptor," *Current Molecular Medicine*, vol. 12, no. 2, pp. 153–162, 2012.
- [182] P. Totta, F. Acconcia, S. Leone, I. Cardillo, and M. Marino, "Mechanisms of naringenin-induced apoptotic cascade in cancer cells: involvement of estrogen receptor α and β signalling," *IUBMB Life*, vol. 56, no. 8, pp. 491–499, 2004.
- [183] H. R. Frydoonfar, D. R. McGrath, and A. D. Spigelman, "The variable effect on proliferation of a colon cancer cell line by the citrus fruit flavonoid Naringenin," *Colorectal Disease*, vol. 5, no. 2, pp. 149–152, 2003.
- [184] S. Aranganathan, J. Panneer Selvam, N. Sangeetha, and N. Nalini, "Modulatory efficacy of hesperetin (citrus flavanone) on xenobiotic-metabolizing enzymes during 1,2-dimethylhydrazine-induced colon carcinogenesis," *Chemico-Biological Interactions*, vol. 180, no. 2, pp. 254–261, 2009.
- [185] S. Aranganathan and N. Nalini, "Antiproliferative efficacy of hesperetin (Citrus Flavanoid) in 1,2-dimethylhydrazine-induced colon cancer," *Phytotherapy Research*, vol. 27, no. 7, pp. 999–1005, 2013.
- [186] K. Manigandan, R. L. Jayaraj, and N. Elangovan, "Taxifolin ameliorates 1,2-dimethylhydrazine induced cell proliferation and redox avulsions in mice colon carcinogenesis," *Biomedicine & Preventive Nutrition*, vol. 4, no. 4, pp. 499–509, 2014.
- [187] S. B. Lee, K. H. Cha, D. Selenge, A. Solongo, and C. W. Nho, "The chemopreventive effect of taxifolin is exerted through ARE-dependent gene regulation," *Biological & Pharmaceutical Bulletin*, vol. 30, no. 6, pp. 1074–1079, 2007.
- [188] K. Manigandan, D. Manimaran, R. L. Jayaraj, N. Elangovan, V. Dhivya, and A. Kaphle, "Taxifolin curbs NF- κ B-mediated Wnt/ β -catenin signaling via up-regulating Nrf2 pathway in experimental colon carcinogenesis," *Biochimie*, vol. 119, pp. 103–112, 2015.
- [189] M. Shimizu, A. Deguchi, J. T. E. Lim, H. Moriwaki, L. Kopelovich, and I. B. Weinstein, "(–)-Epigallocatechin gallate and polyphenon E inhibit growth and activation of the epidermal growth factor receptor and human epidermal growth factor receptor-2 signaling pathways in human colon cancer cells," *Clinical Cancer Research*, vol. 11, no. 7, pp. 2735–2746, 2005.
- [190] N. P. Seeram, Y. Zhang, and M. G. Nair, "Inhibition of proliferation of human cancer cells and cyclooxygenase enzymes by anthocyanidins and catechins," *Nutrition and Cancer*, vol. 46, no. 1, pp. 101–106, 2003.
- [191] J. Ju, J. Hong, J.-N. Zhou et al., "Inhibition of intestinal tumorigenesis in Apcmin/+ mice by (-)-epigallocatechin-3-gallate, the major catechin in green tea," *Cancer Research*, vol. 65, no. 22, pp. 10623–10631, 2005.
- [192] Y. Shirakami, M. Shimizu, H. Tsurumi, Y. Hara, T. Tanaka, and H. Moriwaki, "EGCG and polyphenon E attenuate inflammation-related mouse colon carcinogenesis induced by AOM plus DDS," *Molecular Medicine Reports*, vol. 1, no. 3, pp. 355–361, 2008.
- [193] G.-J. Du, Z. Zhang, X.-D. Wen et al., "Epigallocatechin gallate (EGCG) is the most effective cancer chemopreventive polyphenol in green tea," *Nutrients*, vol. 4, no. 11, pp. 1679–1691, 2012.
- [194] G. Peng, D. A. Dixon, S. J. Muga, T. J. Smith, and M. J. Wargovich, "Green tea polyphenol (-)-epigallocatechin-3-gallate inhibits cyclooxygenase-2 expression in colon carcinogenesis," *Molecular Carcinogenesis*, vol. 45, no. 5, pp. 309–319, 2006.
- [195] X. Zhang, K.-W. Min, J. Wimalasena, and S. J. Baek, "Cyclin D1 degradation and p21 induction contribute to growth inhibition of colorectal cancer cells induced by epigallocatechin-3-gallate," *Journal of Cancer Research and Clinical Oncology*, vol. 138, no. 12, pp. 2051–2060, 2012.
- [196] H. Jin, W. Gong, C. Zhang, and S. Wang, "Epigallocatechin gallate inhibits the proliferation of colorectal cancer cells by regulating Notch signaling," *OncoTargets and Therapy*, vol. 6, pp. 145–153, 2013.
- [197] M. Roomi, V. Ivanov, T. Kalinovsky, A. Niedzwiecki, and M. Rath, "Anticancer effect of lysine, proline, arginine, ascorbic acid and green tea extract on human renal adenocarcinoma line 786-0," *Oncology Reports*, 2006.
- [198] Y. D. Jung, M. S. Kim, B. A. Shin et al., "EGCG, a major component of green tea, inhibits tumour growth by inhibiting VEGF induction in human colon carcinoma cells," *British Journal of Cancer*, vol. 84, no. 6, pp. 844–850, 2001.
- [199] A. Sengupta, S. Ghosh, R. K. Das, S. Bhattacharjee, and S. Bhattacharya, "Chemopreventive potential of diallylsulfide, lycopene and theaflavin during chemically induced colon carcinogenesis in rat colon through modulation of cyclooxygenase-2 and inducible nitric oxide synthase pathways," *European Journal of Cancer Prevention*, vol. 15, no. 4, pp. 301–305, 2006.
- [200] S. Huang, N. Yang, Y. Liu et al., "Grape seed proanthocyanidins inhibit colon cancer-induced angiogenesis through suppressing the expression of VEGF and Ang1," *International Journal of Molecular Medicine*, vol. 30, no. 6, pp. 1410–1416, 2012.
- [201] Y.-J. Kim, H.-J. Park, S.-H. Yoon et al., "Anticancer effects of oligomeric proanthocyanidins on human colorectal cancer cell line, SNU-C4," *World Journal of Gastroenterology*, vol. 11, no. 30, pp. 4674–4678, 2005.

- [202] M. Da Silva, G. K. Jagers, S. V. Verstraeten, A. G. Erleijman, C. G. Fraga, and P. I. Oteiza, "Large procyanidins prevent bile-acid-induced oxidant production and membrane-initiated ERK1/2, p38, and Akt activation in Caco-2 cells," *Free Radical Biology & Medicine*, vol. 52, no. 1, pp. 151–159, 2012.
- [203] Y. Zhang, S. K. Vareed, and M. G. Nair, "Human tumor cell growth inhibition by nontoxic anthocyanidins, the pigments in fruits and vegetables," *Life Sciences*, vol. 76, no. 13, pp. 1465–1472, 2005.
- [204] J. Cvorovic, F. Tramer, M. Granzotto, L. Candussio, G. Decorti, and S. Passamonti, "Oxidative stress-based cytotoxicity of delphinidin and cyanidin in colon cancer cells," *Archives of Biochemistry and Biophysics*, vol. 501, no. 1, pp. 151–157, 2010.
- [205] D. Cooke, M. Schwarz, D. Boocock et al., "Effect of cyanidin-3-glucoside and an anthocyanin mixture from bilberry on adenoma development in the ApcMin mouse model of intestinal carcinogenesis - Relationship with tissue anthocyanin levels," *International Journal of Cancer*, vol. 119, no. 9, pp. 2213–2220, 2006.
- [206] A. Hagiwara, K. Miyashita, T. Nakanishi et al., "Pronounced inhibition by a natural anthocyanin, purple corn color, of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP)-associated colorectal carcinogenesis in male F344 rats pretreated with 1,2-dimethylhydrazine," *Cancer Letters*, vol. 171, no. 1, pp. 17–25, 2001.
- [207] S.-Y. Kang, N. P. Seeram, M. G. Nair, and L. D. Bourquin, "Tart cherry anthocyanins inhibit tumor development in ApcMin mice and reduce proliferation of human colon cancer cells," *Cancer Letters*, vol. 194, no. 1, pp. 13–19, 2003.
- [208] G. K. Harris, A. Gupta, R. G. Nines et al., "Effects of lyophilized black raspberries on azoxymethane-induced colon cancer and 8-hydroxy-2'-deoxyguanosine levels in the Fischer 344 rat," *Nutrition and Cancer*, vol. 40, no. 2, pp. 125–133, 2001.
- [209] G. Lala, M. Malik, C. Zhao et al., "Anthocyanin-rich extracts inhibit multiple biomarkers of colon cancer in rats," *Nutrition and Cancer*, vol. 54, no. 1, pp. 84–93, 2006.
- [210] S. Lim, J. Xu, J. Kim et al., "Role of anthocyanin-enriched purple-fleshed sweet potato p40 in colorectal cancer prevention," *Molecular Nutrition & Food Research*, vol. 57, no. 11, pp. 1908–1917, 2013.
- [211] V. Charepalli, L. Reddivari, R. Vadde, S. Walia, S. Radhakrishnan, and J. K. P. Vanamala, "Eugenia jambolana (Java plum) fruit extract exhibits anti-cancer activity against early stage human HCT-116 colon cancer cells and colon cancer stem cells," *Cancers*, vol. 8, no. 3, article no. 29, 2016.
- [212] W. Yi, J. Fischer, G. Krewer, and C. C. Akoh, "Phenolic compounds from blueberries can inhibit colon cancer cell proliferation and induce apoptosis," *Journal of Agricultural and Food Chemistry*, vol. 53, no. 18, pp. 7320–7329, 2005.
- [213] C. Zhao, M. M. Giusti, M. Malik, M. P. Moyer, and B. A. Magnuson, "Effects of commercial anthocyanin-rich on colonic cancer and nontumorigenic colonic cell growth," *Journal of Agricultural and Food Chemistry*, vol. 52, no. 20, pp. 6122–6128, 2004.
- [214] D. Y. Shin, J. N. Lu, G.-Y. Kim et al., "Anti-invasive activities of anthocyanins through modulation of tight junctions and suppression of matrix metalloproteinase activities in HCT-116 human colon carcinoma cells," *Oncology Reports*, vol. 25, no. 2, pp. 567–572, 2011.
- [215] J. W. Yun, W. S. Lee, M. J. Kim et al., "Characterization of a profile of the anthocyanins isolated from *Vitis coignetiae* Pulliat and their anti-invasive activity on HT-29 human colon cancer cells," *Food and Chemical Toxicology*, vol. 48, no. 3, pp. 903–909, 2010.
- [216] L.-S. Wang, C.-T. Kuo, S.-J. Cho et al., "Black raspberry-derived anthocyanins demethylate tumor suppressor genes through the inhibition of DNMT1 and DNMT3B in colon cancer cells," *Nutrition and Cancer*, vol. 65, no. 1, pp. 118–125, 2013.
- [217] S. Thomasset, D. P. Berry, H. Cai et al., "Pilot study of oral anthocyanins for colorectal cancer chemoprevention," *Cancer Prevention Research*, vol. 2, no. 7, pp. 625–633, 2009.
- [218] H.-F. Lu, Y.-L. Chen, J.-S. Yang et al., "Antitumor activity of capsaicin on human colon cancer cells in vitro and colo 205 tumor xenografts in vivo," *Journal of Agricultural and Food Chemistry*, vol. 58, no. 24, pp. 12999–13005, 2010.
- [219] R. Clark, J. Lee, and S.-H. Lee, "Synergistic anticancer activity of capsaicin and 3,3'-diindolylmethane in human colorectal cancer," *Journal of Agricultural and Food Chemistry*, vol. 63, no. 17, pp. 4297–4304, 2015.
- [220] M. Y. Kim, L. J. Trudel, and G. N. Wogan, "Apoptosis induced by capsaicin and resveratrol in colon carcinoma cells requires nitric oxide production and caspase activation," *Anticancer Research*, vol. 29, no. 10, pp. 3733–3740, 2009.
- [221] M. Y. Kyung, O. P. Jong, G.-Y. Kim et al., "Capsaicin induces apoptosis by generating reactive oxygen species and disrupting mitochondrial transmembrane potential in human colon cancer cell lines," *Cellular & Molecular Biology Letters*, vol. 14, no. 3, pp. 497–510, 2009.
- [222] S.-H. Lee, R. L. Richardson, R. H. Dashwood, and S. J. Baek, "Capsaicin represses transcriptional activity of β -catenin in human colorectal cancer cells," *The Journal of Nutritional Biochemistry*, vol. 23, no. 6, pp. 646–655, 2012.
- [223] H. O. Seon, S. K. Young, C. L. Sung, F. H. Yi, Y. C. In, and J. Y. Ho, "Dihydrocapsaicin (DHC), a saturated structural analog of capsaicin, induces autophagy in human cancer cells in a catalase-regulated manner," *Autophagy*, vol. 4, no. 8, pp. 1009–1019, 2008.
- [224] W. Guo, L. Nie, D. Wu et al., "Avenanthramides inhibit proliferation of human colon cancer cell lines in vitro," *Nutrition and Cancer*, vol. 62, no. 8, pp. 1007–1016, 2010.
- [225] A. Bishayee, "Cancer prevention and treatment with resveratrol: from rodent studies to clinical trials," *Cancer Prevention Research*, vol. 2, no. 5, pp. 409–418, 2009.
- [226] L. Tessitore, A. Davit, I. Sarotto, and G. Caderni, "Resveratrol depresses the growth of colorectal aberrant crypt foci by affecting bax and p21(CIP) expression," *Carcinogenesis*, vol. 21, no. 8, pp. 1619–1622, 2000.
- [227] M. Sengottuvelan, P. Viswanathan, and N. Nalini, "Chemopreventive effect of trans-resveratrol—a phytoalexin against colonic aberrant crypt foci and cell proliferation in 1,2-dimethylhydrazine induced colon carcinogenesis," *Carcinogenesis*, vol. 27, no. 5, pp. 1038–1046, 2006.
- [228] M. Sengottuvelan, R. Senthikumar, and N. Nalini, "Modulatory influence of dietary resveratrol during different phases of 1,2-dimethylhydrazine induced mucosal lipid-peroxidation, antioxidant status and aberrant crypt foci development in rat colon carcinogenesis," *Biochimica et Biophysica Acta (BBA) - General Subjects*, vol. 1760, no. 8, pp. 1175–1183, 2006.
- [229] M. Sengottuvelan and N. Nalini, "Dietary supplementation of resveratrol suppresses colonic tumour incidence in 1,2-dimethylhydrazine-treated rats by modulating biotransforming enzymes and aberrant crypt foci development," *British Journal of Nutrition*, vol. 96, no. 1, pp. 145–153, 2006.

- [230] Y. Schneider, B. Duranton, F. Gossé, R. Schleiffer, N. Seiler, and F. Raul, "Resveratrol inhibits intestinal tumorigenesis and modulates host-defense-related gene expression in an animal model of human familial adenomatous polyposis," *Nutrition and Cancer*, vol. 39, no. 1, pp. 102–107, 2001.
- [231] S. Sale, R. G. Tunstall, K. C. Ruparelia, G. A. Potter, W. P. Steward, and A. J. Gescher, "Comparison of the effects of the chemopreventive agent resveratrol and its synthetic analog trans 3,4,5,4'-tetramethoxystilbene (DMU-212) on adenoma development in the Apc^{Min+} mouse and cyclooxygenase-2 in human-derived colon cancer cells," *International Journal of Cancer*, vol. 115, no. 2, pp. 194–201, 2005.
- [232] C. C. Ziegler, L. Rainwater, J. Whelan, and M. F. McEntee, "Dietary Resveratrol Does Not Affect Intestinal Tumorigenesis in Apc Min/+ Mice," *Journal of Nutrition*, vol. 134, no. 1, pp. 5–10, 2004.
- [233] Z. G. Li, T. Hong, Y. Shimada et al., "Suppression of N-nitrosomethylbenzylamine (NMBA)-induced esophageal tumorigenesis in F344 rats by resveratrol," *Carcinogenesis*, vol. 23, no. 9, pp. 1531–1536, 2002.
- [234] H.-B. Zhou, J.-J. Chen, W.-X. Wang, J.-T. Cai, and Q. Du, "Anticancer activity of resveratrol on implanted human primary gastric carcinoma cells in nude mice," *World Journal of Gastroenterology*, vol. 11, no. 2, pp. 280–284, 2005.
- [235] M. E. Juan, I. Alfaras, and J. M. Planas, "Colorectal cancer chemoprevention by trans-resveratrol," *Pharmacological Research*, vol. 65, no. 6, pp. 584–591, 2012.
- [236] S. M. Saud, W. Li, N. L. Morris et al., "Resveratrol prevents tumorigenesis in mouse model of Kras activated sporadic colorectal cancer by suppressing oncogenic Kras expression," *Carcinogenesis*, vol. 35, no. 12, pp. 2778–2786, 2014.
- [237] Q. Ji, X. Liu, Z. F. Han et al., "Resveratrol suppresses epithelial-to-mesenchymal transition in colorectal cancer through TGF- β 1/Smads signaling pathway mediated Snail/E-cadherin expression," *BMC Cancer*, vol. 15, article 97, 2015.
- [238] C. Buhrmann, P. Shayan, P. Kraehe, B. Popper, A. Goel, and M. Shakibaei, "Resveratrol induces chemosensitization to 5-fluorouracil through up-regulation of intercellular junctions, epithelial-to-mesenchymal transition and apoptosis in colorectal cancer," *Biochemical Pharmacology*, vol. 98, no. 1, pp. 51–58, 2015.
- [239] Q. Ji, X. Liu, X. Fu et al., "Resveratrol inhibits invasion and metastasis of colorectal cancer cells via MALAT1 mediated Wnt/ β -catenin signal pathway," *PLoS ONE*, vol. 8, no. 11, Article ID e78700, 2013.
- [240] Z. Wang, L. Zhang, Z. Ni et al., "Resveratrol induces AMPK-dependent MDRI inhibition in colorectal cancer HCT116/L-OHP cells by preventing activation of NF- κ B signaling and suppressing cAMP-responsive element transcriptional activity," *Tumor Biology*, vol. 36, no. 12, pp. 9499–9510, 2015.
- [241] D. P. Chauhan, "Chemotherapeutic potential of curcumin for colorectal cancer," *Current Pharmaceutical Design*, vol. 8, no. 19, pp. 1695–1706, 2002.
- [242] I. Villegas, S. Sánchez-Fidalgo, and C. A. De La Lastra, "New mechanisms and therapeutic potential of curcumin for colorectal cancer," *Molecular Nutrition & Food Research*, vol. 52, no. 9, pp. 1040–1061, 2008.
- [243] B. Uzzan and R. Benamouzig, "Is Curcumin a Chemopreventive Agent for Colorectal Cancer?" *Current Colorectal Cancer Reports*, vol. 12, no. 1, pp. 35–41, 2016.
- [244] J. T. Piper, S. S. Singhal, M. S. Salameh, R. T. Torman, Y. C. Awasthi, and S. Awasthi, "Mechanisms of anticarcinogenic properties of curcumin: the effect of curcumin on glutathione linked detoxification enzymes in rat liver," *The International Journal of Biochemistry & Cell Biology*, vol. 30, no. 4, pp. 445–456, 1998.
- [245] H. Chen, Z.-S. Zhang, Y.-L. Zhang, and D.-Y. Zhou, "Curcumin inhibits cell proliferation by interfering with the cell cycle and inducing apoptosis in colon carcinoma cells," *Anticancer Research*, vol. 19, no. 5 A, pp. 3675–3680, 1999.
- [246] L. Li, B. Ahmed, K. Mehta, and R. Kurzrock, "Liposomal curcumin with and without oxaliplatin: effects on cell growth, apoptosis, and angiogenesis in colorectal cancer," *Molecular Cancer Therapeutics*, vol. 6, no. 4, pp. 1276–1282, 2007.
- [247] G. Mudduluru, J. N. George-William, S. Muppala et al., "Curcumin regulates miR-21 expression and inhibits invasion and metastasis in colorectal cancer," *Bioscience Reports*, vol. 31, no. 3, pp. 185–197, 2011.
- [248] M. I. James, C. Iwuji, G. Irving et al., "Curcumin inhibits cancer stem cell phenotypes in *ex vivo* models of colorectal liver metastases, and is clinically safe and tolerable in combination with FOLFOX chemotherapy," *Cancer Letters*, vol. 364, no. 2, pp. 135–141, 2015.
- [249] S. K. Sandur, A. Deorukhkar, M. K. Pandey et al., "Curcumin modulates the radiosensitivity of colorectal cancer cells by suppressing constitutive and inducible NF- κ B activity," *International Journal of Radiation Oncology • Biology • Physics*, vol. 75, no. 2, pp. 534–542, 2009.
- [250] M. A. Encalada, K. M. Hoyos, S. Rehecho et al., "Anti-proliferative effect of *Melissa officinalis* on human colon cancer cell line," *Plant Foods for Human Nutrition*, vol. 66, no. 4, pp. 328–334, 2011.
- [251] K. Venkatachalam, S. Gunasekaran, V. A. S. Jesudoss, and N. Namasivayam, "The effect of rosmarinic acid on 1,2-dimethylhydrazine induced colon carcinogenesis," *Experimental and Toxicologic Pathology*, vol. 65, no. 4, pp. 409–418, 2013.
- [252] C. P. R. Xavier, C. F. Lima, M. Fernandes-Ferreira, and C. Pereira-Wilson, "*Salvia fruticosa*, *Salvia officinalis*, and rosmarinic acid induce apoptosis and inhibit proliferation of human colorectal cell lines: the role in MAPK/ERK pathway," *Nutrition and Cancer*, vol. 61, no. 4, pp. 564–571, 2009.
- [253] Y. Xu, G. Xu, L. Liu, D. Xu, and J. Liu, "Anti-invasion effect of rosmarinic acid via the extracellular signal-regulated kinase and oxidation-reduction pathway in Ls174-T cells," *Journal of Cellular Biochemistry*, vol. 111, no. 2, pp. 370–379, 2010.
- [254] C.-B. Lin, C.-C. Lin, and G. J. Tsay, "6-gingerol inhibits growth of colon cancer cell LoVo via induction of G2/M arrest," *Evidence-Based Complementary and Alternative Medicine*, vol. 2012, Article ID 326096, 2012.
- [255] C.-H. Jeong, A. M. Bode, A. Pugliese et al., "[6]-Gingerol suppresses colon cancer growth by targeting leukotriene A₄ hydrolase," *Cancer Research*, vol. 69, no. 13, pp. 5584–5591, 2009.
- [256] S. H. Lee, M. Cekanova, and J. B. Seung, "Multiple mechanisms are involved in 6-gingerol-induced cell growth arrest and apoptosis in human colorectal cancer cells," *Molecular Carcinogenesis*, vol. 47, no. 3, pp. 197–208, 2008.
- [257] M.-H. Pan, M.-C. Hsieh, J.-M. Kuo et al., "6-Shogaol induces apoptosis in human colorectal carcinoma cells via ROS production, caspase activation, and GADD 153 expression," *Molecular Nutrition & Food Research*, vol. 52, no. 5, pp. 527–537, 2008.
- [258] J. N. Losso, R. R. Bansode, A. Trappey II, H. A. Bawadi, and R. Truax, "In vitro anti-proliferative activities of ellagic acid," *The Journal of Nutritional Biochemistry*, vol. 15, no. 11, pp. 672–678, 2004.

- [259] S. Umesalma and G. Sudhandiran, "Differential inhibitory effects of the polyphenol ellagic acid on inflammatory mediators NF- κ B, iNOS, COX-2, TNF- α , and IL-6 in 1,2-dimethylhydrazine-induced rat colon carcinogenesis," *Basic & Clinical Pharmacology & Toxicology*, vol. 107, no. 2, pp. 650–655, 2010.
- [260] S. Umesalma and G. Sudhandiran, "Ellagic acid prevents rat colon carcinogenesis induced by 1, 2 dimethyl hydrazine through inhibition of AKT-phosphoinositide-3 kinase pathway," *European Journal of Pharmacology*, vol. 660, no. 2-3, pp. 249–258, 2011.
- [261] A. González-Sarrías, M. Á. Núñez-Sánchez, J. Tomé-Carneiro, F. A. Tomás-Barberán, M. T. García-Conesa, and J. C. Espín, "Comprehensive characterization of the effects of ellagic acid and urolithins on colorectal cancer and key-associated molecular hallmarks: MicroRNA cell specific induction of CDKN1A (p21) as a common mechanism involved," *Molecular Nutrition & Food Research*, vol. 60, no. 4, pp. 701–716, 2016.
- [262] M. Á. Núñez-Sánchez, A. Karmokar, A. González-Sarrías et al., "In vivo relevant mixed urolithins and ellagic acid inhibit phenotypic and molecular colon cancer stem cell features: A new potentiality for ellagitannin metabolites against cancer," *Food and Chemical Toxicology*, vol. 92, pp. 8–16, 2016.
- [263] M. Serraino and L. U. Thompson, "Flaxseed supplementation and early markers of colon carcinogenesis," *Cancer Letters*, vol. 63, no. 2, pp. 159–165, 1992.
- [264] A.-M. Pajari, A. I. Smeds, S. I. Oikarinen, P. C. Eklund, R. E. Sjöholm, and M. Mutanen, "The plant lignans matairesinol and secoisolariciresinol administered to Min mice do not protect against intestinal tumor formation," *Cancer Letters*, vol. 233, no. 2, pp. 309–314, 2006.
- [265] J.-H. Yoo, H. J. Lee, K. Kang et al., "Lignans inhibit cell growth via regulation of Wnt/ β -catenin signaling," *Food and Chemical Toxicology*, vol. 48, no. 8-9, pp. 2247–2252, 2010.
- [266] T.-C. Hour, J. Chen, C.-Y. Huang, J.-Y. Guan, S.-H. Lu, and Y.-S. Pu, "Curcumin enhances cytotoxicity of chemotherapeutic agents in prostate cancer cells by inducing p21WAF1/CIP1 and C/EBP β expressions and suppressing NF- κ B activation," *The Prostate*, vol. 51, no. 3, pp. 211–218, 2002.
- [267] M. Notarbartolo, P. Poma, D. Perri, L. Dusonchet, M. Cervello, and N. D'Alessandro, "Antitumor effects of curcumin, alone or in combination with cisplatin or doxorubicin, on human hepatic cancer cells. Analysis of their possible relationship to changes in NF- κ B activation levels and in IAP gene expression," *Cancer Letters*, vol. 224, no. 1, pp. 53–65, 2005.
- [268] F. H. Sarkar and Y. Li, "Using chemopreventive agents to enhance the efficacy of cancer therapy," *Cancer Research*, vol. 66, no. 7, pp. 3347–3350, 2006.
- [269] A. C. Bharti, N. Donato, S. Singh, and B. B. Aggarwal, "Curcumin (diferuloylmethane) down-regulates the constitutive activation of nuclear factor- κ B and I κ B α kinase in human multiple myeloma cells, leading to suppression of proliferation and induction of apoptosis," *Blood*, vol. 101, no. 3, pp. 1053–1062, 2003.
- [270] C. K. Singh, J. George, and N. Ahmad, "Resveratrol-based combinatorial strategies for cancer management," *Annals of the New York Academy of Sciences*, vol. 1290, no. 1, pp. 113–121, 2013.
- [271] A. K. Garg, T. A. Buchholz, and B. B. Aggarwal, "Chemosensitization and radiosensitization of tumors by plant polyphenols," *Antioxidants & Redox Signaling*, vol. 7, no. 11-12, pp. 1630–1647, 2005.
- [272] J. Francy-Guilford and J. M. Pezzuto, "Mechanisms of cancer chemopreventive agents: A perspective," *Planta Medica*, vol. 74, no. 13, pp. 1644–1650, 2008.
- [273] M. Fantini, M. Benvenuto, L. Masuelli et al., "In Vitro and in Vivo antitumoural effects of combinations of polyphenols, or polyphenols and anticancer drugs: perspectives on cancer Treatment," *International Journal of Molecular Sciences*, vol. 16, no. 5, pp. 9236–9282, 2015.
- [274] V. Aires, E. Limagne, A. K. Cotte, N. Latruffe, F. Ghiringhelli, and D. Delmas, "Resveratrol metabolites inhibit human metastatic colon cancer cells progression and synergize with chemotherapeutic drugs to induce cell death," *Molecular Nutrition & Food Research*, vol. 57, no. 7, pp. 1170–1181, 2013.
- [275] M. Shakibaei, A. Mobasheri, C. Lueders, F. Busch, P. Shayan, and A. Goel, "Curcumin enhances the effect of chemotherapy against colorectal cancer cells by inhibition of NF- κ B and Src protein kinase signaling pathways," *PLoS ONE*, vol. 8, no. 2, Article ID e57218, 2013.
- [276] B. Du, L. Jiang, Q. Xia, and L. Zhong, "Synergistic inhibitory effects of curcumin and 5-fluorouracil on the growth of the human colon cancer cell line HT-29," *Chemotherapy*, vol. 52, no. 1, pp. 23–28, 2005.
- [277] J.-T. Hwang, J. Ha, and O. J. Park, "Combination of 5-fluorouracil and genistein induces apoptosis synergistically in chemo-resistant cancer cells through the modulation of AMPK and COX-2 signaling pathways," *Biochemical and Biophysical Research Communications*, vol. 332, no. 2, pp. 433–440, 2005.
- [278] G. G. Hillman, Y. Wang, O. Kucuk et al., "Genistein potentiates inhibition of tumor growth by radiation in a prostate cancer orthotopic model," *Molecular Cancer Therapeutics*, vol. 3, no. 10, pp. 1271–1279, 2004.
- [279] R. L. Goodman, A. Grann, P. Saracco, and M. F. Needham, "Genistein, a tyrosine kinase inhibitor, enhanced radiosensitivity in human esophageal cancer cell lines in vitro: Possible involvement of inhibition of survival signal transduction pathways," *International Journal of Radiation Oncology • Biology • Physics*, vol. 50, no. 1, pp. 195–201, 2001.
- [280] I. Zoberi, C. M. Bradbury, H. A. Curry et al., "Radiosensitizing and anti-proliferative effects of resveratrol in two human cervical tumor cell lines," *Cancer Letters*, vol. 175, no. 2, pp. 165–173, 2002.
- [281] D. Chendil, R. S. Ranga, D. Meigooni, S. Sathishkumar, and M. M. Ahmed, "Curcumin confers radiosensitizing effect in prostate cancer cell line PC-3," *Oncogene*, vol. 23, no. 8, pp. 1599–1607, 2004.
- [282] J. George, M. Singh, A. K. Srivastava et al., "Resveratrol and black tea polyphenol combination synergistically suppress mouse skin tumors growth by inhibition of activated MAPKs and p53," *PLoS ONE*, vol. 6, no. 8, Article ID e23395, 2011.
- [283] A. Mallhotra, P. Nair, and D. K. Dhawan, "Curcumin and resveratrol synergistically stimulate p21 and regulate cox-2 by maintaining adequate zinc levels during lung carcinogenesis," *European Journal of Cancer Prevention*, vol. 20, no. 5, pp. 411–416, 2011.
- [284] C. E. Harper, L. M. Cook, B. B. Patel et al., "Genistein and resveratrol, alone and in combination, suppress prostate cancer in SV-40 tag rats," *The Prostate*, vol. 69, no. 15, pp. 1668–1682, 2009.
- [285] A. Khafif, S. P. Schantz, T.-C. Chou, D. Edelstein, and P. G. Sacks, "Quantitation of chemopreventive synergism between

- (-)-epigallocatechin-3-gallate and curcumin in normal, premalignant and malignant human oral epithelial cells," *Carcinogenesis*, vol. 19, no. 3, pp. 419–424, 1998.
- [286] S. U. Mertens-Talcott and S. S. Percival, "Ellagic acid and quercetin interact synergistically with resveratrol in the induction of apoptosis and cause transient cell cycle arrest in human leukemia cells," *Cancer Letters*, vol. 218, no. 2, pp. 141–151, 2005.
- [287] A. P. Majumdar, S. Banerjee, J. Nautiyal et al., "Curcumin synergizes with resveratrol to inhibit colon cancer," *Nutrition and Cancer*, vol. 61, no. 4, pp. 544–553, 2009.
- [288] C. Dimarco-Crook and H. Xiao, "Diet-based strategies for cancer chemoprevention: The role of combination regimens using dietary bioactive components," *Annual Review of Food Science and Technology*, vol. 6, pp. 505–526, 2015.
- [289] U. Lewandowska, S. Gorlach, K. Owczarek, E. Hrabec, and K. Szewczyk, "Synergistic interactions between anticancer chemotherapeutics and phenolic compounds and anticancer synergy between polyphenols," *Postepy Higieny i Medycyny Doswiadczalnej*, vol. 68, pp. 528–540, 2014.
- [290] K. R. Patel, E. Scott, V. A. Brown, A. J. Gescher, W. P. Steward, and K. Brown, "Clinical trials of resveratrol," *Annals of the New York Academy of Sciences*, vol. 1215, no. 1, pp. 161–169, 2011.
- [291] V. Ruiz de Porras, S. Bystrup, A. Martínez-Cardús et al., "Curcumin mediates oxaliplatin-acquired resistance reversion in colorectal cancer cell lines through modulation of CXCL12/Chemokine/NF- κ B signalling pathway," *Scientific Reports*, vol. 6, article 24675, 2016.
- [292] M. Mathonnet, A. Perraud, N. Christou et al., "Hallmarks in colorectal cancer: angiogenesis and cancer stem-like cells," *World Journal of Gastroenterology*, vol. 20, no. 15, pp. 4189–4196, 2014.
- [293] M. Kakarala, D. E. Brenner, H. Korkaya et al., "Targeting breast stem cells with the cancer preventive compounds curcumin and piperine," *Breast Cancer Research and Treatment*, vol. 122, no. 3, pp. 777–785, 2010.
- [294] M. T. E. Montales, O. M. Rahal, J. Kang et al., "Repression of mammosphere formation of human breast cancer cells by soy isoflavone genistein and blueberry polyphenolic acids suggests diet-mediated targeting of cancer stem-like/progenitor cells," *Carcinogenesis*, vol. 33, no. 3, pp. 652–660, 2012.
- [295] S. Shankar, D. Nall, S.-N. Tang et al., "Resveratrol inhibits pancreatic cancer stem cell characteristics in human and KrasG12D transgenic mice by inhibiting pluripotency maintaining factors and epithelial-mesenchymal transition," *PLoS ONE*, vol. 6, no. 1, Article ID e16530, 2011.
- [296] W. Zhou, G. Kallifatidis, B. Baumann et al., "Dietary polyphenol quercetin targets pancreatic cancer stem cells," *International Journal of Oncology*, vol. 37, no. 3, pp. 551–561, 2010.