

# Comparative study on the bioavailability of peptide extracts from Jeju black pigs and three-way crossbred pigs

Jae Won Jeong<sup>1</sup>, Ermie Mariano Jr<sup>1</sup>, Da Young Lee<sup>1</sup>, Seung Yun Lee<sup>2</sup>, Jae Hyeon Kim<sup>1</sup>, Seung Hyeon Yun<sup>1</sup>, Juhyun Lee<sup>1</sup>, Jinmo Park<sup>1</sup>, Yeongwoo Choi<sup>1</sup>, Dahee Han<sup>1</sup>, Jin Soo Kim<sup>1</sup>, Sun Jin Hur<sup>1\*</sup>

<sup>1</sup>Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea

<sup>2</sup>Division of Animal Science, Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea



Received: May 26, 2023

Revised: Jul 5, 2023

Accepted: Aug 10, 2023

## \*Corresponding author

Sun Jin Hur

Department of Animal Science and Technology, Chung-Ang University, Anseong 17546, Korea.

Tel: +82-31-670-4673

E-mail: hursj@cau.ac.kr

Copyright © 2024 Korean Society of Animal Sciences and Technology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

## ORCID

Jae Won Jeong

<https://orcid.org/0000-0001-5240-1875>

Ermie Mariano Jr

<https://orcid.org/0000-0003-2630-4603>

Da Young Lee

<https://orcid.org/0000-0002-3172-0815>

Seung Yun Lee

<https://orcid.org/0000-0002-8861-6517>

Jae Hyeon Kim

<https://orcid.org/0000-0003-1174-4737>

Seung Hyeon Yun

<https://orcid.org/0000-0002-9940-2960>

Juhyun Lee

<https://orcid.org/0000-0001-6777-4447>

Jinmo Park

<https://orcid.org/0009-0004-9626-1025>

## Abstract

This study aimed to compare the changes in the bioactivities of peptide extracts (< 10 kDa) obtained from Jeju black pigs (JBP) and three-way crossbred pigs (Landrace × Yorkshire × Duroc, LYD) before and after digestion. The results showed that the loin peptide extracts of JBP maintained high 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical scavenging activity after *in vitro* digestion. However, the iron chelating activity and antihypertensive activity of all peptide extracts were decreased. This study suggested that the peptide extracts produced through alkaline-AK digestion have sufficiently high antioxidant and antihypertensive activities; however, these activities were reduced after *in vitro* digestion. Meanwhile, the JBP loin and ham peptide extracts promoted high superoxide dismutase (SOD) activity than that of LYD when administered to mice. Furthermore, the ham peptide extracts of JBP showed a relatively high antihypertensive activity in mice. Therefore, it is deemed that these peptide extracts from JBP are more bioactive than that of LYD, and can be used as bioactive materials.

**Keywords:** Jeju black pigs, Peptide extracts, Antioxidant activity, Antihypertensive activity

## INTRODUCTION

Currently, the Korean pork market is dominated by three-way crossbred pigs (Landrace × Yorkshire × Duroc; LYD). On the other hand, the Jeju black pig (JBP) accounts for only 1.2% of the Korean pork market; however, Korean consumers prefer JBP owing to its higher muscle fat content and redness compared with LYD [1]. In addition, JBP has a higher essential fatty acid content and better overall taste compared with LYD [2]. Because JBP, whose breeding size continues to increase, is recognized as high-quality pork and domestic consumers increasingly prefer it, comparing JBP with LYD to identify differentiated characteristics is essential. Furthermore, there is a need to identify the excellence of JBP, which accounts for only 1.2% of the Korean pork market. Previous studies have compared the carcass

Yeongwoo Choi  
<https://orcid.org/0009-0000-1882-4890>  
Dahee Han  
<https://orcid.org/0009-0005-6423-3414>  
Jin Soo Kim  
<https://orcid.org/0009-0007-7974-7885>  
Sun Jin Hur  
<https://orcid.org/0000-0001-9386-5852>

#### Competing interests

No potential conflict of interest relevant to this article was reported.

#### Funding sources

This work was supported by Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET) through High Value-added Food Technology Development Program, funded by Ministry of Agriculture, Food and Rural Affairs (MAFRA) (322008-5). This research was supported by the Chung-Ang University Graduated Research Scholarship in 2023.

#### Acknowledgements

Not applicable.

#### Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

#### Authors' contributions

Conceptualization: Hur SJ.  
Data curation: Jeong JW, Mariano E, Lee J, Han D, Kim JS.  
Formal analysis: Jeong JW, Lee DY, Lee SY, Yun SH.  
Methodology: Jeong JW, Kim JH.  
Software: Choi Y.  
Validation: Jeong JW, Park J, Choi Y, Han D.  
Investigation: Jeong JW, Kim JH, Yun SH.  
Writing - original draft: Jeong JW, Mariano E.  
Writing - review & editing: Jeong JW, Mariano E, Lee DY, Lee SY, Kim JH, Yun SH, Lee J, Park J, Choi Y, Han D, Kim JS, Hur SJ.

#### Ethics approval and consent to participate

All procedures related to animal testing were approved by the Institutional Animal Care and Use Committee (IACUC) of the Chung-Ang University Research Institute (approval number: 2021-00038).

and meat characteristics of JBP and LYD [1,2]; however, comparative studies on the changes in the bioactivities of pork peptides due to various physiological processes, including digestion, are insufficient.

In general, pork contains bioactive peptides that regulate various biological actions and have beneficial health effects [3,4]. Peptides that exhibit various bioactive functions, such as antioxidant, antihypertension, antithrombosis, and antibacterial functions, have been derived from the myofibrillar proteins of pigs [5–8]. These bioactive peptides can be efficiently extracted from proteins using enzymes such as papain, bromelain, ficin, and alkaline protease [9–11]. Interestingly, peptides with a small molecular weight (< 10 kDa) have higher antioxidant and antihypertensive activities than peptides with a large molecular weight [12,13]. However, research on the digestion-induced changes in the bioactivities of pork peptides is insufficient. Therefore, this study was conducted to compare and analyze the *in vitro*-digestion-induced changes in the antioxidant and antihypertensive activities of peptide extracts (< 10 kDa) derived (using alkaline-AK enzymes) from the loin and ham myofibrillar proteins of JBP and LYD. This study also aimed to ascertain and compare the benefits of these peptides to experimental animals (mice) by *in vivo* experiment.

## MATERIALS AND METHODS

### Chemical materials


$\alpha$ -Amylase (hog pancreas), bile extract (porcine), lipase (porcine pancreas), mucin (porcine stomach type II), pepsin (porcine gastric mucosa), uric acid, 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (ABTS), 2,2-diphenyl-1-picrylhydrazyl (DPPH), potassium persulfate, and 3-(2-pyridyl)-5,6-diphenyl-1,2,4-triazine-*p,p'*-disulfonic acid monosodium salt hydrate (ferrozine) were purchased from Sigma-Aldrich (St. Louis, MO, USA). Bovine serum albumin was obtained from Biosesang (Seongnam, Korea) and pancreatin (porcine pancreas) was purchased from Tokyo Chemical Industry (Tokyo, Japan).

### Experimental animals

All procedures related to animal testing were approved by the Institutional Animal Care and Use Committee (IACUC) of the Chung-Ang University Research Institute (Approval number: 2021-00038) and conducted at the CK-II Specialized Animal Experiment Center, Chung-Ang University. A total of 30 ICR mice (female, eight-week-old) were purchased from Orient Bio (Seongnam, Korea). During the two-week adaptation period, a pellet-type general feed was provided *ad libitum* and the breeding environment was maintained at  $22 \pm 2^\circ\text{C}$  temperature,  $60 \pm 5\%$  humidity, and a 12 h photoperiod (07:00–19:00).

### Peptide extraction

Fig. 1 shows the process for obtaining peptide extracts from the loin and ham of JBP and LYD. JBP loin and ham were obtained from pigs raised in Jeju Island, and trademarked JBP products were purchased and used. LYD loin and ham were purchased from three wholesale stores (Asan and Goyang, Korea). Each sample was sampled at least 5 individuals for meat part. After grinding 2 kg of each sample for 1 min, total 10 L of distilled water compare to 2 kg of each sample was added and the samples were washed 10 times to remove blood and fat as much as possible. 400 g of the sample was homogenized using 4 L of 0.04 M phosphate buffer (PBS, pH 7.4) and then centrifuged at  $1,977\times g$  and  $4^\circ\text{C}$  for 15 min. The myofibrillar protein thus obtained was hydrolyzed using 0.2% alkaline-AK at  $60^\circ\text{C}$  and pH 11 for 2 h. Alkaline-AK (180–200 KU/g solid) was obtained by fermenting soybean meal inoculated with *Bacillus methyilotrophicus* and then extracting



**Fig. 1. Schematic diagram of the procedure for obtaining myofibrillar proteins and peptide extracts (< 10 kDa) from JBP and LYD.** JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace × Yorkshire × Duroc).

from the fermented soybean meal. The protein hydrolysate was heated at 80 °C for 15 min to deactivate the enzyme activity and dried at 55 °C for 24 h. After drying, a peptide extract of < 10 kDa was obtained through a central filter unit (Amicon® Ultra; Merck Millipore, Burlington, MA, USA). Finally, the peptide extract was freeze-dried at -70 °C for 72 h.

## Analysis of digestibility

### *In vitro* digestion

The *in vitro* digestion experiment was performed using the method given by Lee et al. [14], with slight modifications. Table 1 shows the constituents and concentrations of digestive enzymes and inorganic and organic solvents used for conducting the *in vitro* digestion experiment [14,15]. To each peptide extract sample (120 mg), 2 mL of saliva was added and the mixture was stirred in a water bath (37 °C, 150 rpm) for 5 min. Subsequently, 4 mL of gastric juice was added to the mixture and allowed to react for 2 h under the same conditions. Then, 4 mL of the small intestine solution and 2 mL of the bile solution were added, respectively, and allowed to react in a water bath at 37 °C and 150 rpm for 2 h to end the *in vitro* digestion process. In all *in vitro* digestion experiments, a control containing only digestive enzymes and a control containing only samples was prepared by replacing the sample or the digestive enzymes with distilled water. After the digestion process, all samples were centrifuged for 20 min at 13,000×g and 4 °C, and the supernatant was taken and used

**Table 1.** Constituents and concentrations of synthetic digestive juices used in the *in vitro* digestion model<sup>1)</sup>

	Saliva (mouth step)	Gastric juice (stomach step)	Duodenal juice (small intestine step)	Bile juice (small intestine step)
Inorganic and organic components	1.7 mL NaCl <sup>2)</sup> (175.3 g/L) <sup>3)</sup> 8 mL Urea (25 g/L) 15 mg Uric acid	6.5 mL HCl (37 g/L) 18 mL CaCl <sub>2</sub> ·2H <sub>2</sub> O (22.2 g/L) 1 g Bovine serum albumin	6.3 mL KCl (89.6 g/L) 9 mL CaCl <sub>2</sub> ·2H <sub>2</sub> O (22.2 g/L) 1 g Bovine serum albumin	69.3 mL NaHCO <sub>3</sub> (84.7 g/L) 10 mL CaCl <sub>2</sub> ·2H <sub>2</sub> O (22.2 g/L) 1.8 g Bovine serum albumin 30 g Bile
Enzymes	290 mg α-Amylase 25 mg Mucin	2.5 g Pepsin 3 g Mucin	9 g Pancreatin 1.5 g Lipase	
pH	6.8 ± 0.2	1.50 ± 0.02	8.0 ± 0.2	7.0 ± 0.2

<sup>1)</sup>After mixing all components (inorganic components, organic components, and enzymes), the volume was made up to 500 mL with distilled water. If necessary, the pH of the digestive juices was adjusted to the appropriate value.

<sup>2)</sup>The numbers indicate the concentrations of chemicals used to make the digestive juices.

<sup>3)</sup>The numbers in parentheses indicate the concentrations of inorganic or organic components per liter of distilled water.

as the final sample.

### ***In vivo* digestion**

A total of 30 eight-week-old female ICR mice were separated into each cages by mass-based randomize block design. After a two-week adaptation period, the *in vivo* digestion experiment was conducted for 21 d; thus, the total duration of the experiment was 35 d. Diets for the *in vivo* digestion experiments of peptide extracts were limited based on the daily feed intake (5 g) of general ICR mice, and peptide extracts (800 mg/kg) were administered orally once a day (at the same time every day).

### **Analysis of molecular weight**

Gel permeation chromatography was conducted using an ACQUITY APC System (Waters Corporation, Milford, MA, USA) with an Xbridge Protein BEH SEC column (150 × 7.8 mm, 3.5 μm) (Waters Corporation). Deionized water (A) and MeOH (B) were used as the mobile phase solvents; the solvent composition was set to 90% A and 10% B. The flow rate was 0.7 mL/min, and a 10 μL sample was injected and analyzed for 15 min using a reactive index (RI) detector.

### **Analysis of free amino acid composition**

Sample pretreatment for the analysis of free amino acid composition was performed using the method given by Enda et al. [16]. Briefly, 1 mL of a 5% TCA solution was added to 1 mL of each sample (diluted at an appropriate concentration) and vortexed, and the protein was precipitated by centrifuging at 13,572×g for 20 min. Thereafter, 2 mL of the supernatant was taken and 4 mL of n-hexane solution was added to it. The mixture was shaken for 10 min and then centrifuged at 1,977×g and 20 °C for 20 min. After centrifugation, the lower layer solution was obtained, and the process of centrifuging at 20 °C for 20 min at 13,572×g was repeated twice to remove the remaining n-hexane solution. Finally, all samples were filtered using a 0.20 μm syringe filter and analyzed using a Hitachi L-8900 amino acid analyzer (Hitachi High-Tech, Tokyo, Japan); the detailed analysis conditions are presented in Table 2.

### ***In vitro* analysis of bioavailability**

#### ***Analysis of 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) radical scavenging activity***

The ABTS assay was performed according to the method given by Re et al. [17], with slight modifications. The ABTS stock solution was prepared by dissolving 38.41 mg of 2,2'-azino-bis

**Table 2.** The conditions maintained in the amino acid analyzer for studying the changes in free amino acid composition

Parameter	Condition
Column	Hitachi HPLC packed ion exchange column (#2622PF)
Mobile phase	L-8900 buffer solution PF-1, 2, 3, 4, RG
Flow rate	Buffer solution: 0.35 mL/min Ninhydrin solution: 0.30 mL/min
Detection wavelength	440 nm, 570 nm
Temperatures	Reaction coil: 135°C Column: 30°C–70°C
Injection volume	20 µL

(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt (7 mM) and 6.615 mg of potassium persulfate (2.45 mM) in 10 mL of distilled water, mixing and incubating the mixture in a dark room for 12 h. The prepared ABTS stock solution was diluted with MeOH and used as the ABTS working solution; the absorbance of the diluted solution was  $0.7 \pm 0.02$  at 734 nm. Twenty microliters of each sample and 180 µL of ABTS working solution were added to a 96-well plate and incubated in a dark room for 10 min. Subsequently, their absorbance was measured at 734 nm using a microplate reader (Spectramax® 190; Molecular Devices, San Jose, CA, USA). The ABTS radical scavenging activity was calculated as follows:

$$\text{ABTS scavenging activity (\%)} = \left( 1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}} \right) \times 100$$

#### ***Analysis of 2,2-diphenyl-1-picrylhydrazyl radical scavenging activity***

The DPPH assay was performed using the method given by Tepe et al. [18], with slight modifications. The DPPH solution was prepared by dissolving 2 mg of 2,2-diphenyl-1-picrylhydrazyl in 25 mL of MeOH. One hundred microliters of each sample and 100 µL of DPPH solution were added to a 96-well plate and incubated in a dark room for 30 min. Subsequently, their absorbance was measured at 517 nm. The DPPH radical scavenging activity was calculated as follows:

$$\text{DPPH scavenging activity (\%)} = \left( 1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}} \right) \times 100$$

#### ***Analysis of iron chelating activity***

The iron chelating assay was performed using the method given by Dinis et al. [19], with slight modifications. Distilled water was added to prepare 2 mM iron (II) chloride and 5 mM ferrozine. The prepared samples (200 µL) were placed in 1.5 mL tubes, and 200 µL of 2 mM iron (II) chloride and 40 µL of 5 mM ferrozine was sequentially added to the tubes. Thereafter, the tubes were vortexed and incubated at 20°C–25°C for 10 min. Finally, the absorbance of the samples was measured at 562 nm. The iron chelating activity was calculated as follows:

$$\text{Iron chelating activity (\%)} = \left( 1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}} \right) \times 100$$

**Analysis of reducing power**

The reducing power assay was conducted according to the method given by Oyaizu [20]. Iron (III) chloride (1 mg/mL), TCA (100 mg/mL), and potassium hexacyanoferrate (III) (10 mg/mL) were prepared using distilled water as a solvent. The samples (100 µL) were taken in 1.5 mL tubes, and 100 µL each of 0.2 M sodium phosphate buffer (pH 6.6) and potassium hexacyanoferrate (III) was added to the tubes and incubated in a dark room at 50 °C for 20 min. Subsequently, 100 µL of TCA was added and the tubes were centrifuged at 9,425×g for 1 min. The supernatant (200 µL) obtained was mixed with 200 µL of distilled water and 40 µL of iron (III) chloride, and the absorbance of the solution was measured at 700 nm.

**Analysis of angiotensin-converting enzyme (ACE) inhibitory activity**

This experiment was conducted using the method given by Cushman & Cheung [21], with some modifications. To prepare the ACE solution, 0.1 M sodium borate buffer (pH 8.3) was prepared by properly mixing sodium tetraborate and boric acid, and 0.5 M sodium chloride was added to it. Subsequently, lung acetone powder from rabbit was extracted by stirring at a concentration of 50 mg/mL (w/v) for 24 h at 4 °C and centrifuging at 4 °C and 9,425×g for 30 min to obtain the supernatant. The ACE substrate was prepared using 0.1 M sodium borate buffer (pH 8.3) containing 8.3 mM N-Hippuryl-His-Leu hydrate. Each sample (50 µL) was taken in a 2 mL tube (distilled water was taken instead of the sample in the control), and 1 M HCl was added initially to prepare additional samples and controls that stopped the reaction. ACE substrate (50 µL) was added to each of the prepared samples and reacted for 10 min at 37 °C. Thereafter, 50 µL of ACE solution (25 mM/mL) was added to the reaction solution and incubated at 37 °C for 30 min. Initially, 250 µL of 1 M HCl was added to stop the reaction in the samples in which 1 M HCl was not previously added; furthermore, 500 µL of ethyl acetate was added and the mixture was vortexed for 1 min. Subsequently, the mixture was centrifuged at 1,977×g for 10 min and then 200 µL of the supernatant was taken and dried at 60 °C for 30 min. Finally, 1 mL of distilled water was added to the dried sample to prepare the final sample. The absorbance of the final sample was measured at 228 nm using a UV/Vis spectrophotometer (Cary® 300; Agilent, Santa Clara, CA, USA). The ACE inhibitory activity was calculated as follows:

$$\text{ACE inhibitory activity (\%)} = \left( 1 - \frac{\text{Absorbance of sample} - \text{Absorbance of sample}_{\text{HCl}}}{\text{Absorbance of control} - \text{Absorbance of control}_{\text{HCl}}} \right) \times 100$$

**In vivo analysis of bioavailability****Collection of plasma and serum from mice**

After completing the peptide extract feeding experiments, all experimental animals (mice) were sacrificed using CO<sub>2</sub> gas, and blood was collected through cardiac puncture method. For analyzing the antioxidant enzyme activities, the collected blood was placed in a plasma separation gel tube and centrifuged for 10 min at 2,000×g and 4 °C, and the obtained plasma was used as a sample. For ACE activity analysis, serum obtained by centrifuging blood in a serum separation gel tube for 15 at 2,000×g and 4 °C was used as a sample.

**Analysis of catalase activity**

The catalase (CAT) activity of each sample was measured using an OxiTec™ Catalase assay kit (Biomax, Seoul, Korea). Briefly, each 25 µL sample was added to a separate well of a 96-well plate, 25 µL of 40 µM H<sub>2</sub>O<sub>2</sub> solution was added to it, and the plate was incubated at 20 °C–25 °C for 30 min. Thereafter, 50 µL of Oxi-Probe/horseradish peroxidase (HRP) working solution was added

to each well, the plate was incubated at 20°C–25°C for 30 min, and the absorbance was measured at 570 nm. One unit (U) of catalase activity refers to the amount of enzyme that will decompose 1 μM of H<sub>2</sub>O<sub>2</sub> per min at pH 7.0 and 25°C.

#### **Analysis of peroxidase activity**

The peroxidase (POD) activity was analyzed using an OxiTec™ Hydrogen peroxide/Peroxidase assay kit (Biomax). Briefly, each 50 μL sample was added to a separate well of a 96-well plate and 50 μL of Oxi-Probe/H<sub>2</sub>O<sub>2</sub> working solution was added to it. Thereafter, the plate was incubated at 20°C–25°C for 30 min and then the absorbance was measured at 560 nm. One unit of HRP activity refers to the amount of enzyme that catalyzes the production of 1 mg of purpurogallin from pyrogallol in 20 s at pH 6.0 and 20°C.

#### **Analysis of superoxide dismutase activity**

The superoxide dismutase (SOD) activity was analyzed using an OxiTec™ SOD assay kit (Biomax). Briefly, each prepared sample and reagent was reacted at 37°C for 30 min, and then the absorbance of samples and blanks was measured at 450 nm using a microplate reader. The SOD activity was calculated as follows:

$$\text{SOD activity (Inhibition rate, \%)} = \left( 1 - \frac{\text{Absorbance of sample} - \text{Absorbance of blank 2}}{\text{Absorbance of blank 1} - \text{Absorbance of blank 3}} \right)$$

#### **Analysis of angiotensin converting enzyme activity**

An ACE activity assay kit (Elabscience, Houston, TX, USA) was used to measure the ACE activity of the samples. The absorbance of samples and blanks was measured at 340 nm using a UV/Vis spectrophotometer. One unit (U) refers to the amount of 1 μM of substrate catalyzed by 1 L of sample per min at 37°C. ACE activity (U/L) was calculated using the following equation: where ε is the absorbance coefficient, d is the path length of the cuvette, and DF is the dilution factor. The ACE activity was calculated as follows:

$$\text{ACE activity (U/L)} = \left( \frac{\text{Absorbance of sample} - \text{Absorbance of blank}}{\text{Reaction time}} \right) \times \frac{1000}{\varepsilon \times d} \times \frac{\text{Total volume}}{\text{Volume of sample}} \times \text{DF}$$

#### **Statistical analysis**

All statistical analyses were conducted using the IBM SPSS Statistics 26 program (IBM, Armonk, NY, USA). The differences between the groups were analyzed using Student's *t*-test and one-way analysis of variance (ANOVA). Using the *t*-test, the average difference between the cuts within the same breed, or the breeds within the same cut, or before and after *in vitro* digestion was compared at a significance level of  $p < 0.05$ . In the case of the one-way ANOVA, the average difference between the control group and loin group of JBP and LYD, or ham group of JBP and LYD (*In vivo* analysis of bioavailability) were compared at a significance level of  $p < 0.05$ . Pairwise comparisons were performed using a post-hoc test (Student–Newman–Keuls; SNK) at a significance level of  $p < 0.05$ .

## **RESULTS AND DISCUSSION**

#### **Analysis of the molecular weight of peptides**

A common pattern was observed in the hydrolysis of peptide extracts into low-molecular-weight

**Table 3.** The molecular weight distribution of loin and ham peptide extracts from JBP and LYD before and after *in vitro* digestion

Contents	Loin				Ham			
	JBP		LYD		JBP		LYD	
	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion
Mn <sup>1)</sup> (Da)	691	287	376	318	437	303	716	287
> 10,000 Da (%)	1.25	0.61	0.28	0.91	-	1.10	1.82	0.73
5,000–10,000 Da (%)	5.55	2.46	1.73	3.74	0.22	4.28	7.12	3.16
3,000–5,000 Da (%)	6.42	3.40	2.97	5.49	1.42	5.88	9.57	4.68
200–3,000 Da (%)	84.93	66.62	86.65	64.85	88.38	65.84	78.54	63.32
< 200 Da (%)	1.85	26.91	8.37	25.01	9.98	22.89	2.96	28.11

<sup>1)</sup> The number averaged molecular weight

JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace × Yorkshire × Duroc).

peptides after *in vitro* digestion (Table 3). The proportion of 200–3,000 Da peptides in the JBP loin and ham peptide extracts before *in vitro* digestion was approximately 85% and 88%, respectively, and that in the LYD loin and ham peptide extracts was approximately 87% and 79%, respectively. However, after *in vitro* digestion, the proportion of 200–3,000 Da peptides in the JBP loin and ham peptide extracts decreased to approximately 67% and 66%, respectively, and that in the LYD loin and ham peptide extracts decreased to approximately 65% and 63%, respectively. Accordingly, it was confirmed that the molecular weight distribution ratio of peptides with < 200 Da after *in vitro* digestion decreased by approximately 13%–25% compared to that before digestion in both JBP and LYD peptide extracts.

These results are believed to be the consequences of peptide bond decomposition and the formation of peptides with smaller molecular weights due to hydrolysis by digestive enzymes during *in vitro* digestion. The results of this study are similar to those obtained by Paoletta et al. [22] and Gallego et al. [23] in studies comparing and analyzing the molecular weight distribution of peptides produced after *in vitro* digestion of pork proteins. According to previous studies, most peptides that are derived from pork proteins and have antioxidant activity have a molecular weight of approximately 303–1,275 Da [3,8,24]. In addition, pork-derived peptides that exhibit ACE inhibitory activity have a molecular weight of approximately 520–950 Da [5,25–28].

### Analysis of free amino acid composition

Table 4 shows the results of the analysis of changes in free amino acid composition after *in vitro* digestion of peptide extracts. The total content of free amino acids detected after *in vitro* digestion of JBP and LYD ham peptide extracts was 154.91 mg/g and 160.01 mg/g, respectively, which is higher than that detected after *in vitro* digestion of JBP and LYD loin peptide extracts. In addition, the main free amino acids of JBP and LYD peptide extracts before *in vitro* digestion were leucine, glutamic acid, isoleucine, and tyrosine, whereas those detected after *in vitro* digestion were arginine, leucine, lysine, phenylalanine and tyrosine.

The composition of free amino acids before and after *in vitro* digestion is believed to have changed due to various factors, including the hydrolysis of peptides and structural changes in amino acids during digestion [29]. According to previous studies, peptides have smaller molecular weights as the enzymatic hydrolysis process progresses, and relatively more free amino acids are produced [30]. Therefore, these results appear to be the consequences of the decomposition of low-molecular-weight peptides in the peptide extracts (obtained from myofibrillar proteins using alkaline-AK enzymes) into large amounts of free amino acids by *in vitro* digestive enzymes.



**Table 4.** Changes in the free amino acid composition of loin and ham peptide extracts from JBP and LYD after *in vitro* digestion

Amino acid	Free amino acid content (mg/g dw)							
	Loin				Ham			
	JBP		LYD		JBP		LYD	
	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion	Before digestion	After digestion
Aspartic acid	0.20	0.13	0.39	0.28	0.15	0.08	0.34	0.26
Threonine	0.46	2.62	0.55	2.50	0.53	2.74	0.47	2.70
Serine	0.53	1.70	0.62	1.65	0.55	1.79	0.53	1.78
Asparagine	0.77	1.93	0.97	2.03	0.79	2.09	0.92	2.32
Glutamic acid	2.61	3.00	4.64	4.64	3.07	3.50	4.31	5.19
Glutamine	0.27	4.26	0.33	3.55	0.41	4.60	0.53	4.52
Glycine	0.26	0.31	0.29	0.32	0.28	0.36	0.24	0.38
Alanine	1.21	3.68	1.47	3.63	1.38	3.78	1.46	3.70
Valine	0.59	2.91	0.88	2.95	0.67	3.05	0.74	3.10
Methionine	0.84	6.26	1.25	5.90	1.07	6.29	1.30	6.66
Isoleucine	1.72	5.33	2.18	5.15	1.77	5.66	2.03	5.85
Leucine	3.72	24.83	5.47	23.93	4.66	26.19	5.44	26.68
Tyrosine	1.56	16.70	1.77	15.31	1.58	17.47	1.88	17.38
Phenylalanine	0.95	16.11	1.10	15.03	0.99	16.54	1.12	16.82
Tryptophan	ND	2.71	ND	2.50	ND	2.90	ND	2.72
Ornithine	0.32	0.51	0.59	0.55	0.31	0.51	0.50	0.67
Lysine	1.16	22.17	1.83	20.99	1.31	23.67	1.40	24.84
Histidine	0.11	2.41	0.16	2.19	0.12	2.50	0.12	2.52
Arginine	1.20	27.85	1.89	28.60	1.43	30.09	1.80	30.78
Proline	0.34	1.14	0.32	0.96	0.43	1.10	0.35	1.14
Carnosine	ND	ND	ND	ND	ND	ND	ND	ND
Total	18.82	146.56	26.70	142.66	21.50	154.91	25.48	160.01


dw, dry weight; JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace × Yorkshire × Duroc); ND, not detected.

According to Xu et al. [31], arginine, lysine, tyrosine, tryptophan, methionine, and histidine have higher antioxidant activities compared with all other amino acids. Arginine significantly increases the activities of SOD, glutathione S-transferase (GST), and glutathione peroxidase in the blood and liver of mice under oxidative stress, whereas lysine upregulates Nrf2, a transcription factor involved in regulating antioxidant gene expression [32,33].


### **In vitro analysis of bioavailability**

#### **In vitro analysis of antioxidative activity**

The JBP and LYD loin peptide extracts showed no significant difference in ABTS radical scavenging activity before and after *in vitro* digestion (Fig. 2A). However, the ABTS radical scavenging activity of the JBP and LYD ham peptide extracts decreased by approximately 17% and 18%, respectively, after *in vitro* digestion ( $p < 0.05$ ) (Fig. 2B). In addition, there was no significant difference between the ABTS radical scavenging activity of the JBP loin peptide extracts after *in vitro* digestion (Fig. 2C); however, the ABTS radical scavenging activity of the LYD loin peptide extract was significantly higher than that of the LYD ham peptide extract ( $p < 0.05$ ) (Fig. 2D). Furthermore, there was no significant difference in the DPPH radical scavenging activity of all peptide extracts after *in vitro* digestion except LYD ham peptide extract (Fig. 3). Before *in vitro*



**Fig. 2.** Comparison of the 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (ABTS) radical scavenging activity between (A) the loin peptide extracts from JBP and LYD, (B) the ham peptide extracts from JBP and LYD, (C) the loin and ham peptide extracts from JBP, and (D) the loin and ham peptide extracts from LYD. Data are presented as mean  $\pm$  standard deviation. <sup>a,b</sup>Lowercase letters indicate a significant difference in the ABTS radical scavenging activity based on breeds and within cuts after digestion ( $p < 0.05$ ). A single asterisk indicates a significant difference in the ABTS radical scavenging activity before and after *in vitro* digestion ( $p < 0.05$ ). JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).




**Fig. 3.** Comparison of the 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity between (A) the loin peptide extracts from JBP and LYD, (B) the ham peptide extracts from JBP and LYD, (C) the loin and ham peptide extracts from JBP, and (D) the loin and ham peptide extracts from LYD. Data are presented as mean  $\pm$  standard deviation. <sup>A,B</sup>Uppercase letters indicate a significant difference in the DPPH radical scavenging activity based on breeds and within cuts before digestion ( $p < 0.05$ ). A single asterisk indicates a significant difference in the DPPH radical scavenging activity before and after *in vitro* digestion ( $p < 0.05$ ). JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).


digestion, the iron chelating activity of JBP loin and ham peptide extracts was approximately 74% and 55%, respectively, which was significantly higher than that of LYD loin and ham peptide extracts ( $p < 0.05$ ) (Fig. 4). However the iron chelating activity was reduced after *in vitro* digestion ( $p < 0.05$ ), and there was no significant difference between after *in vitro* digestion of JBP and LYD loin and ham peptide extracts (Fig. 4). Furthermore, the reducing power assay confirmed that there was no significant difference in reducing power before and after *in vitro* digestion in all treatments (Fig. 5).

According to previous studies, the antioxidant activity of the peptides detected after enzymatic hydrolysis is higher than that of the proteins before decomposition because enzymatic hydrolysis increases antioxidant activities such as free radical removal, reactive oxygen species inactivation, and metal ion chelating by decomposing the protein's peptide bond [34,35]. In addition, peptides with a small molecular weight of  $< 10$  kDa have higher antioxidant activity than peptides with a larger molecular weight [12]. However, in this study, the structure and function of low-molecular-weight peptides changed due to pH changes and hydrolysis by digestive enzymes during *in vitro* digestion, reducing the antioxidant activities of peptide extracts [36]. Although it was thought that peptides extracted from pork can increase antioxidant activities by exposing the hydrophobic amino acid residues of the peptide through hydrolysis by pepsin during *in vitro* digestion, it has been confirmed that antioxidant activities are reduced when the hydrophobic amino acid residues of the peptide are exposed [37,38]. In addition, peptides can be reduced in bioactivity by oxidation, deamidization, or hydrolysis due to pH changes in the digestive system [39]. The pH change may affect the functional properties of the peptide by modifying one or more amino acids. For example, glutamine and asparagine are destroyed under acidic pH conditions, whereas cysteine, serine, and threonine are destroyed under alkaline pH conditions [40].

Meanwhile, the results of this study showed that the proportion of peptides with a molecular



**Fig. 4.** Comparison of the iron chelating activity between (A) the loin peptide extracts from JBP and LYD, (B) the ham peptide extracts from JBP and LYD, (C) the loin and ham peptide extracts from JBP, and (D) the loin and ham peptide extracts from LYD. Data are presented as mean  $\pm$  standard deviation. <sup>A,B</sup>Uppercase letters indicate a significant difference in iron chelating activity based on breeds and within cuts before digestion ( $p < 0.05$ ). A single asterisk indicates a significant difference in iron chelating activity before and after *in vitro* digestion ( $p < 0.05$ ). JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).




**Fig. 5.** Comparison of the reducing power between (A) the loin peptide extracts from JBP and LYD, (B) the ham peptide extracts from JBP and LYD, (C) the loin and ham peptide extracts from JBP, and (D) the loin and ham peptide extracts from LYD. Data are presented as mean  $\pm$  standard deviation. JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).

weight of  $< 200$  Da increased after *in vitro* digestion, which is relatively small compared to the molecular weight of most peptides with antioxidant activities [3,8,24]. In addition, free amino acids may have different antioxidant activities owing to the structural differences in amino acid side chains, and amino acids with side chains containing alkyl groups such as leucine, isoleucine, and phenylalanine have low antioxidant capacity [31]. Thus, free amino acids such as arginine, lysine, and tyrosine, which were detected in large quantities after *in vitro* digestion in the present study, were judged to exhibit high antioxidant activities, whereas leucine and phenylalanine were judged to have significantly low antioxidant activities [31].

Therefore, in the case of ABTS radical scavenging activity, it is judged that bioactive peptides and large amounts of free amino acids contribute to antioxidant activity and can exhibit high radical elimination even after *in vitro* digestion [41]. The ABTS assay is thought to show significantly higher free radical scavenging activity than the DPPH assay because ABTS radicals are more reactive and are less affected by pH than DPPH radicals [42]. In the case of the iron chelating activity, peptides with amino acid residues that can be chelated with  $\text{Fe}^{2+}$  are mostly hydrolyzed through the *in vitro* digestion process and lose their activity [34,43]. Regarding the reducing power, peptides that can act as electron donors are decomposed and lose their activity due to enzymatic hydrolysis; so, it is expected that the reducing power decreases after *in vitro* digestion [44].

### Analysis of angiotensin-converting enzyme inhibitory activity

The ACE inhibitory activity of the loin and ham peptide extracts of JBP and LYD was approximately 94%–97% before *in vitro* digestion; however, after *in vitro* digestion, the ACE inhibitory activity was reduced ( $p < 0.05$ ) (Fig. 6). Nevertheless, the ACE inhibitory activity of the loin peptide extract of JBP was approximately 3% higher than that of the LYD ( $p < 0.05$ ) (Fig. 6). ACE has been confirmed to have an important effect on the renin-angiotensin system (RAS) that regulates blood pressure [45]. ACE converts an inactive decapeptide, angiotensin I, into an octapeptide with vascular contraction activity, angiotensin II, and inactivates bradykinin, which




**Fig. 6.** Comparison of the angiotensin-converting enzyme (ACE) inhibitory activity between (A) the loin peptide extracts from JBP and LYD, (B) the ham peptide extracts from JBP and LYD, (C) the loin and ham peptide extracts from JBP, and (D) the loin and ham peptide extracts from LYD. Data are presented as mean  $\pm$  standard deviation. <sup>A,B</sup>Uppercase letters indicate a significant difference in the ACE inhibitory activity based on breeds and within cuts before digestion ( $p < 0.05$ ). <sup>a,b</sup>Lowercase letters indicate a significant difference in the ACE inhibitory activity based on breeds and within cuts after digestion ( $p < 0.05$ ). A single asterisk indicates a significant difference in the ACE inhibitory activity before and after *in vitro* digestion ( $p < 0.05$ ). JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).

mediates vascular expansion; thus, high ACE activity causes hypertension [46,47]. Therefore, intensive research has been conducted on bioactive peptides derived from food proteins that can suppress ACE activity to reduce high blood pressure while avoiding the side effects caused by synthetic drugs [48]. According to previous papers, peptides derived through enzymatic hydrolysis of pork-derived proteins exhibit strong ACE inhibitory activity [5,25–28]. In addition, peptides with a molecular weight of  $< 10$  kDa were found to have a higher antihypertensive activity than peptides with larger molecular weights [13]. However, these peptides can exhibit ACE inhibitory effects only if they maintain their active form and reach the bloodstream; in this study, it is judged that the peptides were decomposed due to the action of digestive enzymes, resulting in reduced ACE inhibitory activity [23,49]. Most of the dipeptides and tripeptides, which can inhibit ACE activity and also block the active site of ACE, are hydrolyzed into amino acids by cytoplasmic peptides in the small intestine, and only some of them maintain their peptide forms without breaking down into amino acids [50,51].

### ***In vivo* analysis of bioavailability**

#### ***Analysis of antioxidative activity***

The CAT activity of the groups fed with JBP and LYD loin and ham was approximately 260–400 mU/mL higher than the control group ( $p < 0.05$ ) (Fig. 7A). Also, the CAT activity of the groups fed with JBP loin and LYD ham was higher than the JBP ham and LYD loin group, respectively ( $p < 0.05$ ) (Fig. 7A). The POD activity of the groups fed with JBP and LYD loin and ham was approximately 0.31–0.52 mU/mL higher than the control group ( $p < 0.05$ ) (Fig. 7B). The POD activity of the groups fed with LYD loin and ham was significantly higher than that of the JBP group ( $p < 0.05$ ) (Fig. 7B). The SOD activity of the groups fed with JBP loin and ham was  $80.01 \pm 1.51\%$  and  $76.49 \pm 10.19\%$ , respectively, which was relatively high compare to LYD group ( $p <$



**Fig. 7.** Dietary effects of the loin and ham peptide extracts from JBP and LYD on the (A) catalase (CAT), (B) peroxidase (POD), (C) superoxidase dismutase (SOD) activities in mice blood. Data are presented as mean  $\pm$  standard deviation. <sup>A,B</sup>Uppercase letters indicate a significant difference in the antioxidant enzyme activities between the breeds within the same cut ( $p < 0.05$ ). <sup>a,b</sup>Lowercase letters indicate a significant difference in the antioxidant enzyme activities between the cuts within the same breed ( $p < 0.05$ ). A single asterisk indicates a significant difference in the ACE activity between the control and treatment groups ( $p < 0.05$ ). CTL, control; JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).


0.05) (Fig. 7C). However, JBP group was not significantly different to the control group (Fig. 7C).

Antioxidant enzymes such as CAT, POD, and SOD play an important role in protecting the body from oxidative stress caused by free radicals [52]. SOD, an antioxidant enzyme that catalyzes the displacement of free radical  $O^{\cdot 2}$  to  $H_2O_2$  and  $O_2$ , can protect biomolecules from oxidative damage by active oxygen species. In addition, it is judged that the large amount of free amino acids detected after digestion can enhance antioxidant activities [53,54]. For example, arginine has been found to increase the expression of antioxidant-related genes and proteins, leucine increases the activity of antioxidant enzymes such as GST and total superoxide dismutase (T-SOD), and lysine promotes the expression of antioxidant-related genes [33,55,56]. Therefore, when peptide extracts obtained from the loin and ham myofibrillar protein of JBP are consumed, CAT and POD activities increase and SOD activity is triggered to promote antioxidant activity.

### Analysis of angiotensin-converting enzyme activity

The antihypertensive activity was analyzed by measuring the ACE activity after feeding mice with the different peptide extracts; the results of the analysis are presented in Fig. 8. The ACE activity of the groups fed with JBP and LYD loin and ham was higher than that of the control group ( $p < 0.05$ ). The ACE activity of the groups fed with JBP ham and LYD loin was  $287.87 \pm 8.26$  U/L and  $297.28 \pm 3.09$  U/L, respectively, which was significantly lower than that of the groups fed with JBP loin and LYD ham ( $p < 0.05$ ).

In this study, the ACE inhibitory activity of peptide extracts obtained from the loin and ham myofibrillar proteins of JBP and LYD was more than 94% before *in vitro* digestion (Fig. 6). However, the activity of peptides that exhibit ACE inhibitory activity is believed to decrease in peptide extracts owing to enzymatic hydrolysis, changes in pH, and the intestinal residence



**Fig. 8. Dietary effects of the loin and ham peptide extracts from JBP and LYD on the ACE activity in mice blood.** Data are presented as mean  $\pm$  standard deviation. <sup>A,B</sup>Uppercase letters indicate a significant difference in the ACE activity between breeds within the same cut ( $p < 0.05$ ). <sup>a,b</sup>Lowercase letters indicate a significant difference in the ACE activity between cuts within the same breed ( $p < 0.05$ ). A single asterisk indicates a significant difference in the ACE activity between the control and treatment groups ( $p < 0.05$ ). CTL, control; JBP, Jeju black pigs; LYD, three-way crossbred pigs (Landrace  $\times$  Yorkshire  $\times$  Duroc).

time during the digestive process [23,39,57]. Previously, peptides were reported to have low bioavailability when administered orally owing to their weak intestinal mucosal permeability, which can vary depending on the difference in mucosal thickness and surface area of the gastrointestinal tract [57, 58]. In addition, there is little action of protein hydrolyzing enzymes in the large intestine, making it easy to absorb peptides, but peptides can be decomposed by bacteria in the large intestine [59]. According to Lee et al. [60], peptides obtained using alkaline-AK can have relatively reduced digestive stability due to the influence of intestinal microbiome. Thus, various factors in the intestine affect the digestion, absorption, and activity of peptides, resulting in results that may vary from those of the *in vitro* experiments in this study.

Meanwhile, peptides derived from animal protein are believed to impact ACE activity due to the toxins produced from amino acid decomposition metabolites through fermentation by intestinal microbiome during digestion [61,62]. For example, indoxyl sulfate, a toxin produced from the metabolites of an intestinal microbiome, can activate RAS [63]. When RAS is activated, the activity of ACE is promoted and the amount of angiotensin II increases, which results in vascular contraction and high blood pressure [64,65]. In addition, trimethylamine N-oxide (TMAO), which is related to high blood pressure, also promotes an increase in ACE activity [66,67]. It is predicted that the ACE activity of the groups fed with different peptide extracts increased due to these factors. Nevertheless, the ACE activity of the groups fed with JBP ham and LYD loin was lower than those of the groups fed with JBP loin and LYD ham; thus, JBP ham peptide extract believed to have a relatively high ACE inhibitory activity.

## CONCLUSION

In this study, changes in the molecular weight distribution, free amino acid composition, and antioxidant and antihypertensive activities of peptide extracts ( $< 10$  kDa) of JBP and LYD after *in vitro* digestion were compared and analyzed. In addition, the changes in antioxidant and antihypertensive activities of the peptide extracts and their benefits were confirmed through animal

experiments. The loin peptide extract of JBP maintained a relatively high ABTS radical scavenging activity after *in vitro* digestion. The iron chelating activity of JBP loin and ham peptide extracts was significantly higher than that of LYD loin and ham peptide extracts before *in vitro* digestion. However, the iron chelating activity of all peptide extracts decreased after *in vitro* digestion, and there was no significant difference. Although the ACE inhibitory activity of all peptide extracts decreased, the ACE inhibitory activity of JBP loin peptide extract was higher than that of the LYD loin peptide extract. These results are believed to be the consequences of changes in peptide structure and function due to hydrolysis by digestive enzymes and change in pH. Both JBP and LYD peptide extracts promoted antioxidant enzyme activities in the experimental animals; among the different peptide extracts, the JBP loin and ham peptide extracts showed significantly high SOD activity than that of LYD. In addition, the JBP ham peptide extracts showed a relatively higher ACE inhibitory activity in the case of *in vivo* experiment. Therefore, it is deemed that the peptide extracts of JBP are more bioactive than that of LYD, and can be used as bioactive materials; however, additional research is needed to improve their bioavailability.

## REFERENCES

1. Jin SK, Kim CW, Song YM, Jang WH, Kim YB, Yeo JS, et al. Physicochemical characteristics of longissimus muscle between the Korean native pig and Landrace. *Korean J Food Sci Anim Resour.* 2001;21:142-8.
2. Kim GW, Kim HY, Chun JY. Different characteristics between improved breeds and Jeju black pigs. In: 2016 KFN International Symposium and Annual Meeting; 2016; Jeju, Korea. p. 320-1.
3. Arihara K. Functional properties of bioactive peptides derived from meat proteins. In: Nollet LML, Toldrá F, editors. *Advanced technologies for meat processing.* Boca Raton, FL: CRC Press; 2006. p. 245-74.
4. Sánchez A, Vázquez A. Bioactive peptides: a review. *Food Qual Saf.* 2017;1:29-46. <https://doi.org/10.1093/fqsafe/fyx006>
5. Katayama K, Mori T, Kawahara S, Miake K, Kodama Y, Sugiyama M, et al. Angiotensin-I converting enzyme inhibitory peptide derived from porcine skeletal muscle myosin and its antihypertensive activity in spontaneously hypertensive rats. *J Food Sci.* 2007;72:S702-6. <https://doi.org/10.1111/j.1750-3841.2007.00571.x>
6. Keska P, Stadnik J. Antimicrobial peptides of meat origin - an *in silico* and *in vitro* analysis. *Protein Pept Lett.* 2017;24:165-73. <https://doi.org/10.2174/0929866523666161220113230>
7. Kęska P, Rohn S, Halagarda M, Wójciak KM. Peptides from different carcass elements of organic and conventional pork—potential source of antioxidant activity. *Antioxidants.* 2020;9:835. <https://doi.org/10.3390/antiox9090835>
8. Saiga A, Tanabe S, Nishimura T. Antioxidant activity of peptides obtained from porcine myofibrillar proteins by protease treatment. *J Agric Food Chem.* 2003;51:3661-7. <https://doi.org/10.1021/jf021156g>
9. Ryan JT, Ross RP, Bolton D, Fitzgerald GF, Stanton C. Bioactive peptides from muscle sources: meat and fish. *Nutrients.* 2011;3:765-91. <https://doi.org/10.3390/nu3090765>
10. Sharma M, Gat Y, Arya S, Kumar V, Panghal A, Kumar A. A review on microbial alkaline protease: an essential tool for various industrial approaches. *Ind Biotechnol.* 2019;15:69-78. <https://doi.org/10.1089/ind.2018.0032>
11. Singh PK, Shrivastava N, Ojha BK. Enzymes in the meat industry. In: Kuddus M, editor. *Enzymes in food biotechnology: production, applications, and future prospects.* London: Academic Press; 2019. p. 111-28.



12. Ajibola CF, Fashakin JB, Fagbemi TN, Aluko RE. Effect of peptide size on antioxidant properties of African yam bean seed (*Sphenostylis stenocarpa*) protein hydrolysate fractions. *Int J Mol Sci.* 2011;12:6685-702. <https://doi.org/10.3390/ijms12106685>
13. Ruiz-Ruiz J, Dávila-Ortíz G, Chel-Guerrero L, Betancur-Ancona D. Angiotensin I-converting enzyme inhibitory activity in peptide fractions from hard-to-cook bean hydrolysates. *J Biotechnol.* 2010;150:309-10. <https://doi.org/10.1016/j.jbiotec.2010.09.283>
14. Lee SJ, Lee SY, Chung MS, Hur SJ. Development of novel in vitro human digestion systems for screening the bioavailability and digestibility of foods. *J Funct Foods.* 2016;22:113-21. <https://doi.org/10.1016/j.jff.2016.01.005>
15. Hur SJ, Lim BO, Decker EA, McClements DJ. In vitro human digestion models for food applications. *Food Chem.* 2011;125:1-12. <https://doi.org/10.1016/j.foodchem.2010.08.036>
16. Enda H, Sagane Y, Nakazawa Y, Sato H, Yamazaki M. Data on free amino acid contents in Japanese basket clams (*Corbicula japonica*) from Lake Abashiri and Abashirigawa River. *Data Br.* 2018;16:639-43. <https://doi.org/10.1016/j.dib.2017.11.075>
17. Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic Biol Med.* 1999;26:1231-7. [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3)
18. Tepe B, Sokmen M, Akpulat HA, Daferera D, Polissiou M, Sokmen A. Antioxidative activity of the essential oils of *Thymus sipyleus* subsp. *sipyleus* var. *sipyleus* and *Thymus sipyleus* subsp. *sipyleus* var. *rosulans*. *J Food Eng.* 2005;66:447-54. <https://doi.org/10.1016/j.jfoodeng.2004.04.015>
19. Dinis TCP, Madeira VMC, Almeida LM. Action of phenolic derivatives (acetaminophen, salicylate, and 5-aminosalicylate) as inhibitors of membrane lipid peroxidation and as peroxy radical scavengers. *Arch Biochem Biophys.* 1994;315:161-9. <https://doi.org/10.1006/abbi.1994.1485>
20. Oyaizu M. Studies on products of browning reaction: antioxidative activities of products of browning reaction prepared from glucosamine. *Jpn J Nutr.* 1986;44:307-15. <https://doi.org/10.5264/eiyogakuzashi.44.307>
21. Cushman DW, Cheung HS. Spectrophotometric assay and properties of the angiotensin-converting enzyme of rabbit lung. *Biochem Pharmacol.* 1971;20:1637-48. [https://doi.org/10.1016/0006-2952\(71\)90292-9](https://doi.org/10.1016/0006-2952(71)90292-9)
22. Paoletta S, Falavigna C, Faccini A, Virgili R, Sforza S, Dall'Asta C, et al. Effect of dry-cured ham maturation time on simulated gastrointestinal digestion: characterization of the released peptide fraction. *Food Res Int.* 2015;67:136-44. <https://doi.org/10.1016/j.foodres.2014.10.026>
23. Gallego M, Mauri L, Aristoy MC, Toldrá F, Mora L. Antioxidant peptides profile in dry-cured ham as affected by gastrointestinal digestion. *J Funct Foods.* 2020;69:103956. <https://doi.org/10.1016/j.jff.2020.103956>
24. Xing LJ, Hu YY, Hu HY, Ge QF, Zhou GH, Zhang WG. Purification and identification of antioxidative peptides from dry-cured Xuanwei ham. *Food Chem.* 2016;194:951-8. <https://doi.org/10.1016/j.foodchem.2015.08.101>
25. Escudero E, Sentandreu MA, Arihara K, Toldrá F. Angiotensin I-converting enzyme inhibitory peptides generated from in vitro gastrointestinal digestion of pork meat. *J Agric Food Chem.* 2010;58:2895-901. <https://doi.org/10.1021/jf904204n>
26. Escudero E, Toldrá F, Sentandreu MA, Nishimura H, Arihara K. Antihypertensive activity of peptides identified in the in vitro gastrointestinal digest of pork meat. *Meat Sci.* 2012;91:382-4. <https://doi.org/10.1016/j.meatsci.2012.02.007>
27. Muguruma M, Ahmed AM, Katayama K, Kawahara S, Maruyama M, Nakamura T.

- Identification of pro-drug type ACE inhibitory peptide sourced from porcine myosin B: evaluation of its antihypertensive effects in vivo. *Food Chem.* 2009;114:516-22. <https://doi.org/10.1016/j.foodchem.2008.09.081>
28. Nakashima Y, Arihara K, Sasaki A, Mio H, Ishikawa S, Itoh M. Antihypertensive activities of peptides derived from porcine skeletal muscle myosin in spontaneously hypertensive rats. *J Food Sci.* 2002;67:434-7. <https://doi.org/10.1111/j.1365-2621.2002.tb11424.x>
  29. Diep TT, Yoo MJY, Rush E. Effect of in vitro gastrointestinal digestion on amino acids, polyphenols and antioxidant capacity of tamarillo yoghurts. *Int J Mol Sci.* 2022;23:2526. <https://doi.org/10.3390/ijms23052526>
  30. Weng TM, Chen MT. Changes of protein in natto (a fermented soybean food) affected by fermenting time. *Food Sci Technol Res.* 2010;16:537-42. <https://doi.org/10.3136/fstr.16.537>
  31. Xu N, Chen G, Liu H. Antioxidative categorization of twenty amino acids based on experimental evaluation. *Molecules.* 2017;22:2066. <https://doi.org/10.3390/molecules22122066>
  32. Abu-Serie MM, El-Gamal BA, El-Kersh MA, El-Saadani MA. Investigation into the antioxidant role of arginine in the treatment and the protection for intralipid-induced non-alcoholic steatohepatitis. *Lipids Health Dis.* 2015;14:128. <https://doi.org/10.1186/s12944-015-0124-0>
  33. Li XY, Liu Y, Jiang WD, Jiang J, Wu P, Zhao J, et al. Co- and post-treatment with lysine protects primary fish enterocytes against Cu-induced oxidative damage. *PLOS ONE.* 2016;11:e0147408. <https://doi.org/10.1371/journal.pone.0147408>
  34. Elias RJ, Kellerby SS, Decker EA. Antioxidant activity of proteins and peptides. *Crit Rev Food Sci Nutr.* 2008;48:430-41. <https://doi.org/10.1080/10408390701425615>
  35. Park SY, Chin KB. Antioxidant activities of pepsin hydrolysates of water- and salt-soluble protein extracted from pork hams. *Int J Food Sci Technol.* 2011;46:229-35. <https://doi.org/10.1111/j.1365-2621.2010.02454.x>
  36. Segura-Campos M, Chel-Guerrero L, Betancur-Ancona D, Hernandez-Escalante VM. Bioavailability of bioactive peptides. *Food Rev Int.* 2011;27:213-26. <https://doi.org/10.1080/87559129.2011.563395>
  37. Acquah C, Di Stefano E, Udenigwe CC. Role of hydrophobicity in food peptide functionality and bioactivity. *J Food Bioact.* 2018;4:88-98. <https://doi.org/10.31665/JFB.2018.4164>
  38. Zhu CZ, Zhang WG, Kang ZL, Zhou GH, Xu XL. Stability of an antioxidant peptide extracted from Jinhua ham. *Meat Sci.* 2014;96:783-9. <https://doi.org/10.1016/j.meatsci.2013.09.004>
  39. Chen MC, Sonaje K, Chen KJ, Sung HW. A review of the prospects for polymeric nanoparticle platforms in oral insulin delivery. *Biomaterials.* 2011;32:9826-38. <https://doi.org/10.1016/j.biomaterials.2011.08.087>
  40. Korhonen H, Pihlanto-Leppäla A, Rantamäki P, Tupasela T. Impact of processing on bioactive proteins and peptides. *Trends Food Sci Technol.* 1998;9:307-19. [https://doi.org/10.1016/S0924-2244\(98\)00054-5](https://doi.org/10.1016/S0924-2244(98)00054-5)
  41. Clausen MR, Skibsted LH, Stagsted J. Characterization of major radical scavenger species in bovine milk through size exclusion chromatography and functional assays. *J Agric Food Chem.* 2009;57:2912-9. <https://doi.org/10.1021/jf803449t>
  42. Shalaby EA, Shanab SMM. Antioxidant compounds, assays of determination and mode of action. *Afr J Pharm Pharmacol.* 2013;7:528-39. <https://doi.org/10.5897/AJPP2013.3474>
  43. Zhu CZ, Zhang WG, Zhou GH, Xu XL. Identification of antioxidant peptides of Jinhua ham generated in the products and through the simulated gastrointestinal digestion system. *J Sci*

- Food Agric. 2016;96:99-108. <https://doi.org/10.1002/jsfa.7065>
44. Keşka P, Stadnik J. Effect of in vitro gastro-pancreatic digestion on antioxidant activity of low-molecular-weight (<3.5 kDa) peptides from dry-cured pork loins with probiotic strains of LAB. *Int J Food Sci Technol*. 2021;56:6268-78. <https://doi.org/10.1111/ijfs.15066>
  45. Gurley SB, Coffman TM. The renin-angiotensin system and diabetic nephropathy. *Semin Nephrol*. 2007;27:144-52. <https://doi.org/10.1016/j.semnephrol.2007.01.009>
  46. Mora L, Gallego M, Toldrá F. ACEI-inhibitory peptides naturally generated in meat and meat products and their health relevance. *Nutrients*. 2018;10:1259. <https://doi.org/10.3390/nu10091259>
  47. Zhuo JL, Ferrao FM, Zheng Y, Li XC. New frontiers in the intrarenal renin-angiotensin system: a critical review of classical and new paradigms. *Front Endocrinol*. 2013;4:166. <https://doi.org/10.3389/fendo.2013.00166>
  48. Toldrá F, Reig M, Aristoy MC, Mora L. Generation of bioactive peptides during food processing. *Food Chem*. 2018;267:395-404. <https://doi.org/10.1016/j.foodchem.2017.06.119>
  49. Vercruyse L, Van Camp J, Morel N, Rougé P, Herregods G, Smagghe G. Ala-Val-Phe and Val-Phe: ACE inhibitory peptides derived from insect protein with antihypertensive activity in spontaneously hypertensive rats. *Peptides*. 2010;31:482-8. <https://doi.org/10.1016/j.peptides.2009.05.029>
  50. Hernández-Ledesma B, del Mar Contreras M, Recio I. Antihypertensive peptides: production, bioavailability and incorporation into foods. *Adv Colloid Interface Sci*. 2011;165:23-35. <https://doi.org/10.1016/j.cis.2010.11.001>
  51. Shimizu M. Food-derived peptides and intestinal functions. *BioFactors*. 2004;21:43-7. <https://doi.org/10.1002/biof.552210109>
  52. Woo JY, Gu W, Kim KA, Jang SE, Han MJ, Kim DH. *Lactobacillus pentosus* var. *plantarum* C29 ameliorates memory impairment and inflammaging in a D-galactose-induced accelerated aging mouse model. *Anaerobe*. 2014;27:22-6. <https://doi.org/10.1016/j.anaerobe.2014.03.003>
  53. Boguszewska D, Grudkowska M, Zagdańska B. Drought-responsive antioxidant enzymes in potato (*Solanum tuberosum* L.). *Potato Res*. 2010;53:373-82. <https://doi.org/10.1007/s11540-010-9178-6>
  54. Vlasova II. Peroxidase activity of human hemoproteins: keeping the fire under control. *Molecules*. 2018;23:2561. <https://doi.org/10.3390/molecules23102561>
  55. Chen X, Xiang L, Jia G, Liu G, Zhao H, Huang Z. Effects of dietary leucine on antioxidant activity and expression of antioxidant and mitochondrial-related genes in longissimus dorsi muscle and liver of piglets. *Anim Sci J*. 2019;90:990-8. <https://doi.org/10.1111/asj.13249>
  56. Liang M, Wang Z, Li H, Cai L, Pan J, He H, et al. L-Arginine induces antioxidant response to prevent oxidative stress via stimulation of glutathione synthesis and activation of Nrf2 pathway. *Food Chem Toxicol*. 2018;115:315-28. <https://doi.org/10.1016/j.fct.2018.03.029>
  57. Hamman JH, Enslin GM, Kotzé AF. Oral delivery of peptide drugs: barriers and developments. *BioDrugs*. 2005;19:165-77. <https://doi.org/10.2165/00063030-200519030-00003>
  58. Kompella UB, Lee VHL. Delivery systems for penetration enhancement of peptide and protein drugs: design considerations. *Adv Drug Deliv Rev*. 2001;46:211-45. [https://doi.org/10.1016/S0169-409X\(00\)00137-X](https://doi.org/10.1016/S0169-409X(00)00137-X)
  59. Chourasia MK, Jain SK. Pharmaceutical approaches to colon targeted drug delivery systems. *J Pharm Pharm Sci*. 2003;6:33-66.
  60. Lee SY, Lee DY, Hur SJ. Changes in the stability and antioxidant activities of different

- molecular weight bioactive peptide extracts obtained from beef during in vitro human digestion by gut microbiota. *Food Res Int.* 2021;141:110116. <https://doi.org/10.1016/j.foodres.2021.110116>
61. Jaworska K, Koper M, Ufnal M. Gut microbiota and renin-angiotensin system: a complex interplay at local and systemic levels. *Am J Physiol Gastrointest Liver Physiol.* 2021;321:G355-66. <https://doi.org/10.1152/ajpgi.00099.2021>
  62. Mafra D, Borges NA, de Franca Cardozo LFM, Anjos JS, Black AP, Moraes C, et al. Red meat intake in chronic kidney disease patients: two sides of the coin. *Nutrition.* 2018;46:26-32. <https://doi.org/10.1016/j.nut.2017.08.015>
  63. Akbar MF. Inhibition of indoxyl sulfate-induced intrarenal renin-angiotensin system activation: targeting the aryl hydrocarbon receptor. *Transl Clin Pharmacol.* 2017;25:114-6. <https://doi.org/10.12793/tcp.2017.25.3.114>
  64. Aiemratchanee P, Panyawechamontri K, Phaophu P, Reamtong O, Panbangred W. In vitro antihypertensive activity of bioactive peptides derived from porcine blood corpuscle and plasma proteins. *Int J Food Sci Technol.* 2021;56:2315-24. <https://doi.org/10.1111/ijfs.14853>
  65. Volpe M, Tocci G, Pagannone E. Activation of the renin-angiotensin-aldosterone system in heart failure. *Ital Heart J.* 2005;6:16S-23S.
  66. Ge X, Zheng L, Zhuang R, Yu P, Xu Z, Liu G, et al. The gut microbial metabolite trimethylamine N-oxide and hypertension risk: a systematic review and dose-response meta-analysis. *Adv Nutr.* 2020;11:66-76. <https://doi.org/10.1093/advances/nmz064>
  67. Liu M, Han Q, Yang J. Trimethylamine-N-oxide (TMAO) increased aquaporin-2 expression in spontaneously hypertensive rats. *Clin Exp Hypertens.* 2019;41:312-22. <https://doi.org/10.1080/10641963.2018.1481420>