

EDITORIAL COMMENT

Internal Bleeding

Is Intraplaque Hemorrhage a Decoration or a Driver?*



Jean-Baptiste Michel, MD, PhD,^a Peter Libby, MD,^b Grégory Franck, PhD^a

Contemporary advances in high-throughput screening of biological samples have added unbiased screens to the classical candidate approach for implicating mediators in various disease states. In this issue of *JACC: Basic to Translational Science*, Matic et al. (1) report transcriptomic and proteomic profiling of atherosclerotic plaques banked from carotid endarterectomies. In parallel, they looked at blood sampled near the lesion in vivo in comparison to peripheral blood. The investigators found a single molecule of many tested to be overexpressed in this collection of atherosclerotic lesions that received surgical remediation compared with nonatherosclerotic arterial specimens. They identified biliverdin reductase B (BLVRB) as significantly increased at a messenger RNA, local protein, and plasma protein levels, in company with parallel rises in heme oxygenase 1 (HMOX1). BLVRB participates in catabolism of heme.

SEE PAGE 464

These results that emerged from high-throughput screening technologies applied to human atherosclerosis draw our attention to pathways long implicated


in atherogenesis, but remain worthy of renewed attention in light of new data including these from the Stockholm/Karolinska group. The strong regulation of heme-metabolizing enzymes in active human atheromata indicates a substantial local and/or systemic tissue response to heme, at least in carotid artery lesions sufficiently active to warrant surgical treatment.

From whence does heme that could evoke a clearance response within the plaque come? The microvessels that penetrate into plaques during their evolution (inward neo-angiogenesis) exhibit impaired barrier function and display characteristics of fragility that could lead to extravasation of erythrocytes and intraplaque hemorrhage (Figure 1). In the extreme, leaky microvessels in the plaque could give rise to rapid plaque expansion and promote hematoma formation that could embarrass blood flow. Such macroscopic episodes of intraplaque hemorrhage, although uncommon, likely represent the extreme case of much more common microscopic episodes of intraplaque hemorrhage. Matic et al. (1) found that exposure to ferric iron was associated with iron uptake by THP-1 human monocyte cell line-derived macrophage-like cells and stimulated HMOX1 and BLVRB expression, suggesting that local hemolysis and iron derived from heme drive the elevated concentrations of these enzymes in plaques. Intraplaque hemorrhage promotes erythrocyte sequestration and localized hemolysis, hemostasis, and coagulation, including platelet activation, fibrin formation, and trapping of leukocytes, mainly neutrophils, in human plaques (2). Foci of intraplaque hemorrhage colocalize with proteases and markers of oxidative stress, which are possible promoters of plaque rupture (3,4). In contrast, products of local blood coagulation can lead to the release of platelet products, such as transforming growth factor-beta and platelet-derived growth factor, which can foster a local healing response that could augment smooth muscle production of extracellular matrix,

*Editorials published in *JACC: Basic to Translational Science* reflect the views of the authors and do not necessarily represent the views of *JACC: Basic to Translational Science* or the American College of Cardiology.

From the ^aUnité mixte de Recherche 1148 Inserm-Paris7 University, Xavier Bichat Hospital, Paris, France; and the ^bDivision of Cardiovascular Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts. Dr. Libby was supported by grants from the National Heart, Lung, and Blood Institute (HL080472) and the RRM Charitable Fund. Dr. Franck was supported by the Lefoulon-Delalande Foundation and the UE PRESTIGE Fund. Dr. Michel has reported that he has no relationships relevant to the contents of this paper to disclose.

All authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the *JACC: Basic to Translational Science* [author instructions page](#).



smooth muscle proliferation, and thus, growth of the atherosclerotic lesion.

In addition to erythrocyte extravasation from fragile microvessels, entry of erythrocytes from the macrovascular compartment into the intima could arise from breaches in the integrity of the endothelial layer that lines the macrovascular lumen. Altered hydrodynamics at regions of flow disturbance could contribute to endothelial cell desquamation or impaired barrier function. Recent studies suggest accumulation of erythrocytes near the intimal interface with blood even in early atherogenesis (5). Not only do red blood cells (RBCs) accumulate locally within the intima, but hemoglobin, glycophorin A, and iron colocalize with regions of RBC collections that compose small intraplaque hematomas. The sites of local RBC accumulation in the intima also contain HMOX1, ferroportin, and members of the natural resistance-associated macrophage protein family of metal transporters. Furthermore, sites of these intimal hematomas contain markers of oxidation of lipids and proteins. Smooth muscle cells appear capable of erythrophagocytosis in addition to classical scavenging of modified or senescent RBC via scavenger receptors associated with mononuclear phagocytes including CD163 and CD36 (5).

Whether they are derived from microvascular leaks or from breaches in the macrovascular endothelium, redox-active ferrous iron (Fe^{++}) derived from erythrocyte heme prosthetic groups can contribute to local oxidative damage through the Fenton reaction. The local plaque environment favors hemolysis of RBC, releasing heme and iron in the intimal lesions. Recent systems biology analyses point to hydroxyl radical derived through Fenton chemistry as an agonist in atherogenesis (6). In addition, genome-wide association studies have implicated iron metabolism and hepcidin as risk factors for biomarkers of atherosclerosis such as increased intima-media thickness (7). Experimental studies have implicated hepcidin in activating plaque macrophages following erythrophagocytosis. Some experimental data suggest that clearance of hemoglobin by CD163-positive cells and concomitant HMOX1 expression can provide an anti-inflammatory response that can defend against some of the potentially adverse effects of local heme metabolism and accumulation of iron (8). Beyond the role of RBC-derived heme and iron in potentiating atherosclerosis, the membranes of erythrocytes that enter the intima can provide a local source of cholesterol (9).

The finding of increased BLVRB in carotid atheromata shines a bright spotlight on these various pathogenic and potentially counter-regulatory pathways during human atherogenesis. Whereas intraplaque hemorrhage does not likely initiate the atherosclerotic process, operation of the iron-related pathways enumerated herein may sustain the disease and promote properties of plaques implicated in causing clinical complications (Figure 1).

The observations of Matic et al. (1) raise a number of interesting questions that merit further investigation. Is intraplaque hemorrhage a characteristic of carotid atheromata, or does it affect atherosclerotic plaques in other vascular beds? Do particular hemodynamic conditions near the carotid bifurcation predilect to the formation of intraplaque hemorrhage? In this regard, femoral artery atheromata rarely exhibit intraplaque hemorrhage (10). About one-third of acute coronary events arise from superficial erosion (11), a mechanism of plaque disruption that would not deposit RBC in the core of coronary atheromata (12). In contrast, because intraluminal thrombi characteristically contain abundant RBC (13) and abdominal aortic aneurysms usually associate with a burden of mural thrombosis, the heme pathways portrayed in the study by Matic et al. (1) might apply particularly to these manifestations of atherosclerosis.

The focus on heme metabolism stimulated by the results of Matic et al. (1) raises therapeutic questions as well. Does modulation of human atherosclerosis, for example, by aggressive lipid-lowering therapy, reduce RBC accumulation of plaque arising either

from microvessels or intimal breaches? Substantial evidence suggests that contemporary preventive therapies including statins may change the character of human atherosclerotic plaques. “Stabilization” of microvessels and of the integrity of luminal endothelial cells might limit erythrocyte entry into the plaque. Microvascular permeability studies using magnetic resonance techniques could help to test this hypothesis (14).

Animal experiments could explore further mechanisms of intraplaque hemorrhage. Pigs and rodents have some limitations for probing plaque neovascularization (15). But, some preparations in rabbits (16) or mice (17) involve accumulation of RBC near neovessels, suggesting microvascular leakiness and intraplaque hemorrhage. Animal experiments that produce flow disturbance, for example, by the induction of a partial stenosis, might also help to define mechanisms of the entry of RBC from the microcirculation into the artery wall (18).

In sum, the results from the Karolinska group highlight the local responses in plaque to heme and provide support for the operation in human atherosclerosis of many pathways that involve erythrocyte extravasation and iron-mediated oxidative stress as a potential potentiator of human atherogenesis.

ADDRESS FOR CORRESPONDENCE: Dr. Peter Libby, Division of Cardiovascular Medicine, Department of Medicine, Brigham and Women’s Hospital, Harvard Medical School, 77 Avenue Louis Pasteur, Boston, Massachusetts 02115. E-mail: plibby@bwh.harvard.edu.

REFERENCES

1. Matic LP, Iglesias MJ, Vesterlund M, et al. Novel multiomics profiling of human carotid atherosclerotic plaques and plasma reveals biliverdin reductase B as a marker of intraplaque hemorrhage. *J Am Coll Cardiol Basic Trans Science* 2018;3:464-80.
2. Leclercq A, Houard X, Philippe M, et al. Involvement of intraplaque hemorrhage in atherothrombosis evolution via neutrophil protease enrichment. *J Leukoc Biol* 2007;82:1420-9.
3. Michel JB, Virmani R, Arbustini E, Pasterkamp G. Intraplaque haemorrhages as the trigger of plaque vulnerability. *Eur Heart J* 2011; 32:1977-1985a-c.
4. Michel JB, Martin-Ventura JL, Nicoletti A, Ho-Tin-Noe B. Pathology of human plaque vulnerability: mechanisms and consequences of intraplaque haemorrhages. *Atherosclerosis* 2014; 234:311-9.
5. Delbosc S, Bayles RG, Laschet J, et al. Erythrocyte efferocytosis by the arterial wall promotes oxidation in early-stage atheroma in humans. *Front Cardiovasc Med* 2017;4:43.
6. Formanowicz D, Radom M, Rybarczyk A, Formanowicz P. The role of Fenton reaction in ROS-induced toxicity underlying atherosclerosis—modeled and analyzed using a Petri net-based approach. *Biosystems* 2018;165: 71-87.
7. Galesloot TE, Janss LL, Burgess S, et al. Iron and hepcidin as risk factors in atherosclerosis: what do the genes say? *BMC Genet* 2015;16:79.
8. Schaer CA, Schoedon G, Imhof A, Kurrer MO, Schaer DJ. Constitutive endocytosis of CD163 mediates hemoglobin-heme uptake and determines the noninflammatory and protective transcriptional response of macrophages to hemoglobin. *Circ Res* 2006;99:943-50.
9. Tziakas DN, Kaski JC, Chalikias GK, et al. Total cholesterol content of erythrocyte membranes is increased in patients with acute coronary syndrome: a new marker of clinical instability? *J Am Coll Cardiol* 2007;49: 2081-9.
10. Herisson F, Heymann MF, Chetiveaux M, et al. Carotid and femoral atherosclerotic plaques show different morphology. *Atherosclerosis* 2011;216: 348-54.
11. Farb A, Burke AP, Tang AL, et al. Coronary plaque erosion without rupture into a lipid core. A frequent cause of coronary thrombosis in sudden coronary death. *Circulation* 1996;93: 1354-63.
12. Partida RA, Libby P, Crea F, Jang IK. Plaque erosion: a new in vivo diagnosis and a potential major shift in the management of patients with acute coronary syndromes. *Eur Heart J* 2018;39: 2070-6.

13. Martinez-Pinna R, Lindholt JS, Madrigal-Matute J, et al. From tissue iron retention to low systemic haemoglobin levels, new pathophysiological biomarkers of human abdominal aortic aneurysm. *Thromb Haemost* 2014;112:87-95.
14. Taqueti VR, Di Carli MF, Jerosch-Herold M, et al. Increased microvascularization and vessel permeability associate with active inflammation in human atheromata. *Circ Cardiovasc Imaging* 2014;7:920-9.
15. Van der Veken B, De Meyer GRY, Martinet W. Intraplaque neovascularization as a novel therapeutic target in advanced atherosclerosis. *Expert Opin Ther Targets* 2016;20:1247-57.
16. Giannarelli C, Ibanez B, Cimmino G, et al. Contrast-enhanced ultrasound imaging detects intraplaque neovascularization in an experimental model of atherosclerosis. *J Am Coll Cardiol Img* 2010;3:1256-64.
17. de Vries MR, Niessen HW, Lowik CW, Hamming JF, Jukema JW, Quax PH. Plaque rupture complications in murine atherosclerotic vein grafts can be prevented by TIMP-1 overexpression. *PLoS One* 2012;7:e47134.
18. Franck G, Mawson T, Sausen G, et al. Flow perturbation mediates neutrophil recruitment and potentiates endothelial injury via TLR2 in mice: implications for superficial erosion. *Circ Res* 2017;121:31-42.

KEY WORDS atherosclerosis, Fenton reaction, heme oxygenase, iron