



# Subcutaneous delivery of immune checkpoint inhibitors: new route replacing intravenous administration?

Yusuke Inoue<sup>^</sup>

Second Division, Department of Internal Medicine, Hamamatsu University School of Medicine, Shizuoka, Japan

*Correspondence to:* Yusuke Inoue, MD, PhD. Second Division, Department of Internal Medicine, Hamamatsu University School of Medicine, 1-20-1 Handayama, Chuo-ku, Hamamatsu, Shizuoka 431-3192, Japan. Email: yinoue@hama-med.ac.jp.

*Comment on:* Burotto M, Zvirbulė Z, Mochalova A, *et al.* IMscin001 Part 2: a randomised phase III, open-label, multicentre study examining the pharmacokinetics, efficacy, immunogenicity, and safety of atezolizumab subcutaneous versus intravenous administration in previously treated locally advanced or metastatic non-small-cell lung cancer and pharmacokinetics comparison with other approved indications. *Ann Oncol* 2023;34:693-702.

**Keywords:** Subcutaneous; immune checkpoint inhibitor (ICI); atezolizumab; recombinant human hyaluronidase PH20 (rHuPH20); IMscin001

Submitted Jan 17, 2024. Accepted for publication Feb 26, 2024. Published online Apr 18, 2024.

doi: 10.21037/tlcr-24-63

**View this article at:** <https://dx.doi.org/10.21037/tlcr-24-63>

The use of intravenously delivered immune checkpoint inhibitors (ICIs) as antibody therapies has become a central node in the treatment of locally advanced or metastatic non-small cell lung cancer (NSCLC) as well as extensive-stage small cell lung cancer. For NSCLC, the indications for ICIs targeting programmed death-1 (PD-1) and its ligand (PD-L1) have expanded covering resectable disease as neoadjuvant (1), adjuvant (2), and perioperative (3) therapies. Meanwhile, the number of indications has increased substantially across various types of cancers, with marketing approval covering at least 90 uses for more than 11 different ICIs (4). Because ICIs offer durable responses in a subset of patients who may require long-term repeated administrations over years, they are associated with an accumulating treatment burden in cancer patients with numerous indications, particularly within hospital and outpatient treatment settings. Therefore, the development of strategies to improve convenience and efficiency in healthcare by shortening the times for dose preparation and administration and to reduce patient time and discomfort in the clinic has become an important field of investigation.

Supported by patient preferences and healthcare professional satisfaction over intravenous administration for a range of reasons, including ease of administration,

less emotional distress, and shorter treatment duration (5-8), monoclonal antibodies (mAbs) with subcutaneous formulations have been approved in the United States and European Union for several disease areas, including rheumatoid arthritis [e.g., etanercept (anti-TNF $\alpha$ ), tocilizumab (anti-IL-6), anakinra (anti-IL-1), abatacept (selective co-stimulating modulator)], multiple sclerosis [e.g., ofatumumab (anti-CD-20)], and high-dose mAbs in oncology [e.g., trastuzumab (anti-HER2), rituximab (anti-CD20), daratumumab (anti-CD38)], as discussed in a previous report (9). For successful subcutaneous delivery of mAbs, the unique characteristics of the subcutaneous space, consisting of extracellular matrix, adipose tissue, blood and lymph vessels, fibroblasts, and macrophages, as well as the physicochemical characteristics of mAbs, including large molecular size and low permeability that limit diffusion, require consideration (10). Meanwhile, the maximum volume that can be safely injected into the subcutaneous space is considered to be 1.5 mL, meaning that subcutaneous formulations containing high drug doses in a limited injection volume need to be developed (11). This volume limitation for subcutaneous drug delivery together with the limitations for drug dispersion and absorption are, at least in part, mediated by hyaluronan. Hyaluronan forms

<sup>^</sup> ORCID: 0000-0001-8075-0597.

a gel-like substance with water and acts as the main filter for the extracellular matrix (12,13). Consequently, novel drug delivery technologies involving recombinant human hyaluronidases, such as recombinant human hyaluronidase PH20 (rHuPH20), have been developed to overcome the barriers for subcutaneous delivery of co-administered mAbs (12). rHuPH20 induces local depolymerization of hyaluronan and enables temporal subcutaneous administration of reformulated existing intravenous drugs in relatively large volumes (12,13). rHuPH20 has already been approved in the United States and/or global markets as a drug delivery technology for daratumumab, trastuzumab, pertuzumab, trastuzumab, rituximab, and immune globulin.

Burotto and colleagues reported the primary results of a randomized phase III, open-label, multicenter, noninferiority study (IMscin001 Part 2) investigating the pharmacokinetics, efficacy, and safety of subcutaneous atezolizumab, an anti-PD-L1 mAb, compared with approved intravenous administration (1,200 mg every 3 weeks) in patients with previously treated locally advanced or metastatic NSCLC (14). In the study, a total of 371 patients were randomized at 2:1 to receive subcutaneous atezolizumab (N=247) or intravenous atezolizumab (N=124). The total injection volume of the ready-to-use subcutaneous formulation of atezolizumab (1,875 mg) was 15 mL containing 30,000 U of rHuPH20 (2,000 U/mL), with a suggested delivery time of <10 minutes. The subcutaneous atezolizumab regimen of 1,875 mg every 3 weeks was set based on the findings in the prior phase 1b IMscin001 Part 1 study, which demonstrated that this regimen provided similar exposure to an intravenous dosing regimen of 1,200 mg every 3 weeks (15). Despite the inherent limitations of subcutaneous drug delivery, such as interpatient variability in bioavailability, the study by Burotto and colleagues met the co-primary pharmacokinetics endpoints, cycle 1 observed trough serum concentration and model-predicted area under the curve from days 0 to 21, showing comparable levels of atezolizumab in the blood. Although the study was not formally designed to examine the noninferiority of efficacy endpoints, the objective response rates (subcutaneous: 12% versus intravenous: 10%) and progression-free survival durations (median survival: 2.8 months for the subcutaneous group versus 2.9 months for the intravenous group) were similar between the subcutaneous and intravenous arms, consistent with the findings in the pivotal OAK study (16).

Given that antidrug antibody development can be associated with lower serum drug levels and worse clinical

outcomes in cancer patients treated with ICIs (17,18), one of the major concerns for subcutaneous administration of mAbs is the potential risk of increased immunogenicity, as observed for other subcutaneous mAbs in oncology. For example, a higher incidence of antidrug antibodies was reported for subcutaneous trastuzumab compared with its intravenous comparator (19). In the IMscin001 Part 2 study, however, there was no noticeable difference in the positivity rates of treatment-emergent anti-atezolizumab antibodies between the subcutaneous (19.5%) and intravenous (13.9%) groups (14). Unfortunately, information on whether these treatment-emergent anti-atezolizumab antibodies were neutralizing in action was not available. Because ICIs can enhance the nontumor-specific immune system, further immunogenicity analyses are required in patients receiving subcutaneous ICIs.

Regarding safety, no new signals were identified in the subcutaneous atezolizumab group in the IMscin001 Part 2 study (14). Reflecting the different administration routes, injection site reactions occurred specifically in the subcutaneous arm (4.5%), while infusion-related reactions occurred specifically in the intravenous arm (3.2%).

While Burotto and colleagues did not provide data on patient preferences (14), the CheckMate 8KX study investigating subcutaneous nivolumab, an anti-PD-1 antibody, with or without rHuPH20 in patients with advanced solid tumors found that most patients reported high satisfaction with subcutaneous administration and preferred it over intravenous administration, and that limited pain was associated with subcutaneous injection (20), consistent with the results for other subcutaneous mAbs in oncology (5-8). Taken together, these findings should facilitate a shift in ICI administration from the intravenous to subcutaneous route. Indeed, many studies have been investigating the pharmacokinetics, efficacy, and safety of different subcutaneous ICIs (*Table 1*). Among the ICIs listed in *Table 1*, sasanlimab, a humanized, hinge region-stabilized anti-PD-1 antibody, does not require any modification or combination to enable subcutaneous delivery (21). Furthermore, single-domain antibodies that lack the immunoglobulin light chain are more soluble and stable than complete mAbs, and can more rapidly penetrate tissues, thereby enabling subcutaneous administration (22). For example, envafolimab, a humanized single-domain anti-PD-L1 antibody fused to a human immunoglobulin Fc fragment, is the first subcutaneously injectable anti-PD-L1 antibody approved by the National Medical Products Administration in China. Despite the ongoing shift in focus for pharmaceutical companies toward the development of subcutaneous ICI formulations, there

**Table 1** Overview of clinical trials on subcutaneous administration of immune checkpoint inhibitors

Molecule	Target	Trial name or clinical trial No.	Phase	Setting	Tumor type	Experimental treatment arm	Coformulation with hyaluronidase
Nivolumab	PD-1	CheckMate 8KX	1/2	Advanced	NSCLC, melanoma, RCC, CRC	Monotherapy	Yes/no
Pembrolizumab	PD-1	CheckMate 67T	3	Advanced or metastatic	ccRCC	Monotherapy	Yes
		KEYNOTE-555	1	Advanced	Melanoma	Monotherapy	No
		NCT06041802	2	Recurrent or metastatic	cSCC	Monotherapy	Yes
		NCT05722015	3	Metastatic	NSCLC	Pembrolizumab plus chemotherapy	Yes
Sasanlimab	PD-1	NCT02573259	1b/2	Locally advanced or metastatic	NSCLC, UC	Monotherapy	No
		CREST	3	Non-muscle invasive	Bladder cancer	Sasanlimab in combination with BCG	No
Atezolizumab	PD-L1	IMscin001 Part 1	1b	Advanced or metastatic	NSCLC	Monotherapy	Yes
		IMscin001 Part 2	3	Locally advanced or metastatic	NSCLC	Monotherapy	Yes
Durvalumab	PD-L1	NCT04870112	1/2a	Locally advanced or metastatic	NSCLC, SCLC	Monotherapy	Unclear
Envafoлимab	PD-L1	NCT02827968	1	Advanced	Malignant solid tumors	Monotherapy	No
		NCT05024214	1b/2	Advanced	Solid tumors	Envafoлимab plus lenvatinib	No
		NCT03667170	2	Locally advanced or metastatic	dMMR/MSI-H malignant solid tumors	Monotherapy	No
Ipilimumab	CTLA-4	CheckMate 76U	1/2	Advanced	Melanoma, UC, HCC, NSCLC, RCC	Monotherapy or in combination with subcutaneous nivolumab	Yes

PD-1, programmed death-1; PD-L1, programmed death-ligand 1; CTLA-4, cytotoxic T-lymphocyte-associated antigen 4; NSCLC, non-small cell lung cancer; RCC, renal cell carcinoma; CRC, colorectal cancer; ccRCC, clear cell renal cell carcinoma; cSCC, cutaneous squamous cell carcinoma; UC, urothelial carcinoma; SCLC, small cell lung cancer; dMMR, deficient mismatch repair; MSI-H, microsatellite instability-high; HCC, hepatocellular cancer; BCG, Bacillus Calmette-Guerin.

remain some issues to be addressed. First, the amounts of subcutaneous ICI mAbs that can lead to reductions in the patient burden and clinic time as well as the overall cost need to be explored in the current chemo-immunotherapy era. Some patients may prefer not to undergo the additional subcutaneous puncture associated with ICIs in addition to the intravenous administration of other anticancer agents

in chemo-immunotherapy regimens. In terms of costs, subcutaneous atezolizumab delivery required a 56% dose increase compared with intravenous delivery (1,875 versus 1,200 mg) to achieve similar drug exposure because of the lower bioavailability (14). In addition to the use of the drug delivery technology with rHuPH20, subcutaneous atezolizumab would incur an additional manufacturing

burden (23). Furthermore, pharmacoeconomic studies of subcutaneous ICIs will need to be updated following the upcoming arrival of intravenous ICI biosimilars that should reduce the prices of intravenous ICIs. Second, there is no clear relationship between drug exposure and efficacy for ICIs (24). Impressively, for example, the addition of ultra-low-dose nivolumab (20 mg Q3W), representing only approximately 6% of the United States Food and Drug Administration-approved flat dose of 240 mg once every 2 weeks, to triple metronomic chemotherapy for head and neck cancer showed a clinical benefit comparable to that observed with single-agent PD-1/PD-L1 inhibitors at approved doses (25). Therefore, when aiming at cost reduction, it may not be the best strategy to optimize the dose of subcutaneous ICIs based on their noninferiority to pharmacokinetic parameters of corresponding intravenous products. Exploration of alternative approaches to reduce the cost while maintaining treatment benefits is also encouraged, such as administering intravenous ICIs at the approved doses less frequently (23).

In summary, the introduction of subcutaneous mAbs in cancer immunotherapy is one of the notable areas of investigation for improving patient care and healthcare resource utilization. The IMscin001 Part 2 study (14) demonstrated that subcutaneous administration of atezolizumab (1,875 mg Q3W) was noninferior to intravenous administration of atezolizumab (1,200 mg Q3W) in terms of drug exposure, with similar efficacy, safety, and immunogenicity between the two arms. The study has led to the approval of subcutaneous atezolizumab by the Medicines and Healthcare products Regulatory Agency in the UK and the European Commission for use with all of the indications previously approved for the intravenous formulation. The fact that the innovative subcutaneous atezolizumab formulation can cut treatment time by up to 80% with simplified administration should be appreciated and will likely lead to further approvals by other agencies in the future.

### Acknowledgments

The author thanks Alison Sherwin, PhD, from Edanz (<https://jp.edanz.com/ac>) for editing a draft of this manuscript.

*Funding:* None.

### Footnote

*Provenance and Peer Review:* This article was commissioned

by the editorial office, *Translational Lung Cancer Research*. The article has undergone external peer review.

*Peer Review File:* Available at <https://tlcr.amegroups.com/article/view/10.21037/tlcr-24-63/prf>

*Conflicts of Interest:* The author has completed the ICMJE uniform disclosure form (available at <https://tlcr.amegroups.com/article/view/10.21037/tlcr-24-63/coif>). The author has no conflicts of interest to declare.

*Ethical Statement:* The author is accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

### References

1. Forde PM, Spicer J, Lu S, et al. Neoadjuvant Nivolumab plus Chemotherapy in Resectable Lung Cancer. *N Engl J Med* 2022;386:1973-85.
2. Felip E, Altorki N, Zhou C, et al. Adjuvant atezolizumab after adjuvant chemotherapy in resected stage IB-IIIa non-small-cell lung cancer (IMpower010): a randomised, multicentre, open-label, phase 3 trial. *Lancet* 2021;398:1344-57.
3. Wakelee H, Liberman M, Kato T, et al. Perioperative Pembrolizumab for Early-Stage Non-Small-Cell Lung Cancer. *N Engl J Med* 2023;389:491-503.
4. Johnson PC, Gainor JF, Sullivan RJ, et al. Immune Checkpoint Inhibitors - The Need for Innovation. *N Engl J Med* 2023;388:1529-32.
5. Pivot X, Gligorov J, Müller V, et al. Preference for subcutaneous or intravenous administration of trastuzumab in patients with HER2-positive early breast cancer (PrefHer): an open-label randomised study. *Lancet Oncol* 2013;14:962-70.
6. Rummel M, Kim TM, Aversa F, et al. Preference for

- subcutaneous or intravenous administration of rituximab among patients with untreated CD20+ diffuse large B-cell lymphoma or follicular lymphoma: results from a prospective, randomized, open-label, crossover study (PrefMab). *Ann Oncol* 2017;28:836-42.
7. O'Shaughnessy J, Sousa S, Cruz J, et al. Preference for the fixed-dose combination of pertuzumab and trastuzumab for subcutaneous injection in patients with HER2-positive early breast cancer (PHranceSCa): A randomised, open-label phase II study. *Eur J Cancer* 2021;152:223-32.
  8. Pivot X, Gligorov J, Müller V, et al. Patients' preferences for subcutaneous trastuzumab versus conventional intravenous infusion for the adjuvant treatment of HER2-positive early breast cancer: final analysis of 488 patients in the international, randomized, two-cohort PrefHer study. *Ann Oncol* 2014;25:1979-87.
  9. Bittner B, Schmidt J. Advancing Subcutaneous Dosing Regimens for Biotherapeutics: Clinical Strategies for Expedited Market Access. *BioDrugs* 2024;38:23-46.
  10. Homšek A, Spasić J, Nikolić N, et al. Pharmacokinetic characterization, benefits and barriers of subcutaneous administration of monoclonal antibodies in oncology. *J Oncol Pharm Pract* 2023;29:431-40.
  11. Alves NJ. Antibody conjugation and formulation. *Antib Ther* 2019;2:33-9.
  12. Locke KW, Maneval DC, LaBarre MJ. ENHANZE® drug delivery technology: a novel approach to subcutaneous administration using recombinant human hyaluronidase PH20. *Drug Deliv* 2019;26:98-106. Erratum in: *Drug Deliv* 2019;26:1300.
  13. Knowles SP, Printz MA, Kang DW, et al. Safety of recombinant human hyaluronidase PH20 for subcutaneous drug delivery. *Expert Opin Drug Deliv* 2021;18:1673-85.
  14. Burotto M, Zvirbule Z, Mochalova A, et al. IMscin001 Part 2: a randomised phase III, open-label, multicentre study examining the pharmacokinetics, efficacy, immunogenicity, and safety of atezolizumab subcutaneous versus intravenous administration in previously treated locally advanced or metastatic non-small-cell lung cancer and pharmacokinetics comparison with other approved indications. *Ann Oncol* 2023;34:693-702.
  15. Felip E, Burotto M, Zvirbule Z, et al. Results of a Dose-Finding Phase 1b Study of Subcutaneous Atezolizumab in Patients With Locally Advanced or Metastatic Non-Small Cell Lung Cancer. *Clin Pharmacol Drug Dev* 2021;10:1142-55.
  16. Rittmeyer A, Barlesi F, Waterkamp D, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomised controlled trial. *Lancet* 2017;389:255-65.
  17. Kverneland AH, Enevold C, Donia M, et al. Development of anti-drug antibodies is associated with shortened survival in patients with metastatic melanoma treated with ipilimumab. *Oncoimmunology* 2018;7:e1424674.
  18. Kim C, Yang H, Kim I, et al. Association of High Levels of Antidrug Antibodies Against Atezolizumab With Clinical Outcomes and T-Cell Responses in Patients With Hepatocellular Carcinoma. *JAMA Oncol* 2022;8:1825-9.
  19. Jackisch C, Kim SB, Semiglazov V, et al. Subcutaneous versus intravenous formulation of trastuzumab for HER2-positive early breast cancer: updated results from the phase III HannaH study. *Ann Oncol* 2015;26:320-5.
  20. Lonardi S, Ługowska I, O'Donnell A, et al. 616 Pharmacokinetics and safety of a subcutaneous formulation of nivolumab (NIVO SC) monotherapy: updated results from the phase 1/2 CheckMate 8KX study. *J Immunother Cancer* 2023. doi: 10.1136/jitc-2023-SITC2023.0616.
  21. Johnson ML, Braiteh F, Grilley-Olson JE, et al. Assessment of Subcutaneous vs Intravenous Administration of Anti-PD-1 Antibody PF-06801591 in Patients With Advanced Solid Tumors: A Phase 1 Dose-Escalation Trial. *JAMA Oncol* 2019;5:999-1007.
  22. Li J, Deng Y, Zhang W, et al. Subcutaneous envafolimab monotherapy in patients with advanced defective mismatch repair/microsatellite instability high solid tumors. *J Hematol Oncol* 2021;14:95.
  23. Peer CJ, Zimmerman SM, Figg WD, et al. Subcutaneous Atezolizumab: A Jab Without a Benefit. *Clin Pharmacol Drug Dev* 2022;11:134-5.
  24. Feng Y, Wang X, Bajaj G, et al. Nivolumab Exposure-Response Analyses of Efficacy and Safety in Previously Treated Squamous or Nonsquamous Non-Small Cell Lung Cancer. *Clin Cancer Res* 2017;23:5394-405.
  25. Patil VM, Noronha V, Menon N, et al. Low-Dose Immunotherapy in Head and Neck Cancer: A Randomized Study. *J Clin Oncol* 2023;41:222-32.

**Cite this article as:** Inoue Y. Subcutaneous delivery of immune checkpoint inhibitors: new route replacing intravenous administration? *Transl Lung Cancer Res* 2024;13(4):947-951. doi: 10.21037/tlcr-24-63