

Regional drivers of clutch loss reveal important trade-offs for beach-nesting birds

Brooke Maslo¹, Thomas A. Schlacher², Michael A. Weston³,
Chantal M. Huijbers², Chris Anderson², Ben L. Gilby², Andrew D. Olds²,
Rod M. Connolly⁴ and David S. Schoeman²

¹ Ecology, Evolution and Natural Resources, Rutgers, The State University of New Jersey, New Brunswick, NJ, United States

² School of Science and Engineering, University of the Sunshine Coast, Maroochydore, Australia

³ Centre for Integrative Ecology, Deakin University, Geelong, Australia

⁴ Australian Rivers Institute - Coast & Estuaries, and School of Environment, Griffith University, Gold Coast, Queensland, Australia

ABSTRACT

Coastal birds are critical ecosystem constituents on sandy shores, yet are threatened by depressed reproductive success resulting from direct and indirect anthropogenic and natural pressures. Few studies examine clutch fate across the wide range of environments experienced by birds; instead, most focus at the small site scale. We examine survival of model shorebird clutches as an index of true clutch survival at a regional scale (~200 km), encompassing a variety of geomorphologies, predator communities, and human use regimes in southeast Queensland, Australia. Of the 132 model nests deployed and monitored with cameras, 45 (34%) survived the experimental exposure period. Thirty-five (27%) were lost to flooding, 32 (24%) were depredated, nine (7%) buried by sand, seven (5%) destroyed by people, three (2%) failed by unknown causes, and one (1%) was destroyed by a dog. Clutch fate differed substantially among regions, particularly with respect to losses from flooding and predation. 'Topographic' exposure was the main driver of mortality of nests placed close to the drift line near the base of dunes, which were lost to waves (particularly during storms) and to a lesser extent depredation. Predators determined the fate of clutches not lost to waves, with the depredation probability largely influenced by region. Depredation probability declined as nests were backed by higher dunes and were placed closer to vegetation. This study emphasizes the scale at which clutch fate and survival varies within a regional context, the prominence of corvids as egg predators, the significant role of flooding as a source of nest loss, and the multiple trade-offs faced by beach-nesting birds and those that manage them.

Subjects Animal Behavior, Conservation Biology, Ecology, Zoology

Keywords Shorebirds, Sandy shore, Egg loss, Predators, Flood, Seascape

Several iconic, threatened species of the world's coastlines nest on ocean-exposed sandy shores (e.g., turtles, birds) and are thought to use nest-site selection to increase clutch success, hatchling survival, and ultimately fitness (*Refsnider & Janzen, 2010; Spencer, 2002*). Sandy shores include distinct habitat types (dunes, non-vegetated beach, surf-zone), each with variable attributes and subject to a variety of pressures (e.g., predators, people, urban

Submitted 11 May 2016
Accepted 18 August 2016
Published 13 September 2016

Corresponding author
Brooke Maslo,
brooke.maslo@rutgers.edu

Academic editor
Patricia Gandini

Additional Information and
Declarations can be found on
page 16

DOI 10.7717/peerj.2460

© Copyright
2016 Maslo et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

development) (Meager, Schlacher & Nielsen, 2012; Schlacher, Meager & Nielsen, 2014). Heterogeneity of the sandy beach environment across space and time implies that reproductive success for these species might depend on their flexibility in finding suitable nest sites.

Coastal birds are important contributors to sandy beach and dune ecosystems, transferring resources between marine and terrestrial systems (Huijbers et al., 2015), providing critical nutrient inputs (Sekercioglu, 2006), and sometimes filling the role of apex consumers (Brown et al., 2015). They also act as surrogates for beach and dune conservation and may be prime indicators of beach condition (Schlacher, Meager & Nielsen, 2014). Despite their significant role in ecosystem function and management, persistence of many coastal bird populations is severely threatened by direct and indirect anthropogenic impacts (e.g., Brinker et al., 2007; Dowding & Murphy, 2001; Van De Pol et al., 2010). Viability of resident coastal bird populations is limited by failed nesting attempts (i.e., clutch failure) and high chick mortality resulting from predation, flooding, and human disturbance (Erwin et al., 2006; Martín et al., 2015; Tjørve & Underhill, 2008).

For decades, researchers have examined potential factors influencing clutch survival of coastal birds. They have identified a suite of egg and chick predators across several taxonomic groups (Brooks et al., 2014; Ivan & Murphy, 2005), evaluated links between direct and indirect human interference and reproductive failure (Ruhlen et al., 2003; Weston & Elgar, 2007), and quantified the impact of high tides and storms on reproductive success (Brooks et al., 2013; Pol et al., 2010). These studies were generally performed on small sections of coastline under relatively uniform management and human use regimes (e.g., Dutton et al., 2005; Neuman et al., 2004), and were assumed to experience similar conditions (e.g., Hardy & Colwell, 2012; Morse, Powell & Tetreau, 2006). Thus, we have a good understanding of threats coastal ground-nesting birds encounter at local scales. For example, nests on beaches that are used heavily for human recreation may experience higher direct mortality from trampling, crushing, and vandalism (Brooks et al., 2013; Pol et al., 2010); these areas might also support higher predator densities due to the provision of additional food resources from garbage (Yasué & Dearden, 2006). Nests on uninhabited beaches may experience less chronic human disturbance, but be more vulnerable to severe acute human impacts due to a lack of regulatory presence and enforcement (Brown et al., 2015; Del Viejo et al., 2004). The suite of predators may also vary in response to variations in landscape connectivity, habitat types and the presence of invasive species (Brown et al., 2015). While such studies are important in understanding local threats and informing local management, it remains unclear whether outcomes are more broadly applicable.

Conservation practitioners have at their disposal a portfolio of management options that can be applied on a site-by-site basis (Maslo & Lockwood, 2009; Neuman et al., 2004). Management at a given site can improve reproductive success of a target species, although it can be costly (e.g., Hecht & Melvin, 2009). Further, managers must continuously monitor changing site conditions to determine when and how to appropriately intervene (Cohen et al., *in press*). Mitigation of all threats places a heavy burden on beach-nesting bird managers, who are already constrained for time and resources, and such efforts may be fruitless if nests are lost to flooding. Importantly, even the most effective management at the local scale does not guarantee population viability. Understanding the pressures

reducing clutch success at a regional scale may relieve some of the management burden by allowing managers to prioritize threat mitigation within or across regions. Identifying the relative importance of the generalized predictors of clutch failure across a regional scale may also increase population-level benefits of management. Managers would increase both their confidence in deciding appropriately where and when to apply specific interventions, as well as the efficiency with which management is implemented.

Determination of clutch fate of beach-nesting birds historically has been elusive (*Ivan & Murphy, 2005; Mabee, 1997*), preventing the identification of generalized predictors of clutch failure. Few studies investigate clutch survival over spatial scales that are large enough to encompass different environmental conditions or gradients in anthropogenic pressures. To address this gap, we examined beach-nesting bird clutch fate (using a model system of shorebird nests and eggs) along 180 km of coastline selected to encompass a diversity of geomorphological, anthropogenic, and predator environments. We identified the primary causes of clutch loss among a suite of putative factors at a regional scale. By definitively assigning clutch fate to a set of artificial nests distributed across varying beach habitats and examining an exhaustive list of predictor variables hypothesized to influence clutch failure, we determined the relative importance of predation, flooding, and human disturbance on the probability of clutch loss.

STUDY AREA

We purposefully selected four study sites within the region of southeast Queensland, Australia that differed in the types of likely mortality agents present, including two sand-barrier islands (Bribie and Moreton Islands) and two mainland sections of the ocean-exposed coastline (Noosa North Shore and Sunshine Coast; [Fig. 1](#)). Sites were separated between 12 and 27 km. The main differences between sites were the presence of exotic and feral predators (red foxes, *Vulpes vulpes*; dogs, *Canis familiaris*); the intensity of development and human use; and the types of recreational activities (camping, fishing, off-road vehicles; [Table 1](#)). Basic habitat attributes were similar between sites, all being ocean-exposed sandy beaches of intermediate morphodynamic state, backed by generally low (typically 2–4 m) foredunes of 10–40 m width (*Brown et al., 2015; Huijbers et al., 2015; Huijbers et al., 2013; Schlacher & Thompson, 2012; Schlacher & Thompson, 2013a; Schlacher & Thompson, 2013b; Schlacher et al., 2015a*). Local geomorphology varied, however, along these coasts (e.g., width and steepness of dunes and beaches), so the study area also provided an opportunity to examine the influence of these variables on clutch survival.

METHODS

We experimentally mimicked nests of red-capped plovers, *Charadrius ruficapillus*, and monitored with camera traps the fate of clutches (after *Cardilini et al., 2013*). Red-capped plovers are widely distributed across Australia and breed in Queensland throughout the year, with a peak in spring and summer (*Marchant & Higgins, 1993*). Females typically lay two eggs of cryptic colouration in a shallow scrape in the sand. For beach-nesting individuals, most nests are located on the upper beach and in foredunes, either in the open,




Figure 1 Location of study regions in southeast Queensland in eastern Australia where we monitored experimental nests on beaches and dunes on Moreton Island ($n = 38$), Bribie Island ($n = 27$), the Noosa North Shore ($n = 38$), and the Sunshine Coast ($n = 30$) in early 2015. Image: Google, 2015 Digital Globe.

Table 1 Comparison of study regions within SE-Queensland, Australia with respect to likely causes of clutch mortality associated with shore- and dune-nesting birds. Assessments are based on five years of field studies at these locations by TAS and MW.

Region	Foxes, Dogs, Cats	Off-road vehicles	Dune camping	Urban Development
Moreton Island	no	yes	yes	none
Bribie Island	yes	yes	limited	moderate
Noosa North Shore	yes	yes	yes	sparse
Sunshine Coast	yes	no	no	intense

under vegetative cover, or next to flotsam objects (Ekanayake et al., 2015a; Marchant & Higgins, 1993).

To mimic plover clutches, we placed two European quail (*Coturnix japonica*) eggs in shallow sand scrapes; quail eggs are approximately the same size, weight and colour as red-capped plover eggs and have been used previously for these types of experiments

Table 2 Summary of fates after five days of experimental clutches on ocean-exposed beaches and dunes at four study sites in southeast Queensland, Australia, during Jan–May 2015.

Fate of Nest	Moreton Island		Noosa North Shore		Bribie Island		Sunshine Coast		Total	
Carnivores										
Ghost crab			1	(3%)					1	(1%)
Torresian crow	10	(27%)	1	(3%)	12	(44%)	1	(3%)	24	(18%)
Red fox			2	(5%)			4	(13%)	6	(5%)
Snake			1	(3%)					1	(1%)
Carnivores all species	10	(27%)	5	(13%)	12	44%	5	(17%)	32	(24%)
Domestic dog							1	(3%)	1	(1%)
Flooded	17	(46%)	1	(3%)	1	(4%)	16	(53%)	35	(27%)
Intact	7	(19%)	27	(71%)	4	(15%)	7	(23%)	45	(34%)
People			5	(13%)	2	(7%)			7	(5%)
Sand burial	2	(5%)			7	(26%)			9	(7%)
Unknown	1	(3%)			1	(4%)	1	(3%)	3	(2%)
	37		38		27		30		132	

(Maguire, Stojanovic & Weston, 2010). We used survival of the model clutches as an index of real clutch survival, whilst acknowledging that the presence and behaviour of an incubating adult may alter absolute clutch loss rates (Smith, Gilchrist & Smith, 2007). However, there is no a priori expectation that systematic bias amongst regions would result from using experimental (model) clutches. We obtained unwashed quail eggs from a local hatchery 1–2 days prior to field deployment and handled them as little as possible. The quail eggs used in this experiment were 32.5 ± 0.13 (SE) mm long and 24.8 ± 0.07 mm wide, and weighed 10.8 ± 0.10 g (real Red-capped Plover eggs are 30.42 ± 0.15 mm long and 22.42 ± 0.07 mm wide; $n = 73$ MA Weston, 2016, unpublished data).

Model clutch survival experiments

We ran experiments during the late summer and early autumn of 2015 (Moreton Island: 31 Jan–07 Feb; Noosa North Shore: 3–9 March; Bribie Island: 13–19 March; Sunshine Coast: 27 Apr–1 May). We placed nests approximately 800 m apart along the shore to minimize the possibility of the same predator depredating more than one nest sequentially, whilst still achieving adequate replication and dispersion within each region. We aimed for equal replication levels at each site, but due to logistical constraints (travel times, access) the number of experimental units varied between 27 and 38, with a total of 132 nests for the entire study (Table 2).

We positioned model nests to mimic natural plover nesting habitat (Lomas et al., 2014), using the following criteria: (i) position on the beach: on the upper part of the unvegetated beach above the spring tide drift line near the base of the dunes (narrow strip of accumulated wrack deposited by swash), in the fore-dune area (the dune slope above the lowest vegetation line up to the first crest), or in the dune area (landward of the top of the fore-dune); (ii) microhabitat exposure: under vegetative cover, in the open next to flotsam, or in the open at least 2 m away from flotsam; (iii) microhabitat topography: in a hollow/swale/depression, or on a mound/ridge. We randomly assigned nests to each nest

location category using a randomisation function in Microsoft Excel, which was reapplied until we achieved ‘reasonable’ balance of nests between factor levels to enable subsequent analysis. We determined the placement of the clutch perpendicular to the water’s edge by walking to the midpoint of the dune or fore-dune at predetermined GPS coordinates and selecting the nearest location that met the randomly allocated microhabitat criteria. For beach nests, we used the upper beach just below the foredune to most closely mimic the location of real plover nests (Lomas *et al.*, 2014; Maslo, Handel & Pover, 2011). To allow equal chances for predation from both diurnal and nocturnal predators, we deployed half the nests in the early morning and the other half just after sunset. There was no significant difference ($p = 0.79$) in depredation between nests deployed near sunrise (24%) and those deployed near sunset (26%).

We then monitored nests for a 5-day period with a digital passive infrared (PIR) motion sensor camera (Scoutguard SG560Z-8M) concealed within 1–2 m of each nest, checking daily for signs of predation. Camera operations followed protocols developed for beach scavengers (Huijbers *et al.*, 2015; Lomas *et al.*, 2014). All work was conducted under Animal Ethics Permit No. AN/A/14/84 issued by the University of the Sunshine Coast, and the Scientific Purposes Permits WITK14608214 and WISP14609114 issued by the Queensland Government Department of Environment and Heritage Protection.

Environmental variables

We measured a suite of microhabitat attributes for each experimental nest. We recorded vegetation characteristics as three complementary variables: (i) small-scale (1-m² quadrat) plant cover (%) centred at the scrape (digital photograph and Coral Point Count (CPCe) software); (ii) height and distance of vegetation nearest to nest (measuring tape); and (iii) dune-wide vegetation cover (line intercept along transects extending from the base of the foredune to the landward edge of the feasible (potential) plover nesting habitat. We used a theodolite to measure four complementary metrics that described the local geomorphology: (i) distance and elevation of the nest relative to the storm drift-line (visible as an accumulation of wrack on the upper beach near the dunes); (ii) distance and elevation relative to the seaward base of the foredune (defined as a distinct rise in the angle of the beach-face); (iii) ‘exposure’ defined as the elevation of the nest relative to the nearest two profile survey points along the beach-dune transect; and (iv) dune dimensions (max. dune ridge height, width of the dune field). We obtained distances of nests from the nearest creek, rocky headland, and house using Google Earth. To index the type and intensity of human activity, we counted (once every day during nest and camera checks) the number of campsites, swimmers, fishers, and dog walkers during approximately 10 min within 100 m of each experimental nest. Data collection occurred between 0700–1100 hours consistently across all deployments. To account for potential differences in human use of shores with respect to weekends, every site was sampled at least once during the weekend and four times during the week.

We gathered wave data from wave-rider buoys operated by the Queensland Government (<http://www.qld.gov.au/environment/coasts-waterways/beach/monitoring/>), using recordings from the Brisbane buoy because it was close to the sites and contained the

longest and most complete time-series for the region. Wave heights differed significantly (ANOVA, $P < 0.005$) among sites (Fig. 2).

Data analysis

To identify the most important predictors of clutch fate, we first built a conditional inference tree using the party package in the statistical programming environment R (Hothorn, Hornik & Zeileis, 2006; R Core Team, 2014). This routine works by testing the hypothesis that the response variable clutch fate, discrete with five levels: intact; flooded; depredated; disturbed by people; or other (covered by sand, or unknown) is independent of the predictors. If this hypothesis is rejected, the routine selects the predictor that has the strongest association with the response and splits the data in two so as to best represent this association. The routine is then repeated recursively on each of the resulting nodes, stopping in each case only when the null hypothesis of no association cannot be rejected. The strength of this approach lies in the fact that because all predictors are reused at each instance of binary partitioning, the routine can reveal not only important predictors, but also statistical interactions and possible nonlinearity.

We supplemented the results of the conditional inference tree with conventional generalized linear modeling (glm) for predated clutches. Clutches lost to flooding were almost perfectly explained by the inference tree, while there were too few observations for disturbance by people to support a more detailed analysis. Our glm employed a logit link function (binomial family) and modeled clutch fate (depredated = 1; intact = 0) as a function of all available unconfounded predictors. In this sense, we considered geographic coordinates, presence of humans (including off-road vehicles and dogs), and both wave and tide metrics confounded with site, because there was no within-region variation for any of these predictors. We used a forward stepwise model-building approach based on the AIC corrected for finite sample sizes (AICc), considering only main effects. This decision to omit interactions was based on initial inspection of the data, which indicated a lack of replication and/or contrast in data across potential interaction cells (especially those associated with site). Following the forward stepwise selection process, we interrogated the final model by recalculating the AICc for all possible combinations of variables used in model building (Quinn & Keough, 2002) and used multi-model inference to determine the relative importance of predictors based on their summed Akaike weights (Burnham, Anderson & Huyvaert, 2011; Symonds & Moussalli, 2011). Finally, we used standard log-likelihood ratio tests to simplify the model by dropping predictors one at a time, starting with the least important, until only significant terms ($\alpha = 0.05$) remained.

RESULTS

Causes of model clutch loss and spatial variability

Flooding and depredation were the main causes of clutch failure, accounting for 27% and 25% of all nests lost, respectively. Of the 132 nests deployed, 45 (34%) survived the full 5-day experimental exposure period, 32 were depredated, 35 were lost due to flooding, and seven were destroyed by people (Table 3). Crows were the main predator, accounting for 75% of all depredated clutches; red foxes accounted for 19% of depredated nests, whilst




Figure 2 Wave height data recorded offshore (ca. 10 km) from the beaches on which experiments in nest survival of shore- and dune-nesting plovers were conducted in early 2015 (cf. Fig. 1 for region locations), SE-Queensland, Australia. (A) –longer-term record of maximum wave height from 1977 to 2015. (B) –Summary of maximum wave height for the first five months of each year from 2001 to 2015 for which data at comparable recording frequency (30 min) were available; (C) –maximum wave height during the study; (D) –comparison of the frequency of large wave events between 2015 and the period 2001–2014. Large wave events are defined as days on which maximum wave height exceeded the 90th (D), 95th (E) or 99th (F) percentile of the historical record; p values refer to t -tests contrasting the 2015 value with the mean of the preceding 14 years.

Table 3 Summary of final binomial generalized linear model used to predict the probability of depredation for experimental nests placed in the four study regions within SE-Queensland, Australia, 2015. Estimates for model coefficients refer to log-odds ratios and are additive on the model intercept, which represents the log-odds of depredation at Bribie Island.

	Estimate	Std Err	z	p-value
(Intercept)	2.964	1.018	2.912	0.0036
Region Moreton Island	0.078	0.9738	0.081	0.9358
Region Noosa North Shore	-3.660	0.906	-4.041	5.33×10^{-05}
Region Sunshine Coast	-0.604	0.976	-0.619	0.5360
Dune Height (m)	-1.106	0.432	-2.561	0.0104
Dist. to Vegetation (m)	0.011	0.005	2.095	0.0362

Notes.

Null deviance: 99.099 on 73 degrees of freedom.
Residual deviance: 68.765 on 68 degrees of freedom.
AIC: 80.765.

a single clutch each was eaten by a ghost crab and a snake (Table 3). Red foxes visited a further eight nests at the Noosa North Shore and Sunshine Coast, but they did not consume or detect the clutches in these instances. A domestic dog trampled one clutch, and nine clutches were buried by wind-blown sand.

Overall clutch survival differed significantly among sites (Mantel Cox test, Chi square = 20.26, df 3, $P \leq 0.001$; Fig. 3). Clutches placed on the Noosa North Shore survived longest and in highest numbers compared with all other sites (Mantel – Cox min. $P = 0.039$; min. Hazard Ratio –North Shore: Other Site = 0.40, 95% CI [0.19–0.72]). Survival patterns were comparable for nests placed at Bribie Island and the Sunshine Coast (Mantel – Cox $P = 0.96$; Hazard Ratio –Bribie Island: Sunshine Coast = 0.98, 95% CI [0.53–1.81]). Clutches on Moreton Island had significantly lower survival rates than those on the Sunshine Coast ($P = 0.02$, Hazard Ratio –Moreton: Sunshine Coast = 1.97; 95% CI [1.12–3.45]) but were comparable to those on Bribie Island ($P = 0.27$; Hazard Ratio - Moreton: Bribie Island. = 1.37, 95% CI [0.79–2.37]).

The conditional inference tree (Fig. 4) strongly resolved distance from the drift line as a significant ($P < 0.001$) predictor of flooding. Thirty-four of the 35 flooded nests in the study were located on the upper beach above the spring. Site was a strong predictor of fate of the remaining 97 clutches ($P = 0.011$). At mainland sites (Noosa North Shore and Sunshine Coast), most of the 50 clutches not situated low on the shore remained intact ($n = 34$) for the duration of the study, with the remainder depredated ($n = 9$), disturbed by people ($n = 6$), or lost to other causes ($n = 1$). The group of nests from Bribie and Moreton Islands that was not lost to flooding was further split on the basis of dune height ($P = 0.023$). Of the 31 nests located where dunes were smaller than 2.49 m in height, more than half were destroyed by predators ($n = 17$). By contrast, where dunes were taller than 4.5 m, only 3 of the 12 nests were lost to predators (Fig. 4).

The forward stepwise binomial glm identified site, distance to nearest creek, dune height and distance to nearest vegetation as the most important predictors (in order of addition to the additive model) of depredation. Subsequent multi-model inference provided a slightly




Figure 3 Survival curves for: (A) all sites (Moreton Island, Bribe Island, Noosa North Shore, and the Sunshine Coast, Queensland, Australia) and (B–E) individual sites showing identified causes of nest loss over the time span of the experiments in each region (numbers in parentheses are the actual number of experimental clutches lost attributed to a specific cause; shaded areas are 95% confidence intervals of survival).




Figure 4 Conditional inference tree indicating the primary predictors of each clutch fate category for all experimental nests placed with the study region of SE-Queensland, Australia, 2015: intact, flooded, depredated, disturbed by people, and other (buried by sand, unknown). Variables are ranked based upon the strength of their association with specific clutch fates and their quantitative values are split to best represent the relevant association.

different prioritization (relative importance of predictors based on their summed Akaike weights in parentheses): site (1.00); dune height (0.77); distance to nearest vegetation (0.72); and distance to nearest creek (0.69). Log-likelihood ratio tests indicated that removing distance to nearest creek from the model did not cause a significant deterioration in model fit ($\Delta\text{Deviance} = 2.369$, $\Delta\text{DF} = 1$, $p = 0.124$), but that the subsequent removal of distance to nearest vegetation did cause the fit to deteriorate significantly ($\Delta\text{Deviance} = 4.634$, $\Delta\text{DF} = 1$, $p = 0.031$). The final predictive model therefore included main effects for region, dune height, and distance to nearest vegetation, and explained 31.3% of the null deviation.

Coefficients from the final model indicate that at average values for dune height and distance to nearest vegetation for each site, probability of depredation was significantly higher than would be expected by chance at Bribie Island, significantly lower than would be expected by chance at Noosa North Shore, but no different from null expectation at either Moreton Island or the Sunshine Coast. The odds of depredation declined ($P = 0.010$) by a factor of 0.33 for every metre of dune height and increased ($p = 0.036$) by a factor of 1.01 for every additional metre away from the nearest vegetation (Table 3, Fig. 5).

DISCUSSION

Our broad-scale evaluation of the relative importance of multiple drivers of clutch loss on beaches suggests that predation and flooding are important causes of clutch failure in a regional context. Contrary to expectations that predation pressure is particularly




Figure 5 Influence of dune height and distance to nearest vegetation on the probability of experimental clutch depredation across all study sites within SE-Queensland, Australia, 2015. Shaded regions indicate 95% confidence intervals.

strong on human-dominated beaches (*Seneviratne et al., 2012*), predation across the entire study region was high, even at sites with no or little human development. Specifically, corvids were responsible for the majority of predated nests, complementing the suite of local-scale studies that increasingly demonstrate corvids as the primary cause of clutch loss in beach- and ground-nesting birds (*Ekanayake et al., 2015b; Madden, Arroyo & Amar, 2015*). As human commensals, corvids have rapidly expanded their global population, with the highest densities occurring in areas of significant urbanization (*Hardy & Colwell, 2012; Lauro & Tanacredi, 2002; Rees et al., 2015b*). Corvids are highly mobile, moving readily between natural and highly urbanized areas (*Whisson, Weston & Shannon, 2015*), and their impact on reproductive success can be extreme (*Burrell & Colwell, 2012; Lima, 2009*). While corvid densities are typically higher in urbanized areas (*Marzluff et al., 2001; Rees et al., 2015a*), the spillover effect of increased corvid recruitment has cascading effects on the avian species in periurban environments (*Marzluff et al., 2001*). A troubling aspect of intense corvid depredation is the limited and hitherto unconfirmed effectiveness of management responses available. Very few studies examine the effects of corvid management on breeding bird productivity, and they report small or equivocal effects (*Forys et al., 2015; Neatherlin & Marzluff, 2004; Velasco, 2015*).

Our results indicate that predation risk is higher for nests occurring in low-lying dunes away from vegetation, and the biological explanation for this phenomenon remains unclear. Corvids typically land within 1 m of the nest and approach on foot, or they discover nests while walking around open substrate (*Forys et al., 2015; Velasco, 2015*). Therefore, it is possible that corvids, as visual predators (*Ekanayake et al., 2015b*), are less able to detect or access nests in thicker vegetation (*Rees et al., 2015b*). Nest placement in shorebirds is partly about survival of the clutch, but also about survival of the parents (*Gillis et al., 2012; Tieleman, Noordwijk & Williams, 2008*); early detection of incoming predators enables adults to survive to re-clutch multiple times within a season and/or breed over several or many subsequent seasons (*Dowling & Weston, 1999; Lomas et al., 2014*). There may also be sublethal stress effects of incubating nests away from cover (*Amat & Masero, 2004*).

The sizeable proportion of clutches that failed due to flooding emphasized the severity of flooding as a threat to coastal ground-nesting birds (*Pol et al., 2010; Van De Pol et al., 2010*). Although many of the flooded nests in this study occurred during stormy weather on the Sunshine Coast (53%) and Moreton Island (45%), we found little evidence that the type of large wave occurrences that led to clutch losses in 2015 were atypical events. The number of days in which maximum wave height exceeded the 90th percentile of historical records for the first five months of each year ($n = 17$ days) was not significantly different from the mean of 14.6 days in the previous 14 years, with only four years (2001, 2008, 2009, 2013) exceeding values recorded in 2015 (*Fig. 2*). Similarly, wave heights exceeded the top 1% of the historical record on only a single day during the present study. The impact of storm or high tide events can be catastrophic to bird reproduction, particularly if storms occur late in the incubation period when there is little time for re-nesting. There exists considerable uncertainty about when storms will arise, where they will be most severe, and for how long habitats will be impacted (*Hemer et al., 2013; IPCC, 2013; Schlacher et al., 2015b*), suggesting that assessment of flood risk for beach-nesting birds is imperfect (*Lomas*

et al., 2014). Human-mediated flood risk abatement, if feasible, may be of great importance to population viability.

Human disturbance accounted for the loss of ≤ 9 clutches at any site and only 11% across the entire region, illustrating the relatively weak influence of this source of failure on beach-nesting bird reproductive success. Human presence in the vicinity of a nest was not related to the probability of clutch loss to direct human causes, as would be expected. In fact, we found no significant effect of dogs, camping, off-road vehicles or proximity to development on the probability of a nest being disturbed. These results suggest that passive management of human disturbance (i.e., symbolic fencing, signage) is effective in reducing anthropogenic impacts to beach-nesting birds in a regional context. Where human disturbance of nesting birds is severe in a specific location, localized management can address the problem (*Del Viejo et al.*, 2004; *Ruhlen et al.*, 2003; *Weston & Elgar*, 2007).

CAVEATS

The results described here result from the placement of model clutches within suitable plover habitat. Model clutches are used extensively in research and when implemented carefully can reveal important ecological patterns and processes (*Berry & Lill*, 2003). We acknowledge beach-nesting birds theoretically select nesting sites to reduce the risks of clutch loss, perhaps basing their choice in part on previous nesting experiences. We also recognise that model clutches are not associated with incubator behaviour, which either can be protective (i.e., defence) or may render clutches more vulnerable to predation (e.g., visual and scent cues) (*Ekanayake et al.*, 2015a). Indeed, model clutch studies are enlightening in terms of physical destruction of the eggs (e.g., *Buick & Paton*, 1989), but they shed no light on the impacts of disturbance (disruption of incubation which may reduce egg viability), even though disturbance to breeding shorebirds is considered a conservation threat (*Maslo, Burger & Handel*, 2012; *Meager, Schlacher & Nielsen*, 2012; *Powell & Collier*, 2000; *Quinn et al.*, 1996; *Schlacher, Nielsen & Weston*, 2013; *Weston, Schlacher & Lynn*, 2014). In a review of 80 studies, (*Major & Kendal*, 1996) report that artificial nests underestimate actual survival of real nests, while cameras on nests either do not affect or slightly increase clutch survival (*Richardson, Gardali & Jenkins*, 2009; *Sanders & Maloney*, 2002). *Ekanayake et al.* (2015a) and *Ekanayake et al.* (2015b) also used model clutches on a study of red-capped plovers in southern Victoria and confirmed the comparability of natural and model clutches in identifying egg predators.

Model nests also permit otherwise infeasible studies to be performed. While common, widespread nesting species may enable the study of real nests in some areas to address some research questions (none were available on coasts in subtropical Australia). However, this would represent a biased research effort because studies would be confined to species apparently coping well with prevailing conditions; traditional yet abandoned habitats could not be assessed for threats. Artificial nests permit an examination of egg predation risk in areas of suitable but often unoccupied habitat, as we have done here. Calibration of this model egg study with real clutch fate and survival (impossible here because no such data, which is strictly comparable, are available) would not only be confirmatory, but would

shed light on the utility of studies that use model eggs to guide real nest management. We also note that survival estimates of real clutches are often biased (e.g., *Nichols et al., 1984*).

MANAGEMENT IMPLICATIONS

As coastal bird populations continue to decline globally (e.g., *Rodrigues et al., 2004*), the enhancement of reproductive success through active management plays an increasingly critical role in species' persistence. Our analysis suggests that management of human disturbance through symbolic fencing, signage, and regulatory measures across a region is likely enough to effectively mitigate anthropogenic impacts. The significant influence of predation and flooding on clutch loss at the regional scale demonstrates that actively addressing these threats will lead to increased viability of beach-nesting bird populations. Predator management is typically performed at the site scale and can be effective in the short-term, and targeted application may increase clutch success (e.g., exclosures, electric fencing, removal; *Maslo & Lockwood, 2009; Neuman et al., 2004*). However, predator mortality may elicit a compensatory response (i.e., increased reproduction) in affected populations and new individuals can quickly occupy open niche space (*Harding, Doak & Albertson, 2001; Smith et al., 2010*). Both corvids and foxes can disperse great distances from their natal territories (*Dekker, Stein & Heitkönig, 2001; Marzluff et al., 2001*); therefore, targeting source populations within the region may have trickle-down benefits to beach-nesting birds. Predation management, however, may be unnecessary and inefficient if nests are highly vulnerable to flooding.

Managing flood risk of clutches is likely difficult to implement. Possibilities include modifying habitats to maintain nesting areas at higher elevations, for example through dredge spoil or ecologically sensitive beach nourishment. Identification of the sites within a region where habitat modifications have the highest likelihood of providing benefits to the population will be critical. At the location scale, moving nests or raising them before flood events can also be attempted. In flood prone locations, a tractable strategy may be to assess a clutch's vulnerability to flooding (*Haig et al., 2005; Sanders & Maloney, 2002; Thomas, Lanctot & Székely, 2006*) and then manage failure risk for the subset of nests that are not flood prone (this also preserves any learnt, adaptive, nest placement whereby birds learn to avoid flood prone habitats).

In short, conservation of coastal bird species in the presence of multiple threats and measureable uncertainty requires practitioners to make hard choices about management intervention. An understanding of the driving forces of clutch success at a regional scale offers managers some confidence in developing strategies that promote the viability of coastal bird populations (*Meager, Schlacher & Nielsen, 2012*) rather than mitigating impacts on a site-by-site or location-by-location basis.

ACKNOWLEDGEMENTS

We especially thank I Franks for many hours help in the field and contributions by Prof. Ischlberg. Nicholas Horrocks (and two anonymous reviewers) commented on the manuscript in a most helpful and constructive manner.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

This study was supported by the Australian government's Collaborative Research Network (CRN). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:
Australian government's Collaborative Research Network.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Brooke Maslo analyzed the data, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Thomas A. Schlacher conceived and designed the experiments, performed the experiments, analyzed the data, contributed reagents/materials/analysis tools, wrote the paper, prepared figures and/or tables, reviewed drafts of the paper.
- Michael A. Weston conceived and designed the experiments, performed the experiments, contributed reagents/materials/analysis tools, wrote the paper, reviewed drafts of the paper.
- Chantal M. Huijbers, Ben L. Gilby, Andrew D. Olds and Rod M. Connolly wrote the paper, reviewed drafts of the paper.
- Chris Anderson conceived and designed the experiments, performed the experiments.
- David S. Schoeman analyzed the data, prepared figures and/or tables, reviewed drafts of the paper.

Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

All work was conducted under Animal Ethics Permit No. AN/A/14/84 issued by the University of the Sunshine Coast, and the Scientific Purposes Permits WITK14608214 and WISP14609114 issued by the Queensland Government Department of Environment and Heritage Protection.

Data Availability

The following information was supplied regarding data availability:

Maslo B, Schlacher T, Michael W, Huijbers C, Anderson C, Gilby B, Olds A, Connolly R, Schoeman D Data from: Regional Drivers of Model Clutch Loss Reveal Important Trade-offs for Beach-nesting Birds: <http://dx.doi.org/10.5061/dryad.hn132>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.2460#supplemental-information>.

REFERENCES

- Amat JA, Masero JA. 2004.** Predation risk on incubating adults constrains the choice of thermally favourable nest sites in a plover. *Animal Behaviour* **67**:293–300 DOI [10.1016/j.anbehav.2003.06.014](https://doi.org/10.1016/j.anbehav.2003.06.014).
- Berry L, Lill A. 2003.** Do predation rates on artificial nests accurately predict predation rates on natural nests? The effects of nest type, egg type and nest-site characteristics. *Emu* **103**:207–214 DOI [10.1071/MU02054](https://doi.org/10.1071/MU02054).
- Brinker DF, McCann JM, Williams B, Watts BD. 2007.** Colonial-nesting seabirds in the Chesapeake Bay region: where have we been and where are we going? *Waterbirds* **30**:93–104 DOI [10.1675/1524-4695\(2007\)030\[0093:CSITCB\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2007)030[0093:CSITCB]2.0.CO;2).
- Brooks GL, Sanders FJ, Gerard PD, Jodice PG. 2013.** Daily survival rate for nests and chicks of least terns (*Sternula antillarum*) at natural nest sites in South Carolina. *Waterbirds* **36**:1–10 DOI [10.1675/063.036.0101](https://doi.org/10.1675/063.036.0101).
- Brooks GL, Sanders FJ, Gerard PD, Jodice PG. 2014.** Daily survival rate for nests of Black Skimmers from a core breeding area of the southeastern USA. *The Wilson Journal of Ornithology* **126**:443–450 DOI [10.1676/13-136.1](https://doi.org/10.1676/13-136.1).
- Brown MB, Schlacher TA, Schoeman DS, Weston MA, Huijbers CM, Olds AD, Connolly RM. 2015.** Invasive carnivores alter ecological function and enhance complementarity in scavenger assemblages on ocean beaches. *Ecology* **96**:2715–2725 DOI [10.1890/15-0027.1](https://doi.org/10.1890/15-0027.1).
- Buick AM, Paton DC. 1989.** Impact of off-road vehicles on the nesting success of Hooded Plovers (*Charadrius rubricollis*) in the Coorong region of South Australia. *Emu* **89**:159–172.
- Burnham KP, Anderson DR, Huyvaert KP. 2011.** AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology* **65**(1):23–35 DOI [10.1007/s00265-010-1029-6](https://doi.org/10.1007/s00265-010-1029-6).
- Burrell NS, Colwell MA. 2012.** Direct and indirect evidence that productivity of Snowy Plovers *Charadrius nivosus* varies with occurrence of a nest predator. *Wildfowl* **62**:204–223.
- Cardilini AP, Weston MA, Nimmo DG, Dann P, Sherman CD. 2013.** Surviving in sprawling suburbs: suburban environments represent high quality breeding habitat for a widespread shorebird. *Landscape and Urban Planning* **115**:72–80 DOI [10.1016/j.landurbplan.2013.04.001](https://doi.org/10.1016/j.landurbplan.2013.04.001).
- Cohen JB, Hecht A, Robinson KF, Osnas EE, Tyre AJ, Brinker DF, Davis C, Hake M, Johnson L, King E, Kocek A, Maslo B, Melvin SM. 2016.** Structured decision making for the use of predator exclosures to recover a threatened species, the Atlantic Coast piping plover. *Ecosphere* In Press.

- Dekker JJ, Stein A, Heitkönig I. 2001.** A spatial analysis of a population of red fox (*Vulpes vulpes*) in the Dutch coastal dune area. *Journal of Zoology* 255:505–510 DOI 10.1017/S0952836901001583.
- Del Viejo AM, Vega X, González M, Sánchez J. 2004.** Disturbance sources, human predation and reproductive success of seabirds in tropical coastal ecosystems of Sinaloa State, Mexico. *Bird Conservation International* 14:191–202 DOI 10.1017/S0959270904000243.
- Dowding JE, Murphy EC. 2001.** The impact of predation by introduced mammals on endemic shorebirds in New Zealand: a conservation perspective. *Biological Conservation* 99:47–64 DOI 10.1016/S0006-3207(00)00187-7.
- Dowling B, Weston MA. 1999.** Managing a breeding population of the Hooded Plover *Thinornis rubricollis* in a high-use recreational environment. *Bird Conservation International* 9:255–270 DOI 10.1017/S0959270900003440.
- Dutton DL, Dutton PH, Chaloupka M, Boulon RH. 2005.** Increase of a Caribbean leatherback turtle *Dermochelys coriacea* nesting population linked to long-term nest protection. *Biological Conservation* 126:186–194 DOI 10.1016/j.biocon.2005.05.013.
- Ekanayake KB, Weston MA, Nimmo DG, Maguire GS, Endler JA, Küpper C. 2015a.** The bright incubate at night: sexual dichromatism and adaptive incubation division in an open-nesting shorebird. *Proceedings of the Royal Society of London B: Biological Sciences* 282:20143026 DOI 10.1098/rspb.2014.3026.
- Ekanayake KB, Whisson DA, Tan LXL, Weston MA. 2015b.** Intense predation of non-colonial, ground-nesting bird eggs by corvid and mammalian predators. *Wildlife Research* 42:518–528 DOI 10.1071/WR15080.
- Erwin RM, Sanders GM, Prosser DJ, Cahoon DR. 2006.** High tides and rising seas: potential effects on estuarine waterbirds. *Studies in Avian Biology* 32:214–228.
- Forys EA, Hopkins D, Ingham P, Miller M, Gluckman L. 2015.** Do effigies deter fish crows hunting in a black skimmer colony mid-season? *Southeastern Naturalist* 14:635–640 DOI 10.1656/058.014.0405.
- Gillis H, Gauffre B, Huot R, Bretagnolle V. 2012.** Vegetation height and egg coloration differentially affect predation rate and overheating risk: an experimental test mimicking a ground-nesting bird. *Canadian Journal of Zoology* 90:694–703 DOI 10.1139/z2012-035.
- Haig SM, Ferland CL, Cuthbert FJ, Dingleline J, Goossen JP, Hecht A, McPhillips N. 2005.** A complete species census and evidence for regional declines in Piping Plovers. *Journal of Wildlife Management* 69:160–173 DOI 10.2193/0022-541X(2005)069<0160:ACSCAE>2.0.CO;2.
- Harding EK, Doak DF, Albertson JD. 2001.** Evaluating the effectiveness of predator control: the non native red fox as a case study. *Conservation Biology* 15:1114–1122 DOI 10.1046/j.1523-1739.2001.0150041114.x.
- Hardy MA, Colwell MA. 2012.** Factors influencing Snowy Plover nest survival on ocean-fronting beaches in coastal northern California. *Waterbirds* 35:503–656 DOI 10.1675/063.035.0401.

- Hecht A, Melvin SM. 2009.** Expenditures and effort associated with recovery of breeding Atlantic Coast Piping Plovers. *The Journal of Wildlife Management* **73**:1099–1107 DOI [10.2193/2008-061](https://doi.org/10.2193/2008-061).
- Hemer MA, Fan Y, Mori N, Semedo A, Wang XL. 2013.** Projected changes in wave climate from a multi-model ensemble. *Nature Climate Change* **3**:471–476 DOI [10.1038/nclimate1791](https://doi.org/10.1038/nclimate1791).
- Hothorn T, Hornik K, Zeileis A. 2006.** Unbiased recursive partitioning: a conditional inference framework. *Journal of Computational and Graphical Statistics* **15**:651–674 DOI [10.1198/106186006X133933](https://doi.org/10.1198/106186006X133933).
- Huijbers CM, Schlacher TA, Schoeman DS, Olds AD, Weston MA, Connolly RM. 2015.** Limited functional redundancy in vertebrate scavenger guilds fails to compensate for the loss of raptors from urbanized sandy beaches. *Diversity and Distributions* **21**:55–63 DOI [10.1111/ddi.12282](https://doi.org/10.1111/ddi.12282).
- Huijbers CM, Schlacher TA, Schoeman DS, Weston MA, Connolly RM. 2013.** Urbanisation alters processing of marine carrion on sandy beaches. *Landscape and Urban Planning* **119**:1–8 DOI [10.1016/j.landurbplan.2013.06.004](https://doi.org/10.1016/j.landurbplan.2013.06.004).
- IPCC. 2013.** Climate change 2013: the physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change. In: *Long-term climate change: projections, commitments and irreversibility*. Cambridge: Cambridge University Press.
- Ivan JS, Murphy RK. 2005.** What preys on piping plover eggs and chicks? *Wildlife Society Bulletin* **33**:113–119 DOI [10.2193/0091-7648\(2005\)33\[113:WPOPPE\]2.0.CO;2](https://doi.org/10.2193/0091-7648(2005)33[113:WPOPPE]2.0.CO;2).
- Lauro B, Tanacredi J. 2002.** An examination of predatory pressures on Piping Plovers nesting at Breezy Point, New York. *Waterbirds* **25**:401–409 DOI [10.1675/1524-4695\(2002\)025\[0401:AEOPPO\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2002)025[0401:AEOPPO]2.0.CO;2).
- Lima SL. 2009.** Predators and the breeding bird: behavioral and reproductive flexibility under the risk of predation. *Biological Reviews* **84**:485–513 DOI [10.1111/j.1469-185X.2009.00085.x](https://doi.org/10.1111/j.1469-185X.2009.00085.x).
- Lomas SC, Whisson DA, Maguire GS, Tan LX, Guay P-J, Weston MA. 2014.** The influence of cover on nesting red-capped plovers: a trade-off between thermoregulation and predation risk? *The Victorian Naturalist* **131**(4):115–127.
- Mabee TJ. 1997.** Using eggshell evidence to determine nest fate of shorebirds. *The Wilson Bulletin* **109**(2):307–313.
- Madden CF, Arroyo B, Amar A. 2015.** A review of the impacts of corvids on bird productivity and abundance. *Ibis* **157**:1–16 DOI [10.1111/ibi.12223](https://doi.org/10.1111/ibi.12223).
- Maguire GS, Stojanovic D, Weston MA. 2010.** Conditioned taste aversion reduces fox depredation on model eggs on beaches. *Wildlife Research* **36**:702–708 DOI [10.1071/WR09123](https://doi.org/10.1071/WR09123).
- Major RE, Kendal CE. 1996.** The contribution of artificial nest experiments to understanding avian reproductive success: a review of methods and conclusions. *Ibis* **138**:298–307 DOI [10.1111/j.1474-919X.1996.tb04342.x](https://doi.org/10.1111/j.1474-919X.1996.tb04342.x).
- Marchant S, Higgins PJ. 1993.** *Handbook of Australian, New Zealand and Antarctic Birds, Volume 2 (Raptors to Lapwings)*. Melbourne: Oxford University Press.

- Martín B, Delgado S, Adl C, Tirado S, Ferrer M. 2015.** Effects of human presence on the longterm trends of migrant and resident shorebirds: evidence of local population declines. *Animal Conservation* **18**:73–81 DOI [10.1111/acv.12139](https://doi.org/10.1111/acv.12139).
- Marzluff JM, McGowan KJ, Donnelly R, Knight RL. 2001.** Causes and consequences of expanding American Crow populations. In: *Avian ecology and conservation in an urbanizing world*. New York: Springer, 331–363.
- Maslo B, Burger J, Handel SN. 2012.** Modeling foraging behavior of piping plovers to evaluate habitat restoration success. *The Journal of Wildlife Management* **76**:181–188 DOI [10.1002/jwmg.210](https://doi.org/10.1002/jwmg.210).
- Maslo B, Handel SN, Pover T. 2011.** Restoring beaches for Atlantic Coast Piping Plovers (*Charadrius melodus*): a classification and regression tree analysis of nest-site selection. *Restoration Ecology* **19**:194–203 DOI [10.1111/j.1526-100X.2010.00709.x](https://doi.org/10.1111/j.1526-100X.2010.00709.x).
- Maslo B, Lockwood JL. 2009.** Evidence-based decisions on the use of predator enclosures in shorebird conservation. *Biological Conservation* **142**:3213–3218 DOI [10.1016/j.biocon.2009.07.034](https://doi.org/10.1016/j.biocon.2009.07.034).
- Meager JJ, Schlacher TA, Nielsen T. 2012.** Humans alter habitat selection of birds on ocean-exposed sandy beaches. *Diversity & Distributions* **18**:294–306 DOI [10.1111/j.1472-4642.2011.00873.x](https://doi.org/10.1111/j.1472-4642.2011.00873.x).
- Morse JA, Powell AN, Tetreau MD. 2006.** Productivity of Black Oystercatchers: effects of recreational disturbance in a national park. *The Condor* **108**:623–633 DOI [10.1650/0010-5422\(2006\)108\[623:POBOEO\]2.0.CO;2](https://doi.org/10.1650/0010-5422(2006)108[623:POBOEO]2.0.CO;2).
- Neatherlin EA, Marzluff JM. 2004.** Responses of American crow populations to campgrounds in remote native forest landscapes. *Journal of Wildlife Management* **68**:708–718 DOI [10.2193/0022-541X\(2004\)068\[0708:ROACPT\]2.0.CO;2](https://doi.org/10.2193/0022-541X(2004)068[0708:ROACPT]2.0.CO;2).
- Neuman KK, Page GW, Stenzel LE, Warriner JC, Warriner JS. 2004.** Effect of mammalian predator management on Snowy Plover breeding success. *Waterbirds* **27**:257–263 DOI [10.1675/1524-4695\(2004\)027\[0257:EOMPMO\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2004)027[0257:EOMPMO]2.0.CO;2).
- Nichols JD, Percival HF, Coon RA, Conroy MJ, Hensler GL, Hines JA. 1984.** Observer visitation frequency and success of mourning dove nests; a field experiment. *Auk* **101**:398–402.
- Pol MVD, Vindenes Y, Sæther B-E, Engen S, Ens BJ, Oosterbeek K, Tinbergen JM. 2010.** Effects of climate change and variability on population dynamics in a long-lived shorebird. *Ecology* **91**:1192–1204 DOI [10.1890/09-0410.1](https://doi.org/10.1890/09-0410.1).
- Powell AN, Collier CL. 2000.** Habitat use and reproductive success of Western Snowy Plovers at new nesting areas created for California Least Terns. *The Journal of Wildlife Management* **64**:24–33 DOI [10.2307/3802971](https://doi.org/10.2307/3802971).
- Quinn GP, Keough MJ. 2002.** *Experimental design and data analysis for biologists*. New York: Cambridge University Press.
- Quinn J, Morris R, Blokpoel H, Weseloh D, Ewins P. 1996.** Design and management of bird nesting habitat: tactics for conserving colonial waterbird biodiversity on artificial islands in Hamilton Harbour, Ontario. *Canadian Journal of Fisheries and Aquatic Sciences* **53**:45–57 DOI [10.1139/f95-260](https://doi.org/10.1139/f95-260).

- R Core Team.** 2014. *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing. Available at <http://www.r-project.org/>.
- Rees J, Webb J, Crowther M, Letnic M.** 2015a. Carrion subsidies provided by fishermen increase predation of beachnesting bird nests by facultative scavengers. *Animal Conservation* 18:44–49 DOI [10.1111/acv.12133](https://doi.org/10.1111/acv.12133).
- Rees JD, Webb JK, Crowther MS, Letnic M.** 2015b. Ravens are a key threat to beach-nesting birds. *Australian Field Ornithology* 32(2):100–107.
- Refsnider JM, Janzen FJ.** 2010. Putting eggs in one basket: ecological and evolutionary hypotheses for variation in oviposition-site choice. *Annual Review of Ecology, Evolution, and Systematics* 41:39–57 DOI [10.1146/annurev-ecolsys-102209-144712](https://doi.org/10.1146/annurev-ecolsys-102209-144712).
- Richardson T, Gardali T, Jenkins SH.** 2009. Review and metaanalysis of camera effects on avian nest success. *The Journal of Wildlife Management* 73:287–293 DOI [10.2193/2007-566](https://doi.org/10.2193/2007-566).
- Rodrigues AS, Andelman SJ, Bakarr MI, Boitani L, Brooks TM, Cowling RM, Fishpool LD, Da Fonseca GA, Gaston KJ, Hoffmann M.** 2004. Effectiveness of the global protected area network in representing species diversity. *Nature* 428:640–643 DOI [10.1038/nature02422](https://doi.org/10.1038/nature02422).
- Ruhlen TD, Abbott S, Stenzel LE, Page GW.** 2003. Evidence that human disturbance reduces Snowy Plover chick survival. *Journal of Field Ornithology* 74:300–304 DOI [10.1648/0273-8570-74.3.300](https://doi.org/10.1648/0273-8570-74.3.300).
- Sanders MD, Maloney RF.** 2002. Causes of mortality at nests of ground-nesting birds in the Upper Waitaki Basin, South Island, New Zealand: a 5-year video study. *Biological Conservation* 106:225–236 DOI [10.1016/S0006-3207\(01\)00248-8](https://doi.org/10.1016/S0006-3207(01)00248-8).
- Schlacher TA, Meager JJ, Nielsen T.** 2014. Habitat selection in birds feeding on ocean shores: landscape effects are important in the choice of foraging sites by oystercatchers. *Marine Ecology* 35:67–76 DOI [10.1111/maec.12055](https://doi.org/10.1111/maec.12055).
- Schlacher TA, Nielsen T, Weston MA.** 2013. Human recreation alters behaviour profiles of non-breeding birds on open-coast sandy shores. *Estuarine, Coastal and Shelf Science* 118:31–42 DOI [10.1016/j.ecss.2012.12.016](https://doi.org/10.1016/j.ecss.2012.12.016).
- Schlacher TA, Thompson L.** 2012. Beach recreation impacts benthic invertebrates on ocean-exposed sandy shores. *Biological Conservation* 147:123–132 DOI [10.1016/j.biocon.2011.12.022](https://doi.org/10.1016/j.biocon.2011.12.022).
- Schlacher TA, Thompson L.** 2013a. Environmental control of community organisation on ocean-exposed sandy beaches. *Marine and Freshwater Research* 64:119–129 DOI [10.1071/MF12172](https://doi.org/10.1071/MF12172).
- Schlacher TA, Thompson L.** 2013b. Spatial structure on ocean-exposed sandy beaches: faunal zonation metrics and their variability. *Marine Ecology Progress Series* 478:43–55 DOI [10.3354/meps10205](https://doi.org/10.3354/meps10205).
- Schlacher TA, Weston MA, Lynn D, Schoeman DS, Huijbers CM, Olds AD, Masters S, Connolly RM.** 2015a. Conservation gone to the dogs: when canids rule the beach in small coastal reserves. *Biodiversity and Conservation* 24:493–509 DOI [10.1007/s10531-014-0830-3](https://doi.org/10.1007/s10531-014-0830-3).

- Schlacher TA, Weston MA, Schoeman DS, Olds AD, Huijbers CM, Connolly RM. 2015b.** Golden opportunities: a horizon scan to expand sandy beach ecology. *Estuarine, Coastal and Shelf Science* **157**:1–6 DOI [10.1016/j.ecss.2015.02.002](https://doi.org/10.1016/j.ecss.2015.02.002).
- Sekercioglu CH. 2006.** Increasing awareness of avian ecological function. *Trends in Ecology and Evolution* **21**:464–471 DOI [10.1016/j.tree.2006.05.007](https://doi.org/10.1016/j.tree.2006.05.007).
- Seneviratne SI, Nicholls N, Easterling D, Goodess CM, Kanae S, Kossin J, Luo Y, Marengo J, McInnes K, Rahimi M. 2012.** Changes in climate extremes and their impacts on the natural physical environment: an overview of the IPCC SREX report. In: *EGU General Assembly Conference Abstracts. Vol. 14*.
- Smith PA, Gilchrist H, Smith JN. 2007.** Effects of nest habitat, food, and parental behavior on shorebird nest success. *The Condor* **109**:15–31 DOI [10.1650/0010-5422\(2007\)109\[15:EONHFA\]2.0.CO;2](https://doi.org/10.1650/0010-5422(2007)109[15:EONHFA]2.0.CO;2).
- Smith RK, Pullin AS, Stewart GB, Sutherland WJ. 2010.** Effectiveness of predator removal for enhancing bird populations. *Conservation Biology* **24**:820–829 DOI [10.1111/j.1523-1739.2009.01421.x](https://doi.org/10.1111/j.1523-1739.2009.01421.x).
- Spencer R-J. 2002.** Experimentally testing nest site selection: fitness trade-offs and predation risk in turtles. *Ecology* **83**:2136–2144 DOI [10.1890/0012-9658\(2002\)083\[2136:ETNSSF\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2002)083[2136:ETNSSF]2.0.CO;2).
- Symonds MRE, Moussalli A. 2011.** A brief guide to model selection, multimodel inference and model averaging in behavioural ecology using Akaike's information criterion. *Behavioral Ecology and Sociobiology* **65**(1):13–21 DOI [10.1007/s00265-010-1037-6](https://doi.org/10.1007/s00265-010-1037-6).
- Thomas G, Lanctot R, Székely T. 2006.** Can intrinsic factors explain population declines in North American breeding shorebirds? A comparative analysis. *Animal Conservation* **9**:252–258 DOI [10.1111/j.1469-1795.2006.00029.x](https://doi.org/10.1111/j.1469-1795.2006.00029.x).
- Tieleman BI, Van Noordwijk HJ, Williams JB. 2008.** Nest site selection in a hot desert: trade-off between microclimate and predation risk. *The Condor* **110**:116–124 DOI [10.1525/cond.2008.110.1.116](https://doi.org/10.1525/cond.2008.110.1.116).
- Tjørve KMC, Underhill L. 2008.** Influence of disturbance and predation on breeding success of the African Black Oystercatcher, *Haematopus moquini*, on Robben Island, South Africa. *Waterbirds* **31**:83–96 DOI [10.1675/1524-4695\(2008\)31\[83:IODAPO\]2.0.CO;2](https://doi.org/10.1675/1524-4695(2008)31[83:IODAPO]2.0.CO;2).
- Van De Pol M, Ens BJ, Heg D, Brouwer L, Krol J, Maier M, Exo KM, Oosterbeek K, Lok T, Eising CM. 2010.** Do changes in the frequency, magnitude and timing of extreme climatic events threaten the population viability of coastal birds? *Journal of Applied Ecology* **47**:720–730 DOI [10.1111/j.1365-2664.2010.01842.x](https://doi.org/10.1111/j.1365-2664.2010.01842.x).
- Velasco VN. 2015.** Investigation of non-lethal electric shock on American crows as a predator aversion treatment for reducing depredation on California least tern eggs. MS thesis, Seaver College of Science and Engineering.
- Weston MA, Elgar MA. 2007.** Responses of incubating hooded plovers (*Thinornis rubricollis*) to disturbance. *Journal of Coastal Research* **23**(3):569–576 DOI [10.2112/04-0151.1](https://doi.org/10.2112/04-0151.1).

- Weston MA, Schlacher TA, Lynn D. 2014.** Pro-environmental beach driving is uncommon and ineffective in reducing disturbance to beach-dwelling birds. *Environmental Management* **53**:999–1004 DOI [10.1007/s00267-014-0256-4](https://doi.org/10.1007/s00267-014-0256-4).
- Whisson DA, Weston MA, Shannon K. 2015.** Home range, habitat use and movements by the little raven (*Corvus mellori*) in a coastal peri-urban landscape. *Wildlife Research* **42**:500–508 DOI [10.1071/WR15039](https://doi.org/10.1071/WR15039).
- Yasué M, Dearden P. 2006.** The potential impact of tourism development on habitat availability and productivity of Malaysian plovers *Charadrius peronii*. *Journal of Applied Ecology* **43**:978–989 DOI [10.1111/j.1365-2664.2006.01204.x](https://doi.org/10.1111/j.1365-2664.2006.01204.x).