

Response to Alzheimer's research paper

H Mina Tait

Department of Biomedical Science,
The University of Sheffield,
Sheffield, UK

Dear editor

Zhang et al have recently published an interesting article in *Clinical Interventions in Aging* entitled "Atorvastatin attenuates the production of IL-1 β , IL-6, and TNF- α in the hippocampus of an amyloid β 1-42-induced rat model of Alzheimer's disease".¹ The conclusion that "IL-1 β , IL-6, and TNF- α were significantly decreased in the atorvastatin-treated AD group than that in the AD group" is not justified by the data presented in Figures 1, 2, and 3. Given that the AD group and the atorvastatin-treated AD group showed no significant difference in numbers of cells positive for interleukin-1 β , interleukin-6, or tumor necrosis factor- α , and in some cases there may be a trend towards an increase in positive cells in the atorvastatin-treated AD group, it is hard to see how they came to this conclusion.

In their introduction, Li et al state that "Cholesterol-lowering agents, particularly 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase inhibitors (also known as statins), have been reported to be capable of increasing the ratio of alpha to beta secretase activity and then increasing the concentrations of extracellular A β ."

The conventional wisdom, supported by the references that they cite (eg, Buxbaum et al²), is that changing the ratio in this way is likely to lead to a decrease, not an increase, in extracellular A β . I would very much like to hear the authors' and your views on these issues.

Reference

1. Zhang YY, Fan YC, Wang M, Wang D, Li XH. Atorvastatin attenuates the production of IL-1 β , IL-6, and TNF- α in the hippocampus of an amyloid β 1-42-induced rat model of Alzheimer's disease. *Clin Interv Aging*. 2013;8:103–110.
2. Buxbaum JD, Geoghagen NS, Friedhoff LT. Cholesterol depletion with physiological concentrations of a statin decreases the formation of the Alzheimer amyloid A β peptide. *J Alzheimers Dis*. 2001;3:221–229.

Correspondence: H Mina Tait
Group X10, Western Bank,
Sheffield, S10 2TN, UK
Tel +44 011 4222 0000
Fax +44 011 4276 5413
Email hmtait1@sheffield.ac.uk

Response

Xiao-Hong Li

Department of Neurology, Jinan Central Hospital affiliated to Shandong University and School of Medicine, Shandong University, Jinan, People's Republic of China

Correspondence: Xiao-Hong Li
Department of Neurology, Jinan
Central Hospital affiliated to Shandong University and School of
Medicine, Shandong University, Jiefang Road 105,
Jinan 250013, People's Republic of China
Email xiaohongli2011@yeah.net

Dear editor

We thank Ms Tait for her interest in our recently published paper in *Clinical Interventions in Aging*. Regarding the first issue raised, we have checked the original manuscript and raw data of our experiment and found that there is a mis-pairing of the columns for group 2 and group 3 in Figures 3, 4, and 5. Below are the corrected Figures 3E, 4E, and 5E based on the raw data. In these figures, the original column of group

2 has been modified to group 3, and vice versa. From these figures, it is clear that the atorvastatin-treated AD group had a significant reduction in the numbers of cells positive for interleukin-1 β , interleukin-6, or tumor necrosis factor- α in the atorvastatin-treated AD group.

We agree about the point concerning “the changing of ratio of alpha to beta secretase is likely to lead to a decrease in extracellular A β ”. There may be a typographical on this sentence. Thus, the introduction of “Cholesterol-lowering agents, particularly 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase inhibitors (also known as statins), have been reported to be capable of increasing the ratio of alpha to beta secretase activity and then increasing the concentrations of extracellular A β ” should be revised to read “Cholesterol-lowering agents, particularly 3-hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) reductase inhibitors (also known as statins), have been reported to be capable of increasing the ratio of alpha to beta secretase activity and then decreasing the concentrations of extracellular A β ”. We are grateful for the interest shown in this study.





Figure 3 Atorvastatin attenuated interleukin-1 β expression in the hippocampus of a rat treated with A β 1-42. The upper panel shows interleukin-1 β -positive cells in rat hippocampus, detected by immunohistochemistry on day 7 after A β injection (original magnification \times 400). (A) Control group (group 1), (B) atorvastatin control group (group 2), (C) AD group (group 3), and (D) atorvastatin-treated AD group (group 4). (E) The lower panel shows the number of interleukin-1 β -positive cells counted in the rat hippocampus.

Notes: The data are presented as the mean \pm standard deviation. * $P < 0.01$.

Abbreviations: A β , amyloid-beta peptide; IL, interleukin; AD, Alzheimer's disease.





Figure 4 Atorvastatin attenuated interleukin-6 expression in the hippocampus of a rat treated with A β 1-42. The upper panel shows interleukin-6-positive cells in rat hippocampus, detected by immunohistochemistry on day 7 after A β injection (original magnification \times 400). **(A)** Control group (group 1), **(B)** atorvastatin control group (group 2), **(C)** AD group (group 3), and **(D)** atorvastatin-treated AD group (group 4). **(E)** The lower panel shows the number of interleukin-6-positive cells counted in the rat hippocampus.

Notes: The data are presented as the mean \pm standard deviation. * $P < 0.01$.

Abbreviations: A β , amyloid-beta (peptide); IL, interleukin; AD, Alzheimer's disease.




Figure 5 Atorvastatin attenuated TNF- α expression in hippocampus of the A β 1-42-treated rat. The upper panel shows TNF- α -positive cells in rat hippocampus, detected by immunohistochemistry on day 7 after A β injection (original magnification $\times 400$). (A) Control group (group 1), (B) atorvastatin control group (group 2), (C) AD group (group 3), and (D) atorvastatin-treated AD group (group 4). (E) The lower panel shows the number of TNF- α -positive cells counted in the rat hippocampus.

Notes: The data are presented as the mean \pm standard deviation. *P < 0.01.

Abbreviations: A β , amyloid-beta (peptide); IL, interleukin; AD, Alzheimer's disease; TNF- α , tumor necrosis factor alpha.

Clinical Interventions in Aging

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, the American Chemical Society's 'Chemical Abstracts

Submit your manuscript here: <http://www.dovepress.com/clinical-interventions-in-aging-journal>

Dovepress

Service' (CAS), Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.