



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review

Disinfection methods against SARS-CoV-2: a systematic review

C.P. Viana Martins^{a,*}, C.S.F. Xavier^a, L. Coimbra^{b,c,d}

^a Faculty of Medicine, University of Porto, Porto, Portugal

^b Division of Microbiology, Department of Pathology, Faculty of Medicine, University of Porto, Porto, Portugal

^c CINTESIS, Centre for Health Technology and Science Research, Porto, Portugal

^d Burn Unit and Department of Plastic and Reconstructive Surgery, University Hospital Centre of São João, Porto, Portugal

ARTICLE INFO

Article history:

Received 14 April 2021

Accepted 26 July 2021

Available online 18 October 2021

Keywords:

SARS-CoV-2

COVID-19

Coronavirus

Disinfection methods

Systematic review



SUMMARY

Background: Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), the causative agent of coronavirus disease 2019, has caused millions of deaths worldwide. The virus is transmitted by inhalation of infectious particles suspended in the air, direct deposition on mucous membranes and indirect contact via contaminated surfaces. Disinfection methods that can halt such transmission are important in this pandemic and in future viral infections.

Aim: To highlight the efficacy of several disinfection methods against SARS-CoV-2 based on up-to-date evidence found in the literature.

Methods: Two databases were searched to identify studies that assessed disinfection methods used against SARS-CoV-2. In total, 1229 studies were identified and 60 of these were included in this review. Quality assessment was evaluated by the Office of Health Assessment and Translation's risk-of-bias tool.

Findings: Twenty-eight studies investigated disinfection methods on environmental surfaces, 16 studies investigated disinfection methods on biological surfaces, four studies investigated disinfection methods for airborne coronavirus, and 16 studies investigated methods used to recondition personal protective equipment (PPE).

Conclusions: Several household and hospital disinfection agents and ultraviolet-C (UV-C) irradiation were effective for inactivation of SARS-CoV-2 on environmental surfaces. Formulations containing povidone-iodine can provide virucidal action on the skin and mucous membranes. In the case of hand hygiene, typical soap bars and alcohols can inactivate SARS-CoV-2. Air filtration systems incorporated with materials that possess catalytic properties, UV-C devices and heating systems can reduce airborne viral particles effectively. The decontamination of PPE can be conducted safely by heat and ozone treatment.

© 2021 The Healthcare Infection Society. Published by Elsevier Ltd. All rights reserved.

Introduction

The coronavirus disease 2019 (COVID-19) pandemic has become an ongoing global health crisis responsible for causing millions of deaths and has devastated the world's economy

* Corresponding author. Address: Faculty of Medicine, University of Porto, Porto, Portugal.

E-mail address: up201507735@med.up.pt (C.P. Viana Martins).

[1,2]. Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), a novel betacoronavirus, is known to be transmitted through exposure to infectious particles in respiratory droplets of infected individuals [3]. This can take place by inhalation of viral particles suspended in the air, deposition of exhaled infectious droplets directly on mucous membranes, or indirect contact with contaminated secondary surfaces, such as hands or fomites [4]. It is believed that airborne transmission may be the dominant form of transmission that best explains the occurrence of superspreading events, the higher risk of transmission in indoor settings, and the fact that more than half of transmission events are observed in asymptomatic or pre-symptomatic patients [5–8]. Particles emitted from infected individuals can be deposited on environmental surfaces and can remain viable for hours to days; as such, it is possible that transmission occurs due to indirect contact with contaminated surfaces [9–11].

The process of decontaminating surfaces normally uses chemical agents such as alcohol or quaternary ammonium compounds (QACs). There is evidence that these agents are active against viruses including SARS-CoV-2 [12]. The World Health Organization (WHO) recommends alcohol-based formulations to disinfect hands; such formulations have been shown to inactivate SARS-CoV-2 efficiently [13]. Many other accessible formulations with a broad range of application, such as hydrogen peroxide or povidone-iodine (PVP-I), possess antiviral properties, potentially serving as effective alternatives for the disinfection of biological surfaces [14,15].

As recent findings suggest that the airborne route is the most plausible and dominant form of transmission, this matter should be highlighted, and methods that can inactivate viruses suspended in the air may contribute substantially to lower the number of cases. Besides natural and mechanical ventilation, only two methods are available commercially: air cleaners fitted with filters or ultraviolet light; and upper room fixtures of ultraviolet germicidal irradiation (UVGI) [16]. UVGI uses short-wavelength ultraviolet C (UV-C) light which, in turn, has been tested against SARS-CoV-2 and proven to be effective [17].

Personal protective equipment (PPE) is essential to protect healthcare workers (HCWs) from contracting infections. Frontline HCWs are at higher risk of contracting SARS-CoV-2 infection compared with the general public [18]. While it is recommended that PPE should be disposable, in times of crisis, a shortage of PPE can cause more harm than benefit, as observed in many countries during the COVID-19 pandemic [19]. In cases when PPE is scarce, methods that provide proper sterilization, while preserving functionality, can be highly beneficial.

The first vaccines, distributed by the end of 2020, have reduced the number of hospitalizations, deaths and incidence of infection, proving to be the most effective tool to combat the COVID-19 pandemic [20–22]. However, disinfection methods will continue to play a major role and must still be put into practice to control local transmissions, whether from human to human, fomites or airborne. Halting the chain of transmission through the implementation of disinfection methods is not only useful in this present pandemic but also in any future similar pandemic. Therefore, the goal of this systematic review is to highlight the best disinfection methods to eliminate SARS-CoV-2 from environmental surfaces, biological surfaces and the air, and to determine the best methods to recondition PPE adequately.

Methods

Eligibility criteria

This review included original articles and experimental studies. Guidelines, protocols, recommendations and non-experimental studies, such as case reports, case series, cross-sectional, prospective case–control studies, opinions and review articles, were excluded. No limitations were considered regarding language, date or status of publication.

Participants

Studies that mentioned SARS-CoV-2 as the main target of any type of disinfection method tested were included. If the study did not mention SARS-CoV-2 specifically, inclusion of the family of coronaviruses that shares genetic or morphological similarities with SARS-CoV-2, preferentially the betacoronaviruses responsible for previous outbreaks of respiratory diseases such as severe acute respiratory syndrome (SARS-CoV) and/or Middle Eastern respiratory syndrome (MERS-CoV), was mandatory. During an outbreak of a highly contagious viral disease such as COVID-19, the availability of the virus under investigation can be limited. Therefore, surrogate viruses (i.e. enveloped virus references) used to study the efficacy of disinfection methods were also included. Studies that did not meet the participant criteria were excluded.

Interventions

Trials that compared the virucidal effects of disinfection methods with the potential to halt transmission of SARS-CoV-2 on environmental surfaces, biological surfaces, air and PPE were assessed.

Information sources, search and study selection

A search was conducted by two reviewers in two separate databases from January to June 2021. PubMed and Web of Science were searched using the following terms: ('SARS-CoV-2' OR 'Coronavirus' OR 'COVID-19') AND ('Disinfection Methods' OR 'Surface Disinfection' OR 'Hand Disinfection' OR 'Air disinfection' OR 'Environmental disinfection' OR 'Inactivation'). Thirteen articles were identified from other sources and included in the screening process. Two reviewers screened (by title and abstract) the initial 1229 articles found, and the information collected was registered on a shared EndNote Vx9 (Clarivate Analytics, Philadelphia, PA, USA) library and a shared online Microsoft Excel V16.42/2020 (Microsoft Corp., Redmond, WA, USA) document. Eligibility assessment was performed independently in an unblinded standardized manner by two reviewers and disagreements between reviewers were solved by consensus. In total, 60 articles were found to meet the inclusion criteria and were included in this systematic review.

Risk of bias

To determine the risk of bias in the individual studies selected, the Office of Health Assessment and Translation Risk-of-Bias Rating Tool for Human and Animal Studies was used.

This tool includes a questionnaire aimed to study risk of bias in several domains:

- selection bias;
- performance bias;
- attrition/exclusion bias;
- detection bias;
- selective reporting bias; and
- other bias.

Potential source of bias was graded as low risk (++), probable low risk (+), probable high risk or not reported (-), and high risk (-).

Results

Study selection

In total, 1229 articles were identified through a search of two databases, PubMed and Web of Science, from January 2021; 13 of these articles were included from other sources. After eliminating duplicate articles, 1021 articles remained. All articles were screened based on the title and abstract, leaving 83 articles eligible for this review. A further 36 articles were eliminated as they did not meet the inclusion criteria. In total, 60 articles were included in this systematic review. The details of this process are represented in Figure 1. The main characteristics of each individual study included in the systematic review are summarized in Table I.

Study characteristics

The studies included were from 15 countries. Thirty-eight studies used SARS-CoV-2 in their experiments, and the other studies depended on surrogate viruses to represent virucidal

activities of some disinfection methods. Six of the articles included were pre-prints.

Stability and survival of SARS-CoV-2 exposed to heat and high humidity

SARS-CoV-2 can remain viable on glass, stainless steel and plastic for more than 3.5 h at ambient temperature and humidity [23]. Increasing relative humidity alone at a constant temperature of 25°C can reduce the survival of SARS-CoV-2 on non-porous surfaces from approximately 15 h–8 h. When temperature and relative humidity are increased simultaneously, the half-life can be reduced remarkably to approximately 1 h [24]. The findings of another study conducted on other coronaviruses [mouse hepatitis virus (MHV) and transmissible gastroenteritis virus (TGEV)] revealed similar results. However, at low temperatures of 4°C and relative humidity of 20%, viruses can persist for up to 28 days [25]. SARS-CoV-2 can be deactivated at different rates when exposed to distinct heating procedures; one study showed that conditions that block evaporation can speed up virus inactivation rates substantially [26].

Disinfection methods on environmental surfaces

Amongst all the reviewed and included studies, 28 articles were categorized as disinfection methods with potential activity on environmental surfaces [12,23–49]. A summary of this category is provided in Table II.

Several studies demonstrated the virucidal properties of commonly used alcohols, leading to the inactivation of viruses on environmental surfaces. One study illustrated inactivation of human coronavirus (HCoV) on porcelain and ceramic surfaces with different concentrations of ethanol and isopropanol. Ethanol with concentrations ranging from 62% to 80% can cause a 4 log₁₀ reduction of viral titres after exposure ≥ 15 s.




Figure 1. PRISMA flow diagram of included articles. SARS-CoV-2, severe acute respiratory syndrome coronavirus-2.

Table I
Characteristics of included studies (N=60)

Study	Country	Year	Study design	Environmental surfaces	Personal protective equipment (masks/respirators etc.)	Biological surfaces (hands, skin, oral cavity, respiratory tract)	Air
Anderson <i>et al.</i>	UK	2020	<i>In vitro</i>	X			
Bedell <i>et al.</i>	USA	2016	<i>In vitro</i>	X			
Behzadinasab <i>et al.</i>	Hong Kong	2020	<i>In vitro</i>	X			
Biryukov <i>et al.</i>	USA	2020	<i>In vitro</i>	X	x		
Casanova <i>et al.</i>	USA	2010	<i>In vitro</i>	X			
Colnago <i>et al.</i>	Brasil	2020	<i>In vitro</i>	X		x	
Criscuolo <i>et al.</i>	Italy	2020	<i>In vitro</i>	X			
Gamble <i>et al.</i>	USA	2020	<i>In vitro</i>	X			
Gerchman	Israel	2020	<i>In vitro</i>	X			
He <i>et al.</i>	China	2004	<i>In vitro</i>				x
Heilingloh <i>et al.</i>	Germany	2020	<i>In vitro</i>	X			
Hulkower <i>et al.</i>	USA	2011	<i>In vitro</i>	X		x	
Khaiboullina <i>et al.</i>	USA	2020	<i>In vitro</i>	X			
Liu <i>et al.</i>	China	2020	<i>In vitro</i>	X			
Malenovská	Czech Republic	2020	<i>In vitro</i>	X			
Martins <i>et al.</i>	Brasil	2020	<i>In vitro</i>	X			
Meyers <i>et al.</i>	USA	2021	<i>In vitro</i>	X			
Monge <i>et al.</i>	USA	2020	<i>In vitro</i>	X			
Rabenau <i>et al.</i>	Germany	2005	<i>In vitro</i>	X		x	
Ratnesar-Shumate <i>et al.</i>	USA	2020	<i>In vitro</i>	X			
Wood and Payne	UK	1998	<i>In vitro</i>	X			
Blanchard <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Campos <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Buonanno <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Daeschler <i>et al.</i>	Canada	2020	<i>In vitro</i>		X		
Gopal <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Ibanez-Cervantes <i>et al.</i>	Mexico	2020	<i>In vitro</i>		X		
Ludwig-Begall <i>et al.</i>	Belgium	2020	<i>In vitro</i>		X		
Ma <i>et al.</i>	China	2020	<i>In vitro</i>		X		
Mantlo <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Ozog <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Perkins <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Rathnasinghe <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Rockey <i>et al.</i>	USA	2020	<i>In vitro</i>		X		
Bidra <i>et al.</i>	USA	2020	<i>In vitro</i>			X	
Bidra <i>et al.</i>	USA	2020	<i>In vitro</i>			X	
Eggers <i>et al.</i>	Germany	2015	<i>In vitro</i>			X	
Frank <i>et al.</i>	USA	2020	<i>In vitro</i>			X	
Gudmundsdottir <i>et al.</i>	Iceland	2020	<i>In vitro</i>			X	
Kratzel <i>et al.</i>	Germany	2020	<i>In vitro</i>			X	
Leslie <i>et al.</i>	USA	2020	<i>In vitro</i>			X	
Liang <i>et al.</i>	China	2020	<i>In vivo</i> and <i>in vitro</i>			X	
Meister <i>et al.</i>	Germany	2020	<i>In vitro</i>			X	
Mukherjee <i>et al.</i>	India	2020	<i>In vitro</i>			X	
Buonanno <i>et al.</i>	USA	2020	<i>In vitro</i>				x
Qiao <i>et al.</i>	USA	2020	<i>In vitro</i>				x
Yu <i>et al.</i>	USA	2020	<i>In vitro</i>				x
Franke <i>et al.</i>	Germany	2021	<i>In vitro</i>	x			
Gidari <i>et al.</i>	Italy	2021	<i>In vitro</i>	x			
Glasbrenner <i>et al.</i>	USA	2021	<i>In vitro</i>		x		
Hirose <i>et al.</i>	Japan	2020	<i>In vitro</i>			x	
Hu <i>et al.</i>	China	2021	<i>In vitro</i>	x			
Huang <i>et al.</i>	USA	2020	Prospective cohort			x	

(continued on next page)

Table 1 (continued)

Study	Country	Year	Study design	Environmental surfaces	Personal protective equipment (masks/respirators etc.)	Biological surfaces (hands, skin, oral cavity, respiratory tract)	Air
Ijaz <i>et al.</i>	USA	2021	<i>In vitro</i>	x		x	
Messina <i>et al.</i>	Italy	2021	<i>In vitro</i>	x			
Steinhauer <i>et al.</i>	Germany	2020	<i>In vitro</i>	x		x	
Steinhauer <i>et al.</i>	Germany	2020	<i>In vitro</i>			x	
Trivellini <i>et al.</i>	Italy	2021	<i>In vitro</i>	x			
Uppal <i>et al.</i>	USA	2021	<i>In vitro</i>		x		
Valdez-Salas <i>et al.</i>	Mexico	2021	<i>In vitro</i>		x		

Isopropanol 60–70% exposed on surfaces for at least 15 s demonstrated similar results with a 4 log₁₀ reduction of viral titre [40]. Hulkower *et al.* demonstrated the virucidal effects of three products containing different concentrations of alcohol on stainless steel surfaces. Ethanol 62%, 70% and 71% showed approximately 1.98–3.92 log₁₀ reduction of MHV and 3.19–4.04 log₁₀ reduction of TGEV after 1 min of exposure [29]. Hygiene wipes containing water and ethanol (0.6 g/wipe) destined to decontaminate plastic food packaging can reduce alphacoronavirus 1 to undetectable levels after 72 h of refrigeration (4°C) compared with wipes containing 99% water. However, this study showed evidence that hygiene wipes can potentially transfer viral particles to secondary surfaces [38]. Only two studies showed virucidal efficacy with at least 20% ethanol against SARS-CoV-2 in suspension [12,47].

In the case of sodium hypochlorite, one study showed that 0.525% and 0.1% sodium hypochlorite was sufficient to produce a 4 log₁₀ reduction of HCoV after 15 s of exposure on porcelain and ceramic surfaces [40]. Sodium hypochlorite 0.06% caused <1 log₁₀ reduction of MHV and TGEV after 1 min of exposure on stainless steel. This indicates that either a higher concentration of sodium hypochlorite is needed to cause a more significant reduction in viral titre after 1 min of exposure or a longer exposure time should be considered if 0.06% sodium hypochlorite is used [29]. When it comes to SARS-CoV-2, 0.14% sodium hypochlorite has been shown to reduce the viral titre significantly after 30 s of exposure [12].

With reference to aldehydes, one study showed that glutaraldehyde can lead to a >4 log₁₀ reduction in HCoV with contact times as low as 15 s on porcelain and ceramic surfaces [40]. Glutaraldehyde 4% and ethylenedioxy dimethanol at different concentrations were also capable of causing more than 3.5 log₁₀ reduction in SARS-CoV titre after 15 min in a suspension test [28]. Ortho-phthalaldehyde (OPA) 0.55% caused <2.5 log₁₀ reduction of MHV and TGEV after 1 min of exposure, indicating that OPA may need a longer exposure time to reach its total inactivation capacity [29].

QACs are common disinfection agents with a wide range of microbicidal action. Disinfectant wipes containing 0.75% didecyl-dimethyl-ammonium chloride associated with 0.5% hydrogen peroxide can reduce the alphacoronavirus 1 titre by 3.8 log₁₀ on plastic carriers, and can prevent transmission to secondary surfaces [38]. Combined surface disinfection solutions containing 0.5% benzalkonium chloride with laurylamine can reduce the SARS-CoV titre by 6.13 log₁₀ after 30 min of exposure, while 0.5% benzalkonium chloride associated with glutaraldehyde and didecylidmonium chloride showed a 3.75 log₁₀ reduction in the SARS-CoV titre in 30 min [28]. However, a

study conducted in 1997 on suspended HCoV revealed that 1% of benzalkonium chloride and a combination of 5% cetrimide and chlorhexidine gluconate were both ineffective in reducing viral titre after 1 min of exposure [27]. Moreover, QACs were shown to be active against SARS-CoV-2, vaccinia virus Elstree and modified vaccinia virus Ankara with contact times ≤5 min [12,47].

Phenols are another group of disinfectants active against a variety of micro-organisms. Cleaners that consist of 9.09% O-phenylphenol and 7.66% P-tertiary amyphenol showed a moderate reduction in infectivity for MHV and TGEV, revealing approximately 0.8–3.17 log₁₀ reduction on stainless steel surfaces [29]. Chloroxylenol 5% was ineffective for reducing the HCoV titre, but a study conducted in 2020 demonstrated that lower concentrations can efficiently inactivate a number of coronaviruses, including SARS-CoV-2, deposited on glass and in suspension after 1 min of exposure [12,27].

Ozonated water could be an alternative for environmental disinfection as it can cause a 2.0–5.0 log₁₀ reduction in SARS-CoV-2 titre after only 1 min of exposure [39,45].

Other chemical agents, such as magnesium monoperoxyphthalate, lead to a ≥4.5 log₁₀ reduction in SARS-CoV titre after 15 min of exposure [28]. Surface disinfectants based on citric acid, hydrochloric acid or lactic acid were shown to reduce viral titres of coronaviruses (including SARS-CoV-2) efficiently [12]. Virusend (TX-10), a detergent-based disinfectant, was able to reduce infectious SARS-CoV-2 with high titre inoculum by at least 4.0 log₁₀ plaque-forming units (PFU)/mL, and reduce infectious SARS-CoV-2 with low titre inoculum by at least 2.3 log₁₀ PFU/mL on hard surfaces, such as stainless steel, and in solution [31].

UV-C irradiation and ozone exposure

On glass surfaces, UV-C radiation can reduce MHV titre by an average of 2.71 log₁₀ and 6.11 log₁₀ with exposure times of 5 and 10 min, respectively. It is also able to reduce MERS-CoV titre by 5.9 log₁₀ after 5 min of exposure [30]. Findings in two studies indicated that at least 3 min of exposure to UV-C irradiation is able to inactivate SARS-CoV-2 in suspension completely [35,46]. Spherical objects such as footballs, volleyballs and basketballs were completely decontaminated from SARS-CoV-2 after 1 min of exposure to a UV-C-LED device (275 nm) [48]. UV-A, characterized by a longer wavelength (315–400 nm) is less efficient in viral inactivation, revealing only 1 log₁₀ reduction after 9 min of exposure to radiation [35]. It is suggested that peak emission of approximately 286 nm can be effective in inactivating coronaviruses [34]. An in-vitro study provided evidence that UV-B (280–315 nm) levels

Table II
Results of disinfection methods used on environmental surfaces

	Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log10) or (%)	Reduction of viral half-life (t1/2)
1	Anderson <i>et al.</i> (2020)	SARS-CoV-2	ViruSend (TX-10) detergent	Suspension test and stainless steel disc surface	1 min 10 min	4.0 log10 4.0 log10	(-)
2	Bedell <i>et al.</i> (2016)	MHV-A59 MERS-CoV	Multiple-emitter, automated, continuous, whole-room UV-C disinfection system	Glass coverslip surface	5 min 10 min 5 min 10 min	2.71 log10 6.11 log10 5.91 log10 (-)	(-)
3	Behzadinasab <i>et al.</i> (2020)	SARS-CoV-2	Cu2O/PU coating film	Glass surface Stainless steel surface	1 h	3.64 log10 2.97 log10	(-)
4	Biryukov <i>et al.</i> (2020)	SARS-CoV-2	24°C + 20% relative humidity 24°C + 40% relative humidity 24°C + 60% relative humidity 24°C + 80% relative humidity 35°C + 20% relative humidity 35°C + 40% relative humidity 35°C + 60% relative humidity	Stainless steel, ABS plastic and nitrile rubber surfaces	(-)	(-)	15.33 h ± 2.75 11.52 h ± 1.72 9.15 h ± 3.39 8.33 h ± 1.80 7.33 h ± 1.33 7.52 h ± 1.22 2.26 h ± 1.42
5	Casanova <i>et al.</i> (2010)	TGEV and MHV	4°C + 20% relative humidity 4°C + 50% relative humidity 4°C + 80% relative humidity 20°C + 20% relative humidity 20°C + 50% relative humidity 20°C + 80% relative humidity 40°C + 20% relative humidity 40°C + 50% relative humidity 40°C + 80% relative humidity	Stainless steel surface	28 days 21 days 28 days 28 days 3 days (TGEV) and 5 days (MHV) 14 days (TGEV) 11 days (MHV) 5 days (-) 3 h	0.5 log10 3.5 log10 3.2 log10 (TGEV) and 2.5 log10 (MHV) 2 log10 2 log10 (TGEV) and 3 log10 (MHV) 3 log10 (TGEV) and 5 log10 (MHV) 3.5 log10 (TGEV) and 4.7 log10 (MHV) (-) 2.8 log10 (TGEV) and 4.1 log10 (MHV) >4 log10	(-)
6	Colnago <i>et al.</i> (2020)	ACoV	Household dishwashing detergent (2% sodium dodecyl sulfate and 6% linear alkylbenzene sulfonates)	Suspension test	10 min	>4 log10	(-)
7	Criscuolo <i>et al.</i> (2021)	SARS-CoV-2	UV-C	Glass Plastic Gauze Wood Fleece Wool	15 min	>99.9% >99.9% >99.9% 0.0% 90.0% 94.4%	(-)

Table II (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Reduction of viral half-life (t _{1/2})	
		Ozone (0.2 ppm)	Glass Plastic Gauze Wood Fleece	2 h	90.0% 82.2% 96.8% 93.3% >99.9%		
		Ozone (4 ppm)	Glass Plastic Gauze Wood Fleece	2 h	94.4% 90.0% 99.8% (-) 99.7%		
8	Gamble <i>et al.</i> (2020)	SARS-CoV-2	Uncovered plate oven (70°C) Covered plate oven (70°C) Closed vial oven (70°C) Closed vial heat block (70°C)	Suspension test (-)	(-)	~ 37 min ~ 3 min and 56 s ~ 51.6 s ~ 1 min and 55 s	
9	Gerchman <i>et al.</i> (2020)	HCoV-OC43	UV-LED (267 nm wavelength, 6 –7 mJ/cm ²) UV-LED (279 nm wavelength, 6 –7 mJ/cm ²) UV-LED (286 nm wavelength, 13 mJ/cm ²) UV-LED (297 nm wavelength, 32 mJ/cm ²)	Suspension test 60 s 90 s	>3 log ₁₀ >3 log ₁₀ >3 log ₁₀ >3 log ₁₀	(-)	
10	Heilingloh <i>et al.</i> (2020)	SARS-CoV-2	UV-C 1.94 mJ/cm ² /s UV-A 0.54 mJ/cm ² /s combined (UV-C and UV-A)	Suspension test	9 min	Total inactivation 1 log reduction Total inactivation	(-)
11	Hulkower <i>et al.</i> (2011)	TGEV	9.09% O-phenylphenol, 7.66% P-tertiary amylphenol 6% sodium hypochlorite 0.55% ortho-phthalaldehyde 70% ethanol 62% ethanol 71% ethanol	Stainless steel surface	1 min	2.03 log ₁₀ 0.35 log ₁₀ 2.27 log ₁₀ 3.19 log ₁₀ 4.04 log ₁₀ 3.51 log ₁₀ 1.33 log ₁₀	(-)
		MHV	9.09% O-phenylphenol, 7.66% P-tertiary amylphenol 6% sodium hypochlorite 0.55% ortho-phthalaldehyde 70% ethanol 62% ethanol 71% ethanol			0.62 log ₁₀ 1.71 log ₁₀ 3.92 log ₁₀ 2.66 log ₁₀ 1.98 log ₁₀	

12	Khaiboullina <i>et al.</i> (2020)	HCoV	TNP coating + UV-C (254 nm wavelength)	Glass coverslip surface	TNP (20 min to dry or left wet) o and UV-C (30 s and 1 minute)	Reduction in viral copies on both wet and dry surfaces potentiated by the addition of TNP	(-)
13	Liu <i>et al.</i> (2020)	SARS-CoV-2	Ultra-high power UV-C	Suspension test	1 s	100%	(-)
14	Malenovská (2020)	SARS-CoV-2	99% water, caprylyl/capryl glucoside, citric acid, sodium citrate, sodium benzoate	Plastic (4°C)	24 h 48 h 72 h 96 h 120 h	~1.9 log ₁₀ ~2.6 log ₁₀ ~2.2 log ₁₀ >1 log ₁₀ >0.3 log ₁₀	(-)
			Water, ethanol (0.6 g/wipe), glycerine, Aloe barbadensis leaf extract, chlorhexidine digluconate		24 h 48 h 72 h 96 h 120 h	2.4 log ₁₀ 2.2 log ₁₀ >1.8 log ₁₀ >1 log ₁₀ >0.3 log ₁₀	
			0.75% didecyl-dimethyl-ammonium chloride, 0.5% hydrogen peroxide, less than 5% non-ionic surface active agent, cationic surface active agent, bleaching agent based on oxygen, perfume, limonene, iodopropynyl butylcarbamate		24 h 48 h 72 h 96 h 120 h	>3.3 log ₁₀ >3.1 log ₁₀ >2.3 log ₁₀ Not performed Not performed	
15	Martins <i>et al.</i> (2020)	SARS-CoV-2	Ozonated water [0.2–0.8 ppm (mg/L)]	Suspension test	1 min	2 log ₁₀	(-)
16	Meyers <i>et al.</i> (2021)	HCoV	62% ethanol	Porcelain surface	15 s, 30 s, 1 min	>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀	(-)
			70% ethanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀	
			75% ethanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀	
			80% ethanol			>4 log ₁₀ , ≥4 log ₁₀ , >4 log ₁₀	
			95% ethanol			>2 log ₁₀ , 2–3 log ₁₀ , 1–2 log ₁₀	
			70% isopropanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀	
			75% isopropanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀	
			80% isopropanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀	
			95% isopropanol			>4 log ₁₀ , 3–4 log ₁₀ , 3–4 log ₁₀	
			0.0525% sodium hypochlorite			1–2 log ₁₀ , 2–3 log ₁₀ , 2–3 log ₁₀	

(continued on next page)

Table II (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Reduction of viral half-life (t _{1/2})	
		0.525% sodium hypochlorite			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		0.1% sodium hypochlorite			Not performed		
		Glutaraldehyde			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		62% ethanol	Ceramic surface		>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		70% ethanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		75% ethanol			3–4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		80% ethanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		95% ethanol			1–2 log ₁₀ , 1–2 log ₁₀ , 1–2 log ₁₀		
		70% isopropanol			>4 log ₁₀ , 3–4 log ₁₀ , >4 log ₁₀		
		75% isopropanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		80% isopropanol			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		95% isopropanol			3–4 log ₁₀ , 1–2 log ₁₀ , 1–2 log ₁₀		
		0.0525% sodium hypochlorite			1–3 log ₁₀ , 1–2 log ₁₀ , 1–2 log ₁₀		
		0.525% sodium hypochlorite			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		0.1% sodium hypochlorite			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
		Glutaraldehyde			>4 log ₁₀ , >4 log ₁₀ , >4 log ₁₀		
17	Monge <i>et al.</i> (2020)	SARS-CoV-2	Cationic phenylene ethynylene polymers (conjugated electrolytes)	Suspension test	10 min	1–5 log	(-)
			Cationic phenylene ethynylene oligomers (conjugated electrolytes)		20 min	1.5 log	
					60 min	5 log	
18	Rabenau <i>et al.</i> (2005)	SARS-CoV	Mikrobac forte (0.5% benzalkonium chloride and laurylamine)	Suspension test	30 min	≥6.13 log ₁₀	(-)

			Korsolin FF (0.5% benzalkonium chloride, glutaraldehyde and didecyldimonium chloride)			$\geq 3.75 \log_{10}$	
			Dismozon pur (magnesium monoperphthalate)			$\geq 4.5 \log_{10}$	
			Korsolex basic [4% glutaraldehyde and (ethylenedioxy) dimethanol]	15 min		$\geq 3.5 \log_{10}$	
			Korsolex basic [3% glutaraldehyde and (ethylenedioxy) dimethanol]	30 min		$\geq 3.5 \log_{10}$	
			Korsolex basic [2% glutaraldehyde and (ethylenedioxy) dimethanol]	60 min		$\geq 3.5 \log_{10}$	
19	Ratnesar-Shumate et al. (2020)	SARS-CoV-2	37°C + 5% CO ₂	Stainless steel coupons	20 min	1.6 W/m ² UV-B -> ~2.5 log ₁₀ 0.7 W/m ² UV-B -> ~2.2 log ₁₀ 0.3 W/m ² UV-B -> ~2.5 log ₁₀ Darkness -> 0.5 log ₁₀	(-)
20	Wood and Payne (1998)	HCoV	Dettol (5% chloroxylenol) Dettol for hospitals (1% benzalkonium chloride) Savlon (5% cetrimide and chlorhexidine gluconate)	Suspension test	1 min	0.0 log ₁₀ 0.0 log ₁₀ 0.0 log ₁₀	(-)
21	Franke et al. (2021)	Bacteriophage F6 (phi 6)	Ozone (80 ppm) + 90% relative humidity	Melamine-coated solid core panels Ceramic tiles Stainless steel carriers	60 min	4.29 log ₁₀ 6.15 log ₁₀ 5.31 log ₁₀	(-)
		Bovine coronavirus		Melamine-coated solid core panels Ceramic tiles Stainless steel carriers		5.03 log ₁₀ 4.88 log ₁₀ 5.31 log ₁₀	
22	Gidari et al. (2021)	SARS-CoV-2	23–25°C + 40–50 relative humidity	Plastic Stainless steel carriers Glass	(-)	(-)	3.5 h 4.4 h 4.2 h
			UV-C (254 nm)	Plastic Stainless steel carriers Glass	20.06 mJ/cm ² (36 s) 20.06 mJ/cm ² (36 s) 10.25 mJ/cm ² (21 s)	>4.00 log ₁₀ >4.00 log ₁₀ >4.00 log ₁₀	(-)

(continued on next page)

Table II (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Reduction of viral half-life (t _{1/2})	
23	Hu <i>et al.</i> (2021)	SARS-CoV-2	Ozonated water (36 mg/L)	Suspension test	0 min 1 min 5 min 10 min	0.0 log ₁₀ ~5 log ₁₀ ~5 log ₁₀ ~5 log ₁₀	(-)
24	Ijaz <i>et al.</i> (2020)	MHV-1 HCoV-229E SARS-CoV MERS-CoV SARS-CoV-2 HCoV-229E (1), SARS-CoV (2), SARS-CoV-2 (3)	Ozonated water (18 mg/L) 0.12% p-chloro-m-xenolol (PCMX) PCMX (0.125% w/v) Alkyl dimethyl benzyl ammonium chloride QAC (0.19% w/w) Citric acid (2.4% w/w) Ethanol (50% w/w)/QAC (0.082% w/w) Alkyl dimethyl benzyl ammonium chloride (0.0916%) QAC (0.092% w/w) QAC (0.077% w/w) Lactic acid (1.9% w/w) Hydrochloric acid (0.25% w/w) Sodium hypochlorite (0.14% w/w) Benzalkonium chloride (0.45% w/w) Ethanol (44% w/w) Sodium hypochlorite (0.32% w/w)	Glass Suspension test Glass + organic load	0.5 min 10 min 5 min 5 min 1 min 5–10 min 1.75 min 0.5 min 0.5–1.75 min 5 min 2 min 5 min 5 min 0.5 min 0.5 min 5 min 5 min 5 min	~5 log ₁₀ ~5 log ₁₀ ~5 log ₁₀ ~5 log ₁₀ ≥4.2 log ₁₀ ≥4.0 log ₁₀ ≥6.0 log ₁₀ ≥5.0 log ₁₀ ≥5.0 log ₁₀ (1) ≥4.0 log ₁₀ , (2) ≥6.0 log ₁₀ , (3) not performed (1) ≥6.0 log ₁₀ , (2) ≥5.8 log ₁₀ , (3) ≥3.5 log ₁₀ (1) ≥4.3 log ₁₀ , (2) ≥3.0 log ₁₀ , (3) ≥3.0 log ₁₀ (1) ≥5.5 log ₁₀ , (2) not performed, (3) ≥4.5 log ₁₀ (1) ≥3.5 log ₁₀ , (2) ≥4.8 log ₁₀ , (3) not performed (1) ≥3.3 log ₁₀ , (2) ≥3.8 log ₁₀ , (3) ≥4.0 log ₁₀ (1) Not performed (2) ≥4.1 log ₁₀ (1) Not performed (2) ≥5.5 log ₁₀ (1) Not performed (2) ≥4.1 log ₁₀ (1) Not performed (2) ≥5.1 log ₁₀ (1) Not performed (2) ≥4.5 log ₁₀ (1) ≥4.0 log ₁₀ (2) ≥4.1 log ₁₀ (1) Not performed (2) ≥5.1 log ₁₀	(-)

25	Messina <i>et al.</i> (2021)	SARS-CoV-2	UV irradiation chips (265–350 nm) box with lid - reflected light	Suspension test	3 min	4.70 log ₁₀	(-)
			UV irradiation chips (265–350 nm) box with lid		3 min	3.45 log ₁₀	
			UV irradiation chips (265–350 nm) box with lid		6 min	5.53 log ₁₀	
			UV irradiation chips (265–350 nm) box with lid		6 min	5.53 log ₁₀	
			UV irradiation chips (265–350 nm) box with lid		10 min	5.70 log ₁₀	
			UV irradiation chips (265–350 nm) box with lid		10 min	5.70 log ₁₀	
			UV irradiation chips (265–350 nm) box without lid - direct light		3 min	4.62 log ₁₀	
			UV irradiation chips (265–350 nm) box without lid		3 min	5.53 log ₁₀	
			UV irradiation chips (265–350 nm) box without lid		10 min	5.70 log ₁₀	
			UV irradiation chips (265–350 nm) box without lid		10 min	5.70 log ₁₀	
			26	Steinhauer <i>et al.</i> (2020)	Modified vaccinia virus Ankara	20% surface disinfectant - propan-2-ol, ethanol	
90% surface disinfectant - propan-2-ol, ethanol		15 s				≥4.25 log ₁₀	
Vaccinia virus Elstree	80% surface disinfectant - QAC				30 s	≥4.32 log ₁₀	
	80% surface disinfectant - QAC				60 s	≥4.51 log ₁₀	
SARS-CoV-2	20% surface disinfectant - propan-2-ol, ethanol				15 s	≥4.02 log ₁₀	
	80% surface disinfectant - propan-2-ol, ethanol				15 s	≥4.02 log ₁₀	
	20% surface disinfectant - QAC				15 s	≥4.02 log ₁₀	
	20% surface disinfectant - QAC				60 s	≥3.17 log ₁₀	
	80% surface disinfectant - QAC				15 s	≥4.38 log ₁₀	
	80% surface disinfectant - QAC				30 s	≥4.38 log ₁₀	
	80% surface disinfectant - QAC				60 s	≥2.17 log ₁₀	
27	Trivellin <i>et al.</i> (2020)	SARS-CoV-2	UV-C LED (275 nm) spherical irradiation box	Football	1 min	>3 log ₁₀	(-)
					2 min	>3 log ₁₀	
			Basketball	1 min	>3 log ₁₀		
				2 min	>3 log ₁₀		
			Volleyball	1 min	>3 log ₁₀		
				2 min	>3 log ₁₀		
28	Uppal <i>et al.</i> (2021)	HCoV-OC43	Ozone (20 ppm) Ozone (25 ppm)	Glass	10 min	90.71%	(-)
					10 min	92.3245%	

(continued on next page)

Table II (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log10) or (%)	Reduction of viral half-life (t1/2)
		Ozone (50 ppm)		15 min 20 min 10 min 15 min 20 min	99.99% 100.00% 99.987% 99.985% 100.00%	

SARS-CoV-2, severe acute respiratory syndrome coronavirus-2; MHV, mouse hepatitis virus; MERS-COV, Middle East respiratory syndrome coronavirus; TGEV, transmissible gastroenteritis coronavirus; HCoV, human coronavirus; ACOV, avian coronavirus; UV-C, ultraviolet C irradiation; UV-A, ultraviolet A irradiation; UV-LED, ultraviolet light emitting diode; Cu₂O/PU, cuprous oxide/polyurethane; TiO₂, titanium dioxide; TNP, TiO₂ nanoparticle; ABS plastic, acrylonitrile butadiene styrene plastic.

similar to natural sunlight can significantly reduce SARS-CoV-2 titre by 2.5 log₁₀ on stainless steel surfaces after 20 min of exposure [42].

Exposure of glass, plastic and gauze samples infected with SARS-CoV-2 to UV-C irradiation for 15 min led to a 99.99% reduction of viral titre, while a reduction of 90–95% was obtained for fleece and wool samples. No reduction in viral titre was quantified on wood samples with this method [43]. In the same study, 2 h of exposure to ozone 0.2 ppm was able to completely disinfect (99.99% reduction) fleece samples, and to achieve a 96.8% reduction on gauze, 93.3% on wood, 90% on glass and 82.2% on plastic. Exposure of the same materials to higher concentrations of ozone was effective in reducing viral titre in a shorter period. Uppal *et al.* demonstrated that ozone gas of at least 25 ppm can optimally eliminate ≥99% of HCoV deposited on glass in 15 min, while another study showed that ozone 80 ppm and 90% relative humidity obtained significant viral inactivation after 60 min [44,49].

Complete inactivation of HCoV is seen on TiO₂ nanoparticle (TNP)-coated glass coverslips exposed to UV-C for 30 s and 1 min. Viral inactivation was enhanced and accelerated with TNP coating, making viral titres undetectable after shorter time exposures to UV-C irradiation [36].

SARS-CoV-2 can be eliminated completely after only 1 s of exposure to a high-powered deep UV light. The UV light source is an aluminium gallium nitride-based device and can achieve an output power as high as 2 W at a current of 1.3 A allowing the ultra-rapid inactivation of SARS-CoV-2 [37].

Coatings and films

Coating surfaces with cuprous oxide/polyurethane or conjugated electrolytes such as cationic phenylene ethynylene polymers and oligomers was shown to have virucidal activity against SARS-CoV-2, and reduce viral titre significantly after 1 h of exposure on glass, stainless steel and in suspension [32,41]. Films made from an accessible household dishwashing detergent containing 8% surfactant can provide longer virucidal activity on inanimate surfaces, reducing avian coronavirus to undetectable levels after 10 min of exposure. The activity of these films can persist for up to 7 days [33].

Disinfection methods on biological surfaces

Sixteen articles addressed disinfection methods that can be used on biological surfaces (Table III) with application on skin, hands and mucous membranes, such as the oral cavity and upper respiratory tract [12,13,28,47,50–61].

Alcohols were mainly evaluated via suspension tests showing optimal virucidal activity (including SARS-CoV-2) at concentrations >65% and with exposure times of 15–60 s specifically for application to hands and the oral cavity [12,13,28,47,54,55,57,58,61]. One study evaluated the efficacy of ethanol and propanol directly on human skin against SARS-CoV-2 and found that 40% concentrations of these alcohols can cause >4 log₁₀ reduction in viral titre after just 5 s of exposure [59]. Interestingly, WHO-recommended hand rub formulations containing 80% ethanol or propanol showed inferior efficacy compared with modified formulations (with 75% ethanol or propanol and half of the concentration of glycerol from the original formulation) when tested with SARS-CoV-2 [13]. Soap bars evaluated in two studies were shown to reduce the quantity of SARS-CoV-2 significantly, with

optimal results seen with a contact time of 20 s to 1 min [12,58]. QACs, specifically benzalkonium chloride 0.2%, can produce maximum virucidal activity after 60 s of exposure, verified in suspension tests and on human skin [12,59]. Liquids containing chloroxylenol, citric acid, lactic acid or salicylic acid were also effective in reducing coronavirus titres, including SARS-CoV-2 [12].

Oral rinses containing PVP-I 1–3% lead to >4.33 log₁₀ reduction of SARS-CoV-2, MERS-CoV and modified vaccinia virus Ankara titres after 15–30 s of contact time [50–52,57]. The action of hydrogen peroxide oral rinses, on the other hand, is inferior to PVP-I [51]. Chlorhexidine gluconate (oral and skin formulations) seems to provide suboptimal virucidal activity compared with other agents in in-vitro suspension test experiments. However, a prospective cohort study on patients who were initially admitted to hospital with a positive SARS-CoV-2 test indicated that the application of chlorhexidine gluconate mouthwash and nasopharyngeal spray of the same agent can accelerate the clearance of SARS-CoV-2 in these areas, resulting in a negative reverse transcriptase polymerase chain reaction test after 4 days [60]. Other antiseptic oral rinses containing chloride and benzalkonium or ethanol have also been shown to deactivate SARS-CoV-2 [54,57].

PVP-I can also be applied topically on eyes as an additional pre-procedure disinfection as concentrations of 0.9% can reduce SARS-CoV-2 titre significantly after 30 s of exposure [56]. On the other hand, a toxicity study carried out in rabbits revealed that groups exposed to ocular PVP-I 0.6% and 1.0% every day for 7 days showed signs of mild and transient ocular irritation [56]. Nasal cavity formulations consisting of PVP-I 0.54–5% are able to cause >3 log₁₀ reduction in SARS-CoV-2 titre after 15 s of exposure [62].

Disinfection methods against airborne viruses

Regarding disinfection methods against airborne coronaviruses, four articles were identified (Table IV) [63–66]. Wafers containing silver and copper combined with aluminium oxide display catalytic properties and can be incorporated in air conditioning systems to trap and kill viruses. These wafers are active against coronaviruses and can cause complete viral inactivation after 5 min of exposure [63].

UV-C can efficiently inactivate up to 99.9% of aerosolized coronaviruses [64]. Ventilation systems fitted with a UV-C light source that can control its flow rate, control the exposure time of air passage indirectly, as lower flow rates translate into longer exposure times which results in superior viral removal efficacy [65]. As all human coronaviruses have similar genomic size, a key determinant of radiation sensitivity, it is likely that UV-C irradiation will show comparable inactivation efficiency against other human coronaviruses, including SARS-CoV-2 [64].

Methods to decontaminate and recondition personal protective equipment

Methods with potential use to decontaminate and recondition PPE were examined by 16 studies (Table V) [49,67–81]. Most of these studies investigated filtering facepiece respirators (FFRs), especially 3M N95 masks. Heat (70–95°C) combined with different levels of relative humidity is capable of inactivating enveloped viruses, including SARS-CoV-2,

inoculated on N95 level melt-blown polypropylene fabric after at least 20 min of exposure [68,74,79]. Filtration efficacy was maintained after several cycles. However, cycles should be limited to avoid compromising mask function. A limit of 20 disinfection cycles is suggested for treatments under high relative humidity (100%) and temperatures ≤85°C. Treatment should also be limited to five cycles under high relative humidity (100%) and temperatures ≤95°C [68–70,73]. Caution must be taken when a dry oven is utilized to generate dry heat (0% relative humidity) as samples placed on parchment paper prior to heating can result in lower efficacy of viral inactivation [77].

Other than heat treatment, face masks made with cloth fabric, disposable gowns and powered air purifying respirator hoods can all be decontaminated successfully with doses of at least 20 ppm of ozone [67]. N95 respirators inoculated with HCoV were also adequately decontaminated after 10–20 min of exposure to 20–50 ppm ozone gas [49].

Metals such as copper and zinc possess antiviral activity. Zinc ions incorporated in fabrics, such as cotton and polyamide 66 (PA66), can inactivate SARS-CoV-2 while maintaining virucidal activity after 50 washes, supporting the possibility of long-lasting virucidal protection [71]. It is worth noting that cotton and PA66 can trap viruses, as only 56% and 92% of SARS-CoV-2 can be recovered from cotton samples and PA66, respectively, after viral inoculation. This information is relevant as cotton- and PA66-based masks can trap large amounts of SARS-CoV-2, making cross-contamination more probable when masks are reused without decontamination [71]. Copper iodine complex has the potential to be used on non-critical PPE as it has been shown to completely deactivate SARS-CoV-2 in suspension after 30 min of exposure [75]. An innovative formulation that consists of silver and antimicrobial substances (ethanol and QACs) has also been shown to possess antiviral activity when impregnated in the matrix of surgical masks [81].

Hydrogen peroxide vapour can also inactivate SARS-CoV-2 deposited on N95 masks and FFRs. This last process can be conducted in a STERRAD 100NX sterilization system or a V-PRO Max Sterilizer providing exposure cycles of ≤47 min [72,73].

UV irradiation was able to inactivate coronaviruses deposited on surgical masks and FFRs [73,76,78]. Exposure times needed to decontaminate these materials completely ranged from 60 s to 4 min when the models tested were N95 FFRs. It is worth noting that the efficacy of UV-C irradiation is model-dependent, and straps that contain hydrophilic properties seem to cause a lower reduction in viral titre [76]. Exposure to simulated sunlight for 20 min, characterized by UV irradiation with wavelengths ranging between 300 and 400 nm, can reduce SARS-CoV-2 titre significantly on specific models of N95 masks [80].

Risk of bias

It was only possible to evaluate selection bias in two studies as the majority of experiments took place in in-vitro settings. Only one study blinded the personnel, so the other studies may contain performance bias. Thirteen studies were considered to have a probable risk of attrition or exclusion bias, eight studies had probable risk of detection bias, two studies had probable risk of selective reporting bias, and three studies had probable risk of potential threat to internal validity. A summary of the evaluation is provided in Table VI.

Table III
Results of disinfection methods used on biological surfaces

	Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Type
1	Bidra <i>et al.</i> (2020)	SARS-CoV-2	PVP-I 1.0% oral rinse PVP-I 2.5% oral rinse PVP-I 3.0% oral rinse H ₂ O ₂ 3.0% H ₂ O ₂ 6.0%	ST	15 s and 30 s	~4.33 log ₁₀ ~4.33 log ₁₀ ~4.33 log ₁₀ 1.33 log ₁₀ 1 log ₁₀	Oral
2	Bidra <i>et al.</i> (2020)	SARS-CoV-2	PVP-I (3.0%) oral rinse antiseptic PVP-I (1.5%) oral rinse antiseptic PVP-I (1.0%) oral rinse antiseptic	ST	30 s	3.33 log ₁₀ 3.33 log ₁₀ 3.33 log ₁₀	Oral
3	Eggers <i>et al.</i> (2015)	MERS-CoV	PVP-I surgical scrub (7.5 g/L available iodine) PVP-I skin cleanser (4 g/L available iodine) PVP-I gargle and mouthwash (1 g/L available iodine)	ST	15 s	4.64 log ₁₀ 4.97 log ₁₀ 4.30 log ₁₀	Skin and oral
		Modified vaccinia virus Ankara	PVP-I surgical scrub (7.5 g/L available iodine) PVP-I skin cleanser (4 g/L available iodine) PVP-I gargle and mouthwash (1 g/L available iodine)		15 s, 30 s and 60 s	≥4.17 log ₁₀ , ≥4.17 log ₁₀ ≥4.00 log ₁₀ , ≥4.00 log ₁₀ ≥4.00 log ₁₀ , ≥4.00 log ₁₀ ≥6.50 log ₁₀ , ≥6.50 log ₁₀	
4	Frank <i>et al.</i> (2020)	SARS-CoV-2	PVP-I nasal antiseptic 5.0% PVP-I nasal antiseptic 2.5% PVP-I nasal antiseptic 1.0%	Dilution test	15 s and 30 s	3 log ₁₀ (15 s), 3.33 log ₁₀ (30 s) 3 log ₁₀ (15 s), 3.33 log ₁₀ (30 s) 3 log ₁₀ (15 s), 3.33 log ₁₀ (30 s)	Respiratory tract
5	Gudmundsdottir <i>et al.</i> (2020)	SARS-CoV-2 and HCoV	Coldzyme (glycerol, water, cod trypsin, ethanol, calcium chloride, hydroxymethy, and menthol)	ST	20 min	1.76 log ₁₀ (SARS-CoV-2), 2.88 log ₁₀ (HCoV)	Oral
6	Kratzel <i>et al.</i> (2020)	SARS-CoV-2	Original WHO formulation I consists of 80% (vol/vol) ethanol, 1.45% (vol/vol) glycerol, and 0.125% (vol/vol) hydrogen peroxide	ST	30 s	>3.8 log ₁₀	Hands

			Original WHO formulation II consists of 75% (vol/vol) 2-propanol, 1.45% (vol/vol) glycerol, and 0.125% (vol/vol) hydrogen peroxide			>3.8 log ₁₀	
			Modified WHO formulation I consists of 80% (wt/wt) ethanol, 0.725% (vol/vol) glycerol, and 0.125% (vol/vol) hydrogen peroxide			>5.9 log ₁₀	
			Modified isopropyl-based WHO formulation II contains 75% (wt/wt) 2-propanol, 0.725% (vol/vol) glycerol, and 0.125% (vol/vol) hydrogen peroxide			>5.9 log ₁₀	
7	Leslie <i>et al.</i> (2020)	SARS-CoV-2	PURELL hand sanitizer gel, 70% ethanol (vol/vol)	ST	30 s	≥3.22 log ₁₀	Hands
			PURELL hand sanitizer foam, 70% ethanol (vol/vol)			≥3.10 log ₁₀	
8	Liang <i>et al.</i> (2020)	SARS-CoV-2	Povidone-iodine in-situ gel (polyvinylpyrrolidinone-iodine complex) (0.9%)	ST in tear fluid	30 s, 2 min and 10 min	3.5 log ₁₀ (30 s), 2.9 log ₁₀ (2 min), 3.3 log ₁₀ (10 min)	eyes
			Povidone-iodine in-situ gel (polyvinylpyrrolidinone-iodine complex) (0.5%)			3.2 log ₁₀ (30 s), 2.9 log ₁₀ (2 min), 3.3 log ₁₀ (10 min)	
			Povidone-iodine in-situ gel (polyvinylpyrrolidinone-iodine complex) (0.28%)			2.2 log ₁₀ (30 s), 2.6 log ₁₀ (2 min), 3.3 log ₁₀ (10 min)	
			Povidone-iodine in-situ gel (polyvinylpyrrolidinone-iodine complex) (0.09%)			1.2 log ₁₀ (30 s), 0.8 log ₁₀ (2 min), 1.0 log ₁₀ (10 min)	
			Povidone-iodine nasal spray (1-vinyl-2-pyrrolidinone polymers, iodine complex) (0.54%)	ST in nasal fluid		3.1 log ₁₀ (30 s), 2.9 log ₁₀ (2 min), 3.3 log ₁₀ (10 min)	Respiratory tract
			Povidone-iodine nasal spray (1-vinyl-2-pyrrolidinone polymers, iodine complex) (0.3%)			3.1 log ₁₀ (30 s), 2.9 log ₁₀ (2 min), 3.3 log ₁₀ (10 min)	
			Povidone-iodine nasal spray (1-vinyl-2-pyrrolidinone polymers, iodine complex) (0.17%)			2.9 log ₁₀ (30 s), 2.9 log ₁₀ (2 min), 3.3 log ₁₀ (10 min)	
			Povidone-iodine nasal spray (1-vinyl-2-pyrrolidinone polymers, iodine complex) (0.05%)			2.3 log ₁₀ (30 s), 1.9 log ₁₀ (2 min), 1.6 log ₁₀ (10 min)	

(continued on next page)

Table III (continued)

	Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Type
9	Meister <i>et al.</i> (2020)	SARS-CoV-2	Cavex oral pre rinse (hydrogen peroxide)	ST	30 s	0.33–0.78 log	Oral
			Chlorhexamed Forte [chlorhexidinebis (D-gluconate)]			0.78–1.17 log	
			Dequonal (dequalinium chloride, benzalkonium chloride)			≥2.61–3.11 log	
			Dynexidine Forte 0.2% [chlorhexidinebis (D-gluconate)]			0.50–0.56 log	
			Iso-betadine mouthwash 0% (polyvidone-iodine)			≥2.61–3.11 log	
			Listerine cool mint (ethanol, essential oils)			≥2.61–3.11 log	
			Octenident mouthwash (octenidine dihydrochloride)			0.61–1.11 log	
			ProntOral mouthwash (polyaminopropyl biguanide (polyhexanide))			0.61–≥1.78 log	
10	Mukherjee <i>et al.</i> (2020)	SARS-CoV-2	Soap bar with 67 total fatty matter	ST	20 s	≥3.14 log ₁₀	Hands
			Soap bar with 68 total fatty matter		20 s	≥3.06 log ₁₀	
			Soap bar with 72 total fatty matter		20 s	≥4.06 log ₁₀	
			Liquid cleansers with 10% surfactant w/w		20 s	≥3.10 log ₁₀	
			Liquid cleansers with 12% surfactant w/w		10 s	≥3.01 log ₁₀	
			Liquid cleansers with 19% surfactant w/w		10 s	≥3.42 log ₁₀	
			Alcohol-based sanitizers (60.5% alcohol w/w)		10 s	≥3.25 log ₁₀	
			Alcohol-based sanitizers (65% alcohol w/w)		10 s	≥4.01 log ₁₀	
			Alcohol-based sanitizers (95% alcohol w/w)		15 s	≥4.01 log ₁₀	
11	Rabenau <i>et al.</i> (2005)	SARS-CoV	Sterillium (45% iso-propanol, 30% n-propanol and 0.2% mecetronium etilsulphate)	ST	30 s	≥4.25 log ₁₀	Hands

			Sterillium rub (80% ethanol) {Gopal, 2020 #204}			$\geq 4.25 \log_{10}$ $\geq 5.5 \log_{10}$ $\geq 5.5 \log_{10}$	
12	Hirose <i>et al.</i> (2020)	IAV	Sterillium Virugard (95% ethanol)	ST and HS	ST (5 s, 15 s and 60 s)/HS (5 s, 15 s and 60 s)	$> 4.10, > 4.11, > 4.07 \log$ /HS ($> 4.12, > 4.16, > 4.16 \log$)	Skin
			80% EA (ethanol)			ST ($> 4.10, > 4.11, > 4.07 \log$)/HS ($> 4.12, > 4.16, > 4.16 \log$)	
			60% EA			ST ($> 4.10, > 4.11, > 4.07 \log$)/HS ($> 4.12, > 4.16, > 4.16 \log$)	
			40% EA			ST ($> 4.10, > 4.11, > 4.07 \log$)/HS ($> 4.12, > 4.16, > 4.16 \log$)	
			20% EA			ST ($\sim 0.09, \sim 0.07, \sim 0.06 \log$)/HS ($\sim 0.73, \sim 0.85, \sim 0.88 \log$)	
			70% IPA (isopropanol)			ST ($> 4.10, > 4.11, > 4.07 \log$)/HS ($> 4.12, > 4.16, > 4.16 \log$)	
			0.2% CHG (chlorhexidine gluconate)			ST ($\sim 0.08, \sim 0.17, \sim 0.19 \log$)/HS ($\sim 0.74, \sim 0.95, \sim 1.02 \log$)	
			1.0% CHG			ST ($\sim 0.23, \sim 0.24, \sim 0.40 \log$)/HS ($\sim 2.85, \sim 3.25, \sim 3.39 \log$)	
			0.05% BAC (benzalkonium chloride)			ST ($\sim 0.69, \sim 1.78, \sim 2.71 \log$)/HS ($\sim 0.78, \sim 1.04, \sim 1.23 \log$)	
			0.2% BAC			ST ($\sim 2-43, \sim 2.34, > 4.07 \log$)/HS ($\sim 1.64, \sim 2.85, \sim 3.24 \log$)	
		SARS-CoV-2	80% EA			ST ($> 4.50, > 4.50, > 4.50 \log$)/HS ($> 4.19, > 4.17, > 4.14 \log$)	
			60% EA			ST ($> 4.50, > 4.50, > 4.50 \log$)/HS ($> 4.19, > 4.17, > 4.14 \log$)	
			40% EA			ST ($> 4.50, > 4.50, > 4.50 \log$)/HS ($> 4.19, > 4.17, > 4.14 \log$)	
			20% EA			ST ($> 4.50, > 4.50, > 4.50 \log$)/HS ($> 4.19, > 4.17, > 4.14 \log$)	

(continued on next page)

Table III (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Type	
		70% IPA			ST (~0.08, ~0.25, ~0.33 log)/HS (~0.53, ~0.61, ~0.81 log)		
		0.2% CHG			ST (>4.50, >4.50, >4.50 log)/HS (>4.19, >4.17, >4.14 log)		
		1.0% CHG			ST (~0.33, ~0.42, ~0.58 log)/HS (~2.19, ~2.31, ~2.42 log)		
		0.05% BAC			ST (~1.00, ~1.42, ~1.83 log)/HS (~2.62, ~2.94, ~3.17 log)		
		0.2% BAC			ST (~1.33, ~1.75, ~2.17 log)/HS (~2.03, ~2.19, ~2.36 log)		
13	Huang <i>et al.</i> (2020)	Patients with SARS-CoV-2 infection	Chlorhexidine oral rinse (15 mL)	Oral and oropharyngeal cavity	30 s twice a day for 4 days	37.9% positive SARS-CoV-2 test, 62.1% negative test	Oral and oropharyngeal cavity
			without exposure		(-)	94.5% positive SARS-CoV-2 test, 5.5% negative test	
			Chlorhexidine oral rinse (15 mL) + oropharyngeal spray (1.5 mL)		30 s oral rinse + spray, twice a day for 4 days	14.0% positive SARS-CoV-2 test, 80% negative test	
			without exposure		(-)	93.8% positive SARS-CoV-2 test, 6.2% negative test	
14	Ijaz <i>et al.</i> (2020)	HCoV-229E (1), SARS-CoV-2 (2)	Bar soap PCMX - (0.090% w/w)	ST	0.5–1 min	(1) ≥3.3 log ₁₀ , (2) ≥4.1 log ₁₀	Hands
			Liquid gel handwash - salicylic acid (0.025% w/w)		0.5–1 min	(1) ≥3.6 log ₁₀ , (2) ≥3.6 log ₁₀	
			Foaming handwash - benzalkonium chloride (0.025% w/w)		1 min	(1) ≥3.3 log ₁₀ , (2) ≥3.4 log ₁₀	

			Foaming handwash - salicylic acid (0.023% w/w)		0.5–1 min	(1) $\geq 3.5 \log_{10}$, (2) $\geq 5.0 \log_{10}$	
			Antiseptic liquid - PCMX (0.021% w/v)		5 min	(1) $\geq 5.2 \log_{10}$, (2) $\geq 4.7 \log_{10}$	
			Hand sanitizer gel - ethanol (53% w/w)		1 min	(1) $\geq 5.4 \log_{10}$, (2) $\geq 4.2 \log_{10}$	
			Hand sanitizer gel - citric acid (1.5% w/w), lactic acid (0.41% w/w)		0.5–1 min	(1) $\geq 5.2 \log_{10}$, (2) $\geq 4.7 \log_{10}$	
15	Steinhauer <i>et al.</i> (2020)	Modified vaccinia virus Ankara	20% hand disinfectant - propan-2-ol	ST	15 s	$\sim 0.17 \log_{10}$	Hands
			80% hand disinfectant - propan-2-ol		15 s	$\geq 4.19 \log_{10}$	
		SARS-CoV-2	20% hand disinfectant - propan-2-ol		15 s	$\geq 4.02 \log_{10}$	
			20% hand disinfectant - propan-2-ol		30 s	$\geq 3.02 \log_{10}$	
			80% hand disinfectant - propan-2-ol		15 s	$\geq 2.02 \log_{10}$	
			80% hand disinfectant - propan-2-ol		30 s	$\geq 4.38 \log_{10}$	
16	Steinhauer <i>et al.</i> (2020)	SARS-CoV-2	80% chlorhexidine bis-(D-gluconate) 0.1 g	ST	5–10 min	$< 1.00 \log_{10}$	Oral
			80% chlorhexidine bis-(D-gluconate) 0.2 g		1–5 min	$< 1.00 \log_{10}$	
			80% 0.1 g octenidine dihydrochloride, 2 g phenoxyethanol		15 s	$\geq 4.00 \log_{10}$	
			80% 0.1 g octenidine dihydrochloride, 2 g phenoxyethanol		30 s	$\geq 4.00 \log_{10}$	
			80% 0.1 g octenidine dihydrochloride, 2 g phenoxyethanol		1 min	$\geq 4.00 \log_{10}$	
			20% 0.1 g octenidine dihydrochloride, 2 g phenoxyethanol		15 s	$\geq 4.00 \log_{10}$	

SARS-CoV-2, severe acute respiratory syndrome coronavirus-2; MERS-CoV, Middle East respiratory syndrome coronavirus; HCoV, human coronavirus; PVP-I, povidone-iodine; WHO, World Health Organization; ST, suspension test; HS, human skin.

Discussion

Stability and survival of SARS-CoV-2 exposed to heat and high humidity

Under ambient conditions (temperatures of 21–23°C and relative humidity of 40%), SARS-CoV-2 can remain viable on surfaces for hours to days [7,40]. The findings of this review support the evidence that coronaviruses are less viable when exposed to higher temperatures and higher relative humidity. It is not always possible to change the room temperature or humidity in indoor settings. However, rooms with the possibility to set these parameters between a determined range, such as intensive care units, operating rooms or hospital wards, can benefit as the survival of viruses is reduced markedly in warmer and higher humidity conditions.

Disinfection methods on environmental surfaces

Although most chemical agents have demonstrated virucidal activity against the coronavirus family, alcohols with concentrations of at least 60% showed a more constant and significant reduction in viral titres, promoting viral inactivation with shorter time exposures. This suggests that alcohols may be a better option when it comes to choosing a fast-acting and effective agent. Sodium hypochlorite, if preferred, should be used as a 0.1% solution, at least. If using QACs, a minimum exposure time of 30 min is recommended.

As household dishwashing detergent is more accessible compared with the other coatings discussed, it can be an effective alternative in providing long-lasting virucidal protection on surfaces in household settings or in countries that have difficulty in accessing other products, such as alcohols. However, further investigation is still needed to determine the efficacy and practicality of these coatings.

Ozone has virucidal activity targeting proteins on the viral envelope, inhibiting its entry to host cells. Higher concentrations of ozone must be used with caution due to the potential toxicity to humans; therefore, an ozone concentration of 20 ppm and an exposure time of 15 min is considered to be sufficient for optimal disinfection of surfaces [43,82,83].

For surface disinfection, UV-C irradiation seems to be the best alternative, as it is widely available and exceptionally convenient. It may be preferred over ozone as it is safer and less toxic to humans. However, when used with the purpose of whole-room disinfection, other methods, such as surface antimicrobial agents, could complement the strategy, as some surfaces may not be fully decontaminated due to shadowing or the composition of absorbable materials, such as fleece and wood.

Disinfection methods on biological surfaces

Adequate disinfection of hands is an important way to prevent indirect transmission of respiratory infections, especially during the era of SARS-CoV-2. Based on the review findings and

Table IV

Results of disinfection methods against airborne viruses

	Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀ or (%))
1	Buonanno <i>et al.</i> (2020)	Alphacoronavirus HCoV-229E Betacoronavirus HCoV-OC43	Far-UV-C light at 222 nm (0.5, 1 and 2 mJ/cm ²)	Dynamic aerosol/virus irradiation chamber	~20 s	1.7 mJ/cm ² produce 99.9% inactivation (3-log reduction) of aerosolized alpha HCoV-229E 1.2 mJ/cm ² produce 99.9% inactivation (3-log reduction) of aerosolized beta HCoV-OC43
2	Qiao <i>et al.</i> (2020)	PRCV	UV-C light 200–850 nm (13.9 mJ/cm ²) UV-C light 253±1 nm (49.6 mJ/cm ²)	Wind tunnel (high flow rate of 2439 L/min) Wind tunnel (low flow rate of 684 L/min)	1.3 s 5.1 s	2.2 log ₁₀ (99.4% removal efficiency) 3.7 log ₁₀ (99.98% removal efficiency)
3	Yu <i>et al.</i> (2020)	SARS-CoV-2	Novel Ni-foam-based filter (up to 200°C)	Aerosolized SARS-CoV-2	Single pass	99.8% reduction
4	He <i>et al.</i> (2004)	SARS coronavirus	Ag/Al ₂ O ₃ (Ag 5 wt%) catalytic oxidation Cu/Al ₂ O ₃ (Cu 10 wt%) catalytic oxidation	Ag/Al ₂ O ₃ and Cu/Al ₂ O ₃ wafers	5 min and 20 min	Virus undetectable Virus undetectable

SARS-CoV-2, severe acute respiratory syndrome coronavirus-2; PRCV, porcine respiratory coronavirus; HCoV, human coronavirus; UV-C, ultraviolet C irradiation; Ni, nickel; Ag, silver; Cu, copper.

Table V
Results of methods to recondition personal protective equipment

	Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Effect on material properties and functionality
1	Blanchard <i>et al.</i> (2020)	IAV and RSV	Ozone (20 ppm) + 50–70% RH	Surgical facemasks (1 cm x 1 cm sample swatches) Tyvek (disposable gown) 1 cm x 1 cm N95 respirators 1 cm x 1 cm Bunny suits 1 cm x 1 cm PAPR hoods 1 cm x 1 cm	40 min	Equal to 70% ethanol inactivation	Material properties were preserved and filtration capacity of masks was maintained.
2	Campos <i>et al.</i> (2020)	SARS-CoV-2	Ambient humidity (60%) without BSA Ambient humidity (60%) without BSA Ambient humidity (60%) without BSA Ambient humidity (60%) without BSA Ambient humidity (60%) with BSA Ambient humidity (60%) with BSA Ambient humidity (60%) with BSA Ambient humidity (60%) with BSA 100% humidity without BSA 100% humidity without BSA 100% humidity without BSA 100% humidity without BSA 100% humidity with BSA 100% humidity with BSA	Meltblown fabric from N95-grade FFRs	60°C for 30 min 75°C for 30 min 85°C for 20 min 95°C for 5 min 60°C for 30 min 75°C for 30 min 85°C for 20 min 95°C for 5 min 60°C for 30 min 75°C for 30 min 85°C for 20 min 95°C for 5 min 60°C for 30 min 75°C for 30 min 85°C for 20 min	2.16 ± 0.23 log ₁₀ 3.69 ± 0.32 log ₁₀ >4.77 log ₁₀ >4.77 log ₁₀ 1.07 ± 0.06 log ₁₀ 2.89 ± 0.31 log ₁₀ 4.3 ± 0.55 log ₁₀ 4.8 ± 0.44 log ₁₀ 2.82 ± 0.09 log ₁₀ >4.97 log ₁₀ >4.97 log ₁₀ >4.97 log ₁₀ 2.27 ± 0.09 log ₁₀ 4.92 ± 0.12 log ₁₀ >5.02 log ₁₀	Temperatures of 75–85 °C are able to efficiently inactivate the virus in 20–30 min under 100% RH, without lowering filtration efficiency. Filtration efficacy started to decrease significantly after 10 cycles with temperature of 95°C probably due to the absorption of water or other mechanisms that can decay the electrostatic charge.

(continued on next page)

Table V (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Effect on material properties and functionality	
3	Choi <i>et al.</i> (2020)	SARS-CoV-2	100% humidity with BSA				All FFRs absorbed <1 g of water when in a paper bag. Collection efficacy exceeded 95% and inhalation resistance was preserved. After five cycles of moist heat treatment, 3M 8210 and NS 721 had no change in strap elasticity, while 3M 1860 and 3M 8511 showed a small change (<10%). Masks maintained fibre diameters similar to untreated masks and continued to meet standards for fit, filtration efficiency and breathing resistance. Cotton and polyamide 66 (PA66) can strongly trap viruses as only 56% of SARS-CoV-2 can be recovered from cotton samples and 92% from PA66 after viral inoculation. PPP is poor at trapping viruses.
			100% humidity with BSA		95°C for 5 min	>5.02 log ₁₀	
			Moist heat generated by multi-cooker	FFRs 3M Model 1860 in simulated saliva	65°C for 30 min	~1.5 log ₁₀	
				FFRs 3M Model 1860 in simulated lung fluid		~3.2 log ₁₀	
				FFRs 3M Model 8511 in simulated saliva		~2.5 log ₁₀	
				FFRs 3M Model 8511 in simulated lung fluid		~3.2 log ₁₀	
		FFRs 3M Model 8210 in simulated saliva		~2.2 log ₁₀			
		FFRs NS Model 7210 in simulated saliva		~2.2 log ₁₀			
4	Daeschler <i>et al.</i> (2020)	SARS-Cov-2	70°C + 50% RH	N95 respirators	2–18 min	Reduced to undetectable levels	Masks maintained fibre diameters similar to untreated masks and continued to meet standards for fit, filtration efficiency and breathing resistance.
5	Gopal <i>et al.</i> (2020)	SARS-Cov-2	Zinc oxide embedded into fabrics (only tested on PA66)	Cotton, polypropylene (PPP) fabrics and polyamide (PA66)	60 min	2 log	Cotton and polyamide 66 (PA66) can strongly trap viruses as only 56% of SARS-CoV-2 can be recovered from cotton samples and 92% from PA66 after viral inoculation. PPP is poor at trapping viruses.
6	Ibanez-Cervantes <i>et al.</i> (2020)	SARS-CoV-2	Hydrogen peroxide plasma	N95 3M Model 8210	47 min	Undetectable by RT-PCR	Not tested
7	Ludwig-Begall <i>et al.</i> (2020)	PRCV	UV irradiation Vaporized H ₂ O ₂ (59% liquid H ₂ O ₂) 750 ppm	Surgical mask coupons	2 min 28 min	~5 log ₁₀ ~5 log ₁₀	Not tested

			Dry heat (102°C)		60 min	~5.5 log ₁₀	
			UV irradiation	Surgical mask straps	2 min	~2.9 log ₁₀	
			Vaporized H ₂ O ₂ (59% liquid H ₂ O ₂) 750 ppm		28 min	Non-significant	
			Dry heat (102°C)		60 min	~1.2 log ₁₀	
			UV irradiation	FFR coupons	4 min	~3.2 log ₁₀	
			Vaporized H ₂ O ₂ (59% liquid H ₂ O ₂) 750 ppm		28 min	~4 log ₁₀	
			Dry heat (102°C)		60 min	~2.5 log ₁₀	
			UV irradiation	FFR straps	4 min	(-)	
			Vaporized H ₂ O ₂ (59% liquid H ₂ O ₂) 750 ppm		28 min	~1.2 log ₁₀	
			Dry heat (102°C)		60 min	(-)	
8	Ma <i>et al.</i> (2020)	Avian infectious bronchitis virus	Steam	N95 FFR masks	5 min	Undetectable by RT-PCR	Blocking efficacy of 99% verified in all masks except for one model that seemed to have thinner layers compared with other models. Therefore, masks with thinner layers can have reduced blocking efficacy.
9	Mantlo <i>et al.</i> (2020)	SARS-CoV-2	Clyraguard copper iodine complex undiluted	Suspension test	10 min 30 min 60 min	2 log Below limit of detection (<75 TCID ₅₀ per mL) Below limit of detection (<75 TCID ₅₀ per mL)	Not tested
10	Ozog <i>et al.</i> (2020)	SARS-CoV-2	UV-C irradiation (1.5 J/cm ² to each side)	N95 FFR models (3M 1860, 8210, 8511, 9211; Moldex 1511)	60–70 s (for each side)	Below limit of detection (10 ^{1.3} TCID ₅₀ /4 mm punch)	Not tested
11	Perkins <i>et al.</i> (2020)	SARS-CoV-2	Dry heat (60°C)	N95 respirator coupons + parchment paper	60 min	All samples were positive analysed by microscopy for cytopathic effect	Not tested
			Dry heat (70°C)		60 min	All samples were positive analysed by microscopy for cytopathic effect	
			Dry heat (75°C)		60 min	All samples were positive analysed by microscopy for cytopathic effect	

(continued on next page)

Table V (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Effect on material properties and functionality	
		Dry heat (60°C) Dry heat (70°C)	N95 respirator coupons + tissue culture	(-) 60 min	(-) All samples were positive analysed by microscopy for cytopathic effect		
		Dry heat (75°C)		60 min	All samples were positive analysed by microscopy for cytopathic effect		
		Dry heat (60°C) Dry heat (70°C)	Intact N95 respirators	(-) 60 min	(-) All samples were positive analysed by microscopy for cytopathic effect		
		Dry heat (75°C)		60 min	Most samples were positive analysed by microscopy for cytopathic effect		
		Ambient temperature		5 days	5/9 samples were positive analysed by microscopy for cytopathic effect		
12	Rathnasinghe <i>et al.</i> (2020)	SARS-CoV-2	UV-C irradiation (5.43 mW/cm ²)	N95 mask squares	120 s per side	3.5 log	Not tested
13	Rockey <i>et al.</i> (2020)	Bacteriophage MS2	Temperature (72°C and 82°C) + PBS	N95 respirator coupons	30 min	0.24 log ₁₀ (72°C + 1% RH), 0.19 log ₁₀ (82°C + 1% RH) 6.87 log ₁₀ (72°C + 89% RH), 6.90 log ₁₀ (82°C + 89% RH) 1.44 log ₁₀ (72°C + 1% RH), 2.77 log ₁₀ (82°C + 1% RH) 6.56 log ₁₀ (72°C + 89% RH), 7.16 log ₁₀ (82°C + 89% RH) 0.99 log ₁₀ (72°C + 13% RH), 0.88 log ₁₀ (82°C + 1% RH) 1.45 log ₁₀ (72°C + 25% RH), 1.74 log ₁₀ (82°C + 13% RH)	Not tested
		Temperature (72°C and 82°C) + DMEM-A					
		Temperature (72°C and 82°C) + saliva					

			Temperature (72°C and 82°C) + (PBS + BSA)			1.5 log ₁₀ (72°C + 13% RH), 0.77 log ₁₀ (82°C + 1% RH) 2.72 log ₁₀ (72°C + 25% RH), 3.56 log ₁₀ (82°C + 13% RH) 0.99 log ₁₀ (72°C + 1% RH), 1.48 log ₁₀ (82°C + 1% RH) 6.79 log ₁₀ (72°C + 89% RH), 6.70 log ₁₀ (82°C + 89% RH) 2.58 log ₁₀ (72°C + 1% RH), 3.87 log ₁₀ (82°C + 1% RH) 6.81 log ₁₀ (72°C + 89% RH), 7.63 log ₁₀ (82°C + 89% RH) 0.95 log ₁₀ (72°C + 13% RH), 1.09 log ₁₀ (82°C + 1% RH) 1.69 log ₁₀ (72°C + 25% RH), 2.62 log ₁₀ (82°C + 13% RH) 1.33 log ₁₀ (72°C + 13% RH), 0.76 log ₁₀ (82°C + 1% RH) 1.34 log ₁₀ (72°C + 25% RH), 1.98 log ₁₀ (82°C + 13% RH) 2.51 log ₁₀ (72°C + 1% RH), 3.30 log ₁₀ (82°C + 1% RH) 4.19 log ₁₀ (72°C + 89% RH), 4.38 log ₁₀ (82°C + 89% RH) 1.25 log ₁₀ (72°C + 1% RH), 2.71 log ₁₀ (82°C + 1% RH) 3.71 log ₁₀ (72°C + 89% RH), 3.37 log ₁₀ (82°C + 89% RH)	
			Temperature (72°C and 82°C) + PBS				
			Temperature (72°C and 82°C) + DMEM-A				
			Temperature (72°C and 82°C) + saliva				
			Temperature (72°C and 82°C) + (PBS + BSA)				
		MHV	Temperature (72°C and 82°C) + DMEM-A				
		IAV	Temperature (72°C and 82°C) + DMEM-A				
14	Glasbrenner <i>et al.</i> (2021)	TGEV	UV (300–400 nm) simulated sunlight	FFR 3M 1860 FFR 3M 8210 FFR 3M 8511	(-) (-) (-)	(-) (-) Inactivation less efficient	All FFRs maintained collection efficacy and breathing resistance after one

(continued on next page)

Table V (continued)

Study	Virus	Disinfectant	Disinfection phase	Exposure time	Reduction of viral infectivity (log ₁₀) or (%)	Effect on material properties and functionality
			FFR NS 7210	(-)	Inactivation below level of detection	and five cycles ((Model 3M 8210 not tested for five cycles).
	SARS-CoV-2		FFR 3M 1860 + SS and LF	20 min (13.3 J cm ²) SS/40 min (26.5 J cm ²) FL	Inactivation below level of detection	Reduced strap elasticity from NS 7210 model with 19% change in stress).
			FFR 3M 8210 + SS and LF	(-)	(-)	
			FFR 3M 8511 + SS and LF	60 min (37.8 J cm ²) for SS and FL	Complete inactivation	
			FFR NS 7210 + SS and LF	20 min (13.3 J cm ²) for SS and LF	Inactivation below level of detection	
15	Uppal <i>et al.</i> (2021)	HCoV-OC43	N95 FFRs	10 min	98.1411%	Not tested
		Ozone (20 ppm) Ozone (25 ppm)		10 min 15 min 20 min	97.4138% 99.9947% 99.9966%	
		Ozone (50 ppm)		10 min 15 min 20 min	99.9860% 99.9956% 99.9925%	
16	Valdez-Salas <i>et al.</i> (2021)	Enveloped H5N1 avian influenza virus	Formulated disinfectant - 0.2% benzalkonium chloride, 85% ethanol-water, 0.03% triclosan, 10% silver nanoparticles, 0.3% lauryl alcohol ethoxylate, 0.2% Triton X-100, 2% citric acid, microdacyn	Suspension test 15 min	No presence of haemagglutinine - complete inactivation	Not tested

IAV, avian influenza virus; RSV, respiratory syncytial virus; SARS-CoV-2, severe acute respiratory syndrome coronavirus-2; PRCV, porcine respiratory coronavirus; IBV, avian infectious bronchitis virus; MHV, mouse hepatitis coronavirus; PAPR, powered air purifying respirator; FFR, filtering facepiece respirator; RH, relative humidity; UV-C, ultraviolet C irradiation; PSB, phosphate-buffered saline; BSA, bovine serum albumin; DMEM, Dulbecco's modified Eagle medium; SS, simulated saliva; FF, lung fluid.

Table VI

Risk of bias assessment using the Office of Health Assessment and Translation (OHAT) Risk of Bias Rating Tool for Human and Animal Studies Potential source of bias was graded as low risk (++) , probable low risk (+) , probable high risk or not reported (-) and high risk (--)

Study	Study design	Was administered dose or exposure level adequately randomized?	Was allocation to study groups adequately concealed?	Were experimental conditions identical across study groups?	Were research personnel blinded to the study group during the study?	Were outcome data complete without attrition or exclusion from analysis?	Can we be confident in the exposure characterization?	Can we be confident in the outcome assessment (including blinding of assessors)?	Were all measured outcomes reported?	Were there no other potential threats to internal validity?
Anderson <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(NR)	(NR)	(NR)	(+)	(++)	(+)
Bedell <i>et al.</i> (2016)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(NR)	(NR)	(-)	(++)	(-)
Behzadinasab <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Bidra <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(NR)	(++)	(++)	(++)	(+)
Bidra <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(NR)	(++)	(++)	(++)	(+)
Biryukov <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(+)	(++)
Blanchard <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(-)	(-)	(NR)	(++)	(++)	(+)	(+)
Buonanno <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(NR)	(++)	(++)	(++)	(++)
Campos <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Casanova <i>et al.</i> (2010)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(++)	(+)	(++)	(++)	(++)
Choi. <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(++)	(+)	(++)	(++)	(+)
Colnago <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(-)	(-)	(++)	(++)	(+)
Criscuolo <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(+)	(++)	(+)
Daeschler <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(+)	(++)	(++)	(+)	(++)	(++)
Eggers <i>et al.</i> (2015)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(++)	(+)
Frank <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(+)	(++)	(++)	(+)

(continued on next page)

Table VI (continued)

Study	Study design	Was administered dose or exposure level adequately randomized?	Was allocation to study groups adequately concealed?	Were experimental conditions identical across study groups?	Were research personnel blinded to the study group during the study?	Were outcome data complete without attrition or exclusion from analysis?	Can we be confident in the exposure characterization?	Can we be confident in the outcome assessment (including blinding of assessors)?	Were all measured outcomes reported?	Were there no other potential threats to internal validity?
Gamble <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(+)	(++)	(++)	(++)	(+)
Gerchman <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(-)	(++)	(+)	(++)	(+)
Gopal <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(++)	(+)
Gudmundsdottir <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(-)	(++)	(++)	(+)
He <i>et al.</i> (2004)	<i>In vitro</i>	Not applicable	Not applicable	(-)	(-)	(+)	(++)	(++)	(-)	(+)
Heilingloh <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(+)	(++)
Hulkower <i>et al.</i> (2011)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Ibanez-Cervantes <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(+)	(++)	(++)	(++)
Khaiboullina <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(+)	(++)
Kratzel <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(NR)	(-)	(++)	(++)	(++)	(++)	(++)
Leslie <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(NR)	(-)	(+)	(+)	(++)	(+)	(-)
Liang <i>et al.</i> (2020)	<i>In vivo</i> and <i>in vitro</i>	(+)	(NR)	(++)	(NR)	(++)	(++)	(+)	(++)	(+)
Liu <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(+)	(++)	(+)	(++)	(+)
Ludwig-Begall <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Ma <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(-)	(-)	(NR)	(+)	(+)	(++)	(+)
Malenovská (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(++)	(++)
Mantlo <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(++)	(++)
Martins <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(+)	(++)
Meister <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(-)	(++)	(+)	(+)
Meyers <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(-)	(++)	(++)	(+)	(-)
Monge <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(NR)	(-)	(+)	(+)	(+)	(++)	(+)

Mukherjee <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(+)	(-)	(+)	(++)	(+)
Ozog <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Perkins <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(-)	(+)	(++)	(+)
Qiao <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(+)
Rabenau <i>et al.</i> (2005)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(+)	(+)	(++)
Rathnasinghe <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(+)	(++)	(++)
Ratnesar-Shumate <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(+)	(++)	(++)	(++)	(++)	(++)
Rockey <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Wood and Payne (1998)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(+)	(+)	(+)
Yu <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(NR)	(-)	(NR)	(++)	(+)	(NR)	(+)
Franke <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Gidari <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Glasbrenner <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(-)	(-)	(++)	(+)	(+)
Hirose <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Hu <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(+)	(++)	(++)	(+)	(+)
Huang <i>et al.</i> (2020)	Prospective cohort	(++)	(-)	(-)	(-)	(++)	(++)	(++)	(++)	(+)
Ijaz <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Messina <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Steinhauer <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(+)	(-)	(+)	(++)	(++)	(+)	(+)
Steinhauer <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Trivellini <i>et al.</i> (2020)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Uppal <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(++)	(-)	(++)	(++)	(++)	(++)	(++)
Valdez-Salas <i>et al.</i> (2021)	<i>In vitro</i>	Not applicable	Not applicable	(-)	(-)	(-)	(++)	(+)	(+)	(+)

evidence in the literature, the original formulations of WHO-recommended hand rubs seem to be less active against SARS-CoV-2 compared with modified formulations [13,84]. This is significant as many companies seek standard recommendations from WHO to produce disinfectants with the adequate proportion of ethanol/isopropanol and glycerol. These formulations could be updated to ensure optimal disinfection efficacy of formulations against SARS-CoV-2. Commercially available personal care products, such as soap bars, liquid cleansers (containing surfactant) and alcohol-based hand sanitizers (at least 30% ethanol or propanol), were all able to reduce SARS-CoV-2 titre after 10–20 s of exposure [13,55,58,85]. This suggests that the current procedure for handwashing is effective against SARS-CoV-2 at the established concentrations and duration.

At present, no methods are in place regarding eye or respiratory tract disinfection in order to stop the transmission of SARS-CoV-2, and this deserves further investigation due to potential toxicity. However, there are viable options in specific settings, such as during ophthalmologic procedures or interventions where aerosols may be generated. While very low concentrations of PVP-I showed in-vitro viral inactivation, in-vivo conditions must be taken into account due to the fact that biological debris such as physiological buffers in nasal secretions can lower the effective concentration of PVP-I. Therefore, a concentration of at least 1.25% PVP-I is recommended for in-vivo application [53].

In summary, for oral rinses and skin cleansers, products containing PVP-I should be preferred, as its action is rapid and efficient. Soap bars, surfactant and alcohol-based hand sanitizers are all excellent alternatives for hand hygiene.

Disinfection methods against airborne viruses

Recent evidence indicates that airborne transfer is the main route of transmission of SARS-CoV-2, being more evident in indoor spaces with poor ventilation. Considering that coronaviruses cannot tolerate high temperatures, filtration or ventilation systems coupled with heatable metal filters may be an effective option. It is also evident that SARS-CoV-2 is susceptible to UV-C irradiation. As the latter is the only commercially available option at present, the installation of an upper room germicidal UV-C irradiation device, for example, in healthcare facilities, indoor spaces that accommodate a large number of people, or even in household settings, can be beneficial. Other than UV irradiation, the remaining methods in this section provide preliminary evidence of effective ways to decontaminate the air, indicating the future of more sophisticated and efficient air conditioning systems.

Methods to decontaminate and recondition personal protective equipment

The COVID-19 pandemic has had a significant impact on the environment and mass production of PPE to meet the world's rapid and urgent demand, creating major challenges in waste management on a global scale [86–88]. Surgical masks, for instance, are composed of plastic that is not biodegradable and may end up in waterbeds, causing harm to the environment and the fauna of these areas. Methods that aim to decontaminate and recondition PPE for reuse can be beneficial not only for the

environment but also in cases of shortages of PPE, as experienced by many countries during the COVID-19 pandemic.

Based on these studies, there is still insufficient evidence to support the virucidal efficacy of metal-embedded fabrics. Moreover, as it is important to preserve the functionality of PPE after decontamination, the only methods that provided evidence of effective sterilization without compromising the integrity of PPE (with a limited number of cycles) were heat and ozone treatment, making these methods better and safer options at the present time.

Deposition solutions

Three of the studies included in this review addressed how different deposition solutions can change the viral inactivation rate. It was found that the viral load of SARS-CoV-2 and bacteriophages MS2 and Phi6 deposited in DMEM-A (cell culture medium formulations) showed, under different temperature and humidity exposures, a more significant reduction in viral titre compared with the viral load deposited in phosphate-buffered saline (PBS) [40]. Interestingly, the viral load deposited in freshly collected human saliva demonstrated a log₁₀ reduction trend more similar to PBS compared with DMEM-A. Bovine serum albumin containing higher concentrations of protein can be used to mimic body fluids, particularly sputum [31,42]. This may suggest that laboratory-made solutions may not fully represent the behaviour of biological fluids.

Limitations of this review

One major limitation of this systematic review is that all the studies included are based on in-vitro findings, with some extensive experiments trying to mimic in-vivo conditions. However, the real efficacy in in-vivo settings needs further investigation.

Nineteen of the studies included in this review used surrogate viruses to mimic the behaviour of SARS-CoV-2. Surrogate viruses were included due to the biosafety level of SARS-CoV-2 that may hinder the use of this virus in some experiments. It may also have been unavailable in some laboratories, especially at the beginning of the pandemic when little was known about SARS-CoV-2. To evaluate the efficacy of disinfectants, vaccinia virus, in particular, is a reference virus used in Europe as a surrogate for enveloped viruses (EN 14476) [89]. As SARS-CoV-2 is an enveloped virus easily susceptible to disinfection, as verified in the review findings, methods that can effectively target more resilient surrogate enveloped viruses translate into efficacy against SARS-CoV-2.

In conclusion, the results demonstrate that several household and hospital disinfection agents, UV-C irradiation, ozone and surface coatings are effective for inactivation of the coronavirus family, including SARS-CoV-2, on environmental surfaces. While SARS-CoV-2 can survive for hours to days depending on the surface, high temperature and humidity are key factors in viral decay. Decontamination of PPE can be performed effectively using heat treatment, UV-C irradiation and hydrogen peroxide vapour. Zinc ions can potentially provide prolonged disinfection when embedded into fabrics. Formulations containing PVP-I at different concentrations can provide virucidal action in the form of oral rinses, topical eye disinfection and skin cleansers. In the case of hand hygiene, typical soap bars, ethanol and propanol can inactivate SARS-CoV-2. Regarding disinfection methods against airborne

particles, air filtration systems with materials that possess catalytic properties, UV-C devices and heating systems can reduce viral particles effectively. This review supports improved selection of the most effective disinfection method for each specific setting, potentially resulting in better outcomes during the present pandemic, and also the prevention of viral healthcare-associated infections.

Acknowledgements

The first author wishes to thank Dr. Luís Cobrado for his expertise and guidance throughout this work, and Carolina Xavier for support, effort and assistance regarding the writing and language of this work.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jhin.2021.07.014>.

Conflicts of interest statement

None declared.

Funding sources

None.

References

- [1] Mofijur M, Fattah IMR, Alam MA, Islam ABMS, Ong HC, Rahman SMA, et al. Impact of COVID-19 on the social, economic, environmental and energy domains: lessons learnt from a global pandemic. *Sustain Prod Consum* 2021;26:343–59.
- [2] World Health Organization. COVID-19 weekly epidemiological update. Geneva: WHO; 2021.
- [3] Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020;395:565–74.
- [4] Centers for Disease Control and Prevention. Scientific brief: SARS-CoV-2 transmission. Atlanta, GA: CDC; 2021.
- [5] Bulfone TC, Malekinejad M, Rutherford GW, Razani N. Outdoor transmission of SARS-CoV-2 and other respiratory viruses: a systematic review. *J Infect Dis* 2021;223:550–61.
- [6] Greenhalgh T, Jimenez JL, Prather KA, Tufekci Z, Fisman D, Schooley R. Ten scientific reasons in support of airborne transmission of SARS-CoV-2. *Lancet* 2021;397:1603–5.
- [7] Johansson MA, Quandelacy TM, Kada S, Prasad PV, Steele M, Brooks JT, et al. SARS-CoV-2 transmission from people without COVID-19 symptoms. *JAMA Network Open* 2021;4:e2035057.
- [8] Lewis D. Superspreading drives the COVID pandemic – and could help to tame it. *Nature* 2021;590:544–6.
- [9] World Health Organization. Transmission of SARS-CoV-2: implications for infection prevention precautions. Geneva: WHO; 2020.
- [10] van Doremalen N, Bushmaker T, Morris DH, Holbrook MG, Gamble A, Williamson BN, et al. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *N Engl J Med* 2020;382:1564–7.
- [11] Mondelli MU, Colaneri M, Seminari EM, Baldanti F, Bruno R. Low risk of SARS-CoV-2 transmission by fomites in real-life conditions. *Lancet Infect Dis* 2021;21:e112.
- [12] Ijaz MK, Nims RW, Zhou SS, Whitehead K, Srinivasan V, Kapes T, et al. Microbicidal actives with virucidal efficacy against SARS-CoV-2 and other beta- and alpha-coronaviruses and implications for future emerging coronaviruses and other enveloped viruses. *Sci Rep* 2021;11:5626.
- [13] Kratzel A, Todt D, V'Kovski P, Steiner S, Gultom M, Thao TTN, et al. Inactivation of severe acute respiratory syndrome coronavirus 2 by WHO-recommended hand rub formulations and alcohols. *Emerg Infect Dis* 2020;26:1592–5.
- [14] Mentel R, Shirmakher R, Kevich A, Dreizin RS, Shmidt I. Virus inactivation by hydrogen peroxide. *Vopr Virusol* 1977;731–3.
- [15] Sriwilajaroen N, Wilairat P, Hiramatsu H, Takahashi T, Suzuki T, Ito M, et al. Mechanisms of the action of povidone-iodine against human and avian influenza A viruses: its effects on hemagglutination and sialidase activities. *Viol J* 2009;6:124.
- [16] Nardell EA, Nathavitharana RR. Airborne spread of SARS-CoV-2 and a potential role for air disinfection. *JAMA* 2020;324:141–2.
- [17] Storm N, McKay LGA, Downs SN, Johnson RI, Birru D, de Samber M, et al. Rapid and complete inactivation of SARS-CoV-2 by ultraviolet-C irradiation. *Sci Rep* 2020;10:22421.
- [18] Nguyen LH, Drew DA, Graham MS, Joshi AD, Guo C, Ma W, et al. Risk of COVID-19 among front-line health-care workers and the general community: a prospective cohort study. *Lancet Publ Health* 2020;5:e475–83.
- [19] Burki T. Global shortage of personal protective equipment. *Lancet Infect Dis* 2020;20:785–6.
- [20] Haas EJ, Angulo FJ, McLaughlin JM, Anis E, Singer SR, Khan F, et al. Impact and effectiveness of mRNA BNT162b2 vaccine against SARS-CoV-2 infections and COVID-19 cases, hospitalisations, and deaths following a nationwide vaccination campaign in Israel: an observational study using national surveillance data. *Lancet* 2021;397:1819–29.
- [21] Moghadas SM, Vilches TN, Zhang K, Wells CR, Shoukat A, Singer BH, et al. The impact of vaccination on COVID-19 outbreaks in the United States. *Clin Infect Dis* 2021. <https://doi.org/10.1093/cid/ciab079>.
- [22] Tenforde MW, Olson SM, Self WH, Talbot HK, Lindsell CJ, Steingrub JS, et al. *MMWR Morb Mortal Wkly Rep* 2021;70:674–9.
- [23] Gidari A, Sabbatini S, Bastianelli S, Pierucci S, Busti C, Bartolini D, et al. SARS-CoV-2 survival on surfaces and the effect of UV-C light. *Viruses* 2021;13.
- [24] Biryukov J, Boydston JA, Dunning RA, Yeager JJ, Wood S, Reese AL, et al. Increasing temperature and relative humidity accelerates inactivation of SARS-CoV-2 on surfaces. *mSphere* 2020;5.
- [25] Casanova LM, Jeon S, Rutala WA, Weber DJ, Sobsey MD. Effects of air temperature and relative humidity on coronavirus survival on surfaces. *Appl Environ Microbiol* 2010;76:2712–7.
- [26] Gamble A, Fischer RJ, Morris DH, Yinda KC, Munster VJ, Lloyd-Smith JO. Heat-treated virus inactivation rate depends strongly on treatment procedure. *bioRxiv* 2020. <https://doi.org/10.1101/2020.08.10.242206>.
- [27] Wood A, Payne D. The action of three antiseptics/disinfectants against enveloped and non-enveloped viruses. *J Hosp Infect* 1998;38:283–95.
- [28] Rabenau HF, Kampf G, Cinatl J, Doerr HW. Efficacy of various disinfectants against SARS coronavirus. *J Hosp Infect* 2005;61:107–11.
- [29] Hulkower RL, Casanova LM, Rutala WA, Weber DJ, Sobsey MD. Inactivation of surrogate coronaviruses on hard surfaces by health care germicides. *Am J Infect Control* 2011;39:401–7.
- [30] Bedell K, Buchaklian AH, Perlman S. Efficacy of an automated multiple emitter whole-room ultraviolet-C disinfection system against coronaviruses MHV and MERS-CoV. *Infect Control Hosp Epidemiol* 2016;37:598–9.
- [31] Anderson ER, Hughes GL, Patterson EI. Inactivation of SARS-CoV-2 on surfaces and in solution with Virusend (TX-10), a novel disinfectant. *bioRxiv* 2020. <https://doi.org/10.1101/2020.11.25.394288>.
- [32] Behzadinasab S, Chin A, Hosseini M, Poon L, Ducker WA. A surface coating that rapidly inactivates SARS-CoV-2. *ACS Appl Mater Interfaces* 2020;12:34723–7.

- [33] Colnago LA, Trevisol IM, Voss Rech D, Forato LA, Igreja do Nascimento Mitre C, Gagliardi LJP, et al. Simple, low-cost and long-lasting film for virus inactivation using avian coronavirus model as Challenge. *Int J Environ Res Public Health* 2020;17.
- [34] Gerchman Y, Mamane H, Friedman N, Mandelboim M. UV-LED disinfection of coronavirus: wavelength effect. *J Photochem Photobiol B-Biol* 2020;212:7.
- [35] Heilingloh CS, Aufderhorst UW, Schipper L, Dittmer U, Witzke O, Yang D, et al. Susceptibility of SARS-CoV-2 to UV irradiation. *Am J Infect Control* 2020;48:1273–5.
- [36] Khaiboullina S, Uppal T, Dhabarde N, Subramanian VR, Verma SC. Inactivation of human coronavirus by titania nanoparticle coatings and UVC radiation: throwing light on SARS-CoV-2. *Viruses* 2020;13.
- [37] Liu SF, Luo W, Li D, Yuan Y, Tong W, Kang JJ, et al. Sec-eliminating the SARS-CoV-2 by AlGaIn based high power deep ultraviolet light source. *Adv Funct Mater* 2020. <https://doi.org/10.1002/adfm.202008452>. 7.
- [38] Malenovská H. Coronavirus persistence on a plastic carrier under refrigeration conditions and its reduction using wet wiping technique, with respect to food safety. *Food Environ Virol* 2020;12:361–6.
- [39] Martins RB, Castro IA, Pontelli M, Souza JP, Lima TM, Melo SR, et al. SARS-CoV-2 inactivation by ozonated water: a preliminary alternative for environmental disinfection. *Ozone Sci Eng* 2021;43:108–11.
- [40] Meyers C, Kass R, Goldenberg D, Milici J, Alam S, Robison R. Ethanol and isopropanol inactivation of human coronavirus on hard surfaces. *J Hosp Infect* 2020. <https://doi.org/10.1016/j.jhin.2020.09.026>.
- [41] Monge FA, Jagadesan P, Bondu V, Donabedian PL, Ista L, Chi EY, et al. Highly effective inactivation of SARS-CoV-2 by conjugated polymers and oligomers. *ACS Appl Mater Interfaces* 2020;12:55688–95.
- [42] Ratnesar-Shumate S, Williams G, Green B, Krause M, Holland B, Wood S, et al. Simulated sunlight rapidly inactivates SARS-CoV-2 on surfaces. *J Infect Dis* 2020;222:214–22.
- [43] Criscuolo E, Diotti RA, Ferrarese R, Alippi C, Viscardi G, Signorelli C, et al. Fast inactivation of SARS-CoV-2 by UV-C and ozone exposure on different materials. *Emerg Microbes Infect* 2021. <https://doi.org/10.1080/22221751.2021.1872354>.
- [44] Franke G, Knobling B, Brill FH, Becker B, Klupp EM, Belmar Campos C, et al. An automated room disinfection system using ozone is highly active against surrogates for SARS-CoV-2. *J Hosp Infect* 2021;112:108–13.
- [45] Hu X, Chen Z, Su Z, Deng F, Chen X, Yang Q, et al. Ozone water is an effective disinfectant for SARS-CoV-2. *Virol Sin* 2021. <https://doi.org/10.1007/s12250-021-00379-7>.
- [46] Messina G, Della Camera A, Ferraro P, Amodeo D, Corazza A, Nante N, et al. *Int J Environ Res Public Health* 2021;18.
- [47] Steinhauer K, Meister TL, Todt D, Krawczyk A, Paßvogel L, Becker B, et al. Comparison of the in-vitro efficacy of different mouthwash solutions targeting SARS-CoV-2 based on the European Standard EN 14476. *J Hosp Infect* 2021;111:180–3.
- [48] Trivellini N, Buffolo M, Onelia F, Pizzolato A, Barbato M, Orlandi VT, et al. Inactivating SARS-CoV-2 using 275 nm UV-C LEDs through a spherical irradiation box: design, characterization and validation. *Materials (Basel)* 2021;14.
- [49] Uppal T, Khazaeili A, Snijders AM, Verma SC. Inactivation of human coronavirus by FATHHOME's dry sanitizer device: rapid and eco-friendly ozone-based disinfection of SARS-CoV-2. *Pathogens* 2021;10.
- [50] Eggers M, Eickmann M, Zorn J. Rapid and effective virucidal activity of povidone-iodine products against Middle East respiratory syndrome coronavirus (MERS-CoV) and modified vaccinia virus Ankara (MVA). *Infect Dis Ther* 2015;4:491–501.
- [51] Bidra AS, Pelletier JS, Westover JB, Frank S, Brown SM, Tessema B. Comparison of in vitro inactivation of SARS CoV-2 with hydrogen peroxide and povidone-iodine oral antiseptic rinses. *J Prosthodont* 2020;29:599–603.
- [52] Bidra AS, Pelletier JS, Westover JB, Frank S, Brown SM, Tessema B. Rapid in-vitro inactivation of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) using povidone-iodine oral antiseptic rinse. *J Prosthodont* 2020;29:529–33.
- [53] Frank S, Brown SM, Capriotti JA, Westover JB, Pelletier JS, Tessema B. In vitro efficacy of a povidone-iodine nasal antiseptic for rapid inactivation of SARS-CoV-2. *JAMA Otolaryngol Head Neck Surg* 2020;146:1–5.
- [54] Gudmundsdottir Á, Scheving R, Lindberg F, Stefansson B. Inactivation of SARS-CoV-2 and HCoV-229E in vitro by ColdZyme® a medical device mouth spray against the common cold. *J Med Virol* 2020. <https://doi.org/10.1002/jmv.26554>.
- [55] Leslie RA, Zhou SS, Macinga DR. Inactivation of SARS-CoV-2 by commercially available alcohol-based hand sanitizers. *Am J Infect Control* 2020. <https://doi.org/10.1016/j.ajic.2020.08.020>.
- [56] Liang B, Yuan X, Wei G, Wang W, Zhang M, Peng H, et al. In-vivo toxicity studies and in-vitro inactivation of SARS-CoV-2 by povidone-iodine in-situ gel forming formulations. *bioRxiv* 2020. <https://doi.org/10.1101/2020.05.18.103184>.
- [57] Meister TL, Brüggemann Y, Todt D, Conzelmann C, Müller JA, Groß R, et al. Virucidal efficacy of different oral rinses against severe acute respiratory syndrome coronavirus 2. *J Infect Dis* 2020;222:1289–92.
- [58] Mukherjee S, Vincent CK, Jayasekera HW, Yekhe AS. Antiviral efficacy of personal care formulations against severe acute respiratory syndrome coronavirus 2. *Infect Dis Health* 2020;26:63–6.
- [59] Hirose R, Bandou R, Ikegaya H, Watanabe N, Yoshida T, Daidoji T, et al. Disinfectant effectiveness against SARS-CoV-2 and influenza viruses present on human skin: model-based evaluation. *Clin Microbiol Infect* 2021. <https://doi.org/10.1016/j.cmi.2021.04.009>.
- [60] Huang YH, Huang JT. Use of chlorhexidine to eradicate oropharyngeal SARS-CoV-2 in COVID-19 patients. *J Med Virol* 2021;93:4370–3.
- [61] Steinhauer K, Meister TL, Todt D, Krawczyk A, Paßvogel L, Becker B, et al. Virucidal efficacy of different formulations for hand and surface disinfection targeting SARS CoV-2. *J Hosp Infect* 2021;112:27–30.
- [62] Frank S, Capriotti J, Brown SM, Tessema B. Povidone-iodine use in Sinonasal and oral cavities: a review of safety in the COVID-19 era. *Ear Nose Throat J* 2020;99:586–93.
- [63] He H, Dong X, Yang M, Yang Q, Duan S, Yu Y, et al. Catalytic inactivation of SARS coronavirus, *Escherichia coli* and yeast on solid surface. *Catal Commun* 2004;5:170–2.
- [64] Buonanno M, Welch D, Shuryak I, Brenner DJ. Far-UVC light (222nm) efficiently and safely inactivates airborne human coronaviruses. *Sci Rep* 2020;10:10285.
- [65] Qiao Y, Yang M, Marabella IA, McGee DAJ, Aboubakr H, Goyal S, et al. Greater than 3-log reduction in viable coronavirus aerosol concentration in ducted ultraviolet-C (UV-C) systems. *Environ Sci Technol* 2020. <https://doi.org/10.1021/acs.est.0c05763>.
- [66] Yu L, Peel GK, Cheema FH, Lawrence WS, Bukreyeva N, Jinks CW, et al. Catching and killing of airborne SARS-CoV-2 to control spread of COVID-19 by a heated air disinfection system. *Mater Today Phys* 2020;15:5.
- [67] Blanchard EL, Lawrence JD, Noble JA, Xu M, Joo T, Ng NL, et al. Enveloped virus inactivation on personal protective equipment by exposure to ozone. *medRxiv* 2020. <https://doi.org/10.1101/2020.05.23.20111435>.
- [68] Campos RK, Jin J, Rafael GH, Zhao M, Liao L, Simmons G, et al. Decontamination of SARS-CoV-2 and other RNA viruses from N95 level meltblown polypropylene fabric using heat under different humidities. *ACS Nano* 2020;14:14017–25.
- [69] Choi YW, Richardson AW, Sunderman M, Mladineo MJ, Keyes PH, Hofacre KC, et al. Decontamination of SARS-CoV-2 contaminated

- N95 filtering facepiece respirators (FFRs) with moist heat generated by a multicooker. *Lett Appl Microbiol* 2020. <https://doi.org/10.1111/lam.13443>.
- [70] Daeschler SC, Manson N, Joachim K, Chin AWH, Chan K, Chen PZ, et al. Effect of moist heat reprocessing of N95 respirators on SARS-CoV-2 inactivation and respirator function. *CMAJ* 2020;192:E1189–97.
- [71] Gopal V, Nilsson-Payant BE, French H, Siegers JY, Yung WS, Hardwick M, et al. Zinc-embedded fabrics inactivate SARS-CoV-2 and influenza A virus. *bioRxiv* 2020. <https://doi.org/10.1101/2020.11.02.365833>.
- [72] Ibanez-Cervantes G, Bravata-Alcantara JC, Najera-Cortes AS, Meneses-Cruz S, Delgado-Balbuena L, Cruz-Cruz C, et al. Disinfection of N95 masks artificially contaminated with SARS-CoV-2 and ESKAPE bacteria using hydrogen peroxide plasma: impact on the reutilization of disposable devices. *Am J Infect Control* 2020;48:1037–41.
- [73] Ludwig-Begall LF, Wielick C, Dams L, Nauwynck H, Demeuldre PF, Napp A, et al. The use of germicidal ultraviolet light, vaporized hydrogen peroxide and dry heat to decontaminate face masks and filtering respirators contaminated with a SARS-CoV-2 surrogate virus. *J Hosp Infect* 2020;106:577–84.
- [74] Ma QX, Shan H, Zhang CM, Zhang HL, Li GM, Yang RM, et al. Decontamination of face masks with steam for mask reuse in fighting the pandemic COVID-19: experimental supports. *J Med Virol* 2020;92:1971–4.
- [75] Mantlo E, Rhodes T, Boutros J, Patterson-Fortin L, Evans A, Paessler S. In vitro efficacy of a copper iodine complex PPE disinfectant for SARS-CoV-2 inactivation. *F1000Res* 2020;9:674.
- [76] Ozog DM, Sexton JZ, Narla S, Pretto-Kernahan CD, Mirabelli C, Lim HW, et al. The effect of ultraviolet C radiation against different N95 respirators inoculated with SARS-CoV-2. *Int J Infect Dis* 2020;100:224–9.
- [77] Perkins DJ, Nofchissey RA, Ye C, Donart N, Kell A, Foo-Hurwitz I, et al. COVID-19 global pandemic planning: dry heat incubation and ambient temperature fail to consistently inactivate SARS-CoV-2 on N95 respirators. *Exp Biol Med* 2020. <https://doi.org/10.1177/1535370220977819>.
- [78] Rathnasinghe R, Karlicek RF, Schotsaert M, Koffas MA, Arduini B, Jangra S, et al. Scalable, effective, and rapid decontamination of SARS-CoV-2 contaminated N95 respirators using germicidal ultraviolet C (UVC) irradiation device. *medRxiv* 2020. <https://doi.org/10.1101/2020.10.05.20206953>.
- [79] Rockey N, Arts PJ, Li L, Harrison KR, Langenfeld K, Fitzsimmons WJ, et al. Humidity and deposition solution play a critical role in virus inactivation by heat treatment of N95 respirators. *mSphere* 2020;5.
- [80] Glasbrenner DC, Choi YW, Richardson AW, Edwards EW, Mladineo MJ, Sunderman M, et al. Decontamination of SARS-CoV-2 contaminated N95 filtering facepiece respirators using artificial sun lamps. *J Appl Microbiol* 2021. <https://doi.org/10.1111/jam.15106>.
- [81] Valdez-Salas B, Beltran-Partida E, Cheng N, Salvador-Carlos J, Valdez-Salas EA, Curiel-Alvarez M, et al. Promotion of surgical masks antimicrobial activity by disinfection and impregnation with disinfectant silver nanoparticles. *Int J Nanomed* 2021;16:2689–702.
- [82] Rojas-Valencia MN. Research on ozone application as disinfectant and action mechanisms on wastewater microorganisms. In: Méndez-Vilas A, editor. *Science against microbial pathogens: communicating current research and technological advances*. Mexico: National Autonomous University of Mexico, Institute of Engineering, Coordination of Environmental Engineering; 2011.
- [83] Nuvolone D, Petri D, Voller F. The effects of ozone on human health. *Environ Sci Pollut Res Int* 2018;25:8074–88.
- [84] Suchomel M, Steinmann J, Kampf G. Efficacies of the original and modified World Health Organization-recommended hand-rub formulations. *J Hosp Infect* 2020;106:264–70.
- [85] Rabenau HF, Cinatl J, Morgenstern B, Bauer G, Preiser W, Doerr HW. Stability and inactivation of SARS coronavirus. *Med Microbiol Immunol* 2005;194:1–6.
- [86] Binda G, Bellasi A, Spanu D, Pozzi A, Cavallo D, Bettinetti R. Evaluating the environmental impacts of personal protective equipment use by the general population during the COVID-19 pandemic: a case study of Lombardy (Northern Italy). *Environments* 2021;8:33.
- [87] Rizan C, Reed M, Bhutta MF. Environmental impact of personal protective equipment distributed for use by health and social care services in England in the first six months of the COVID-19 pandemic. *J R Soc Med* 2021;114:250–63.
- [88] Zhang EJ, Aitchison LP, Phillips N, Shaban RZ, Kam AW. Protecting the environment from plastic PPE. *BMJ* 2021;372:n109.
- [89] Eggers M, Schwebke I, Suchomel M, Fotheringham V, Gebel J, Meyer B, et al. The European tiered approach for virucidal efficacy testing – rationale for rapidly selecting disinfectants against emerging and re-emerging viral diseases. *Euro Surveill* 2021;26.