

Clinical practice guideline for breast-conserving surgery in patients with early-stage breast cancer: Chinese Society of Breast Surgery (CSBrS) practice guidelines 2021

Kai Chen^{1,2}, Jie-Qiong Liu^{1,2}, Wei Wu^{1,2}, Feng-Xi Su^{1,2}, Qiang Zou³, Er-Wei Song^{1,2}; Chinese Society of Breast Surgery

¹Guangdong Provincial Key Laboratory of Malignant Tumor Epigenetics and Gene Regulation, Sun Yat-Sen Memorial Hospital, Sun Yat-Sen University, Guangzhou, Guangdong 510120, China;

²Department of Breast Surgery, Yat-sen Breast Tumor Hospital, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, Guangdong 510120, China;

³Department of Surgery, Huashan Hospital, Fudan University, Shanghai 200040, China.

Breast-conserving surgery (BCS) with radiotherapy is the primarily agreed surgical choice for eligible patients with an early diagnosis of breast cancer. In 2019, the panel of Chinese experts assembled by the Chinese Society for Breast Surgery (CSBrS) developed a Chinese experts' consensus on BCS for early-stage breast cancer (Version 2019) with the aim of standardizing BCS in China. Subsequently, the CSBrS conducted a review of published reports and discussions between experts to determine the key clinical questions for the Clinical Practice Guideline for BCS in Patients with Early-Stage Breast Cancer. The group evaluated the relevant evidence using the grading of recommendations assessment, development, and evaluation system, and developed a Clinical Practice Guideline for BCS in Patients with Early-Stage Breast Cancer (Version 2021), with the aim of providing guidance on clinical practice to breast surgeons in China.

Level of Evidence and Recommendation Strength

Level of evidence standard^[1]

Recommendation strength standard^[1]

Recommendation strength review committee

The panel for this guideline is comprised of 85 experts, including 70 (82.3%) breast surgeons, four (4.7%) medical oncologists, four (4.7%) diagnostic radiologists, two (2.4%) pathologists, one (1.2%) obstetrician, two (2.4%) radiation oncologists, and two (2.4%) epidemiologists.

Target Audience

Clinicians specializing in breast diseases in China.

Access this article online	
Quick Response Code: 	Website: www.cmj.org
	DOI: 10.1097/CM9.0000000000001518

Recommendations

Recommendation 1: Indications (all of the indications should be met).

No.	Indications	Level of evidence	Recommended strength
1.1	Patient wishes to preserve her breast	I ^[2,3]	A
1.2	Clinical Stage I, II, or ≤cT2	I ^[2,5]	A
1.3	Able to achieve acceptable cosmetic outcomes after BCS	I ^[2,3]	A

Recommendation 2: Contraindications (any one of the indications is sufficient).

No.	Contraindications	Level of evidence	Recommended strength
2.1	Cannot receive radiotherapy after BCS	I ^[4,6]	A
2.2	Unable to achieve negative surgical margins	I ^[2,3,7-9]	A
2.3	Extensive microcalcification	I ^[2,3,8]	A
2.4	Inflammatory breast cancer	I ^[2,3]	A
2.5	Patient refusal to undergo BCS	I ^[2,3]	A

Recommendation 3: Surgery.

No.	Surgeries	Level of evidence	Recommended strength
3.1	Incorporation of oncoplastic techniques is able to improve the cosmetic outcomes after BCS	II ^[9-11]	A
3.2	It is recommended that inert metal clips (eg, titanium clips) be placed in the surgical bed after BCS as a localization marker for radiation boosting	I ^[2,3]	A

Correspondence to: Dr. Er-Wei Song, Department of Breast Surgery, Yat-sen Breast Tumor Hospital, Sun Yat-sen Memorial Hospital, Sun Yat-sen University, Guangzhou, Guangdong 510120, China
E-Mail: songew@mail.sysu.edu.cn
Dr. Qiang Zou, Department of Surgery, Huashan Hospital, Fudan University, Shanghai 200040, China
E-Mail: zouqiang003@aliyun.com

Copyright © 2021 The Chinese Medical Association, produced by Wolters Kluwer, Inc. under the CC-BY-NC-ND license. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Chinese Medical Journal 2021;134(18)

Received: 14-01-2021 Edited by: Yan-Jie Yin and Xiu-Yuan Hao

Recommendation 4: Pathology.

No.	Pathologies	Level of evidence	Recommended strength
4.1	Margin assessment after BCS is mandatory	I ^[2,3,8]	A
4.2	Intraoperative frozen section analysis for margin assessment is recommended	I ^[2,3,12]	A
4.3	Post-operative formalin-fixed, paraffin-embedded tissue analysis is recommended for margin assessment	I ^[2,3,8]	A
4.4	Methods for margin assessment: Lumpectomy margin assessment (perpendicular inked method)	I ^[2,4,13]	A
	Lumpectomy margin assessment (tangential shaved method)	II ^[2,14]	A
	Cavity wall (tumor bed) sampling	II ^[3,15-21]	A

Recommendation 5: Radiotherapy.

No.	Radiotherapy	Level of evidence	Recommended strength
5.1	Whole-breast irradiation is recommended after BCS*	I ^[2,4,6]	A

* Whole-breast irradiation can be waived in patients aged >65 years with Stage I breast cancer, hormone receptor-positive tumors, and negative surgical margins (CALGB9343 trial).

Discussion

A considerable amount of evidence from the scientific literature supports the safety and efficacy of combining BCS with radiotherapy in the early stages of breast cancer. In the National Surgical Adjuvant Breast and Bowel Project B-06 trial, a total of 1851 patients diagnosed with Stage I or II breast cancer were randomly allocated to three study arms, namely total mastectomy, BCS, and BCS with radiotherapy. According to results of the long-term follow-up that lasted 20 years, it was found that disease-free survival, metastasis-free survival, and overall survival did not differ significantly between the three arms.^[4] During the same period, in the Milan I trial, seven hundred and one patients with <T2 breast cancers (tumor size measuring <2 cm) were randomized to two arms: radical mastectomy and BCS with radiotherapy. The 20-year follow-up results indicated that there was no significant difference between the overall survival of the two arms. However, the cumulative recurrence rate was higher in the BCS + radiation arm (8.8%) when compared with the mastectomy arm (2.3%).^[5] A short-term follow-up study of 95 Chinese patients by Li *et al*^[22] revealed that, after an average follow-up of 17 months, the 2-year local recurrence rate of Stage I or II breast cancer after BCS was only 1.4%, with no metastases or deaths. Chen *et al*^[16] used propensity-score matching to compare the clinical outcomes after a median follow-up of 67 months of 2866 patients with early-stage breast cancer who had undergone BCS or mastectomy in China; they confirmed the safety and efficacy of BCS. Recently, BCS has been recommended internationally as the standard surgical treatment for eligible breast cancer patients in the early stages.^[2,3,8] The

panel members agreed that, provided acceptable cosmetic outcomes could be expected, BCS is suitable for patients with clinical Stage I or II disease or ≤T2 tumors who wish to preserve their breasts.

The result of Early Breast Cancer Trialists' Collaborative Group meta-analysis revealed that neoadjuvant chemotherapy can significantly increase the BCS rate. The 10-year cumulative local recurrence rate was slightly, but not significantly, higher in the neoadjuvant chemotherapy than in the untreated group (15.1% *vs.* 11.9%, *P* = 0.10). No significant differences were observed in the 10-year cumulative rates of breast cancer-related deaths between patients who did and did not receive neoadjuvant chemotherapy (27.5% *vs.* 24.8%, *P* = 0.15).^[23] Therefore, for patients who were clinically diagnosed with Stage III breast cancer or >T2 tumors, it is possible to administer neoadjuvant chemotherapy to downstage the tumor, and thus increase the chances of being eligible for BCS. The panel members considered that, in clinical practice, it is not always easy to accurately measure the degree of shrinkage of a tumor after neoadjuvant chemotherapy. Thus, there was a lack of consensus in the panel regarding the optimal extent of surgery in patients with breast cancer who have received neoadjuvant chemotherapy. However, there was agreement that achieving negative surgical margins is mandatory in this situation.

The panel considered that the following factors are potential risk factors for local recurrence after BCS: breast cancer located in the central portion; bloody discharge from the nipple; large tumor (eg, >T2); multifocal breast cancer; multicentric breast cancer; young age (<35 years); and radiotherapy contraindicated (eg, active connective tissue disease). However, there is no high level evidence that the abovementioned factors are contraindications to BCS.

Achieving negative surgical margins is mandatory for successful BCS. There is evidence that positive surgical margins are closely associated with local recurrence.^[2,8,9,24] Intra-operative gross inspection,^[25] imprint cytology,^[26] intra-operative specimen imaging,^[27] and novel devices^[28,29] can also reportedly reduce the rates of positive margins. However, the panel does not recommend these approaches because of the lack of high-level evidence. Intra-operative frozen section analysis (FSA) is reportedly capable of reducing the rates of margin positivity and second surgeries.^[30] The panel considers that FSA is widely used in clinical practice in China and supports its use for intra-operative margin assessment. The panel take a cautious attitude toward performing margin assessment only by post-operative formalin-fixed, paraffin-embedded examination.

Both lumpectomy margins and cavity margins are suitable for the margin assessment of BCS. Lumpectomy margins are assessed on the surface of the tumor-containing specimen. There are two techniques that can be used for margin assessment: the perpendicular inked and tangential shaved techniques.^[2,14] Cavity margins are assessed by performing a biopsy (tissue sampling) of the residual cavity (or the wall of the tumor bed) after tumor removal.

Well-designed studies suggesting that cavity margin assessment alone is capable of achieving excellent local control have been published.^[15,19]

A meta-analysis has demonstrated that the “no-ink on tumor” is significantly associated with a reduced local recurrence rate and that wider surgical margins do not further improve the local control rate.^[31] Furthermore, the increased risk of local recurrence associated with positive surgical margins is not nullified by post-operative radiotherapy. There is no evidence that different margin widths should be considered for patients of different ages or with different molecular subtypes. A real-world study (CSBrS-005) conducted by the CSBrS in 2019 revealed that 88.2% (1530/1734) of patients who had undergone BCS had margin widths >5 mm.^[32] Others have reached consensus on diagnosing negative surgical margins for infiltrating ductal carcinoma and ductal carcinoma *in situ* by “no-ink on tumor” and “≥2 mm,” respectively.^[2,3,7] The panel considered these standards inappropriate for China and has not recommended them for routine use.

The panel agreed that whole-breast irradiation (RT) is necessary after BCS. However, the CALGB9493 trial showed a small improvement of locoregional recurrence rate in BCS patients who received RT (RT: 2% (95% confidence interval [CI]: 1% to 4%) vs. no-RT: 10% (95% CI: 7% to 15%)), but no statistically significant differences in 10-year distant metastasis and overall survival between patients with low-risk BCS who did and did not undergo RT.^[33] In contrast, the PRIME II trial showed that, in patients aged ≥65 years with early-stage and hormone receptor-positive disease who underwent BCS, the 5-year ipsilateral recurrence rates were 4.1% and 1.3% ($P = 0.0002$) in the no-RT and RT groups, respectively.^[34] Although this difference is statistically significant, the benefit is not clinically important. The panel suggested that RT might be forgone after BCS in certain situations related to the patients' preferences and comorbidities.

The International Breast Cancer Study Group VI-VII trial^[35] aimed to analyze how the timing of RT affects the local failure rate and disease-free survival (DFS) in breast cancer patients. Among pre/perimenopausal patients, the 15-year DFSs were 48.2% vs. 44.9% (hazard ratio [HR] = 1.12, 95% CI: 0.87–1.45) in patients allocated 3 and 6 months of initial chemotherapy (CT). Among post-menopausal patients, the 15-year DFS was 46.1% and 43.3% in the group that did not receive CT initially and the group allocated 3 months of CT, respectively (HR 1.11, 95% CI: 0.82–1.51). The results of this clinical trial suggest that delaying RT until after completion of CT is safe and reasonable. The panel recommended that whole-breast RT is indicated for eligible breast cancer patients after BCS and CT.

The CO-HO-RT trial^[36] revealed that the risk of developing grade ≥2 radiation-induced subcutaneous fibrosis is similar in patients who receive concurrent vs. sequential RT and endocrine therapy, suggesting that concurrent use of these treatment modalities is safe. According to the 3.7-year follow-up data of the N9831

trial,^[37] concurrent use of RT and trastuzumab did not significantly increase cardiotoxicity, supporting the feasibility of the concurrent use of RT and anti-human epidermal growth factor receptor 2 (HER2) treatments. The panel recommended the concurrent use of RT and endocrine therapy, as well as anti-HER2 therapy if indicated.

Acknowledgements

The authors thank Dr. Trish Reynolds, MBBS, FRACP, from Liwen Bianji, Edanz Group China (www.liwenbianji.cn/ac), for editing the English draft of this manuscript. The authors thank Jia-Fan Ma for revision of this manuscript.

List of compiling committee members (In alphabetical order by surname): Zhong-Wei Cao, De-Dian Chen, Kai Chen, Yuan-Jia Cheng, Xue-Ning Duan, Zhi-Min Fan, Pei-Fen Fu, Jian Huang, Jun Jiang, Hong-Chuan Jiang, Feng Jin, Hua Kang, Rui Ling, Jin-Ping Liu, Jie-Qiong Liu, Ke Liu, Qian Liu, Yin-Hua Liu, Yun-Jiang Liu, Zhen-Zhen Liu, Yong-Hui Luo, Rong Ma, Da-Hua Mao, Jiang-Hua Ou, Xiang Qu, Guo-Sheng Ren, Ai-Lin Song, Er-Wei Song, Feng-Xi Su, Li-Li Tang, Xing-Song Tian, Chuan Wang, Fei Wang, Jian-Dong Wang, Shu Wang, Shui Wang, Xiang Wang, Jiong Wu, Ke-Jin Wu, Wei Wu, Fei Xie, Ling Xin, Zhi-Gang Yu, Jian-Guo Zhang, Jin Zhang, Jing-Hua Zhang, Yi Zhao, Zuo-Wei Zhao, Ang Zheng, Wei Zhu, Qiang Zou

Conflicts of interest

The expert committee for these guidelines declares no conflict of interest.

These guidelines are a reference for breast disease specialists in clinical practice. However, the guidelines are not to be used as the basis for medical evaluation, and do not play an arbitrating role in the handling of any medical disputes. The guidelines are not a reference for patients or non-breast specialists. The Chinese Society of Breast Surgery assumes no responsibility for results involving the inappropriate application of these guidelines, and reserves the right to interpret and revise the guidelines.

References

- Ye JM, Guo BL, Liu Q, Ma F, Liu HJ, Wu Q, *et al.* Clinical practice guidelines for sentinel lymph node biopsy in patients with early-stage breast cancer: Chinese Society of Breast Surgery (CSBrS) Practice Guidelines 2021. *Chin Med J* 2021;134:886–894. doi: 10.1097/CM9.0000000000001410.
- Committee of Breast Cancer Society of Chinese Anti-Cancer Association. Guidelines for clinical diagnosis and treatment of breast cancer: Chinese Anti-Cancer Association guidelines (2019 Edition) (in Chinese). *Chin Oncol* 2019;29:609–679. doi: 10.19401/j.cnki.1007-3639.2019.08.009.
- Chinese Association of Breast Surgery. A consensus statement on the breast-conserving surgery of early-stage breast cancer (2019). *Chin J Surg* 2019;57:81–84. doi: 10.3760/cma.j.issn.0529-5815.2019.02.001.
- Fisher B, Anderson S, Bryant J, Margolese RG, Deutsch M, Fisher ER, *et al.* Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med* 2002;347:1233–1241. doi: 10.1056/NEJMoa022152.

5. Veronesi U, Cascinelli N, Mariani L, Greco M, Saccozzi R, Luini A, *et al.* Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med* 2002;347:1227–1232. doi: 10.1056/NEJMoa020989.
6. Clarke M, Collins R, Darby S, Davies C, Elphinstone P, Evans V, *et al.* Effects of radiotherapy and of differences in the extent of surgery for early breast cancer on local recurrence and 15-year survival: an overview of the randomised trials. *Lancet* 2005;366:2087–2106. doi: 10.1016/S0140-6736(05)67887-7.
7. Moran MS, Schnitt SJ, Giuliano AE, Harris JR, Khan SA, Horton J, *et al.* Society of Surgical Oncology-American Society for Radiation Oncology consensus guideline on margins for breast-conserving surgery with whole-breast irradiation in stages I and II invasive breast cancer. *J Clin Oncol* 2014;32:1507–1515. doi: 10.1200/jco.2013.53.3935.
8. Gradishar WJ, Anderson BO, Abraham J, Aft R, Agnese D, Allison KH, *et al.* Breast cancer, version 3.2020, NCCN clinical practice guidelines in oncology. *J Natl Compr Canc Netw* 2020;18:452–478. doi: 10.6004/jnccn.2020.0016.
9. Cardoso F, Kyriakides S, Ohno S, Penault-Llorca F, Poortmans P, Rubio IT, *et al.* Early breast cancer: ESMO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2019;30:1194–1220. doi: 10.1093/annonc/mdz173.
10. Association of Breast Surgery at Baso 2009. Surgical guidelines for the management of breast cancer. *Eur J Surg Oncol* 2009;35 (Suppl 1): 1–22. doi: 10.1016/j.ejso.2009.01.008.
11. Committee of Breast Cancer Society of Chinese Anti-Cancer Association; Committee of Chinese Society of Breast Surgeon. Consensus of oncoplastic breast surgery and breast reconstruction. *Chin Oncol* 2018;28:439–480. doi: 10.19401/j.cnki.1007-3639.2018.06.008.
12. (ASBrS) TAsoBS. Official Statement: Toolbox to Reduce Lumpectomy Reoperations and Improve Cosmetic Outcome in Breast Cancer Patients: The American Society of Breast Surgeons Consensus Conference; 2015. Available from: <https://www.breastsurgeons.org/resources/statements>. Accessed February 3, 2021.
13. Fisher ER, Sass R, Fisher B, Gregorio R, Brown R, Wickerham L. Pathologic findings from the National Surgical Adjuvant Breast Project (protocol 6). II. Relation of local breast recurrence to multicentricity. *Cancer* 1986;57:1717–1724. doi: 10.1002/1097-0142(19860501)57:9<1717::aid-cnrc2820570902>3.0.co;2-h.
14. Wright MJ, Park J, Fey JV, Park A, O'Neill A, Tan LK, *et al.* Perpendicular inked versus tangential shaved margins in breast-conserving surgery: does the method matter? *J Am Coll Surg* 2007;204:541–549. doi: 10.1016/j.jamcollsurg.2007.01.031.
15. Chen K, Zeng Y, Jia H, Jia W, Yang H, Rao N, *et al.* Clinical outcomes of breast-conserving surgery in patients using a modified method for cavity margin assessment. *Ann Surg Oncol* 2012;19:3386–3394. doi: 10.1245/s10434-012-2331-5.
16. Chen K, Pan Z, Zhu L, Hu T, Peng M, Jia W, *et al.* Comparison of breast-conserving surgery and mastectomy in early breast cancer using observational data revisited: a propensity score-matched analysis. *Sci China Life Sci* 2018;61:1528–1536. doi: 10.1007/s11427-018-9396-x.
17. Chen K, Zhu L, Chen L, Li Q, Li S, Qiu N, *et al.* Circumferential shaving of the cavity in breast-conserving surgery: a randomized controlled trial. *Ann Surg Oncol* 2019;26:4256–4263. doi: 10.1245/s10434-019-07725-w.
18. Hequet D, Bricou A, Koual M, Ziou M, Feron JG, Rouzier R, *et al.* Systematic cavity shaving: modifications of breast cancer management and long-term local recurrence, a multicentre study. *Eur J Surg Oncol* 2013;39:899–905. doi: 10.1016/j.ejso.2013.05.012.
19. Hewes JC, Imkampe A, Haji A, Bates T. Importance of routine cavity sampling in breast conservation surgery. *Br J Surg* 2009;96:47–53. doi: 10.1002/bjs.6435.
20. Barthelmes L, Al Awa A, Crawford DJ. Effect of cavity margin shavings to ensure completeness of excision on local recurrence rates following breast conserving surgery. *Eur J Surg Oncol* 2003;29:644–648. doi: 10.1016/s0748-7983(03)00122-7.
21. Malik HZ, George WD, Mallon EA, Harnett AN, Macmillan RD, Purushotham AD. Margin assessment by cavity shaving after breast-conserving surgery: analysis and follow-up of 543 patients. *Eur J Surg Oncol* 1999;25:464–469. doi: 10.1053/ejso.1999.0680.
22. Li JF, Ou YT, Wang TF, Lin BY. The early results of breast-conserving therapy in ninety-five patients of primary breast cancer. *Chin J Surg* 2004;42:282–284. doi: 10.3760/j.issn.0529-5815.2004.05.008.
23. Early Breast Cancer Trialists' Collaborative Group. Long-term outcomes for neoadjuvant versus adjuvant chemotherapy in early breast cancer: meta-analysis of individual patient data from ten randomised trials. *Lancet Oncol* 2018;19:27–39. doi: 10.1016/S1470-2045(17)30777-5.
24. (ASBrS) TAsoBS. Official Statement: Performance and Practice Guidelines for Breast-Conserving Surgery/Partial Mastectomy; 2015. Available from: <https://www.breastsurgeons.org/resources/statements>. Accessed February 3, 2021.
25. Nunez A, Jones V, Schulz-Costello K, Schmolze D. Accuracy of gross intraoperative margin assessment for breast cancer: experience since the SSO-ASTRO margin consensus guidelines. *Sci Rep* 2020;10: 17344. doi: 10.1038/s41598-020-74373-6.
26. Tamanuki T, Namura M, Aoyagi T, Shimizu S, Suwa T, Matsuzaki H. Effect of intraoperative imprint cytology followed by frozen section on margin assessment in breast-conserving surgery. *Ann Surg Oncol* 2020;28:1338–1346. doi: 10.1245/s10434-020-08955-z.
27. Mariscotti G, Durando M, Pavan LJ, Tagliafico A, Campanino PP, Castellano I, *et al.* Intraoperative breast specimen assessment in breast conserving surgery: comparison between standard mammography imaging and a remote radiological system. *Br J Radiol* 2020;93:20190785. doi: 10.1259/bjr.20190785.
28. Zysk AM, Chen K, Gabrielson E, Taft L, May Gonzalez EA, Canner JK, *et al.* Intraoperative assessment of final margins with a handheld optical imaging probe during breast-conserving surgery may reduce the reoperation rate: results of a multicenter study. *Ann Surg Oncol* 2015;22:3356–3362. doi: 10.1245/s10434-015-4665-2.
29. Allweis TM, Kaufman Z, Lelcuk S, Pappo I, Karni T, Schneebaum S, *et al.* A prospective, randomized, controlled, multicenter study of a real-time, intraoperative probe for positive margin detection in breast-conserving surgery. *Am J Sur* 2008;196:483–489. doi: 10.1016/j.amjsurg.2008.06.024.
30. Esbona K, Li Z, Wilke LG. Intraoperative imprint cytology and frozen section pathology for margin assessment in breast conservation surgery: a systematic review. *Ann Surg Oncol* 2012;19:3236–3245. doi: 10.1245/s10434-012-2492-2.
31. Houssami N, Macaskill P, Marinovich ML, Morrow M. The association of surgical margins and local recurrence in women with early-stage invasive breast cancer treated with breast-conserving therapy: a meta-analysis. *Ann Surg Oncol* 2014;21:717–730. doi: 10.1245/s10434-014-3480-5.
32. Yu LX, Shi P, Tian XS, Yu ZG. Chinese Society of Breast Surgery. A multicenter investigation of breast-conserving surgery based on data from the Chinese Society of Breast Surgery (CSBrS-005). *Chin Med J (Engl)* 2020;133:2660–2664. doi: 10.1097/CM9.0000000000001152.
33. Hughes KS, Schnaper LA, Bellon JR, Cirincione CT, Berry DA, McCormick B, *et al.* Lumpectomy plus tamoxifen with or without irradiation in women age 70 years or older with early breast cancer: long-term follow-up of CALGB 9343. *J Clin Oncol* 2013;31:2382–2387. doi: 10.1200/JCO.2012.45.2615.
34. Kunkler IH, Williams LJ, Jack WJ, Cameron DA, Dixon JM. PRIME II investigators. Breast-conserving surgery with or without irradiation in women aged 65 years or older with early breast cancer (PRIME II): a randomised controlled trial. *Lancet Oncol* 2015;16:266–273. doi: 10.1016/S1470-2045(14)71221-5.
35. Karlsson P, Cole BF, Price KN, Gelber RD, Coates AS, Goldhirsch A, *et al.* Timing of radiation therapy and chemotherapy after breast-conserving surgery for node-positive breast cancer: long-term results from International Breast Cancer Study Group Trials VI and VII. *Int J Radiat Oncol Biol Phys* 2016;96:273–279. doi: 10.1016/j.ijrobp.2016.06.2448.
36. Bourcier C, Kerns S, Gourgou S, Lemanski C, Gutowski M, Fenoglio P, *et al.* Concurrent or sequential letrozole with adjuvant breast radiotherapy: final results of the CO-HO-RT phase II randomized trial. *Ann Oncol* 2016;27:474–480. doi: 10.1093/annonc/mdv602.
37. Halyard MY, Pisansky TM, Dueck AC, Suman V, Pierce L, Solin L, *et al.* Radiotherapy and adjuvant trastuzumab in operable breast cancer: tolerability and adverse event data from the NCCTG Phase III Trial N9831. *J Clin Oncol* 2009;27:2638–2644. doi: 10.1200/JCO.2008.17.9549.

How to cite this article: Chen K, Liu JQ, Wu W, Su FX, Zou Q, Song EW, Chinese Society of Breast Surgery. Clinical practice guideline for breast-conserving surgery in patients with early-stage breast cancer: Chinese Society of Breast Surgery (CSBrS) practice guidelines 2021. *Chin Med J* 2021;134:2143–2146. doi: 10.1097/CM9.0000000000001518