

Drosophila eugracilis – Akt

Ashley Morgan¹, Cole A. Kiser¹, Isabel Acosta-Coley², Hanjun Lin², Nicholas Guillette³, Robert McMahon³, Jennifer A. Kennell², Lindsey J Long³, Laura K. Reed¹, Chinmay P. Rele^{1§}

¹The University of Alabama, Tuscaloosa, AL USA

²Vassar College, Poughkeepsie, NY USA

³Oklahoma Christian University, Edmond, OK USA

[§]To whom correspondence should be addressed: cprele@ua.edu

Abstract

Gene Model for *Akt* in the *D. eugracilis* (DeugGB2) assembly (GCA_000236325.2).



Figure 1.

7/2/2022 - Open Access

(A) Synteny of genomic neighborhood of Akt in both D. melanoqaster as well as D. eugracilis. Gene arrows pointing in the same direction as Akt in both D. eugracilis and D. melanogaster are on the same strand as Akt; gene arrows pointing in the opposite direction are on the opposite strand. The thin underlying arrow in D. melanogaster that points to the right indicates Akt is on the + strand and the thin underlying arrow pointing to the left in D. eugracilis indicates Akt is on the – strand. White arrows in *D. eugracilis* indicate orthology to the corresponding gene in *D. melanogaster*. CG32855 and sxe2 are nested within an intron of *Mhc1* in both species. The gene names given in the *D. eugracilis* gene arrows indicate the orthologous gene in *D*. melanogaster, while the locus identifiers are specific to *D. eugracilis*; (B) Gene Model in UCSC Track Hub (Raney et al., 2014): The gene model in D. eugracilis (black), Spaln of D. melanogaster Proteins (purple, alignment of refseq proteins from D. melanogaster), BLAT alignments of NCBI RefSeq Genes (blue, alignment of refseq genes for D. eugracilis), RNA-Seq from adult females (red), adult males (blue), and mixed embryos (purple) (alignment of Illumina RNAseq reads from D. eugracilis), and Transcripts (green) including coding regions predicted by TransDecoder and Splice Junctions Predicted by regtools using D. eugracilis RNA-Seq (Chen et al., 2014; PRJNA63469). Splice junctions shown have a minimum read-depth of 10 with 10-49, 50-99, 100-499, 500-999, >1000 supporting reads in blue, green, pink, brown, and red respectively. The custom gene model (User Supplied Track) is indicated in black with exons depicted with boxes and introns with narrow lines (arrows indicate direction of transcription). (C) Dot Plot of amino acid identity for Akt-PC in D. melanogaster (x-axis) vs. Akt-PC in D. eugracilis (y-axis). Amino acid number is indicated along the left and bottom; exon number is indicated along the top and right. Each colored rectangle represents an exon. Region 1 indicates a lack of sequence similarity between the two sequences; (D) ROAST Alignments Conservation track within the UCSC Genome Browser. The ROAST Alignments for 28 Drosophila Species track within the UCSC Genome Browser shows the genomic region surrounding the beginning of the Akt-PC isoform in all 28 species shown, demonstrating that the non-canonical start codon for Akt-PC is conserved across the genus.

Description

Introduction

The insulin signaling pathway is a highly conserved pathway in animals, and is central to nutrient uptake (Hietakangas and Cohen 2009, Grewal 2009). Akt kinase (*Akt* also known as *Akt1, Protein Kinase B, PKB*; FBgn0010379) regulates stress response, aging, and cell growth and survival in *Drosophila* (Stavely *et al.*, 1998; Verdu *et al.*, 1999). It is involved in signal transduction pathways in physiological and neurological pathways in *Drosophila* (Guo and Zhong 2006). It encodes a core serine-threonine kinase (Bellacosa et al. 1991) component of the Insulin-like growth factor pathway that functions downstream of, and following its activation by the *Pi3K92E* product in *Drosophila* (Andjelkovic et al., 1995). It is activated by phosphatidylinositol binding and phosphorylation (Potter *et al.*, 2002). The gene model reported (*Deug_Akt*) was determined in the Apr. 2013 (BCM-HGSC/Deug_2.0; GCA_000236325.2) of *D. eugracilis* and compared to the ortholog *dmel_Akt* (GCA_000001215.4, FB2021_02; Larkin *et al.*, 2021). *D. eugracilis* is part of the *melanogaster* species group within the subgenus *Sophophora* of *Drosophila* (Pélandakis et al., 1993). It was first described as *Tanygastrella gracilis* by Duda (1924) and revised to *Drosophila eugracilis* by Bock and Wheeler (1972). *D. eugracilis* is found in humid tropical and subtropical forests across southeast Asia (https://www.taxodros.uzh.ch). The methods and dataset versions used to establish the gene model are described in Rele *et al.* (2020). The Genomics Education Partnership maintains a mirror of the UCSC Genome Browser (Kent WJ *et al.*, 2002; Gonzalez *et al.*, 2021), which is available at http://gander.wustl.edu.The predicted gene model in *D. eugracilis* for *Akt* was found in NCBI RefSeq Accession XM_017228041.1 and Locus ID LOC108116251.

<u>Synteny</u>

Akt is located on chromosome 3R (Muller element D) in D. melanogaster and is surrounded by sxe2, CG32855, and Mhcl (upstream) and *Sb* and *CG5903* (downstream). After performing a *tblastn*, the putative *Akt* ortholog (LOC10811625/XM_017228041.1/XP_017083530.1, e-value of 0.0 and percent identity of 90.39%) in D. eugracilis was found to be on scaffold KB465333 (mapped to Muller element D) and is surrounded by orthologs to sxe2 (LOC108116355/XM_017228178.2/XP_017083667.2, e-value of 0.0 and percent identity of 87.08%), Mhcl (LOC108116352/XM 017228167.2/XP 017083656.2, e-value of 0.0 and percent identity of 97.13%), Sb (LOC108116404/XM_017228281.2/XP_017083770.2, e-value of 0.0 and percent identity of 87.02%), and CG5903 (Mic26-27/LOC108116407/XM_017228287.1/XP_017083776.1, e-value of 7e-151 and percent identity of 88.26%) as determined by *blastp* (Figure 1A, Altschul *et al.*, 1990). The annotated model is likely to be the true ortholog of *Akt* due to the high level of synteny that exists between *D. melanogaster* and *D. eugracilis*, and the reciprocal best blast hits between the two genes.

Protein Model

There are five RNA isoforms of *Akt: Akt-RA, Akt-RB, Akt-RC, Akt-RD,* and *Akt-RE*. The RA, RB, RD, and RE isoforms have identical protein coding sequences, represented by the Akt-PE protein isoform here. The Akt-PC coding sequence is unique.

7/2/2022 - Open Access

Both the Akt-PC and Akt-PE isoforms in *D. melanogaster* are encoded by six coding exons. The Akt-PC and Akt-PE isoforms in *D. eugracilis* are encoded by six coding exons (Figure 1B). The coordinates of the curated gene models can be found in NCBI at GenBank/BankIt using the accessions BK059589, BK059590, BK059591, BK059592, and BK059593, one for each protein-coding isoform of Akt. These data are also available in Extended Data files below, which are archived in CaltechData.

Special characteristics of the gene model

Non-canonical Start Codon: The *Akt-RC* isoform has a non-canonical ACG start codon. This is well conserved across the 28 *Drosophila* species as shown in Figure 1D. This non-canonical start codon is used for the translation of the Akt-PC isoform in *D. melanogaster*, (Figure 1D). There is a high level of conservation of the non-canonical ACG start codon (encoding threonine) across all 28 *Drosophila* species, as well as high conservation of the region surrounding the non-canonical start codon. This provides evidence for the existence of a non-canonical start codon in the Akt-PC isoform in *D. eugracilis*.

Methods

Detailed methods including algorithms, database versions, and citations for the complete annotation process can be found in Rele *et al.* (2020).

Reagents

NA

Acknowledgements: We would like to thank Wilson Leung, who created and maintain the GEP technological infrastructure. We would also like to thank Rachael A. Cowan for helping us submit the microPublication and data to NCBI, and Alyssa C. Koehler and Abigail R. Myers for their input on drafting the microPublication.

Extended Data

Description: GTF. Resource Type: Model. File: <u>DeugGB2.Akt1.gff</u>. DOI: <u>10.22002/D1.20201</u>

Description: FAA. Resource Type: Model. File: <u>DeugGB2.Akt1.fasta</u>. DOI: <u>10.22002/D1.20202</u>

Description: FNA. Resource Type: Model. File: <u>DeugGB2.Akt1.pep</u>. DOI: <u>10.22002/D1.20203</u>

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. J Mol Biol 215: 403-10. PubMed ID: <u>2231712</u>

Andjelković M, Jones PF, Grossniklaus U, Cron P, Schier AF, Dick M, Bilbe G, Hemmings BA. 1995. Developmental regulation of expression and activity of multiple forms of the Drosophila RAC protein kinase. J Biol Chem 270: 4066-75. PubMed ID: <u>7876156</u>

Bellacosa A, Testa JR, Staal SP, Tsichlis PN. 1991. A retroviral oncogene, akt, encoding a serine-threonine kinase containing an SH2-like region. Science 254: 274-7. PubMed ID: <u>1833819</u>

Bock, I.R., Wheeler, M.R. (1972). The Drosophila melanogaster species group. Univ. Texas Publs Stud. Genet. 7(7213): 1--102.

Chen ZX; Sturgill D; Qu J; Jiang H; Park S; Boley N; et al.; Richards S. 2014. Comparative validation of the D. melanogaster modENCODE transcriptome annotation. Genome Res 24: 1209-23. PubMed ID: <u>24985915</u>

Duda, O. 1924. Revision der europäischen u. grönländischen sowie einiger sudostasiat. Arten der Gattung Piophila Fallén (Dipteren) [part]. Konowia 3: 97-113 [1924.07.10]

Navarro Gonzalez J; Zweig AS; Speir ML; Schmelter D; Rosenbloom KR; Raney BJ; et al.; Kent WJ. 2021. The UCSC Genome Browser database: 2021 update. Nucleic Acids Res 49: D1046-D1057. PubMed ID: <u>33221922</u>

Grewal SS. 2009. Insulin/TOR signaling in growth and homeostasis: a view from the fly world. Int J Biochem Cell Biol 41: 1006-10. PubMed ID: <u>18992839</u>

Guo HF, Zhong Y. 2006. Requirement of Akt to mediate long-term synaptic depression in Drosophila. J Neurosci 26: 4004-14. PubMed ID: <u>16611817</u>

Hietakangas V, Cohen SM. 2009. Regulation of tissue growth through nutrient sensing. Annu Rev Genet 43: 389-410. PubMed ID: <u>19694515</u>



7/2/2022 - Open Access

Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D. 2002. The human genome browser at UCSC. Genome Res 12: 996-1006. PubMed ID: <u>12045153</u>

Larkin A; Marygold SJ; Antonazzo G; Attrill H; Dos Santos G; Garapati PV; et al.; FlyBase Consortium.. 2021. FlyBase: updates to the Drosophila melanogaster knowledge base. Nucleic Acids Res 49: D899-D907. PubMed ID: <u>33219682</u>

Potter CJ, Pedraza LG, Xu T. 2002. Akt regulates growth by directly phosphorylating Tsc2. Nat Cell Biol 4: 658-65. PubMed ID: <u>12172554</u>

Pélandakis, M., Solignac, M. Molecular phylogeny of Drosophila based on ribosomal RNA sequences. J Mol Evol 37, 525–543 (1993). https://doi.org/10.1007/BF00160433 DOI: <u>doi.org/10.1007/BF00160433</u>

Raney BJ; Dreszer TR; Barber GP; Clawson H; Fujita PA; Wang T; et al.; Kent WJ. 2014. Track data hubs enable visualization of user-defined genome-wide annotations on the UCSC Genome Browser. Bioinformatics 30: 1003-5. PubMed ID: <u>24227676</u>

Rele CP, Sandlin KM, Leung W, Reed LK. 2020. Manual Annotation of Genes within Drosophila Species: the Genomics Education Partnership protocol. bioRxiv 2020.12.10.420521 doi: 10.1101/2020.12.12.420521 DOI: 10.1101/2020.12.12.420521

Staveley BE, Ruel L, Jin J, Stambolic V, Mastronardi FG, Heitzler P, Woodgett JR, Manoukian AS. 1998. Genetic analysis of protein kinase B (AKT) in Drosophila. Curr Biol 8: 599-602. PubMed ID: <u>9601646</u>

Verdu J, Buratovich MA, Wilder EL, Birnbaum MJ. 1999. Cell-autonomous regulation of cell and organ growth in Drosophila by Akt/PKB. Nat Cell Biol 1: 500-6. PubMed ID: <u>10587646</u>

Funding: This material is based upon work supported by the National Science Foundation under Grant No. IUSE-1915544 to LKR and the National Institute of General Medical Sciences of the National Institute of Health Award R25GM130517 to LKR. The Genomics Education Partnership is fully financed by Federal moneys. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Author Contributions: Ashley Morgan: formal analysis, writing - original draft, writing - review editing. Cole A. Kiser: writing - review editing, writing - original draft, formal analysis. Isabel Acosta-Coley: formal analysis, writing - original draft. Hanjun Lin: formal analysis, writing - original draft. Nicholas Guillette: formal analysis, writing - original draft. Robert McMahon: formal analysis, writing - original draft. Jennifer A. Kennell: supervision, writing - review editing. Lindsey J Long: supervision, writing - review editing. Laura K. Reed: supervision, conceptualization, writing - review editing. Chinmay P. Rele: formal analysis, data curation, methodology, supervision, writing - review editing.

Reviewed By: GEP Review Panel

History: Received October 1, 2021 Revision Received June 15, 2022 Accepted June 17, 2022 Published Online July 2, 2022 Indexed July 16, 2022

Copyright: © 2022 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Morgan, A; Kiser, CA; Acosta-Coley, I; Lin, H; Guillette, N; McMahon, R; et al.; Rele, CP (2022). *Drosophila eugracilis – Akt*. microPublication Biology. <u>10.17912/micropub.biology.000544</u>