



Atopic dermatitis and autoimmunity: the occurrence of autoantibodies and their association with disease severity

James Holmes¹ · Lucy C. Fairclough¹ · Ian Todd¹

Received: 3 July 2018 / Revised: 5 December 2018 / Accepted: 9 February 2019 / Published online: 23 February 2019
© The Author(s) 2019

Abstract

Atopic dermatitis (AD) is a widespread condition that appears to be increasing in prevalence and severity worldwide, yet the underlying mechanisms are not well understood. Recent research has identified various similarities between AD and autoimmune conditions, as well as indicating that there may be an association between AD and autoimmunity. This systematic review evaluates the association between AD and autoimmunity, as well as between severity of disease in AD and autoimmunity, with an emphasis on the associations with autoantibodies. MEDLINE (1946 to December 2017) and Embase (1974 to December 2017) databases were searched. Further relevant articles were retrieved from reference lists. Only studies measuring direct indicators of autoimmunity, in humans, were included. Qualitative analysis was carried out for all studies. In addition, quantitative analysis was used to evaluate prevalence of IgE autoantibodies and anti-nuclear antibodies (ANAs) in AD patients and control subjects. The Mantel–Haenszel method was used with a random-effects model. 28 studies assessed the occurrence of autoantibodies in AD patients and 16 studies were used to evaluate association between disease severity and autoantibodies. Pooled analysis from 14 studies, involving 986 AD patients and 441 control subjects, showed that IgE autoantibodies were significantly more prevalent in patients with AD ($P < 0.00001$) than control subjects. Similar analysis was carried out for ANAs, with eight studies that involved 1045 AD patients and 1273 control subjects. ANAs were significantly more prevalent in patients with AD ($P = 0.003$). This quantitative analysis supported an association between AD and IgE autoantibodies, as well as between AD and ANAs. There was insufficient data to make similar conclusions for other indicators of autoimmunity. The weight of evidence also suggests an association between IgE autoantibodies and disease severity. There was insufficient evidence to make this link for other indicators of autoimmunity.

Keywords Atopic dermatitis · Autoimmunity · Autoantibody · IgE · IgG · IgA · Anti-nuclear autoantibodies

Introduction

The term Atopic Dermatitis (AD) was first used by Wise and Sulzberger in 1933 [1]. The World Allergy Organisation defines eczema (also known as atopic eczema dermatitis syndrome (AEDS) and as AD throughout this review) as ‘an inflammatory, chronically relapsing, non-contagious and extremely pruritic skin disease’ [2].

Prevalence of AD worldwide is high, with 7.9% of 6–7 years presenting with the condition in the ISAAC Phase 3 study [3]. AD appears to be increasing in prevalence

globally, with centres reporting increases in self-reported AD and severe AD [3]. There was significant variance in the prevalence of AD between centres, both within and across regions, which has been attributed, in part, to environmental differences [4]. The increasing prevalence and severity of AD makes it essential to understand the underlying mechanisms of the disease, enabling more effective future treatments.

AD is traditionally thought of as an allergic condition with extrinsic allergens driving the immune response. The discovery of a deficiency in the stratum corneum caused by filaggrin gene mutations supports this, as impaired skin barrier function allows for allergens to infiltrate the body, inducing an immune response [5]. AD patients with these mutations are more likely to have persistent, severe, allergen-driven symptoms, but some patients continue to suffer from AD despite removal of stimuli such as dust mites or

✉ Ian Todd
ian.todd@nottingham.ac.uk

¹ School of Life Sciences, University of Nottingham, Life Sciences Building, University Park, Nottingham NG7 2RD, UK

pollen [6]. Overall, as many as two-thirds of AD patients demonstrate ‘no measurable allergen-specific IgE antibody sensitization’ [7].

One hypothesis as to why chronic inflammation persists, despite removal of environmental allergens, is infection by *Staphylococcus aureus*. *S. aureus* colonization was found in more than 90% of patients and was found to correlate significantly with severity [8]. *S. aureus* has a contribution to most AD lesions, both by worsening inflammation through direct action of toxins and by increasing production of IgE antibodies [9]. It would follow that other skin based microorganisms, such as the fungus *Malassezia sympodialis*, could also cause this type of exacerbation, but this does not account for patients that suffer AD lesions without allergic sensitisation.

Despite the traditional understanding of AD, there has been evidence to suggest there might be an autoimmune component to the disease. Disease progression in AD is similar to that in known autoimmune conditions, involving periods of relapse and remission. Furthermore, a significant association has been found between AD and 11 autoimmune conditions, as well as the fact that AD patients are more likely to present with ‘multiple autoimmune comorbidities’ [10]. Finally, immediate skin reactions to some human proteins indicate that, for some patients, IgE autoantibodies are produced [11] which could contribute to disease progression. Researchers have also recently claimed that they have proven that ‘atopic dermatitis ... is an immune-driven (autoimmune) disease’ [12] by treating AD with Dupilumab. This blocked IL-4 and IL-13 [13], to prevent Th2 cytokines from incorrectly targeting autologous tissues.

Methods

This systematic review investigates the link between AD and autoimmunity (with an emphasis on autoantibodies), providing an update to, and widening of scope of, a systematic review from 6 years earlier [14]. The present review evaluates the strength of the link between AD and autoantibodies, and whether there is an association between AD severity and autoantibodies.

Identification of relevant studies

Two modifications were made to the search strategy employed in 2012 by Tang et al. [14]: ‘autologous’ was added as a search term and the Embase database was searched in addition to MEDLINE. These alterations ensured the search strategy reflected current terminology and that as many relevant papers as possible were found.

The search terms were as follows:

1. exp Dermatitis, Atopic/
2. atopic dermatitis.mp.
3. dermatitis atopic.mp.
4. exp Eczema/
5. childhood eczema.mp.
6. infantile eczema.mp.
7. 1 or 2 or 3 or 4 or 5 or 6
8. exp Autoimmunity/
9. exp Autoantigens/
10. Autoallerg\$.mp.
11. autoreactivity.mp.
12. Profilins/
13. ribosomal P2.mp.
14. exp Thioredoxins/
15. exp immunoglobulin G/
16. exp immunoglobulin E/
17. superoxide Dismutase/
18. autoantibod\$.mp.
19. self-antigen.mp.
20. self-react\$.mp.
21. autologous.mp.
22. 8 or 9 or 10 or 11 or 12 or 13 or 14 or 15 or 16 or 17 or 18 or 19 or 20 or 21
23. 7 and 22
24. limit 23 to (male and female and humans and humans)

Database searching was carried out during October to December 2017.

The PRISMA diagram [15] in Fig. 1 gives an overview of the number of papers at each stage of the literature search. Titles and abstracts from this search were checked, yielding 165 papers that warranted full-text analysis. These were scanned in full, with 28 studies providing evidence relating to an association between AD and autoantibodies, and 16




Fig. 1 PRISMA diagram (adapted from [16])

Table 1 Studies examining the association of autoantibodies of IgE class with atopic dermatitis

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Aichberger et al. [17]	Case-control study 12 patients with eczema 11 patients with rhinoconjunctivitis 6 healthy controls 6 control subjects with chronic dermatoses (All adults. Patients with AD ranged from 20 to 72 years, with a mean age of 38.4. Control subjects had a mean age of 38.6. Insufficient data to calculate SD)	<i>Clear</i> Hanifin and Rajka	<i>Western Blot</i> Serum IgE against epithelial cell line A431 and rHom s 4	0% No control subjects produced serum IgE against epithelial cells (A431) or Hom s 4	91.7% 11/12 patients produced serum IgE against epithelial cells (A431) <i>16.7%</i> 2/12 patients produced serum IgE against Hom s 4
Alrichier et al. [18]	<i>Case series</i> 192 patients with eczema 5 healthy control subjects (All adults. AD patients ranged from 18 to 80 years, with a mean age of 39. Control subjects ranged from 25 to 64 years, with a mean age of 31. Insufficient data to calculate SD)	<i>Clear</i> Hanifin and Rajka	<i>Western blot</i> Serum IgE against epithelial cell line A431 + epidermis	0% 0/26 control subjects displayed any autoreactivity	28% 54/192 eczema patients displayed autoreactivity against A431 and/or epidermis
Guarneri et al. [19]	Case-control study 27 patients with AD 27 control subjects (10 children and 17 adults. All participants ranged from 5 to 49 years. AD patients and controls had a mean age of 22.2 ± 12.2)	<i>Clear</i> Hanifin and Rajka	<i>Skin prick test</i> against hMnSOD	0% 0/27 control patients	14.8% 4/27 patients
Hiragun et al. [20]	IgE and IgG Case-control study 63 patients with AD 24 patients with Cholinergic Urticaria 32 patients with bronchial asthma 14 patients with allergic rhinitis 23 normal control patients (Patients with AD ranged from 0 to 65 years, with a mean age of 29.9 ± 11.5. Other participants in this study varied from 0 to 82)	<i>Moderately Clear</i> 'severity of AD was evaluated using severity index of Japanese guideline for AD'	<i>ELISA</i> Specific to QRX-specific IgE and rMGL_1304-specific IgE	26.1% 6/23 controls were QRX-specific IgE positive 39.1% 9/23 controls were rMGL_1304-specific IgE Positive	66.7% 42/63 AD patients were QRX-specific IgE positive <i>P=0.0012</i> 61.9% 39/63 AD patients were rMGL_1304-specific IgE Positive <i>Positive (P=0.0857)</i>

Table 1 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Iives, Virolainen and Harvima [21]	Case-control study 50 patients with AD 24 control subjects (All adults. Patients with AD ranged from 21 to 64 years, with a mean age of 38. Control subjects ranged from 21 to 68 years, with a mean age of 42. Insufficient data to calculate SD)	Clear Rajka and Langeland	<i>Intracutaneous injection and ImmunoCAP</i> against autologous sweat	50% 1/24 positive 11/24 weakly positive 12/24 negative	72% 19/50 positive 17/50 weakly positive 14/50 negative (difference $P < 0.008$)
Kawamoto et al. [22] IgE and IgG	Case-control study 40 patients with AD 26 control patients (no clinical history of allergic diseases and serum IgE within normal levels) (Patients with AD ranged from 12 to 44 years, with a mean age of 26.4. Control subjects ranged from 19 to 54 years, with a mean age of 29.2. No SD provided and insufficient data to calculate)	<i>Clear</i> UK Working Party's Diagnostic Criteria for Atopic Dermatitis	<i>ELISA</i> Peptide Specific	—	No significant difference in IgG levels or number of positive samples for 5 peptides
Kortekangas-Savolainen et al. [23]	Case-control study 27 patients with IgE-mediated AEDS 13 control subjects (6 healthy controls, 4 patients with urticaria, 3 patients with psoriasis) (All adults. Patients with AD had a mean age of 33 ± 11 . Control subjects had a mean age of 47 but insufficient data to calculate SD.)	<i>Not clear</i>	<i>Western blot</i> Serum IgE against keratinocytes	0% 0/13 control subjects displayed IgE autoantibodies against keratinocytes	37% 10/27 patients displayed IgE autoantibodies against keratinocytes

Table 1 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Mittermann et al. [24]	Case-control study 11 patients with AD 7 patients with rhinoconjunctivitis 5 patients with contact allergy 9 non-atopic individuals (All adults, except 1 child with AD. Patients with AD ranged from 13 to 59 years, with a mean of 31.9 ± 14.9. Control subjects ranged from 26 to 63 years, with a mean of 35.2 ± 10.9.)	<i>Clear</i> Hanfin and Rajka	<i>Immunoblotting</i> Against human epithelial cell extracts	0% 0/12 control subjects	72.7% 8/11 patients
Mittermann et al. (2016) [25]	Case-control study 179 patients with AD (53 severe and 126 moderate) 140 patient control group (43 patients with seborrhic eczema and 97 individuals with no previous allergies or skin conditions) (All adults. All participants ranged from 18 to 65 years. AD patients had a mean age of 28, control subjects had a mean age of 36. No SD provided and insufficient data to calculate)	<i>Clear</i> UK Working Party's Diagnostic Criteria for Atopic Dermatitis	<i>Immunoblotting</i> Against human A431 extract	0% 0/140 control subjects	18% 32/179 patients with AD
Mothes et al. [26]	Case-control study 174 patients with eczema 10 patients with psoriasis 26 healthy controls (All adults. Eczema patients had a mean age of 35.4 ± 14.8. Control subjects had a mean age of 36.5 ± 16.5.)	<i>Clear</i> Hanfin and Rajka	<i>Immunoblotting</i> IgE reactivity to a variety of human epithelial antigens	0% 0/10 patients with psoriasis and 0/26 healthy controls displayed autoreactivity	23% 40/174 patients displayed autoreactivity
Natter et al. [27]	Case series 51 patients with eczema (14 children and 37 adults. Patients with eczema ranged from 1 to 63 years, with a mean age of 26.9 ± 18.6)	<i>Clear</i> Hanfin and Rajka	<i>Western Blot</i> Whole serum IgE against endothelial cells and A431	— —	43.1% 22/51 patients

Table 1 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Schmid-Grendelmeier <i>et al.</i> (2005) [28]	Case-control study 69 patients with eczema 5 healthy controls 13 patients with psoriasis 11 patients with ABPA 13 patients with allergy to A fumigatus (All adults. Eczema patients had a mean age of 29.3 ± 5.6 . Other participants had a mean age of 25.3, but insufficient data to calculate SD)	<i>Clear</i> Hanifin and Rajka	<i>ELISA and SPT</i> against rhMnSOD	100% 11/11 patients with ABPA 0% 0/31 other control subjects	42% 29/69 patients with eczema
Valenta et al. [29]	Case-control study 20 patients with AD 28 control patients including patients with various allergic conditions and non-allergic healthy individuals (Patients with AD ranged from 17 to 52 years with a mean age of 35.3 ± 12.3 . Control subjects ranged from 18 to 64 years with a mean age of 33.0 ± 11.4)	<i>Clear</i> Hanifin and Rajka	<i>Western Blot</i> IgE antibodies against endothelial cells, platelets, fibroblasts and epithelial cells	0% 0/28 tested positive for IgE against human proteins from most cell extracts	60% 12/20 tested positive for IgE against human proteins from most cell extracts
Zeller et al. [30]	Case-control study 71 patients with AE 12 patients with psoriasis 24 healthy control subjects (Patients with AE had a mean age of 33.35 ± 12.70 . Patients with non-atopic AE had a mean age of 39.35 ± 13.5 . Patients with psoriasis had a mean age of 45.17 ± 9.20 . Healthy control subjects had a mean age of 29.00 ± 3.71)	<i>Clear</i> Hanifin and Rajka and European Academy of Allergology and Clinical Immunology	<i>Immunoblotting and ELISA</i> Specific to actin- α , eIF6, RP1, HLA-DR- α and tubulin- α	0% 51/71 of all patients with Atopic Eczema 72.2%	71.8% 13/18 patients with the non-atopic form of AD

that evaluated severity of AD in relation with the occurrence of autoantibodies. During this full-text analysis, the reference sections of papers were checked to identify relevant papers that had not appeared in the search results. Most papers excluded at this point were conference proceedings or included results not linked to autoimmunity and AD.

Inclusion criteria

Language

All literatures evaluated in this systematic review were in English (including papers that had already been translated in to English).

Types of studies

Due to the wide range of study designs present in the literature, all the types of study were included. The main types of study evaluated were case-control studies, case series, and individual case reports.

Types of participants

This systematic review evaluated all studies involving human participants, irrespective of age or gender. Studies varied in the age ranges of the participants (details in Tables 1, 3, 4, 5, 7): most included adults only or a mixture of children and adults, whereas a few studies included children only. However, there was no apparent effect of participants' age on the relationship between autoreactivity and AD (Table 2) or severity of AD (Table 6). All the literature was included regardless of whether they had clear criteria for selecting patients with AD. These criteria have been noted for all

studies in appendices, but have only been referred to in the main text when these criteria are unclear or missing.

Types of publication

In this systematic review, all papers published in journals were included, with the exception of reviews and editorials. Letters to journals were included, as they were a key source of smaller studies. Grey literature was not included.

Outcome measures

The papers evaluated in the systematic review were heterogeneous in both study design and outcome measures. The only restriction was that studies had to provide direct evidence of autoantibodies. When evaluating the association between autoantibodies and AD, any tests that indicated autoantibodies, for example, ELISA and skin prick tests (SPTs), were included. When evaluating the association between severity of AD and autoantibodies, all measures of severity were included. The most common measures of severity were SCORAD (SCORing Atopic Dermatitis) [44] and Rajka and Langeland [45] systems, although some authors used other assessments of severity.

Statistical analysis

Where appropriate, a quantitative analysis was carried out using Review Manager 5.3 [46]. The Mantel-Haenszel method was used with a random-effects model to account for the variation in study design and resultant difference in effect size.




Fig. 2 Forest plot showing data from all studies into IgE antibodies. 'Events' are instances of autoimmunity, and 'total' refers to the total number of people in that group of a study

Results

Association of autoantibodies with atopic dermatitis

28 papers were identified that fulfilled the inclusion criteria and were relevant to this part of the review. The main indicators of autoimmunity investigated were specific IgE and IgG antibodies, as well as IgG ANAs.

Autoantibodies of IgE class

14 studies investigated the presence of specific IgE autoantibodies in patients with AD; the details of these studies are summarised in Table 1 (in alphabetical order of the first authors). Twelve were case-control studies, of which 11 indicated that certain IgE autoantibodies were significantly more common in patients with AD. The study

quality was generally good, with well-described protocols, often using western blot, ELISA or SPT techniques, and robustly defined criteria for selection of AD patients in most cases. Most studies evaluated the presence of IgE autoantigens to epithelial cells, epithelial cell extracts, and other proposed autoantigens associated with the skin. Furthermore, the control groups were comprised predominantly of healthy individuals which makes comparison between, and pooling of, individual studies much easier and more useful. This meant an analysis could be carried out on the combined data, which established an odds ratio of 12.97 (with 95% confidence limits of 5.14–32.68) and $P < 0.00001$, indicating that IgE autoantibodies are significantly more likely to be identified in patients with AD compared to the control groups (Fig. 2). This effect seemed to be uniform across all tested antigens, although the majority of evidence concerns specific antibodies to epithelial cell extracts. A qualitative analysis of the studies also indicated that all but one (13/14) concluded that

Table 2 Association of autoantibodies with atopic dermatitis

Marker of autoimmunity	Papers that indicate autoreactivity is more common in patients with AD	Papers that do not indicate autoreactivity is more common in patients with AD
IgE autoantibodies	13 Aichberger et al. [17] Altrichter et al. [18] Guarneri et al. [19] Hiragun et al. (for QRX) [20] Ilves, Virolainen and Harvima [21]* Kortekangas-Savolainen et al. [23] Mittermann et al. [24, 25] Mothes et al. [26] Natter et al. [27] Schmid-Grendelmeier et al. [28] Valenta et al. [29] Zeller et al. [30]	2 Hiragun et al. (for rMGL_1304) [20] [†] Kawamoto et al. [22] [§]
IgG autoantibodies	4 Bergman et al. [31]* (for CCL-3) Neuber et al. [32]* Ochs et al. [33] Szakos et al. [34]*	5 Ambrozic et al. [35] Bergman et al. [31] (for 5 autoantigens) [†] Du Toit et al. [16] El-Rachkidy et al. [36] Kawamoto et al. [22] (for 5 peptides)**, (for 3 peptides) [§]
IgA autoantibodies	0	1 Ress et al. [37]
ANAs	7 Dhar, Kanwar and Deodhar [38] Higashi et al. [39] Ochs et al. [33] Ohkouchi et al. [40]* Szakos et al. [34] Tada et al. [41] Taniguchi et al. [42]	1 Ress et al. [43] [†]

*Differences were statistically significant

[†]Differences were not statistically significant

[§]Autoreactivity was significantly lower in AD patients

Table 3 Studies examining the association of autoantibodies of IgG class with atopic dermatitis

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Ambrožič et al. [35]	Case–control study 45 patients with AD 26 patients with other atopic conditions (controls) (All children. Patients with AD ranged from 2 months to 16.8 years, with a mean age of 3.7. Control subjects ranged from 2.1 to 17.8 years, with a mean age of 11.0. Insufficient data to calculate SD)	<i>Moderately Clear</i> 'Sera selected from the serum banks of the Pediatrics Clinic and Department of Rheumatology (University Medical Centre, Ljubljana)'	<i>ELISA</i> Specific to Anti-β2GPI	38% 10/26 control patients	42% 19/45 patients with AD <i>No significant difference</i>
Bergman et al. [31]	Case–control study 37 Patients with Psoriasis 18 patients with AD 56 healthy controls (All adults. Patients with psoriasis ranged from 20 to 76 years, with a median age of 43.5. Patients with AD ranged from 19 to 69 years, with a median age of 30. Healthy subjects ranged from 27 to 62 years, with a median age of 45.5)	<i>Clear</i> Hanifin and Rajka	<i>ELISA</i> Against a variety of autoantigens	Mean Log2Ab Titer of autoantibody (HCs)	Mean Log2Ab Titer of autoantibody (HCs)
Du Toit et al. (2006) [16]	Case–control study 33 patients with AEDS 78 patients with Chronic Urticaria (All children. Patients with AEDS ranged from 1.25 to 19 years, with a mean age of 7.6. Patients with Chronic Urticaria were age matched)	<i>Not clear</i>	<i>Anti-Fce receptor assays</i> Specific to IgG FcεR1α autoantibodies	47% 37/78 patients with CU	0% 0/33 patients with AEDS
El-Rachkidy et al. [36]	Case–control study Patients with psoriasis (number varies based on test) 5 pooled patients with AD (Age of patients with AD not specified. Psoriasis patients ranged from 22 to 66 years)	<i>Not clear</i>	<i>Immunoblotting and ELISA</i> Specific to stratum corneum antigens	Psoriasis patients were autoreactive against stratum corneum antigens	AD patients were not autoreactive against stratum corneum antigens

Table 3 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Kawamoto et al. [22] (IgE and IgG)	Case–control study 40 patients with AD 26 control patients (no clinical history of allergic diseases and serum IgE within normal levels) (Patients with AD ranged from 12 to 44 years, with a mean age of 26.4. Control subjects ranged from 19 to 54 years, with a mean age of 29.2). Insufficient data to calculate SD	<i>Clear</i> UK Working Party's Diagnostic Criteria for Atopic Dermatitis	ELISA Peptide Specific	—	No significant difference in IgG levels or number of positive samples for 5 peptides
Neuber et al. [32]	Case–control study 16 patients with AD 72 healthy control subjects (All adults. Patients with anti-CD28 autoantibodies had a mean age of 52.3 ± 18.8 , patients without anti-CD28 autoantibodies were had a mean age of 58.7 ± 19.9)	<i>Moderately clear</i> Treated at the Department of Dermatology, University Hospital Eppendorf, Hamburg	ELISA Specific to Anti-CD28	11.1% 8/72 control patients	87.5% 14/16 AD patients ($P < 0.0001$)
Ochs et al. [33] ANA and IgG	Case–control study 64 patients with eczema 39 control patients (study also included a range of other atopic conditions) (Patients with eczema ranged from 4 to 43 years, with a mean age of 24.4. No ages provided for control subjects. Insufficient data to calculate SD)	<i>Clear</i> Hanifin and Rajka	Indirect immunofluorescence For ANAs Western blot Against anti-DFS70 IgG	ANA: not stated WB-DFS70: 0% 0/39 healthy control subjects WB-DFS70: 29.6% 19/64 patients	ANA: 41% WB-DFS70: 0% 0/39 healthy control subjects WB-DFS70: 29.6% 19/64 patients

Table 3 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Szakos et al. [34] IgG and IgM (including ANA)	Case–control study 72 patients with AEDS 22 healthy control patients (All children. Patients with AEDS ranged from 2 to 17 years, with a mean age of 8 years. Control subjects ranged from 1.5 to 14 years, with a mean age of 8.6 years. Insufficient data to calculate SD)	Not Clear	Indirect immunofluorescence For ANAs ELISA Specific to APLs	9.1% 2/22 healthy control patients had non-allergen-specific IgG and/or IgM antibodies 9.1% 2/22 healthy control patients displayed ANAs	29.3% 21/72 patients had non-allergen-specific IgG and/or IgM antibodies 13.9% 10/72 AD patients displayed ANAs <i>APL antibodies occurred significantly more often in AD patients</i>

IgE class autoantibodies are more common in AD patients than controls (Table 2).

Autoantibodies of IgG class

Eight studies investigated the presence of specific IgG autoantibodies in AD (Table 3). However, it was not possible to carry out a quantitative analysis: whilst the studies predominantly used ELISA techniques, the study design was too varied for a pooled analysis. Some studies only compared AD patients against patients with other conditions, for example psoriasis, whilst others only provided mean titres. Furthermore, only three of the studies had clear definitions for AD diagnosis. Because of the more variable study design, it was more appropriate to carry out a qualitative review. Levels of IgG autoantibodies specific to SART2₁₆₁, SART2₈₉₉, ART4₁₃ [22], CCL3 [31] and CD28 [32], as well as APL antibodies [34] were significantly higher than in their respective control groups, but levels of autoantibodies specific to β_2 GPI [35], TNF- α , IFN- α , CCL-5, CCL-2, IL-17 [31], and five other peptides [22] were not significantly higher. Autoantibodies against DFS70 [33] and Fc_eR1₁ autoantibodies [16] were more commonly identified in AD patients than the respective control groups, but no statistical analysis was carried out. Finally, AD patients were found not to display an autoreactive response to stratum corneum antigens, but psoriasis patients did (although no statistical analysis was carried out). Overall, there is no clear picture as to the presence of IgG autoantibodies in AD patients (Table 2). The mixed results, combined with more variable study techniques and less clarity with respect to inclusion criteria, means that more research is required into the relationship between AD and IgG autoantibodies.

Autoantibodies of IgA class


Only one study solely investigated IgA autoantibodies in AD: Ress et al. [37] carried out a large case–control study, comprising 297 patients with AD and 52 healthy control subjects. ELISA techniques were used to determine the presence of IgA-anti-transglutaminase 1 (anti-TG1) and IgA-anti-transglutaminase 3 (anti-TG3) in the control group and AD group. However, this study did not indicate an association between IgA autoantibodies and AD (Table 2), but there are insufficient studies to draw a clear conclusion.

Table 4 Studies examining the association of anti-nuclear autoantibodies with atopic dermatitis

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Dhar et al. [38]	Case-control study 76 children with AD 58 age matched control subjects (All children. Patients with AD ranged from 6 months – 12 years, with a mean age of 3.4 years. Control subjects were age matched. Insufficient data to calculate SD.)	<i>Clear</i> Hanifin and Rajka	<i>FITC-conjugated goat anti-human IgG /sic/IgG</i>	0.0% 0/58 control subjects displayed ANAs at a titre of 1:40	2.6% 2/76 patients with AD displayed ANAs at a titre of 1:40
Higashi et al. [39]	Case-control study 100 patients with AD 1004 control subjects (Patients with AD ranged from 2 to 64 years, with a mean age of 28.2 years. Control subjects ranged from 20 to 70 years, with a mean age of 48.8 years. Insufficient data to calculate SD.)	<i>Clear</i> Hanifin and Rajka	<i>Indirect immunofluorescence</i> Against ANA	19% 19/100 patients with AD (between titres of 40X–640X)	6.8% 68/1004 control patients (between titres of 40X–640X)
Ochs et al. [33]	Case-control study 64 patients with eczema 39 control patients (study also included a range of other atopic conditions) (Patients with eczema ranged from 4 to 43 years, with a mean age of 24.4 years. Insufficient data to calculate SD. No ages provided for control subjects.)	<i>Clear</i> Hanifin and Rajka	<i>Indirect immunofluorescence</i> Against ANA	41% 26/64 patients	Not stated
Ohkouchi et al. [40]	Case-control study 256 patients with eczema 60 control subjects (All adults. Patients with eczema had a mean age of 23 years. Control subjects were age matched. Insufficient data to calculate SD.)	<i>Clear</i> Hanifin and Rajka	<i>Indirect immunofluorescence</i> Against ANA	11.7% 7/60 control patients at a 40 fold dilution	36.3% 93/256 AD patients at a 40 fold dilution ($P < 0.001$)

Table 4 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Ress et al. (2015) [43]	<i>Cross-sectional study</i> 346 children with active AD 117 hospital controls without known skin diseases, at similar ages (All children. Patients with AD ranged from 0.5 to 18.8 years, with a mean age of 5.8 years. Control subjects ranged from 0.5 to 17.7 years, with a mean age of 7.9 years. Insufficient data to calculate SD.)	<i>Clear</i> UK Working Party's Diagnostic Criteria for Atopic Dermatitis	<i>Indirect immunofluorescence</i> Against ANA	12.8% 15/117 control patients were ANA Positive at a titre of 1:10	13.6% 47/346 patients with AD were ANA Positive at a titre of 1:10 ($P > 0.05$)
Szakos et al. [34] IgG and IgM (including ANA)	Case-control study 72 patients with AEDS 22 healthy control subjects (All children. Patients with AEDS ranged from 2 to 17 years, with a mean age of 8.0 years. Control subjects ranged from 1.5 to 14 years, with a mean age of 8.6 years. Insufficient data to calculate SD.)	<i>Not clear</i>	<i>Indirect immunofluorescence</i> Against ANA	9.1% 2/22 healthy control patients displayed ANAs	13.9% 10/72 AD patients displayed ANAs
Tada et al. [41]	<i>Case series</i> 89 patients with AD (Patients with AD ranged from 5 to 49 years, with a mean age of 19.0 ± 7.4 years)	<i>Clear</i> Hanifin and Rajka	<i>Indirect immunofluorescence</i> Against ANA of the IgG subclass	-	25.8% 23/89 displayed ANAs at dilutions of 1:40–1:640
Taniguchi et al. [42]	Case-control study 47 AD patient sera analysed using method 1 57 AD patient sera analysed using method 2 12 control subject sera analysed using methods 1 and 2 (all adults) All adults. Patients with AD ranged from 15 to 30 years. Insufficient data to calculate mean and SD. No data provided on control subjects	<i>Clear</i> Hanifin and Rajka	<i>FITC-conjugated polyvalent immunoglobulins (method 1)</i> <i>FITC-conjugated anti-IgG (method 2)</i>	25% 3/12 healthy control subjects had ANAs identified using method 1 16.7% 2/12 healthy control subjects had ANAs identified using method 2	34.0% 16/47 AD patients with ANA identified using method 1 26.3% 15/57 AD patients with ANA identified using method 2



N.B. 'Ochs et al., 2000' – this study did not indicate the number of control subjects that displayed auto-reactivity. For this reason, although there were 39 control subjects, these were excluded from the forest plot.

Fig. 3 Forest plot showing data from all studies into ANAs. 'Events' are instances of autoimmunity, 'Total' refers to the total number of people in that group of a study

Anti-nuclear autoantibodies

Eight studies investigated ANAs in AD (Table 4), with two of these including ANAs as part of a wider study. Six of the eight were case-control studies. Most of these papers applied clear, commonly used definitions of AD and most used well-described methods—particularly indirect immunofluorescence—to identify ANAs. Seven of the eight papers indicate that ANAs occurred more commonly in patients with AD than in the control subjects (Table 2). Furthermore, the pooling of data showed that patients with AD have a significantly higher frequency of ANAs than the control subjects with an odds ratio of 2.18 (95% confidence intervals 1.31–3.64) and $P=0.003$ (Fig. 3).

Association of autoantibodies with severity of atopic dermatitis

The second part of this systematic review considers whether there is an association between severity of disease in patients with AD and the presence of an autoimmune response. From the papers identified using the search strategy, 16 papers evaluated the presence of autoimmunity in combination with disease severity. 13 papers provide evidence of the relationship between IgE autoantibodies and disease severity (Tables 5, 6), and two provide research relating to ANAs (Table 6). The final paper investigated ANAs, APL antibodies, and ACL antibodies and their relationship with severity (Table 6).

Two main methods of assessing severity were used, SCORAD [44] and Rajka and Langeland [45] scores, although some researchers used other measures of severity. SCORAD was developed as a standardised method to assess severity of AD that could be used in outpatient clinics. It relies

on three components: intensity (responsible for 60% of the total score), extent and subjective symptoms (each responsible $\approx 20\%$ of the score). Rajka and Langeland also put forward a simple scoring index for AD using three parameters, each scored out of 3: extent, course (how much remission in a year) and intensity (scored according to how much it impacts sleep). The sum of these scores is calculated, with 3–4 being defined as mild, 4.5–7.5 being moderate, and 8–9 being severe.

It is difficult to produce pooled data for the studies assessing a link between IgE autoantibodies and severity of AD due to wide variation in the measures of severity and outcome measures but, as shown in Table 6; the weight of evidence indicates that there is an association between the frequency of IgE autoantibodies and disease severity in AD. Furthermore, two studies [48, 50] looked at individual patients over time, assessed levels of IgE autoantibodies and SCORAD throughout treatment, and established a temporal link between SCORAD and IgE autoantibodies. Whilst it is impossible to determine from these studies whether the decrease in IgE autoreactivity as severity of disease declines is a separate response to the cyclosporine A treatment, an unrelated biomarker resulting from a decrease in disease severity, or indeed the cause of the reduced severity, it provides strong support for a link between AD disease severity and autoimmunity. This association did not seem to be present with respect to ANAs, as neither study that looked exclusively at ANAs demonstrated a significant association (Table 6), but more studies are required to reach a definitive conclusion. Szakos et al. [34] also indicated that there was no significant difference in disease severity in patients that displayed ANAs, ACL, APL or a combination of ANAs and APL (Table 6). Overall, the available evidence indicates that severity of AD is associated with autoimmunity, but that this response might be limited to IgE-mediated

Table 5 Association between autoantibodies of IgE class and the severity of atopic dermatitis

References	Study design and size	Diagnostic criteria for eczema	Measure of severity	Auto-reactivity measurement methods	Results
Altrichter et al. [18]	<i>Case series</i> 192 patients with eczema (All adults. Patients with eczema ranged from 18 to 80 years, with a mean age of 39 years. Control subjects ranged from 25 to 64 years, with a mean age of 31 years. Insufficient data to calculate SD.)	<i>Clear</i> Hanifin and Rajka	<i>IGA Score Eczema Area and Severity Index</i>	<i>Western blot</i> Serum IgE against epithelial cell line A431 + epidermis	'Disease activity...' was also significantly higher in autoreactive patients than in non-autoreactive patients' ($P < 0.001$)
Hide et al. [47]	Case-control study* 66 patients with eczema (17 mild, 20 moderate, 26 severe and 3 undefined) (Patients with eczema ranged from 13 to 37 years, with a mean age of 24.7 ± 5.2 years.)	Moderately clear Severity based on grading by Rajka and Langeland	<i>Rajka and Langeland</i>	<i>Skin test</i> Intradermal injection of autologous sweat	<i>No association</i> between frequency of reaction to autologous sweat and severity of AD
Hiragun et al. [20]	Case-control study* 63 patients with AD (Patients with AD ranged from 0 to 65 years, with a mean age of 29.9 ± 11.5 years.)	Moderately clear 'severity of AD was evaluated using severity index of Japanese guideline for AD'	Severity index of Japanese guidelines for AD	<i>ELISA</i> Specific to QRX and rMGL_1304	<i>Spearman's Correlation Coefficient</i> $R = 0.5468, P < 0.0001$ rMGL_1304-specific IgE: $R = 0.448, p < 0.0001$
Ilves, Virolainen and Harvima [21]	Case-control study* 50 patients with AD (Patients with AD ranged from 21 to 64 years, with a mean age of 38 years. Insufficient data to calculate SD.)	<i>Clear</i> Rajka and Langeland	<i>Own definition</i> 1=symptomless 2=almost symptomless 3=mild eczema 4=moderate eczema	<i>Intracutaneous injection</i> Of autologous sweat	<i>Linear trend in Chi-squared test</i> ($P = 0.009$)
Kinaciyan et al. [48]	Case report 1 patient with eczema (The patient was 26 years)	Not clear	SCORAD	<i>Western blot</i> Whole serum IgE against human epithelial cell extracts	<i>Moderately positive correlation</i> between disease severity and positive intracutaneous test (rs = 0.37, $p = 0.008$)
Kohsaka et al. [49]	<i>Case series</i> 18 patients with mild AD 23 patients with moderate AD 12 patients with severe AD 11 patients with most severe AD (Patients with AD had a mean age of 28.2 ± 11.3. years.)	<i>Clear</i> Japanese Dermatological Association (JDA) criteria for AD	Severity index of Japanese guidelines for AD	<i>Western blot</i> Levels of IgE binding against MGL_1304 and its homologs	<i>Spearman's Correlation coefficient</i> significant for all tested antigens $r = 0.3995, P < 0.005$ TF-rMala s 8 $r = 0.3094, P < 0.05$ TF-rMala r 8 $r = 0.3588, P < 0.005$

Table 5 (continued)

References	Study design and size	Diagnostic criteria for eczema	Measure of severity	Autoreactivity measurement methods	Results
Lucas <i>et al.</i> [50]	<i>Case reports</i> 4 Patients with AD underwent treatment with Cyclosporine C (All adults. Patients with AD ranged from 31 to 54 years. Insufficient data to calculate mean and SD.)	<i>Clear</i> Hanifin and Rajka	SCORAD	<i>Western blot</i> IgE against Human Epithelial Cell Extracts	'The intensity of IgE autoreactivity...seemed to reflect skin inflammation'
Mittermann <i>et al.</i> (2016) [25]	Case-control study* 179 patients with AD (53 severe + 126 moderate) (All adults. Patients ranged from 18 to 65 years, with a median age of 28 years. Insufficient data to calculate mean and SD.)	<i>Clear</i> UK Working Party's Diagnostic Criteria for Atopic Dermatitis	SCORAD	Immunoblot assay Specific to human A431 extract	30% of patients with severe AD displayed autoreactivity compared to 13% of patients with moderate AD <i>Significant Association</i> ($P < 0.01$)
Mothes <i>et al.</i> [26]	Case-control study* 174 patients with eczema (All adults. Eczema patients had a mean age of 35.4 ± 14.8 years. Control subjects had a mean age of 36.5 ± 16.5 years.)	<i>Clear</i> Hanifin and Rajka	<i>Diepgen score</i>	<i>Western blot</i> IgE reactivity to a variety of human epithelial antigens	Significantly higher total Diepgen score ($P = 0.036$) in AA + vs AA- patients
Natter <i>et al.</i> (1998) [27]	Case series 51 patients with eczema (14 children and 37 adults. Patients with eczema ranged from 1 to 63 years, with a mean age of 26.9 ± 18.6 years)	<i>Clear</i> Hanifin and Rajka	<i>Own score</i>	<i>Western blot</i> Whole serum IgE against endothelial cells and A431 AD	12/12 patients with 'intensive IgE reactivity to human proteins' had severe or moderate AD 16/29 patients without IgE autoreactivity displayed severe or moderate AD
Schmid-Grendelmeier <i>et al.</i> [28]	Case-control study* 69 patients with eczema (All adults. Patients with eczema had a mean age of 29.3 ± 5.6 years. Other participants had a mean age of 25.3 years, but insufficient data to calculate SD)	<i>Clear</i> Hanifin and Rajka	SCORAD	ELISA Allergen-specific against rhMnSOD	Correlation between SCO-RAD and anti-human MnSOD IgE level $R = 0.756$ $P < 0.0001$

Table 5 (continued)

References	Study design and size	Diagnostic criteria for eczema	Measure of severity	Autoreactivity measurement methods	Results
Tanaka et al. [51]	Case-control study* 61 patients with AD (Patients with AD ranged from 2 to 43 years, with a mean age of 24.0 ± 7.5 years)	Moderately clear Severity based on grading by Rajka and Langeland	Rajka and Langeland	Histamine release By basophils upon stimulation by semi-purified sweat antigen	'The extents of histamine release were not statistically correlated with disease severity'
Zeller et al. [30]	Case-control study* 71 patients with AE (Patients with AE had a mean age of 33.35 ± 12.70 years. Patients with non-atopic AE had a mean age of 39.35 ± 13.5 years.)	Clear Hanifin and Rajka and European Academy of Allergology and Clinical Immunology	SCORAD	Western blot and ELISA Against a variety of self-antigens	No significant difference in SCORAD between patients with and without IgE against self-antigens ($P < 0.7625$)

autoreactivity. However, further research is required into other indicators of autoimmunity, such as IgG autoantibodies, ANAs and IgA autoantibodies to confirm this, as current evidence is limited in these areas.

Discussion

The weight of evidence presented in this review indicates that there is an association between autoreactivity and AD, as well as disease severity in AD, particularly with regard to autoantibodies of IgE class. However, this review cannot provide evidence for a causal link between the two, as the exact mechanisms for how autoimmunity occurs in patients with AD are unclear. Some papers hypothesise that cross reactivity between *Malassezia* spp. and the autoantigen hMnSOD could underlie the pathogenic response seen in AD [21, 25], whilst many others have identified cells in the skin as a source of autoantigens, potentially indicating that skin damage could lead to an autoimmune response to components within the skin, giving rise to autoimmunity. However, these hypotheses rest on assumptions that have not been proven. Taking all the available evidence into account, it is not possible to declare that autoimmunity is a pathogenic factor underlying the natural history of AD, but it does seem plausible and certainly worthy of future research.

This review points to several areas, where future research into autoimmunity and AD would be most valuable. First, some areas of autoimmunity have been less thoroughly investigated when looking at an association with AD. Whilst IgE autoantibodies and, to some extent, ANAs have been covered extensively by research, other markers of autoimmunity have less evidence available. Studies investigating IgG autoantibodies, for example, were varied in terms of control groups, often comparing against groups that have other diseases, and were less clear about the selection criteria for the patients with AD. More research into IgG autoantibodies in AD, across a wide range of autoantigens and in comparison to healthy control groups, would be useful in drawing more robust conclusions.

Five other studies, not specifically covered by this review, have examined aspects of autoimmunity other than autoantibodies (Table 7). Collectively, these studies do indicate that autoimmunity is related to AD, with three of the five indicating statistically significantly higher rates of autoimmune response in patients with AD compared to their control groups, be it through increased histamine release upon stimulation by semi-purified sweat antigens [51], activation of iNKT cells without the corresponding ligand [52], or significantly higher stimulation index of PBMCs in response to hsp60 [53]. In addition, Hide et al.

Table 6 Association of autoantibodies with severity of atopic dermatitis

Marker of autoimmunity	Papers that indicate a link between AD disease severity and autoimmunity	Papers that do not indicate a link between AD disease severity and autoimmunity
IgE autoantibodies	10 Altrichter et al. [18]* Hiragun et al. [20]* Ilves, Virolainen and Harvima [21]* Kinaciyan et al. [48] Kohsaka et al. [49]* Lucae et al. [50] Mittermann et al. [25]* Mothes et al. [26]* Natter et al. [27] Schmid-Grendelmeier et al. [28]*	3 Hide et al. [47] Tanaka et al. [51]† Zeller et al. [30]†
ANAs	0	2 Higashi et al. [39]† Ress et al. [43]†
Multiple indicators	0	1 Szakos et al., 2004 [34]

*Differences were statistically significant

†Differences were not statistically significant

[47] show a greater frequency of response to intradermal injection of autologous sweat in AD patients, albeit without statistical analysis. The final study [54] does not seem to indicate an increased autoreactive T-cell response in patients with AD, but does indicate that there may be an alteration to the balance of CLA+/CLA- and CCR4+/CCR4- cells. Overall, these less traditional indicators of autoimmunity show promising results in terms of a link with AD, but further research is required to come to definitive conclusions.

There are also new indicators of autoimmunity that, whilst not meeting the inclusion criteria of this review, could provide insight into AD. Some interesting papers have investigated the function of regulatory T cells (Tregs) in AD with research indicating that the suppressive effect of Tregs on CD8⁺CLA⁺ T-cell proliferation is inhibited in AD [55]. This would be a similar mechanism to other autoimmune/inflammatory diseases such as psoriasis, which shows impaired suppression of CD4⁺CD25⁺ T cells [56]. Zhang et al. [55] note that further experiments are needed to address the theory that the CD8⁺CLA⁺ T cells in patients with AD might resist suppression, as opposed to there being dysregulation of Tregs, but further research into this field could provide a valuable insight into the mechanism behind autoimmunity. Later research corroborated this [57], with both papers suggesting that reduced inhibition of Teff cells could cause impaired self-tolerance in patients with AD, but further research would assist in elucidating specific mechanisms underlying the autoimmune response.

Another area that could provide useful research into autoimmunity and AD is analysis of genetic variants and

potential associations with AD. For example, copy-number variations of the human histamine H4 receptor (HRH4) gene have been shown to be associated with AD [58], as well as a number of other autoimmune conditions such as SLE [59]. Whilst the exact mechanisms involved with the human histamine H4 receptor gene and AD are unclear, Chen et al. [58] suggest that the expression of HRH4 could be enhanced because of inflammatory stimuli found in autoimmune diseases, leading to a stronger response to histamine. This is an example of another potential component to the autoimmune pathogenesis in AD and is an area that warrants future research.

How autoimmunity is related to AD in animal models also warrants further research. Using animal models could lead to greater understanding of whether autoimmunity causes AD, perpetuates AD, or whether it is just an associated factor. Differences in ethical considerations for humans and animals allows for different studies to be carried out in animals, allowing more flexibility in study design and an ability to more easily identify causation.

Finally, further research into how different phenotypes of AD are associated with autoimmunity could provide a strong basis for future clinical research. Patients with AD can be divided into intrinsic and extrinsic categories, as well as other groups, including age of diagnosis and ethnic origin [60]. These groups have different disease characteristics, so more research into the underlying pathogenic mechanisms within these groups, be it related to autoimmunity or other factors, would be useful to aid understanding and potentially improve future treatments.

Table 7 Association between markers of autoimmunity (other than autoantibodies) and atopic dermatitis

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Heratizadeh et al. [54]	Case–control study 30 adult patients with eczema 12 healthy individuals 3 non-atopic patients with chronic plaque psoriasis (Patients with eczema ranged from 18 to 65 years. Insufficient data to calculate mean and SD. No information given about control subjects.)	<i>Clear</i> Hamifin and Rajka	Analysis of peripheral blood mononuclear cells via proliferation rate of blood-derived lymphocytes in response to α -NAC	Median: 9.3% with α -NAC, 0.8% with medium control (both statistically significantly increased with α -NAC compared to control medium ($P < 0.001$), no analysis compared to control group)	Median: 8.4% with α -NAC, with medium control (both statistically significantly increased with α -NAC compared to control medium ($P < 0.001$), no analysis compared to control group)
Hide et al. [47]	Skin Patch Test Case–control study 66 patients with eczema 7 patients with allergic rhinitis 27 healthy control patients (Patients with AD ranged from 1.3 to 37 years, with a mean age of 24.7 ± 5.2 years. Patients with allergic rhinitis ranged from 20 to 35 years, with a mean age of 24.6 ± 4.8 years. Control subjects ranged from 20 to 42 years, with a mean age of 27.8 ± 6.7 years.)	<i>Moderately clear</i> Severity based on grading by Rajka and Langeland	<i>Intradermal injection</i> of autologous sweat	11.1% 3/27 control patients 71.4% 5/7 patients with allergic rhinitis	85% 56/66 eczema patients
Kapitein et al. [53]	Case–control study 28 child patients with AD (of 55) 18 healthy control children (of 30) (Patients with AD ranged from 1.5 to 17.5 years, with a mean age of 8.8 years. Healthy controls ranged from 1.2 to 17.3 years, with a mean age of 8.6 years. Insufficient data to calculate SD.)	<i>Clear</i> Hamifin and Rajka	<i>Stimulation index</i> of peripheral blood mononuclear cells in response to hsp60	— <i>'Significantly higher'</i> ($P = 0.004$)	

Table 7 (continued)

References	Study design and size	Diagnostic criteria for AD	Methods	Prevalence of AR in control subjects	Prevalence of AR in patients with AD
Lind et al. [52] IL-18 and iNKT cells	Case-control study 78 patients with eczema 45 healthy subjects (Patients with eczema ranged from 18 to 65 years, with a mean age of 29 years. Healthy controls ranged from 20 to 64 years with a mean age of 39 years. Insufficient data to calculate SD.)	<i>Clear</i> UK Working Party's Diagnostic Criteria for Atopic Dermatitis (Patients with eczema ranged from 18 to 65 years, with a mean age of 29 years. Healthy controls ranged from 20 to 64 years with a mean age of 39 years. Insufficient data to calculate SD.)	<i>Plasma IL-18 levels and iNKT auto-reactivity</i> 2 part hypothesis that IL-18 can activate iNKT cells 'independently of exogenous ligands' and that IL-18 correlates with AE disease severity	kawamom-	<i>Levels of IL-18 significantly higher (P=0.003) in AE patients than healthy controls</i>
Tanaka et al. [51] Histamine release	Case-control study 61 patients with AD 13 patients with psoriasis 46 healthy control patients (Patients with AD ranged from 2 to 43 years, with a mean age of 24.0 ± 7.5 years. Patients with psoriasis ranged from 29 to 73 years, with a mean age of 53.4 ± 17 years. Healthy controls ranged from 1 to 52 years, with a mean age of 28.4 ± 8.9 years.)	<i>Moderately Clear</i> Severity based on grading by Rajka and Langeland (Patients with AD ranged from 2 to 43 years, with a mean age of 24.0 ± 7.5 years. Patients with psoriasis ranged from 29 to 73 years, with a mean age of 53.4 ± 17 years. Healthy controls ranged from 1 to 52 years, with a mean age of 28.4 ± 8.9 years.)	Measurement of <i>histamine released</i> by basophils upon stimulation by <i>semi-purified sweat antigen</i>	8.7% 4/46 healthy control subjects 0% 0/13 patients with psoriasis	77% 4/761 patients with AD ($P < 0.001$ compared to healthy controls)

Funding There is no funding source.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Wise F, S M (1933) The 1933 year book of dermatology and syphilology. Year Book Publishers, Inc., Chicago
- Darow U, Eyerich K, Ring J (2014) Eczema, Atopic Eczema and Atopic Dermatitis. World Allergy Organisation. http://www.worldallergy.org/professional/allergic_diseases_center/atopic_eczema/. Accessed 26 October 2017
- Williams H, Stewart A, von Mutius E, Cookson W, Anderson HR, Three Study G (2008) International study of A, allergies in childhood phase O. Is eczema really on the increase worldwide? *J Allergy Clin Immunol* 121:947–954
- Odhiambo JA, Williams HC, Clayton TO, Robertson CF, Asher MI, Group IPTS (2009) Global variations in prevalence of eczema symptoms in children from ISAAC Phase Three. *J Allergy Clin Immunol* 124:1251–1258
- Boguniewicz M, Leung DY (2011) Atopic dermatitis: a disease of altered skin barrier and immune dysregulation. *Immunol Rev* 242:233–246
- Hong J, Buddenkotte J, Berger TG, Steinhoff M (2011) Management of itch in atopic dermatitis. *Semin Cutan Med Surg* 30:71–86
- Flohr C, Johansson SG, Wahlgren CF, Williams H (2004) How atopic is atopic dermatitis? *J Allergy Clin Immunol* 114:150–158
- Abeck D, Mempel M (1998) *Staphylococcus aureus* colonization in atopic dermatitis and its therapeutic implications. *Br J Dermatol* 139(Suppl 53):13–16
- Travers JB (2014) Toxic interaction between Th2 cytokines and *Staphylococcus aureus* in atopic dermatitis. *J Invest Dermatol* 134:2069–2071
- Andersen YM, Egeberg A, Gislason GH, Skov L, Thyssen JP (2017) Autoimmune diseases in adults with atopic dermatitis. *J Am Acad Dermatol* 76:274–280
- Bieber T (2010) Atopic dermatitis. *Ann Dermatol* 22:125–137
- Mount S (2014) Atopic Dermatitis Found To Be an Immune-Driven Disease, News Releases. <http://www.mountsinai.org/about-us/newsroom/press-releases/atopic-dermatitis-found-to-be-an-immune-driven-disease>. Accessed 28 October 2017
- Beck LA, Thaci D, Hamilton JD, Graham NM, Bieber T, Rocklin R, Ming JE, Ren H, Kao R, Simpson E, Ardeleanu M, Weinstein SP, Pirozzi G, Guttman-Yassky E, Suarez-Farinias M, Hager MD, Stahl N, Yancopoulos GD, Radin AR (2014) Dupilumab treatment in adults with moderate-to-severe atopic dermatitis. *N Engl J Med* 371:130–139
- Tang TS, Bieber T, Williams HC (2012) Does “autoreactivity” play a role in atopic dermatitis? *J Allergy Clin Immunol* 129:1209–1215
- Moher D, Liberati A, Tetzlaff J, Altman DG, Group P (2009) Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ* 339:b2535
- Du Toit G, Prescott R, Lawrence P, Johar A, Brown G, Weinberg EG, Motala C, Potter PC (2006) Autoantibodies to the high-affinity IgE receptor in children with chronic urticaria. *Ann Allergy Asthma Immunol* 96:341–344
- Aichberger KJ, Mittermann I, Reininger R, Seiberler S, Swoboda I, Spitzauer S, Kopp T, Stingl G, Sperr WR, Valent P, Repa A, Bohle B, Kraft D, Valenta R (2005) Hom s 4, an IgE-reactive autoantigen belonging to a new subfamily of calcium-binding proteins, can induce Th cell type 1-mediated autoreactivity. *J Immunol* 175:1286–1294
- Altrichter S, Kriehuber E, Moser J, Valenta R, Kopp T, Stingl G (2008) Serum IgE autoantibodies target keratinocytes in patients with atopic dermatitis. *J Invest Dermatol* 128:2232–2239
- Guarneri F, Costa C, Foti C, Hansel K, Guarneri C, Guarneri B, Lisi P, Stingeni L (2015) Frequency of autoallergy to manganese superoxide dismutase in patients with atopic dermatitis: experience of three Italian dermatology centres. *Br J Dermatol* 173:559–562
- Hiragun M, Hiragun T, Ishii K, Suzuki H, Tanaka A, Yanase Y, Miura S, Haruta Y, Kohno N, Hide M (2014) Elevated serum IgE against MGL_1304 in patients with atopic dermatitis and cholinergic urticaria. *Allergol Int* 63:83–93
- Ilves T, Virolainen A, Harvima IT (2016) Immediate wheal reactivity to autologous sweat in atopic dermatitis is associated with clinical severity, serum total and specific IgE and sweat tryptase activity. *Int Arch Allergy Immunol* 170:84–91
- Kawamoto N, Yamada A, Ohkouchi S, Maeda T, Tanaka S, Hashimoto T, Saijo Y, Saijo S, Nukiwa T, Shichijo S, Aizawa H, Itoh K (2003) IgG reactive to CTL-directed epitopes of self-antigens is either lacking or unbalanced in atopic dermatitis patients. *Tissue Antigens* 61:352–361
- Kortekangas-Savolainen O, Peltonen S, Pummi K, Kalimo K, Savolainen J (2004) IgE-binding components of cultured human keratinocytes in atopic eczema/dermatitis syndrome and their crossreactivity with *Malassezia furfur*. *Allergy* 59:168–173
- Mittermann I, Reininger R, Zimmermann M, Gangl K, Reisinger J, Aichberger KJ, Greisenegger EK, Niederberger V, Seipelt J, Bohle B, Kopp T, Akdis CA, Spitzauer S, Valent P, Valenta R (2008) The IgE-reactive autoantigen Hom s 2 induces damage of respiratory epithelial cells and keratinocytes via induction of IFN-gamma. *J Invest Dermatol* 128:1451–1459
- Mittermann I, Wikberg G, Johansson C, Lupinek C, Lundeberg L, Cramer R, Valenta R, Scheynius A (2016) IgE sensitization profiles differ between adult patients with severe and moderate atopic dermatitis. *PLoS One* 11:e0156077
- Mothes N, Niggemann B, Jennecke C, Hagemann T, Weidinger S, Bieber T, Valenta R, Novak N (2005) The cradle of IgE autoreactivity in atopic eczema lies in early infancy. *J Allergy Clin Immunol* 116:706–709
- Natter S, Seiberler S, Hufnagl P, Binder BR, Hirschl AM, Ring J, Abeck D, Schmidt T, Valent P, Valenta R (1998) Isolation of cDNA clones coding for IgE autoantigens with serum IgE from atopic dermatitis patients. *FASEB J* 12:1559–1569
- Schmid-Grendelmeier P, Fluckiger S, Disch R, Trautmann A, Wuthrich B, Blaser K, Scheynius A, Cramer R (2005) IgE-mediated and T cell-mediated autoimmunity against manganese superoxide dismutase in atopic dermatitis. *J Allergy Clin Immunol* 115:1068–1075
- Valenta R, Maurer D, Steiner R, Seiberler S, Sperr WR, Valent P, Spitzauer S, Kapiotis S, Smolen J, Stingl G (1996)

- Immunoglobulin E response to human proteins in atopic patients. *J Invest Dermatol* 107:203–208
30. Zeller S, Rhyner C, Meyer N, Schmid-Grendelmeier P, Akdis CA, Crameri R (2009) Exploring the repertoire of IgE-binding self-antigens associated with atopic eczema. *J Allergy Clin Immunol* 124:278–285
31. Bergman R, Ramon M, Wildbaum G, Avitan-Hersh E, Mayer E, Shemer A, Karin N (2009) Psoriasis patients generate increased serum levels of autoantibodies to tumor necrosis factor-alpha and interferon-alpha. *J Dermatol Sci* 56:163–167
32. Neuber K, Mahnss B, Hubner C, Gergely H, Weichenthal M (2006) Autoantibodies against CD28 are associated with atopic diseases. *Clin Exp Immunol* 146:262–269
33. Ochs RL, Muro Y, Si Y, Ge H, Chan EK, Tan EM (2000) Autoantibodies to DFS 70 kd/transcription coactivator p75 in atopic dermatitis and other conditions. *J Allergy Clin Immunol* 105:1211–1220
34. Szakos E, Lakos G, Aleksza M, Hunyadi J, Farkas M, Solyom E, Sipka S (2004) Relationship between skin bacterial colonization and the occurrence of allergen-specific and non-allergen-specific antibodies in sera of children with atopic eczema/dermatitis syndrome. *Acta Derm Venereol* 84:32–36
35. Ambrozic A, Avicin T, Ichikawa K, Kveder T, Matsuura E, Hojnik M, Atsumi T, Rozman B, Koike T (2002) Anti-beta(2)-glycoprotein I antibodies in children with atopic dermatitis. *Int Immunol* 14:823–830
36. El-Rachid RG, Young HS, Griffiths CE, Camp RD (2008) Humoral autoimmune responses to the squamous cell carcinoma antigen protein family in psoriasis. *J Invest Dermatol* 128:2219–2224
37. Ress K, Teesalu K, Annus T, Putnik U, Lepik K, Luts K, Uibo O, Uibo R (2014) Low prevalence of IgA anti-transglutaminase 1, 2, and 3 autoantibodies in children with atopic dermatitis. *BMC Res Notes* 7:310
38. Dhar S, Kanwar AJ, Deodhar SD (1997) Lack of antinuclear antibody in children with atopic dermatitis. *Indian J Dermatol Venereol Leprol* 63:5–8
39. Higashi N, Niimi Y, Aoki M, Kawana S (2009) Clinical features of antinuclear antibody-positive patients with atopic dermatitis. *J Nippon Med Sch* 76:300–307
40. Ohkouchi K, Mizutani H, Tanaka M, Takahashi M, Nakashima K, Shimizu M (1999) Anti-elongation factor-1alpha autoantibody in adult atopic dermatitis patients. *Int Immunol* 11:1635–1640
41. Tada J, Toi Y, Yoshioka T, Fujiwara H, Arata J (1994) Antinuclear antibodies in patients with atopic dermatitis and severe facial lesions. *Dermatology* 189:38–40
42. Taniguchi Y, Yamakami A, Sakamoto T, Nakamura Y, Okada H, Tanaka H, Mizutani H, Shimizu M (1992) Positive antinuclear antibody in atopic dermatitis. *Acta Derm Venereol Suppl (Stockh)* 176:62–64
43. Ress K, Metsküla K, Annus T, Putnik U, Lepik K, Luts K, Uibo O, Uibo R (2015) Antinuclear antibodies in atopic dermatitis: a cross-sectional study on 346 children. *Int J Dermatol* 54:24–28
44. Stalder JF, Taëb A, Atherton DJ, Bieber P, Bonifazi E, Broberg A, Calza A, Coleman R, De Prost Y, Stalder JF, Gelmetti C, Cuanatti A, Harper J, Künz B, Lachapelle JM, Langeland T, Lever R, Oranje AP, Oueille-Roussel C, Revuz J et al (1993) Severity scoring of atopic dermatitis: the SCORAD index. Consensus report of the European task force on atopic dermatitis. *Dermatology* 186:23–31
45. Rajka G, Langeland T (1989) Grading of the severity of atopic dermatitis. *Acta Derm Venereol Suppl (Stockh)* 144:13–14
46. The Cochrane Community (2014) Review Manager (RevMan5). <https://community.cochrane.org/help/tools-and-software/revman-5>. Accessed 26 October 2017
47. Hide M, Tanaka T, Yamamura Y, Koro O, Yamamoto S (2002) IgE-mediated hypersensitivity against human sweat antigen in patients with atopic dermatitis. *Acta Derm Venereol* 82:335–340
48. Kinaciyan T, Natter S, Kraft D, Stingl G, Valenta R (2002) IgE autoantibodies monitored in a patient with atopic dermatitis under cyclosporin A treatment reflect tissue damage. *J Allergy Clin Immunol* 109:717–719
49. Kohsaka T, Hiragun T, Ishii K, Hiragun M, Kamegashira A, Hide M (2018) Different hypersensitivities against homologous proteins of MGL_1304 in patients with atopic dermatitis. *Allergol Int* 67:103–108
50. Lucae S, Schmid-Grendelmeier P, Wuthrich B, Kraft D, Valenta R, Linhart B (2016) IgE responses to exogenous and endogenous allergens in atopic dermatitis patients under long-term systemic cyclosporine A treatment. *Allergy* 71:115–118
51. Tanaka A, Tanaka T, Suzuki H, Ishii K, Kameyoshi Y, Hide M (2006) Semi-purification of the immunoglobulin E-sweat antigen acting on mast cells and basophils in atopic dermatitis. *Exp Dermatol* 15:283–290
52. Lind SM, Kyulenstierna C, Moll M, Winqvist EDJ, Lundeberg O, Karlsson L, Johansson MA, MTL, Scheynius C, Sandberg A, Karlsson JK MC (2009) IL-18 skews the invariant NKT-cell population via autoreactive activation in atopic eczema. *Eur J Immunol* 39:2293–2301
53. Kapitein B, Aalberse JA, Klein MR, de Jager W, Hoekstra MO, Knol EF, Prakken BJ (2013) Recognition of self-heat shock protein 60 by T cells from patients with atopic dermatitis. *Cell Stress Chaperones* 18:87–95
54. Heratizadeh A, Mittermann I, Balaji H, Wichmann K, Niebuhr M, Valenta R, Werfel T (2011) The role of T-cell reactivity towards the autoantigen alpha-NAC in atopic dermatitis. *Br J Dermatol* 164:316–324
55. Zhang BX, Lyu JC, Liu HB, Feng DQ, Zhang DC, Bi XJ, Duan ZW, Ding G (2015) Attenuation of peripheral regulatory T-cell suppression of skin-homing CD8(+)T cells in atopic dermatitis. *Yonsei Med J* 56:196–203
56. Sugiyama H, Gyulai R, Toichi E, Garaczi E, Shimada S, Stevens SR, McCormick TS, Cooper KD (2005) Dysfunctional blood and target tissue CD4 + CD25high regulatory T cells in psoriasis: mechanism underlying unrestrained pathogenic effector T cell proliferation. *J Immunol* 174:164–173
57. Zhang YY, Wang AX, Xu L, Shen N, Zhu J, Tu CX (2016) Characteristics of peripheral blood CD4 + CD25 + regulatory T cells and related cytokines in severe atopic dermatitis. *Eur J Dermatol* 26:240–246
58. Chen B, Ye T, Shao Y, Zhang J, Zhong Q, Hu X, Zhang W, Yu B (2013) Association between copy-number variations of the human histamine H4 receptor gene and atopic dermatitis in a Chinese population. *Clin Exp Dermatol* 38:295–300; quiz 300 – 291
59. Zhang M, Venable JD, Thurmond RL (2006) The histamine H4 receptor in autoimmune disease. *Expert Opin Investig Drugs* 15:1443–1452
60. Bieber T, D'Erme AM, Akdis CA, Traidl-Hoffmann C, Lauener R, Schappi G, Schmid-Grendelmeier P (2017) Clinical phenotypes and endophenotypes of atopic dermatitis: Where are we, and where should we go? *J Allergy Clin Immunol* 139:S58–S64

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.