

RESEARCH ARTICLE

Open Access



Association between plasma fluorescent oxidation products and erectile dysfunction: A prospective study

Shuman Yang¹, Edward Giovannucci^{5,6}, Bruce Bracken², Shuk-Mei Ho^{3,4} and Tianying Wu^{1*}

Abstract

Background: Existing epidemiological studies of the association between oxidative stress and erectile dysfunction (ED) are sparse and inconclusive, which is likely due to cross-sectional design and small sample size. Therefore, we investigated the association between biomarkers of oxidative stress and ED in prospective setting among a relatively large sample size of men.

Methods: We conducted the prospective study among 917 men ages between 47 and 80 years at the time of blood draw, which is a part of nested prospective case-control study of prostate cancer in the Health Professionals Follow-up Study. Plasma fluorescent oxidation products (FLOPs), a global biomarker for oxidative stress, were measured at three excitation/emission wavelengths (360/420 nm named as FLOP_360; 320/420 nm named as FLOP_320 and 400/475 nm named as FLOP_400).

Results: Approximately 35 % of men developed ED during follow-up. We did not find an independent association between FLOP_360, FLOP_320, FLOP_400 and risk of ED in the multivariable adjusted model (Tertile 3 vs. tertile 1: odds ratio [OR] = 0.90, 95 % confidence interval [CI] = 0.61-1.34, $P_{\text{trend}} = 0.54$ for FLOP_360; OR = 0.73, 95 % CI = 0.49-1.07, $P_{\text{trend}} = 0.27$ for FLOP_320; and OR = 0.98, 95 % CI = 0.66-1.45, $P_{\text{trend}} = 0.72$ for FLOP_400). Further analysis of the association between FLOPs and ED in the fasting samples or controls only (free of prostate cancer incidence) did not change the results appreciably.

Conclusions: Plasma FLOPs were not associated with the risk of ED, suggesting oxidative stress may not be an independent risk factor for ED.

Background

Oxidative stress reflects an imbalance between systemic levels of reactive oxygen species (ROS) and host antioxidant defense systems that are able to counteract (detoxify) these ROS. Insufficient antioxidant defense systems against ROS can result in damage to proteins, lipids and DNA in cells and organs in humans. High level of oxidative stress is an important risk factor for many prevalent diseases including cardiovascular disease, breast cancer and reduced renal function [1–3].

The level of plasma fluorescent oxidation products (FLOPs) is a reliable and convenient approach to assess

circulating oxidative stress in large epidemiological studies. One advantage of this marker as compared to other traditional specific oxidation markers (i.e., F₂-isoprostanes and malondialdehyde) is that FLOP assay reflects oxidation pathways from multiple sources including lipid, protein and DNA [4, 5], whereas traditional specific oxidation markers reflect only a portion of oxidative stress. In large observational studies, we found that the level of plasma FLOPs is increased with hypertension, smoking and reduced renal function as defined by reduced levels of glomerular filtration rate [1, 3, 5, 6]. Furthermore, we have documented that the FLOP assay is robust in epidemiologic and clinical setting in which the collection and processing of blood samples cannot be well-controlled. We have found that FLOPs are stable in the blood samples with delayed processing up to 48 h at 4 °C, stable for more than 10 years in plasma samples in liquid-nitrogen freezers, and

* Correspondence: tianying.wu@uc.edu

¹Division of Epidemiology and Biostatistics, Department of Environmental Health, University of Cincinnati Medical Center, Kettering Complex, 3223 Eden Ave, Cincinnati, OH, USA, 45267-0056

Full list of author information is available at the end of the article

highly reproducible over 1–2 year among the same individuals [5, 7].

The association between oxidative stress and erectile dysfunction (ED) is sparse and inconclusive, which is likely due to cross-sectional design and small sample size. It is well known that oxidative stress plays an important role in the development of atherosclerotic diseases [8, 9]. Atherosclerosis reduces cavernosal blood flow, leading to vasculogenic ED [10, 11]. However, existing epidemiological studies of the association between oxidative stress and ED either had small sample size ($N \leq 60$) or were cross-sectional [12–14]. Further, none of above studies adjusted for important potential confounders such as diabetes, hypertension and cigarette smoking which are the risk factors for ED and are important determinants of oxidative stress [5, 15, 16]. Large and prospective studies are warranted to examine the independent relationship between oxidative stress and ED. Therefore, we investigated the association between plasma FLOPs and ED in prospective settings among a relatively large sample size of men.

Methods

Study participants and blood collection

The Health Professionals Follow-up Study (HPFS) initiated in 1986 is an ongoing prospective study of 51,529 men. Between 1993 and 1995, blood collection kits were sent to participants and 18,140 men returned specimens on ice by using an overnight courier. All returned blood samples were processed within 36 h after blood draw and stored in liquid nitrogen freezers. Based on the participants who

donated blood samples, a 1:1 matched nested prospective case–control study of prostate cancer was performed from the time of blood draw [17]. All participants were free of diagnosed cardiovascular diseases and cancers at the time of blood draw. After excluding the ineligible participants (Fig. 1), we finally included 917 men ages between 47 and 80 years (median = 62 years) at the time of blood draw in the prospective study. Among 917 men, 457 and 460 men were subsequent incident prostate cancer cases and controls, respectively. Written informed consent was obtained from all participants. This investigation was approved by Institutional Review Board of the Brigham and Women’s Hospital, the Harvard School of Public Health and the University of Cincinnati.

Measurement of FLOPs

Assay procedure

We measured plasma FLOPs using previously described procedures [5]. In brief, plasma was extracted with ethanol/ether (3:1, v/v) and centrifuged to obtain supernatant. We measured fluorescence of the supernatant at three wavelengths (360/420 nm [excitation/emission] named as FLOP_360, 320/420 nm named as FLOP_320 and 400/475 nm named as FLOP_400). FLOP_360 represents the interaction between lipid oxidation products and proteins, DNA and carbohydrates. FLOP_320 can be produced when oxidation products such as lipid hydroperoxides, aldehydes, and ketones react with DNA in the presence of metals, and FLOP_360 reflects the interaction between malondialdehyde, proteins and phospholipids [18]. The




Fig. 1 Flowchart of the participants excluded from the nested prospective case–control study of prostate cancer in the Health Professional Follow-up Study 1993–2001

within-run average coefficient of variations for FLOP measurements were < 13 %.

Assay stability in blood samples with delayed processing

All blood samples were processed within 36 h after receiving the samples. The delay in processing blood samples up to 36 h appeared to have minimal influence on the measurement of FLOPs. The overall intraclass correlation coefficients (ICCs) of FLOPs were all greater than 0.95 in the shorter- (0 to 24 h) and longer-delayed processing (0 to 36 h) [7].

Assay between- and within- person reproducibility

We conducted a pilot study in 40 participants who donated two blood samples from the Nurse Health Study. After adjusting for fasting status, the ICC of the between- and within- person variations of the FLOPs for repeated measurements over 1.4 year apart (range: 0.8-2.2 years) was 0.44 for FLOP_360, 0.55 for FLOP_320, and 0.70 for FLOP_400 [3].

ED Ascertainment

Between 2000 and 2001, a recall questionnaire in HPFS participants was initiated to rate their ability to have and maintain an erection sufficient for sexual intercourse. The question was "Please rate your ability (without treatment) to have and maintain an erection good enough for intercourse for the following time periods:". Time periods included "before 1986", "1986-1989", "1990-1994", "1995 or later" and "in the last 3 months". Responses of the questionnaire included "very poor", "poor", "fair", "good" and "very good". ED was defined if participants answered "very poor" or "poor". Any ED cases ascertained clearly after blood draw (1993-1995) were considered incident ED. If year of blood draw and year of ED ascertainment may be overlapping (i.e., year of blood draw was 1995 and period of ED ascertainment was '1995 or later'), these ED cases were not defined as incident ED. If the ED cases were not defined as incidence above, these ED cases were used in the cross-sectional analysis only (Fig. 1). Regardless ED cases recovered at a later time or not, they were defined as ED cases in the current study. Whether these cases were included in the cross-sectional or prospective analysis was dependent on the time period of ED ascertainment.

Statistical analyses

To make a comprehensive understanding on relationship between plasma FLOPs and ED, we performed both prospective and cross-sectional analysis (Fig. 1). In the prospective analysis, we analyzed the association between plasma FLOPs and incidence of ED with logistic regression model, but not with Cox proportional hazard model as we were not exactly sure the year of incident ED. The risk of incident ED in lowest tertile of FLOPs was compared with

that in the second and third tertile of FLOPs. The covariates included in the prospective analysis were age (continuous), body mass index (BMI, continuous), alcohol intake (in quartiles: < 0.9, \geq 0.9 and < 6.6, \geq 6.6 and < 17.2, \geq 17.2 g/day), physical activity (in quartiles: < 13.85, \geq 13.85 and < 28.9, \geq 28.9 and < 51.95, \geq 51.95 MET-hours/week), Caucasian (yes/no), fasting hours (continuous), benign prostatic hyperplasia with surgery (yes/no), history of hypertension (yes/no), history of diabetes (yes/no), smoking status (current smokers, past smokers and non-smokers), month of blood draw (in seasons: Spring [March, April and May], Summer [June, July and August], Fall [September, October and November], Winter [December, January and February]) and year of blood draw (1993, 1994 and 1995). To further rule out the potential confounding by fasting status and subclinical factors of prostate cancer incidence, we also examined the association between FLOPs and ED in fasting samples and controls only (free of prostate cancer incidence), respectively. In the cross-sectional analysis, we examined the association between plasma FLOPs and ED at baseline with logistic regression model. The covariates included in the cross-sectional analysis were same as the prospective analysis. All analyses were performed with Statistical Analysis System (Version 9, SAS Institute Inc., Cary, NC).

Results

Baseline characteristics according to plasma FLOP levels

In the prospective analysis, higher levels of plasma FLOP_360, FLOP_320 and FLOP_400 were associated with older age, greater alcohol intake, lower proportion of men who had fasted \geq 8 h before blood draw and great proportion of current and past smokers (Table 1). Higher levels of FLOP_360 were correlated with greater proportion of history of hypertension. Higher levels of FLOP_400 were correlated with greater BMI. When similar analysis was performed in prostate cancer cases and controls separately, the relationship between baseline characteristics and FLOPs in either group was similar to that in the overall samples (Additional file 1: Table S1 and Additional file 2: Table S2). Moreover, the relationship between baseline characteristics in prospective analysis (Table 1) was comparable to that in cross-sectional analysis (Additional file 3: Table S3).

Association between plasma FLOPs and ED

In the prospective analysis, 35 % ($N = 323$) of men were identified having incident ED after blood draw. Although the proportion of incident ED appeared to be higher among men with higher levels of FLOPs (Table 2), we did not find an independent association between FLOP_360, FLOP_320, FLOP_400 and risk of ED in the multivariable adjusted model (Tertile 3 vs. tertile 1: OR = 0.90, 95 % CI = 0.61-1.34, $P_{\text{trend}} = 0.54$ for FLOP_360; OR = 0.73,

Table 1 Baseline characteristics according to tertiles of plasma fluorescent oxidation products (FOPs) ($N = 917$): prospective analysis in the Health Professional Follow-up Study, 1993-1995

Variables	FOP_360			FOP_320			FOP_400		
	1	2	3	1	2	3	1	2	3
Range (FI/ml)	< 184	≥ 184 ; < 233	≥ 233	< 356	≥ 356 ; < 524	≥ 524	< 49.1	≥ 49.1 ; < 62.6	≥ 62.6
N	305	306	306	305	306	306	305	306	306
Age (years)	61.1	61.9	62.3	60.4	62.2	62.6	60.7	62.3	62.2
Body mass index (kg/m ²)	25.8	25.8	25.7	25.5	25.9	25.9	25.4	25.9	26.0
Alcohol intake (g/day) ^a	2.9	8.2	10.0	3.5	8.8	8.6	2.4	8.2	10.4
Physical activity (MET-hours/week) ^a	26.0	33.6	29.1	27.9	34.2	26.5	28.3	29.8	28.1
Caucasians (%)	92	92	95	91	95	93	93	91	95
Fasting status (≥ 8 h; %)	76.4	56.2	50.3	75.7	56.5	50.7	71.2	59.5	52.3
History of BPH with surgery (%)	1.6	3.6	3.3	1.3	3.3	3.9	2.0	3.6	2.9
History of hypertension (%)	23.3	23.9	31.4	20.3	30.4	27.8	21.3	31.1	26.1
History of diabetes (%)	3.6	2.6	2.9	2.6	3.3	3.3	2.3	3.9	2.9
Current smokers (%)	1.1	6.7	13.0	1.1	9.6	9.4	1.1	4.2	16.5
Past smokers (%)	37.8	49.8	60.8	38.6	55.4	54.4	35.1	50.9	62.7

Variables with normal distribution are shown in mean, unless otherwise specified

^aVariables with skew distribution are shown in median

Abbreviations: FOP = Fluorescent oxidation products, FI = Fluorescent intensity unites, MET = Metabolic equivalent, BPH = Benign prostatic hyperplasia

Bold-faced values indicate statistical significance at $P < 0.05$ across tertiles of FOPs

95 % CI = 0.49-1.07, $P_{\text{trend}} = 0.27$ for FOP_320; and OR = 0.98, 95 % CI = 0.66-1.45, $P_{\text{trend}} = 0.72$ for FOP_400).

Further analysis of the association between FOPs and ED in the fasting samples (≥ 8 h) did not change the results appreciably (Tertile 3 vs. tertile 1: OR = 0.91, 95 % CI = 0.56-1.47, $P_{\text{trend}} = 0.69$ for FOP_360; OR = 0.75, 95 % CI = 0.46-1.21, $P_{\text{trend}} = 0.52$ for FOP_320; and OR = 0.83, 95 % CI = 0.51-1.35, $P_{\text{trend}} = 0.53$ for FOP_400).

When we analyzed the relationship between plasma FOPs and risk of ED in controls only (free of prostate cancer incidence), levels of FOP_360, FOP_320 and FOP_400 were again not associated with increased risk of ED (Table 3).

In the cross-sectional analysis, we found 3.7 % ($N = 37$) of men with ED at the time of blood draw. Higher levels of FOP_360 were associated with increased risk of baseline ED (Tertile 3 vs. tertile 1: odds ratio [OR] = 2.68, 95 % confidence interval [CI] = 1.01-7.12), and the relationship had a significant trend ($P_{\text{trend}} = 0.03$). However, higher levels of FOP_320 and FOP_400 were not associated with baseline ED (Additional file 4: Table S4).

Discussion

To our knowledge, this is the first study that comprehensively assessed the association between oxidative stress and ED in both prospective and cross-sectional designs. Although higher levels of FOP_360 were associated with increased risk of ED in the cross-sectional design, none of FOP_360, FOP_320 or FOP_400 was associated with incidence of ED in the prospective design. Since the

relationship between biomarkers of oxidative stress and ED is largely derived from animal models [10], the results of our study have challenged the traditional understandings on the independent detrimental effects of oxidative stress on ED in human.

We only found FOP_360, but not FOP_320 or FOP_400, was positively associated with ED in cross-sectional design. Several reasons may be responsible: First, we cannot fully exclude a possibility of false-positive findings due to the fact that cross-sectional analysis had a small number of ED cases ($N = 37$) and only one type of FOPs (FOP_360 only) was associated with ED. Second, because of the cross-sectional design, the positive relationship between FOP_360 and ED in cross-sectional design may be due to ED-related diseases that are correlated with oxidative stress. Certainly, further studies are warranted to confirm this.

In contrast to the positive association between FOP_360 and ED in the cross-sectional study, we did not show consistent evidence of a positive association between oxidative stress and risk of ED in the prospective design, which had approximately 9 times as many cases as cross-sectional analysis. Besides the much larger number of cases, the prospective design limited reverse causation. However, it is possible that the underlying mechanism of oxidative stress on ED was not due to a global oxidative burden but a specific type of ROSs. There is animal and tissue evidence suggesting that superoxide with nitric oxide can result in acute impairment of cavernosal relaxation but also long-term penile vasculopathy [19-21].

Table 2 Association between plasma fluorescent oxidation products (FIOPs) and erectile dysfunction in prostate cancer cases ($N = 457$) and controls ($N = 460$): prospective analysis in the Health Professional Follow-up Study, 1993-2001

Tertile Variables	1	2	3	<i>P</i> for trend
	FIOP_360			
Range (FI/ml)	< 184	≥ 184; < 233	≥ 233	—
N	305	306	306	—
Median (FI/ml)	160	207	280	—
Erectile dysfunction incidence (n, %)	100 (33 %)	112 (37 %)	111 (36 %)	—
Age adjusted	1 (ref)	1.13 (0.79, 1.61)	1.07 (0.75, 1.52)	0.84
Multivariable adjusted ^a	1 (ref)	1.03 (0.71, 1.51)	0.90 (0.61, 1.34)	0.54
	FIOP_320			
Range (FI/ml)	< 356	≥ 356; < 524	≥ 524	—
N	305	306	306	—
Median (FI/ml)	302	410	1837	—
Erectile dysfunction incidence (n, %)	102 (33 %)	114 (37 %)	107 (35 %)	—
Age adjusted	1 (ref)	1.00 (0.70, 1.42)	0.86 (0.60, 1.23)	0.30
Multivariable adjusted ^a	1 (ref)	0.75 (0.51, 1.11)	0.73 (0.49, 1.07)	0.27
	FIOP_400			
Range (FI/ml)	< 49.1	≥ 49.1; < 62.6	≥ 62.6	—
N	305	306	306	—
Median (FI/ml)	44.0	55.1	72.6	—
Erectile dysfunction incidence (n, %)	100 (33 %)	99 (32 %)	124 (41 %)	—
Age adjusted	1 (ref)	0.82 (0.57, 1.17)	1.24 (0.87, 1.77)	0.14
Multivariable adjusted ^a	1 (ref)	0.67 (0.45, 0.99)	0.98 (0.66, 1.45)	0.72

Values are odds ratio (95 % confidence interval), unless otherwise specified. FI = Fluorescent intensity units

^aRisk factors include age (continuous), body mass index (continuous), alcohol intake (in quartiles: < 0.88, ≥ 0.88 and < 5.58, ≥ 5.58 and < 16.38, ≥ 16.38 g/day), physical activity (in quartiles: < 15.6, ≥ 15.6 and < 30.6, ≥ 30.6 and < 56.9, ≥ 56.9 MET-hours/week), Caucasian (yes/no), fasting hours (continuous), benign prostatic hyperplasia with surgery (yes/no), history of hypertension (yes/no), history of diabetes (yes/no), smoking status (current smokers, past smokers and non-smokers), month of blood draw (in seasons: Spring [March, April and May], Summer [June, July and August], Fall [September, October and November], Winter [December, January and February]) and year of blood draw (1993, 1994 and 1995)

The study has several limitations. First, a single measurement of FIOPs may not accurately reflect the average levels of the biomarker over a prolonged period of time. However, we have assessed their reproducibility over approximately a one-year period, and high ICCs suggest that this marker can be used as a marker for chronic exposure. Second, since we only ascertained ED via a single self-reported questionnaire on ED onset during approximately 7 years after blood draw, we cannot exclude the recall bias as a possible explanation of our results. Furthermore, our assessment of ED has not been validated; however, we note that a prior report using these data [22] were consistent with what has been found in other studies [15, 23]. Nonetheless, these two methodological weaknesses regarding the ascertainment of ED may lead to misclassification between incident ED and healthy controls. Third, our study is limited because the specific oxidative stress level at penile site was not available. Fourth, as our study only included

history of BPH with surgery, some BPH cases with less severe clinical conditions might not be included. Fifth, the bias due to other known (i.e., duration of hypertension and smoking) or unknown residual confounding factors that are related to plasma FIOPs and ED is likely present. Sixth, our study was derived from another study designed to study prostate cancer but not ED. The sampling of that nested case-control study may introduce biases that affected our study and may limit the generation of our results. All the above limitations may cause the null association between plasma FIOPs and incident ED.

The strength of the study is that it contained relatively large sample size of men and incident EDs. In addition, as mentioned above, the prospective study design is a better design than cross-sectional design to reduce the possibility of the reverse causation between oxidative stress and ED, although an effect of subclinical disease on biomarkers cannot be excluded.

Table 3 Association between plasma fluorescent oxidation products (FLOPs) and erectile dysfunction in controls only ($N = 460$): prospective analysis in the Health Professional Follow-up Study, 1993-2001

Tertile	1	2	3	<i>P</i> for trend
Variables	FLOP_360			
Range (FI/ml)	< 184	≥ 184; < 233	≥ 233	—
N	164	144	152	—
Median (FI/ml)	159	206	279	—
Erectile dysfunction incidence (n, %)	55 (34 %)	51 (35 %)	51 (32 %)	—
Age adjusted	1 (ref)	1.06 (0.64, 1.75)	0.90 (0.54, 1.48)	0.59
Multivariable adjusted ^a	1 (ref)	1.02 (0.58, 1.79)	0.83 (0.47, 1.47)	0.46
Variables	FLOP_320			
Range (FI/ml)	< 356	≥ 356; < 524	≥ 524	—
N	160	142	158	—
Median (FI/ml)	297	411	1858	—
Erectile dysfunction incidence (n, %)	52 (33 %)	53 (37 %)	52 (33 %)	—
Age adjusted	1 (ref)	0.96 (0.57, 1.60)	0.73 (0.44, 1.22)	0.17
Multivariable adjusted ^a	1 (ref)	0.75 (0.42, 1.35)	0.55 (0.31, 0.99)	0.10
Variables	FLOP_400			
Range (FI/ml)	< 49.1	≥ 49.1; < 62.6	≥ 62.6	—
N	150	160	150	—
Median (FI/ml)	43.4	54.9	73.3	—
Erectile dysfunction incidence (n, %)	50 (33 %)	48 (31 %)	59 (39 %)	—
Age adjusted	1 (ref)	0.77 (0.46, 1.28)	1.23 (0.74, 2.03)	0.29
Multivariable adjusted ^a	1 (ref)	0.72 (0.41, 1.28)	1.07 (0.60, 1.88)	0.68

Values are odds ratio (95 % confidence interval), unless otherwise specified. FI = Fluorescent intensity units

^aRisk factors include age (continuous), body mass index (continuous), alcohol intake (in quartiles: < 0.88, ≥ 0.88 and < 5.58, ≥ 5.58 and < 16.38, ≥ 16.38 g/day), physical activity (in quartiles: < 15.6, ≥ 15.6 and < 30.6, ≥ 30.6 and < 56.9, ≥ 56.9 MET-hours/week), Caucasian (yes/no), fasting hours (continuous), benign prostatic hyperplasia with surgery (yes/no), history of hypertension (yes/no), history of diabetes (yes/no), smoking status (current smokers, past smokers and non-smokers), month of blood draw (in seasons: Spring [March, April and May], Summer [June, July and August], Fall [September, October and November], Winter [December, January and February]) and year of blood draw (1993, 1994 and 1995)

Conclusions

In conclusion, we found no overall association between plasma FLOPs and risk of ED. This has raised an important message that systemic oxidative stress markers overall may not be a relevant measure for assessing the risk of incident ED. Therefore, the necessity of oxidative stress measurement in the risk of ED assessment is questionable. Since this is the first prospective study on biomarkers of global oxidation only, further investigation of other biomarkers of oxidative stress in relation to ED is warranted.

Additional files

Additional file 1: Table S1. Baseline characteristics according to tertiles of plasma fluorescent oxidation products in the Health Professional Follow-up Study in prostate cancer cases ($N = 457$), 1993–1995. (DOCX 14 kb)

Additional file 2: Table S2. Baseline characteristics according to tertiles of plasma fluorescent oxidation products in the Health Professional Follow-up Study in controls ($N = 460$), 1993–1995. (DOCX 15 kb)

Additional file 3: Table S3. Baseline characteristics according to tertiles of plasma fluorescent oxidation products (FLOPs) ($N = 1,000$): cross-sectional analysis in the Health Professional Follow-up Study, 1993–1995. (DOCX 13 kb)

Additional file 4: Table S4. Association between plasma fluorescent oxidation products (FLOPs) and erectile dysfunction ($N = 1,000$): cross-sectional analysis in the Health Professional Follow-up Study, 1993–1995. (DOCX 15 kb)

Competing interests

The authors declare that they have no competing interests.

Authors' contribution

S.Y. carried out the study design, drafted the manuscript, and analyzed the data. T.W. supervised the study design, manuscript drafting, data collection, data analyses and interpretation. E.G. participated data collection and interpretation, and manuscript revision. B.B. and S.H. helped with data

collection, data interpretation and manuscript revision. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by funded by Dr. Wu's K07-CA138714 from the National Institute of Health and by American Heart Association grant 0430202 N. The HPFS cohort and prostate cancer case-control study was supported by grant Nos. P01CA055075 and P01CA133891 from the National Institute of Health (NIH)/National Cancer Institute.

Author details

¹Division of Epidemiology and Biostatistics, Department of Environmental Health, University of Cincinnati Medical Center, Kettering Complex, 3223 Eden Ave, Cincinnati, OH, USA, 45267-0056. ²Department of Surgery, University of Cincinnati Medical Center, Cincinnati, OH, USA. ³Division of Environmental Genetics and Molecular Toxicology, Cincinnati, OH, USA. ⁴Center for Environmental Genetics, University of Cincinnati Medical Center, Cincinnati, OH, USA. ⁵Departments of Nutrition and Epidemiology, Harvard School of Public Health, Boston, MA, USA. ⁶The Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA.

Received: 22 August 2014 Accepted: 6 August 2015

Published online: 14 August 2015

References

- Rebholz CM, Wu T, Hamm LL, Arora R, Khan IE, Liu Y, et al. The association of plasma fluorescent oxidation products and chronic kidney disease: a case-control study. *Am J Nephrol*. 2012;36(4):297-304.
- Fortner RT, Tworoger SS, Wu T, Eliassen AH. Plasma fluorescent oxidation products and breast cancer risk: repeated measures in the Nurses' Health Study. *Breast Cancer Res Treat*. 2013;141(2):307-16.
- Jensen MK, Wang YS, Rimm EB, Townsend MK, Willett W, Wu TY. Fluorescent Oxidation Products and Risk of Coronary Heart Disease: A Prospective Study in Women. *J Am Heart Assoc*. 2013;2(5).
- Frankel EN. *Lipid oxidation* 2nd edition. Dundee, Scotland: The Oily Press LTD; 2005.
- Wu TY, Willett WC, Rifai N, Rimm EB. Plasma fluorescent oxidation products as potential markers of oxidative stress for epidemiologic studies. *Am J Epidemiol*. 2007;166(5):552-60.
- Wu T, Rifai N, Willett WC, Rimm EB. Plasma fluorescent oxidation products: independent predictors of coronary heart disease in men. *Am J Epidemiol*. 2007;166(5):544-51.
- Wu TY, Rifai N, Roberts LJ, Willett WC, Rimm EB. Stability of measurements of biomarkers of oxidative stress in blood over 36 h. *Cancer Epidem Biomar*. 2004;13(8):1399-402.
- Singh U, Jialal I. Oxidative stress and atherosclerosis. *Pathophysiology*. 2006;13(3):129-42.
- Harrison D, Griendling KK, Landmesser U, Hornig B, Drexler H. Role of oxidative stress in atherosclerosis. *Am J Cardiol*. 2003;91(3):7A-11A.
- Agarwal A, Nandipati KC, Sharma RK, Zippe CD, Raina R. Role of oxidative stress in the pathophysiological mechanism of erectile dysfunction. *J Androl*. 2006;27(3):335-47.
- Azadzoi KM, Schulman RN, Aviram M, Siroky MB. Oxidative stress in arteriogenic erectile dysfunction: prophylactic role of antioxidants. *J Urology*. 2005;174(1):386-93.
- Aldemir M, Okulu E, Neselioglu S, Erel O, Ener K, Kayigil O. Evaluation of serum oxidative and antioxidative status in patients with erectile dysfunction. *Andrologia*. 2012;44 Suppl 1:266-71.
- Barassi A, Colpi GM, Piediferro G, Dogliotti G, D'Eril GV, Corsi MM. Oxidative stress and antioxidant status in patients with erectile dysfunction. *J Sex Med*. 2009;6(10):2820-5.
- Ciftci H, Yeni E, Savas M, Verit A, Celik H. Paraoxonase activity in patients with erectile dysfunction. *Int J Impot Res*. 2007;19(5):517-20.
- Weber MF, Smith DP, O'Connell DL, Patel MI, de Souza PL, Sitas F, et al. Risk factors for erectile dysfunction in a cohort of 108 477 Australian men. *Med J Aust*. 2013;199(2):107-11.
- Feldman HA, Goldstein I, Hatzichristou DG, Krane RJ, Mckinlay JB. Impotence and Its Medical and Psychosocial Correlates - Results of the Massachusetts Male Aging Study. *J Urology*. 1994;151(1):54-61.
- Wu T, Wang Y, Ho SM, Giovannucci E. Plasma levels of nitrate and risk of prostate cancer: a prospective study. *Cancer epidemiol, biomarkers prev* : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology. 2013;22(7):1210-8.
- Farankel EN. *Lipid oxidation*. Dundee: The Oily Press; 1998.
- Jeremy JY, Jones RA, Koupparis AJ, Hotston M, Persad R, Angelini GD, et al. Reactive oxygen species and erectile dysfunction: possible role of NADPH oxidase. *Int J Impot Res*. 2007;19(3):265-80.
- Burnett AL, Lowenstein CJ, Bredt DS, Chang TS, Snyder SH. Nitric oxide: a physiologic mediator of penile erection. *Science*. 1992;257(5068):401-3.
- Jones RW, Rees RW, Minhas S, Ralph D, Persad RA, Jeremy JY. Oxygen free radicals and the penis. *Expert Opin Pharmacother*. 2002;3(7):889-97.
- Li Y, Batoool-Anwar S, Kim S, Rimm EB, Ascherio A, Gao X. Prospective study of restless legs syndrome and risk of erectile dysfunction. *Am J Epidemiol*. 2013;177(10):1097-105.
- Selvin E, Burnett AL, Platz EA. Prevalence and risk factors for erectile dysfunction in the US. *Am J Med*. 2007;120(2):151-7.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

