

openheart Efficacy of interventions to increase physical activity for people with heart failure: a meta-analysis

Aliya Amirova ^{1,2}, Theodora Fteropoulli,^{3,4} Paul Williams,³ Mark Haddad⁵

► Additional online supplemental material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/openhrt-2021-001687>).

To cite: Amirova A, Fteropoulli T, Williams P, *et al.* Efficacy of interventions to increase physical activity for people with heart failure: a meta-analysis. *Open Heart* 2021;**8**:e001687. doi:10.1136/openhrt-2021-001687

Received 13 April 2021
Accepted 29 April 2021



© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹School of Health Sciences, City University of London, London, UK

²Department of Practice and Policy, Centre for Behavioural Medicine, UCL, London, UK

³Health Services Research and Management, School of Health Sciences, City University of London, London, UK

⁴Medical School, University of Cyprus, Nicosia, Cyprus

⁵Health Services Research and Management, City University of London, London, UK

Correspondence to

Dr Aliya Amirova; a.amirova@ucl.ac.uk

ABSTRACT

Objectives This meta-analysis aims to (1) evaluate the efficacy of physical activity interventions in heart failure and (2) to identify intervention characteristics significantly associated with the interventions' efficacy.

Methods Randomised controlled trials reporting intervention effects on physical activity in heart failure were combined in a meta-analysis using a random-effect model. Exploratory meta-analysis was performed by specifying the general approach (eg, cardiac rehabilitation), strategies used (eg, action planning), setting (eg, centre based), mode of delivery (eg, face to face or online), facilitator (eg, nurse), contact time and behavioural change theory use as predictors in the random-effect model.

Results Interventions (n=21) had a significant overall effect (SMD=0.54, 95% CI (0.13 to 0.95), p<0.0005). Combining an exercise programme with behavioural change intervention was found efficacious (SMD=1.26, 95% CI (0.26 to 2.26), p<0.05). Centre-based (SMD=0.98, 95% CI (0.35 to 1.62), and group-based (SMD=0.89, 95% CI (0.29 to 1.50),) delivery by a physiotherapist (SMD=0.84, 95% CI (0.03 to 1.65),) were significantly associated with efficacy. The following strategies were identified efficacious: prompts/cues (SMD=3.29, 95% CI (1.97 to 4.62)), credible source (standardised mean difference, SMD=2.08, 95% CI (0.95;3.22)), adding objects to the environment (SMD=1.47, 95% CI (0.41 to 2.53)), generalisation of the target behaviour (SMD=1.32, 95% CI (0.22 to 2.41)), monitoring of behaviour by others without feedback (SMD=1.02, 95% CI (0.05 to 1.98)), self-monitoring of outcome(s) of behaviour (SMD=0.79, 95% CI (0.06 to 1.52)), graded tasks (SMD=0.73, 95% CI (0.22 to 1.24)), behavioural practice/rehearsal (SMD=0.72, 95% CI (0.26 to 1.18)), action planning (SMD=0.62, 95% CI (0.03 to 1.21)) and goal setting (behaviour) (SMD=0.56, 95% CI (0.03 to 1.08)).

Conclusion The meta-analysis suggests intervention characteristics that may be suitable for promoting physical activity in heart failure. There is moderate evidence in support of an exercise programme combined with a behavioural change intervention delivered by a physiotherapist in a group-based and centre-based settings.

PROSPERO registration CRD42015015280.

Key questions

What is already known about this subject?

► Individuals diagnosed with heart failure (HF) are advised to engage in physical activity. However, physical activity levels remain extremely low in this population group. Cardiac rehabilitation (CR) is routinely offered to newly diagnosed HF patients. CR is multifaceted; It is unknown which specific components result in physical activity improvements once the programme has ended. It is essential to understand how best to improve everyday physical activity engagement in HF.

What does this study add?

► This meta-analysis assessed what constitutes a successful physical activity intervention designed for individuals living with HF. The findings pinpoint specific intervention features and components that contribute to physical activity improvements in HF. Centre-based interventions that are delivered by a physiotherapist, in group format, which combine exercise with behavioural change intervention are promising for attaining physical activity improvements.

How might this impact on clinical practice?

► The findings of this meta-analysis may inform physical activity intervention designed for individuals diagnosed with HF. There is a need for additional training for physiotherapists in delivering behavioural change interventions alongside an exercise programme that includes the identified efficacious strategies.

INTRODUCTION

The levels of engagement in physical activity of medically stable individuals diagnosed with heart failure (HF) are low.¹ Physical activity is a treatment strategy.² Cardiac rehabilitation (CR) and other exercise-based programmes have been shown to improve quality of life (QoL)^{3,4} and reduce hospitalisation in HF.^{4,5} However, a recent meta-analysis suggested that CR is less likely to be efficacious in sustaining physical activity in HF in particular compared with other cardiovascular diseases (CVD).^{6,7} The uptake of CR remains suboptimal.⁷

Therefore, it is essential to evaluate the efficacy of alternative interventions as well as CR and identify content and features that are likely to be successful in promoting physical activity.

CR is a complex intervention. It is unclear which components are responsible for what outcomes and for which patient group.⁸ There is a need to explore this intervention complexity and identify what makes an intervention successful.⁹ Past reviews have suggested that short-term intervention effects are associated with strategies such as exercise prescription; goal setting; feedback and problem solving; and the use of a behavioural change theory.¹⁰

CR might be missing some efficacious elements. Clark *et al* pointed out that previous healthcare services research has not emphasised CR's goal: how best to ensure that CVD patients benefit from a healthy lifestyle, including physical activity. Clark *et al* also made a call to evaluate a range of potentially effective interventions that are facilitated by various professionals and make use of a diverse set of methods (eg, remote monitoring). Evaluation of home-based and remote interventions is especially vital, given the recent restriction following the SARS-CoV-2 outbreak. It is also essential to understand what features of centre-based, group-based interventions contribute to physical activity improvement. The present meta-analysis of randomised controlled trials (RCTs) reviewed physical activity interventions, including CR, to identify intervention features that contribute to efficacy in improving physical activity.

METHODS

Information sources

The review protocol was registered on PROSPERO database (CRD42015015280). Cochrane Library, MEDLINE, CINAHL, EMBASE, AMED, HEED, PsycArticles, PsycINFO, Global Health, Web of Science: Conference Proceedings, 'Be Part of Research' and ClinicalTrials.gov were searched from inception to 20 February 2020. The search strategy is described in online supplemental material 1.

Eligibility criteria and study selection

Titles, abstracts and full texts were independently screened by two reviewers (AA and PW). The criteria for considering RCTs were: (1) adults diagnosed with HF, (2) intervention targeting physical activity (compared with usual care and/or education), and (3) report of a numerical result for physical activity outcome at intervention completion for both groups. Physical activity outcome was defined as any bodily movement produced by skeletal muscles that requires energy expenditure. Exercise is a subset of physical activity defined as structured physical activity.¹¹ Exercise, in the context of HF, is defined as selfcare behaviour (ie, 'I exercise regularly').

Data collection process

Relevant information was extracted from trial reports (article, online supplemental materials and protocols) using a standardised Cochrane data extraction form.¹²

Risk of bias in individual studies

The risk of bias at the study level was assessed using the Cochrane Collaboration Risk of Bias tool^{2 13} and informed sensitivity analysis.

Data items

Interventions were classified in terms of their general approach to physical activity promotion (eg, exercise), setting (eg, home vs centre), mode of delivery (eg, group vs individual) and facilitator (eg, nurse). The Theory Coding Scheme (TCS)¹⁴ was used to describe the extent to which trials employed a behavioural change theory in the intervention design. TCS scores range from 0 (no theory) to 8 (most extensive theory use). The intervention and comparator treatment were described in terms of the included behavioural change techniques. Interventions' content was independently annotated by AA (100%) and TF (61.90%) using the Behaviour Change Techniques Taxonomy (BCITv1).¹⁵

Statistical analysis

Meta-analysis was performed using the metafor library in R.¹⁶ A random-effect model was used to estimate the overall efficacy of interventions using restricted maximum likelihood. The standardised mean difference (SMD) in physical activity levels between the main intervention and the comparator group was selected as the estimate of efficacy. Heterogeneity index (I^2) was reported as the total unexplained variability in effect. Assessments at the 3 months, 6 months (short-term) and 12 months (long-term) follow-up were included. Meta-regression was performed to explore whether the efficacy was associated with the following: general approach (eg, exercise programme), setting, mode of delivery (eg, home-based), facilitator (eg, nurse), behavioural change strategies (eg, goal setting) and participant characteristics (ie, mean age, New York Heart Association (NYHA) class, proportion of males, mean ejection fraction (EF, %), aetiological aetiology (%)) were specified as predictors in the model.¹⁷ We accounted for the fact that a small number of trials were presenting a particular intervention characteristic using Hartung-Knapp-Sidik adjustment as recommended by Debray *et al*.¹⁸

Risk of bias across studies

The small study bias was evaluated using a funnel plot assessment and Egger's test.


Patient and public involvement

No patients were involved in formulating the research question, the outcome measures or findings interpretation. Patients were not involved in planning or designing of the meta-analysis. This is due to the lack of funding available to include patients as partners in this meta-analysis. Results of this meta-analysis will be disseminated to the relevant patient organisations.

RESULTS

Search results

Search results and reasons for exclusion are listed in the Preferred Reporting Items for Systematic Reviews and



*Authors were contacted with a request to share numerical results on physical activity outcome

Figure 1 The study flow chart (PRISMA, 2009). HF, heart failure; PRISMA, Preferred Reporting Items for Systematic Reviews and Meta-Analyses.

Meta-Analyses diagram (figure 1). A total of 20 trials evaluating 22 interventions postcompletion (n=21,^{19–37}), at 6 months (n=5,^{29 32 33 35 38}) and 12 months (n=5,^{26 27 29 36–38}) follow-up were included in the meta-analysis.

Study characteristics

The trials were conducted between 1999 and 2018. The trials included a total of 6277 participants, and the median sample size was 100 (IQR: 60–204). A large proportion (37%) of participants were drawn from the HF-ACTION trial (n=2331).³⁷

RISK OF BIAS

The overall risk of bias is summarised in figure 2. Six out of 20 trials reported low risk of bias.^{20 22 26 27 29 37} A high risk of bias was present in two trials.^{19 30} The sources of

bias for each trial are summarised in online supplemental material 2. Five trials evaluated the intervention against an active comparator: education.^{19 21 22 27 38}

Participant characteristics

Mean age ranged from 54¹⁹ to 80 years old³³ (SD=7.28; IQR 62–70), and the majority of the sample was male 69.49% (table 1).

Postcompletion efficacy

The present meta-analysis found a significant overall effect as assessed at postcompletion (SMD=0.54, 95% CI (0.13 to 0.95), p<0.005). There was significant high heterogeneity in the estimated effect, I²=95.8%, (Q=1531.74, p<0.001) (figure 3). The following intervention characteristics contributed to the heterogeneity in efficacy:




Figure 2 The risk of bias summary. The Cochrane Collaboration Risk of Bias tool (V.2).¹⁵

general approach of the interventions, setting (ie, centre based vs home based), facilitator and several strategies.

General approach

The included trials delivered interventions that were classified as exercise (k=10, 47.62%), exercise and behavioural change (k=3, 14.29%), motivational interviewing (k=2, 9.42%), remote communication and treatment (k=3, 14.29%), cognitive-behavioural therapy (k=1, 4.76%), disease management (k=1, 4.76%) and self-management (k=1, 4.76%) (table 2). Exercise combined with behavioural change is an efficacious approach (figure 4).

Intervention strategies

A total of 38 strategies (ie, behaviour change techniques, (BCTs)) were present across included trials (online supplemental material 4). Interventions included a mean of 8.90 (SD=3.77; IQR=8–10) strategies. The following strategies were associated with moderate to large effects: prompts/cues, credible source, adding objects to the environment, generalisation of target behaviour, monitoring of behaviour by others without feedback, self-monitoring of outcome(s) of behaviour, graded tasks, behavioural practice/rehearsal, action planning, goal setting (behaviour) (SMD: 0.56–3.29) (table 3).

Settings, facilitator and duration

Interventions were delivered at home (n=8, 38%), in a hospital/clinic (n=8, 38%), or both (n=5, 24%). Only centre-based delivery significantly moderated the efficacy of the included interventions (table 3). Interventions were facilitated by general practice nurses (n=9, 42.85%), physiotherapists (n=6, 28.6%), HF nurses (n=4, 19%), exercise instructors (n=3, 14.29%), researchers (n=2, 9.42%), lay leaders (n=1, 4.76%), advanced practice nurse (n=1, 4.76%), psychologists (n=1, 4.76%), and clinical psychology trainees (n=1, 4.76%). Intervention delivery by a physiotherapist was associated with efficacy

(table 3). Intervention duration varied from 1 day to 72 weeks. Mean contact time was 1849.38 min (SD=1716.40) and was not associated with intervention efficacy.

Theory use

Seven interventions were based on a behavioural change theory (online supplemental material 4). The extent of theory use (TCS) was not associated with efficacy (SMD=0.13, p=0.059, 95% CI (–0.006 to 0.27)).

Sample characteristics, including mean age, gender, mean left-ventricular ejection fraction (LVEF, %), New York Heart Association (NYHA) class, and aetiology, were not significantly associated with intervention efficacy (online supplemental material 5). Likewise, the differences in efficacy between trials using self-reports and trials using accelerometer or pedometer were non-significant (online supplemental material 5).

Long-term efficacy

The included interventions assessed physical activity at a 2-month, 6-month, 12-month and 30-month follow-up. The overall short-term effect was non-significant at the 6 month, (SMD=0.06, 95% CI (–0.49 to 0.38), p=0.80) and 12-month follow-up, (SMD=–0.11, 95% CI (–0.77 to 0.55), p=0.80). Due to the small number of interventions reporting follow-up assessment, it was not feasible to evaluate the long-term effects associated with the individual intervention characteristics.

Sensitivity analysis

Interventions were compared with usual care,^{20 23–26 28–30 33–37} education delivered by an HF specialist nurse^{21 22 27 38} or unspecified health professional,¹⁹ and discouragement to exercise.³² The comparator treatments included a mean of 1.15 (SD=1.49) strategies (online supplemental material 4). When trials comparing the main intervention to education were excluded, the effects of exercise and behavioural change, remote monitoring and treatment, and

Table 1 Study and participant characteristics

Author, year	Country	No of participants		Intervention	Assessment time points		Mean age, years (SD)	Male, %	LVEF, % (SD)	NYHA II-III, %	Physical activity† outcome
		Control	Intervention		Postintervention*	Follow-up†					
Ajiboye <i>et al.</i> ¹⁹ 2015	Nigeria	23	28	–	12 weeks	–	54 (1.6)	53.7	–	–	Self-reported physical activity (diary)
Bernocchi <i>et al.</i> ²⁰ 2018	Italy	56	56	2 months	4 months	2 months	71(9)	88	44.5 (12.4)	45	Accelerometer (Average energy activity counts)
Boynse <i>et al.</i> ²¹ 2014	Netherlands	185	197	1 year	–	–	71 (11.9)	58	<40	29; 21	Exercise: Heart Failure Selfcare, Behaviour Scale (Jaarsma, Strömberg, Mårtensson, & Dracup, 2003)
Brodie <i>et al.</i> ²² 2005; 2008	UK	32	30 (MI and UC);30 (MI)	8 weeks	–	–	79 (6.9)‡	–	31.3 (5.9)§	28; 58	Self-reported general physical activity (Booth <i>et al.</i> , 1996)
Collins <i>et al.</i> ²³ 2004	UK	16	15	24 weeks	–	–	62.7 (11.2)	100	31.7 (6.9)	–	Self-reported physical activity (diary)
Convera-Tindell <i>et al.</i> ²⁴ 2004	USA	42	37	12 weeks	–	–	63.8 (10.1)	99	29.1 (8.5)	80; 20	Pedometer
Cowie <i>et al.</i> ²⁵ 2013	UK	20	20 (Home); 20 (Hospital)	8 weeks	–	–	66 (35–85)¶	85	<40	62; 38	Accelerometer (Average energy activity counts)
Dalal <i>et al.</i> ²⁶ 2019	UK	92	93	12 weeks	–	–	69.7 (10.9)	78	34.5 (25–39)¶¶	59	Accelerometer (Average energy activity counts)
Freedland <i>et al.</i> ²⁷ 2015	USA	60	58	6 months	12 months	12 months	55.8 (11.2)‡	53.8	38.9 (15.5)‡	42.4	Accelerometer (Average energy activity counts)
Jolly <i>et al.</i> ²⁸ 2009	UK	85	84	24 weeks	12 months	12 months	65.9 (12.5)	76	<40	75; 20	Self-reported physical activity (minutes per week, Godin Leisure-Time week, Godin Leisure-Time Exercise Questionnaire, (Godin & Shephard, 1985))
Meng <i>et al.</i> ²⁹ 2013; 2016	Germany	227	248	–	6; 12 months	6; 12 months	61.2 (11.7)	75	31.7 (7.0)	54.7	Self-reported physical activity (minutes per week, Godin Leisure-Time Exercise Questionnaire, (Godin & Shephard, 1985))
O'Connor <i>et al.</i> ³⁰ 2009	USA (88.72%) Canada (8.07%) France (3.33%)	1172	1159	24 weeks	12; 30 months	12; 30 months	59.2 (51.2–67.8)¶¶	72	<35	62; 36	Self-reported physical activity (diary)
Pozzelli <i>et al.</i> ³¹ 2018	USA	102	102	18 months	–	–	60.4 (11.5)‡	56.0	39.4 (12.7)‡	91.2	Self-reported physical activity
Smeulders <i>et al.</i> ²⁹ 2009	Netherlands	131	186	6 weeks	6; 12 months	6; 12 months	66.6 (11.0)	75.8	<40	64; 36	Self-reported physical activity
Tomita <i>et al.</i> ³⁰ 2008	USA	13	19	1 year	–	–	74.2 (9.7)	32.5	–	75; 25	Self-reported physical activity
van den Berg-Emons <i>et al.</i> ³¹ 2004	Netherlands	16	18	12 weeks	–	–	58.6 (12.1)	81	23.9 (9.4)	56; 44	Accelerometer (Average energy activity counts)

Continued

Table 1 Continued

Author, year	Country	No of participants		Assessment time points			Mean age, years (SD)	Male, %	LVEF, % (SD)	NYHA II-III, %	Physical activity† outcome
		Control	Intervention	Postintervention*	Follow-up†						
Willenheimer <i>et al.</i> ³² 1998	Sweden	27	23	16 weeks	6 months	64 (5)	70	0.35 (0.11)	50; 36	Self-reported physical activity (The total physical activity score calculated using the following formula: Time spent on each activity per week intensity*2/100)	
Witham <i>et al.</i> ³³ 2005	UK	41	41	12 weeks	6 months	80 (6)	63	–	61; 39	Accelerometer (Average energy activity counts)	
Yeh <i>et al.</i> ³⁴ 2004	USA	50	50	12 weeks	–	68.1 (11.9)	56	28.3 (8.0)	62; 18	Self-reported physical activity (CHAMPS; Stewart 2001)	
Young <i>et al.</i> ³⁵ 2015; 2016	USA	49	51	3 months	6 months	68.7 (11.8)	47.1	53.4 (12.9)	29.4; 56.9	Accelerometer (Average energy activity counts)	
Meta-analysis sample:		N (Control)=2555	N (intervention)=3722			66 years IQR:(62–70)	Male (69.49 % IQR:(56–78)				

Physical activity outcome was defined as any bodily movement produced by skeletal muscles that requires energy expenditure (WHO, 2018). Exercise is a subset of physical activity defined as structured physical activity (WHO, 2018). In the context of HF, exercise is also defined as self-care behaviour (ie, 'I exercise regularly').

*Time from baseline.
 †Time from intervention completion.
 ‡Mean for the total sample.
 §Group 1: MI + UC; - not reported.
 ¶Median age (range).
 HF, heart failure; MI, Motivational Interviewing; UC, usual care.




Figure 3 Forest plot illustrating overall estimated effect (SMD) and 95% CI and SMD and 95% CI for component trials. SMD, standardised mean difference.

exercise were significant (online supplemental material 6). The exclusion of a large (N=2331) trial with a younger (56 years old) sample³⁷ resulted in a significant decrease in the overall effect. The effect estimates for exercise and behavioural change approach, prompts/cues, credible source, adding objects to the environment, generalisation of the target behaviour, monitoring of behaviour by others without feedback, self-monitoring of outcome(s) of behaviour, action planning, and goal setting (behaviour) were sensitive to the inclusion of the trial (online supplemental material 6). The exclusion of interventions with a high risk of bias indicated that the efficacy of *Exercise* approach was overestimated. The effects of the following strategies were underestimated: social support (emotional), social support (practical), theory use (TCS score), information about health consequences and information on how to perform the behaviour.

Small study bias

A funnel plot for SMD against SE is available in online supplemental material 7. The Egger's test suggested a lack of publication and small study bias (test for funnel plot asymmetry: $Z = 0.46$, $p = 0.46$)

DISCUSSION

The present meta-analysis found moderate evidence in support of existing physical activity interventions designed for individuals living with HF. Centre-based interventions that are delivered by a physiotherapist, in group format, which combine exercise with behavioural change intervention are promising for attaining physical activity improvements. Intervention strategies identified as efficacious are: prompts/cues, credible source, adding objects to the environment, generalisation of the target behaviour, monitoring of behaviour by others without feedback, self-monitoring of outcome(s) of behaviour, graded tasks, behavioural practice/rehearsal, action planning, and goal setting (behaviour). To our knowledge, this is the first meta-analysis evaluating the components of behavioural interventions that are associated with increased physical activity in HF. Interventions that were delivered by a physiotherapist in a centre-based setting were more promising in attaining physical activity improvement than home-based interventions or those delivered by facilitators other than physiotherapist (ie, nurse, lay leader, researcher). This is in contrast to the findings of a previous meta-analysis suggesting that

Table 2 Intervention characteristics

Author, year		Intervention description provided by authors	General approach
Ajiboye <i>et al</i> , ¹⁹ 2015	Main intervention	Aerobic and resistance training and education	Exercise
	Comparator treatment	Usual care and education	
Bernocchi <i>et al</i> , ²⁰ 2018	Main intervention	Telerehabilitation and home-based personalised exercise maintenance programme	Remote communication and treatment
	Comparator treatment	Usual care	
van den Berg-Emons <i>et al</i> , ³¹ 2004	Main intervention	Aerobic exercise training	Exercise
	Comparator treatment	Usual care without particular advice for exercise	
Boyne <i>et al</i> , ²¹ 2014	Main intervention	Individually tailored e-health intervention 'Health Buddy'.	Remote communication and treatment
	Comparator treatment	Education	
Brodie <i>et al</i> , ²² 2005	Main intervention 1	Motivational Interviewing	Motivational interviewing
	Main intervention 2	Motivational Interviewing and education	
	Comparator treatment	Education	
Collins <i>et al</i> , ²³ 2004	Main intervention	Aerobic exercise training	Exercise
	Comparator treatment	Usual care	
Corvera-Tindel <i>et al</i> , ²⁴ 2004	Main intervention	A home walking exercise programme	Exercise
	Comparator treatment	Usual care	
Cowie <i>et al</i> , ²⁵ 2011	Main intervention 1	Hospital-based aerobic exercise training	Exercise
	Main intervention 2	Home-based exercise training	
	Comparator treatment	Usual care	
Dalal <i>et al</i> , ²⁶ 2019 (REACH-HF)	Main intervention	Rehabilitation enablement in HF: self-care and rehabilitation	Exercise and behavioural change
	Comparator treatment	Usual care	
Freedland <i>et al</i> , ²⁷ 2018	Main intervention	Integrative cognitive-behavioural therapy and education and usual care	Cognitive-behavioural therapy
	Comparator treatment	Enhanced (with education) usual care	
O'Connor <i>et al</i> , ³⁷ 2009 (HF-ACTION)	Main intervention	Aerobic exercise training and Exercise adherence facilitation intervention	Exercise and behavioural change
	Comparator treatment	Usual care	
Jolly <i>et al</i> , ³⁶ 2009	Main intervention	Aerobic and resistance exercise training	Exercise
	Comparator treatment	HF specialist nurse care	
Meng <i>et al</i> , ³⁸ 2016	Main intervention	Self-management patient education programme and CR	Self-management
	Comparator treatment	Education	
Pozehl <i>et al</i> , ²⁸ 2018 (HEART Camp)	Main intervention	Multicomponent intervention and resistance exercise training	Exercise and behavioural change
	Comparator treatment	Enhanced (nine exercise sessions for 3 months)	
Smeulders <i>et al</i> , ²⁹ 2009	Main intervention	Chronic disease management programme	Disease management
	Comparator treatment	Usual care	
Tomita <i>et al</i> , ³⁰ 2008	Main intervention	Multidisciplinary internet-based programme on HF management	Remote communication and treatment
	Comparator treatment	Usual care	
Willenheimer <i>et al</i> , ³² 2001	Main intervention	Aerobic exercise training	Exercise
	Comparator treatment	Usual care and discouragement to exercise	
Witham <i>et al</i> , ³³ 2005	Main intervention	Seated aerobic and resistance exercise training	Exercise

Continued

Table 2 Continued

Author, year	Intervention description provided by authors	General approach
	Comparator treatment	Usual care
Yeh <i>et al.</i> ³⁴ 2011	Main intervention	Exercise training (Tai Chi Mind-Body movement)
	Comparator treatment	Usual care
Young <i>et al.</i> ⁸⁵ 2015; 2016	Main intervention	Patient Activation Intervention on self-management in HF
	Comparator treatment	Usual care

CR, cardiac rehabilitation; HF, heart failure.




Figure 4 Forest plot illustrating the standardised mean differences (SMD, 95% CI) moderated by the general approach.

Table 3 Intervention characteristics associated with efficacy

Intervention characteristics	SMD	95% CI
Behavioural change techniques:		
Prompts/cues	3.29	(1.97 to 4.62)
Definition: Introduce or define environmental or social stimulus to promote or cue the behaviour. Examples: frequent phone calls by a health professional/ post or email reminders		
Credible source	2.08	(0.95 to 3.22)
Definition: resent verbal or visual communication from a credible source in favour of or against the behaviour. Examples: Explicit, detailed and salient advice from a health professional to engage in physical activity.		
Adding objects to the environment	1.47	(0.41 to 2.53)
Definition: Add objects to the environment in order to facilitate the performance of the behaviour. Examples: Provision of a treadmill, weights, step, or stationary bicycle.		
Generalisation of the target behaviour	1.32	(0.22 to 2.41)
Definition: Advice to perform the desired behaviour, which is already performed in a particular situation, in another situation. Examples: Encouragement to engage in an exercise in home settings.		
Monitoring of behaviour by others without feedback	1.02	(0.05 to 1.98)
Definition: Observe or record behaviour with the person's knowledge as part of a behavioural change strategy. Examples: The physiotherapist informs participants that their physical activity levels will be monitored using accelerometers and telemonitoring devices.		
Self-monitoring of outcome(s) of behaviour	0.79	(0.06 to 1.52)
Definition: Establish a method for the person to monitor and record the outcome(s) of their behaviour as part of a behavioural change strategy. Examples: Monitoring reduced pain symptoms and dyspnoea as a result of physical activity.		
Graded tasks	0.73	(0.22 to 1.24)
Definition: Set easy-to-perform tasks, making them increasingly difficult, but achievable until the behaviour is performed. Examples: Gradual increase in the level of exertion as assessed using the Borg scale.		
Behavioural practice/rehearsal	0.72	(0.26 to 1.18)
Definition: Prompt practice or rehearsal of the performance of the behaviour one or more times in a context or at a time when the performance may not be necessary. Examples: Exercise training (individual or in a group).		
Action planning	0.62	(0.03 to 1.21)
Definition: prompt, detailed planning of performance of the behaviour (must include at least one of context, frequency, duration and intensity). Examples: plan when, where, how much and at what intensity the participant will perform the exercise.		
Goal setting (behaviour)	0.56	(0.03 to 1.08)
Definition: set or agree on a goal defined in terms of the behaviour to be achieved. Examples: Set a goal to complete 30 min of exercise (brisk walking) at the vigorous intensity in future.		
Setting: Centre-based interventions	0.98	(0.35 to 1.62)
Mode of delivery: Group-based interventions	0.89	(0.29 to 1.50)
Facilitator: Physiotherapist	0.84	(0.03 to 1.65)

Definitions are from Michie *et al.*¹⁵ Intervention characteristics are described in [table 2](#) and online supplemental material 4. SMD and 95% CI for characteristics that were not suggested to be significantly associated with efficacy are summarised in online supplemental material 4.

SMD, standardised mean difference.

centre-based and home-based programmes delivered to individuals post-myocardial infarction or revascularisation, and with HF are equivalent in their efficacy in improving survival, QoL and exercise capacity.³⁹ The present meta-analysis found that delivery of an intervention to a group contributed to efficacy. However, given the ongoing pandemic, it is essential to optimise delivery of physical activity interventions in home settings. Group-based interventions contribute to behavioural change via social comparison, changes in normative beliefs about health behaviour and group member identity.⁴⁰ These factors can also be considered when designing

home-based, contact-free physical activity interventions for older adults with HF.

A previous systematic review of CR programmes found that, in general, educational and behavioural elements of CR did not result in physical activity improvements beyond those achieved by exercise-based programmes.⁶ However, behavioural elements are diverse and vary in their efficacy. The present meta-analysis evaluated a range of such elements and outlined those that are efficacious. A combination of an exercise and behavioural change approach was found to be more efficacious than other approaches, including exercise alone. Several

strategies to improve physical activity appear promising (table 3). Theoretical explanations for the efficacy of these strategies were previously offered.⁴¹ Graded tasks exert an effect on physical activity by fostering positive beliefs about capability through skill mastery (eg, exercise training).⁴¹ Self-monitoring, monitoring by others, planning, goal-setting and feedback are theorised to improve control and regulation of behaviour.⁴² Finally, the efficacy of adding an object associated with physical activity (eg, treadmill) indicates the relevance of cueing (ie, automatic association and non-deliberate regulation of behaviour).⁴¹

Implications for clinical practice and future research

The present meta-analysis found moderate evidence in support of combining exercise programme with behavioural change intervention, delivered by a physiotherapist. Thus, there is a need for additional training for physiotherapists in delivering behavioural change interventions that will include the identified efficacious strategies. Practical limitations of the identified efficacious strategy need to be considered when designing interventions. Adding objects to the environment to support physically active lifestyle (eg, a treadmill) may not be affordable or practical, and does not satisfy the principle of health equity.⁴³ In addition, further research investigating how best to promote a physically active lifestyle in the older HF population is encouraged. The clinical profiles of older adults differ from younger adults, with a significantly worse prognosis and a larger number of comorbidities in the former.⁴⁴ Older adults may also differ in their beliefs about physical activity; and strategies that are suited for promoting an active lifestyle in older adults are different to those that are efficacious for the general population.⁴⁵ Investigation of which behavioural change theory should form the basis for an intervention is also warranted. Only five trials assessed physical activity at 6-month and 12-month follow-ups. Long-term efficacy was not supported. Thus, it is important to investigate how sustained physical activity improvements can be established.

Study-level limitations

High risk of bias was observed in two trials.^{21 32} The sensitivity analysis indicated that the inclusion of these trials may overestimate the efficacy of exercise programmes and underestimate the efficacy of remote monitoring and treatment. Remote communication and feedback interventions that include strategies such as biological feedback (eg, symptom monitoring and feedback) delivered by a nurse using telehealth device, as well as self-monitoring of the behaviour and information about health consequences^{22 23} are identified as efficacious when high risk of bias trials are excluded. The HF-AC-TION³⁷ trial constituted the majority of the meta-analysis sample and when it was excluded in the sensitivity analysis, only a small non-significant effect of exercise combined with behavioural change was observed. High-quality trials

assessing the short and long-term effects of behavioural change; remote communication and treatment; and exercise programmes on physical activity in older adults (>70 years old) with HF are required.

Strengths and limitations of the review

A Cochrane overview of reviews recommended exploring intervention complexity using meta-regression to evaluate the association between intervention characteristics and efficacy.⁹ This meta-analysis identified, annotated and classified behavioural change interventions in terms of their general approach, strategies, settings, facilitator, delivery mode, duration and use of theory; and using meta-regression assessed the association between these characteristics and the efficacy. The clear, consistent and systematic description of the interventions facilitated the reliable grouping and analysis. This helped pinpoint specific efficacious features and elements that can be applied, either as part of CR or otherwise, to improve physical activity outcomes in HF. However, there are a few limitations. Intervention features were present in clusters across the included trials. Given the small number of RCTs evaluating any single included characteristic, multiple comparisons were not feasible. It is not possible to ascertain whether each of the evaluated features is efficacious on their own or only in combination. These features need to be evaluated in a multiarm trial comparing their effects.

CONCLUSIONS

This meta-analysis explored intervention complexity and identified some features of potentially promising physical activity interventions designed for people living with HF. The present review provides moderate evidence that an exercise programme combined with a behavioural change intervention is a promising approach to increasing physical activity in HF. The meta-analysis suggests behavioural change strategies that may be useful in promoting physical activity in HF.

Twitter Theodora Fteropoulli @tzellofouska

Acknowledgements We would like to thank the authors of the included studies for providing the information required for the detailed description of the interventions and data analysis. Many thanks to Professor Gary Morgan at City, the University of London, for his support throughout the review. We thank Dr Raissa Jessica Aquino for her comments on the earlier version of data extraction (excluded at the later stage of the review). We also thank Dr Michael J. Shoemaker (Grand Valley State University) for sharing details on one of the pilot studies he and his colleagues carried out (excluded at the later stage of the review).

Contributors All authors contributed to design and planning. AA and PW contributed by screening articles for inclusion. AA and TF contributed to data extraction and risk of bias assessment. AA implemented the analysis. AA, TF and MH contributed to reporting. TF and MH contributed to the critical revision of the manuscript.

Funding Funding The review was supported by a City, University of London Postgraduate Studentship Grant.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement All data relevant to the study are included in the article or uploaded as online supplemental information.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Aliya Amirova <http://orcid.org/0000-0001-6476-7604>

REFERENCES

- Jaarsma T, Strömberg A, Ben Gal T, *et al.* Comparison of self-care behaviors of heart failure patients in 15 countries worldwide. *Patient Educ Couns* 2013;92:114–20.
- National Institute for Healthcare and Excellence. Overview: chronic heart failure in adults: diagnosis and management guidance, 2018. Available: <https://www.nice.org.uk/guidance/ng106> [Accessed 23 Mar 2020].
- Sagar VA, Davies EJ, Briscoe S, *et al.* Exercise-based rehabilitation for heart failure: systematic review and meta-analysis. *Open Heart* 2015;2:e000163.
- Taylor RS, Long L, Mordi IR, *et al.* Exercise-Based Rehabilitation for Heart Failure: Cochrane Systematic Review, Meta-Analysis, and Trial Sequential Analysis. *JACC Heart Fail* 2019;7:691–705.
- Lewinter C, Doherty P, Gale CP, *et al.* Exercise-based cardiac rehabilitation in patients with heart failure: a meta-analysis of randomised controlled trials between 1999 and 2013. *Eur J Prev Cardiol* 2015;22:1504–12.
- Dibben GO, Dalal HM, Taylor RS, *et al.* Cardiac rehabilitation and physical activity: systematic review and meta-analysis. *Heart* 2018;104:1394–402.
- Santiago de Araújo Pio C, Chaves GS, Davies P, *et al.* Interventions to promote patient utilisation of cardiac rehabilitation. *Cochrane Database Syst Rev* 2019;2:CD007131.
- Clark AM, Redfern J, Briffa T. Cardiac rehabilitation: fit to face the future? *Heart* 2014;100:355–6.
- Anderson L, Taylor RS. Cardiac rehabilitation for people with heart disease: an overview of Cochrane systematic reviews. *Cochrane Database Syst Rev* 2014:CD011273.
- Tierney S, Mamas M, Woods S, *et al.* What strategies are effective for exercise adherence in heart failure? A systematic review of controlled studies. *Heart Fail Rev* 2012;17:107–15.
- Physical activity. Available: <https://www.who.int/news-room/fact-sheets/detail/physical-activity> [Accessed 01 Mar 2020].
- Higgins JP, Green S. *Cochrane Handbook for systematic reviews of interventions, version 5.1.0*. 2011. London: The Cochrane Collaboration, 2017.
- Cochrane Methods. Risk of bias 2 (rob 2) tool. Available: <https://methods.cochrane.org/risk-bias-2> [Accessed 01 Mar 2020].
- Michie S, Prestwich A. Are interventions theory-based? Development of a theory coding scheme. *Health Psychol* 2010;29:1–8.
- Michie S, Richardson M, Johnston M, *et al.* The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: building an international consensus for the reporting of behavior change interventions. *Ann Behav Med* 2013;46:81–95.
- Viechtbauer W. Conducting meta-analyses in R with the metafor package. *J Stat Softw* 2010;36.
- Borenstein M, Hedges LV, Higgins JPT. *Introduction to meta-analysis*. Chichester, UK: John Wiley & Sons, Ltd, 2009.
- Debray TPA, Damen JAAG, Snell KIE, *et al.* A guide to systematic review and meta-analysis of prediction model performance. *BMJ* 2017;356:i6460.
- Ajiboye OA, Anigbogu CN, Ajuluchukwu JN. Exercise training improves functional walking capacity and activity level of Nigerians with chronic biventricular heart failure. *Hong Kong Physiotherapy Journal* 2015;33:42–9.
- Bernocchi P, Vitacca M, La Rovere MT, *et al.* Home-based telerehabilitation in older patients with chronic obstructive pulmonary disease and heart failure: a randomised controlled trial. *Age Ageing* 2018;47:82–8.
- Boyne JJJ, Vrijhoef HJM, Spreeuwenberg M, *et al.* Effects of tailored telemonitoring on heart failure patients' knowledge, self-care, self-efficacy and adherence: a randomized controlled trial. *Eur J Cardiovasc Nurs* 2014;13:243–52.
- Brodie DA, Inoue A. Motivational interviewing to promote physical activity for people with chronic heart failure. *J Adv Nurs* 2005;50:518–27.
- Collins E, Langbein WE, Dilan-Koetje J, *et al.* Effects of exercise training on aerobic capacity and quality of life in individuals with heart failure. *Heart Lung* 2004;33:154–61.
- Corvera-Tindel T, Doering LV, Woo MA, *et al.* Effects of a home walking exercise program on functional status and symptoms in heart failure. *Am Heart J* 2004;147:339–46.
- Cowie A, Thow MK, Granat MH, *et al.* A comparison of home and hospital-based exercise training in heart failure: immediate and long-term effects upon physical activity level. *Eur J Cardiovasc Prev Rehabil* 2011;18:158–66.
- Dalal HM, Taylor RS, Jolly K, *et al.* The effects and costs of home-based rehabilitation for heart failure with reduced ejection fraction: the REACH-HF multicentre randomized controlled trial. *Eur J Prev Cardiol* 2019;26:pp.:262–2.
- Freedland KE, Carney RM, Rich MW, *et al.* Cognitive behavior therapy for depression and self-care in heart failure patients: a randomized clinical trial. *JAMA Intern Med* 2015;175:1773–82.
- Pozehl BJ, McGuire R, Duncan K, *et al.* Effects of the HEART cAMP trial on adherence to exercise in patients with heart failure. *J Card Fail* 2018;24:654–60.
- Smeulders ESTF, van Haastregt JCM, Ambergen T, *et al.* The impact of a self-management group programme on health behaviour and healthcare utilization among congestive heart failure patients. *Eur J Heart Fail* 2009;11:609–16.
- Tomita MR, Tsai B-M, Fisher NM, *et al.* Improving adherence to exercise in patients with heart failure through internet-based self-management. *J Am Geriatr Soc* 2008;56:1981–3.
- van den Berg-Emons R, Balk A, Bussmann H, *et al.* Does aerobic training lead to a more active lifestyle and improved quality of life in patients with chronic heart failure? *Eur J Heart Fail* 2004;6:95–100.
- Willenheimer R, Erhardt L, Cline C, *et al.* Exercise training in heart failure improves quality of life and exercise capacity. *Eur Heart J* 1998;19:774–81.
- Witham MD, Gray JM, Argo IS, *et al.* Effect of a seated exercise program to improve physical function and health status in frail patients > or = 70 years of age with heart failure. *Am J Cardiol* 2005;95:1120–4.
- Yeh GY, McCarthy EP, Wayne PM, *et al.* Tai chi exercise in patients with chronic heart failure: a randomized clinical trial. *Arch Intern Med* 2011;171:750–7.
- Young L, Hertzog M, Barnason S. Effects of a home-based activation intervention on self-management adherence and readmission in rural heart failure patients: the PATCH randomized controlled trial. *BMC Cardiovasc Disord* 2016;16:176.
- Jolly K, Taylor RS, Lip GYH, *et al.* A randomized trial of the addition of home-based exercise to specialist heart failure nurse care: the Birmingham rehabilitation uptake Maximisation study for patients with congestive heart failure (BRUM-CHF) study. *Eur J Heart Fail* 2009;11:205–13.
- O'Connor CM, Whellan DJ, Lee KL, *et al.* Efficacy and safety of exercise training in patients with chronic heart failure: HF-ACTION randomized controlled trial. *JAMA* 2009;301:1439–50.
- Meng K, Musekamp G, Schuler M, *et al.* The impact of a self-management patient education program for patients with chronic heart failure undergoing inpatient cardiac rehabilitation. *Patient Educ Couns* 2016;99:1190–7.
- Anderson L, Sharp GA, Norton RJ, *et al.* Home-based versus centre-based cardiac rehabilitation. *Cochrane Database Syst Rev* 2017;6:CD007130.
- Borek AJ, Abraham C, Greaves CJ, *et al.* Identifying change processes in group-based health behaviour-change interventions: development of the mechanisms of action in group-based interventions (MAGI) framework. *Health Psychol Rev* 2019:1–21.
- Connell LE, Carey RN, de Bruin M, *et al.* Links between behavior change techniques and mechanisms of action: an expert consensus study. *Ann Behav Med* 2019;53:708–20.
- Michie S, Abraham C, Whittington C, *et al.* Effective techniques in healthy eating and physical activity interventions: a meta-regression. *Health Psychol* 2009;28:690–701.
- WHO. World conference on social determinants of health. Available: <https://www.who.int/sdhconference/en/> [Accessed 28 Dec 2020].
- Butrous H, Hummel SL. Heart failure in older adults. *Can J Cardiol* 2016;32:1140–7.
- French DP, Olander EK, Chisholm A, *et al.* Which behaviour change techniques are most effective at increasing older adults' self-efficacy and physical activity behaviour? A systematic review. *Ann Behav Med* 2014;48:225–34.