

Article

Serum 25-Hydroxyvitamin D Concentrations Are Inversely Correlated with Hepatic Lipid Content in Male Collegiate Football Athletes

Xiaomin Sun ^{1,2,3}, Zhen-Bo Cao ^{4,*} , Kumpei Tanisawa ^{3,5}, Satomi Oshima ³ and Mitsuru Higuchi ³

¹ School of Public Health, Xi'an Jiaotong University Health Science Center, Xi'an 710061, China; gzhtxiaomin@xjtu.edu.cn

² Global Health Institute, Xi'an Jiaotong University Health Science Center, Xi'an 710061, China

³ Faculty of Sport Sciences, Waseda University, Tokorozawa, Saitama 359-1192, Japan; tanisawa@nibiohn.go.jp (K.T.); msatomi3939@yahoo.co.jp (S.O.); mhiguchi@waseda.jp (M.H.)

⁴ School of Kinesiology, Shanghai University of Sport, 399 Chang Hai Road, Shanghai 200438, China

⁵ Department of Physical Activity Research, National Institutes of Biomedical Innovation, Health and Nutrition, Tokyo 162-8636, Japan

* Correspondence: caozb_edu@yahoo.co.jp; Tel.: +86-21-5125-3257; Fax: +86-21-5125-3380

Received: 1 June 2018; Accepted: 20 July 2018; Published: 21 July 2018



Abstract: Lower serum 25-hydroxyvitamin D (25(OH)D) concentrations are associated with more weight and fat mass gain in adults in the general population, but it is unknown whether this is the case in collegiate football athletes with greater body weight. This study aimed to investigate associations of serum 25(OH)D concentrations with body fat and ectopic fat accumulation, and to determine which fat indicators are closely related to serum 25(OH)D in male collegiate football athletes. Thirty-four collegiate athletes aged 21 years were recruited. Serum 25(OH)D concentrations and the levels of visceral fat area (VFA), vastus lateralis intramyocellular lipid (IMCL), extramyocellular lipid (EMCL), and intrahepatic lipid (IHCL) were measured. Serum 25(OH)D concentrations were negatively associated with the IHCL values ($r = -0.372$, $p = 0.030$), and the relationship remained after adjustment for several factors ($r = -0.378$, $p = 0.047$). Additionally, multiple stepwise regression analysis of IHCL content as the dependent variable indicated that 25(OH)D concentrations were a stronger predictor of IHCL content ($\beta = -0.363$, $p = 0.030$) than % body fat and $VO_{2peakFFM}$. Higher serum 25(OH)D concentrations are more closely related to lower IHCL content rather than any other fat indicators, suggesting that increasing serum 25(OH)D concentrations may have some effect that inhibits lipid accumulation in hepatic tissue, especially in heavy athletes.

Keywords: abdominal body fat; American football; ectopic fat; rugby football; vitamin D

1. Introduction

Obesity has long been officially recognized as a key factor for the incidence of cardiovascular diseases [1]. In addition to the lipid accumulation in adipose tissue, the accumulation of lipid in non-adipose tissues (such as liver and muscle), also known as ectopic lipid accumulation, also has been associated with the development of cardiovascular diseases [2].

Young collegiate athletes undertaking regular and intensive exercise have presumably been considered to have a very low risk of cardiovascular and metabolic diseases; however, the clustering of metabolic risk factors has been recognized, especially in young heavy athletes [3–5], who often intentionally try to gain weight by increasing energy intake, where weight gain is associated with increases in body fat content and in risks of cardiovascular diseases [6,7]. Guo et al. [3] found that

professional athletes of strength sports in the heavier weight class usually have a higher prevalence of metabolic syndrome and its individual components, including central obesity. The same results were also observed in Japanese collegiate heavyweight Judo athletes and National Football League players [4,5].

Circulating 25-hydroxyvitamin D (25(OH)D) is the main circulating metabolite of vitamin D, as well as the form used for the assessment of vitamin D status. Over the last few decades, vitamin D receptors (VDRs) were also detected in many different tissues, including skeletal muscle, adipose tissue, cardiac muscle, and immune tissue [8]. Available data indicate that low circulating 25(OH)D concentrations are common in obesity, and negative relationships with body mass index (BMI), percent body fat (% body fat), waist circumference (WC), visceral fat area (VFA), and subcutaneous fat area (SFA) have been reported in adolescents, young and healthy middle-aged individuals, and old adults [9–12]. Numerous studies found that higher levels of body fat usually leads to lower 25(OH)D concentrations [13,14]. Recently, it was observed that vitamin D may inhibit lipid accumulation through VDRs or calcium ions [15,16]. Epidemiologic evidence indicates that higher serum 25(OH)D concentrations are associated with less weight and fat mass gain in the future [17], and vitamin D treatments could reduce hepatic triglyceride accumulation in mice [18,19].

However, limited evidence shows that these associations also exist in athletes [20]. Heller et al. [21] found that serum 25(OH)D concentrations were negatively associated with total body fat indicators, such as BMI and % body fat, even after controlling for sex in collegiate athletes. Additionally, Barcal et al. [22] found a negative relationship between serum 25(OH)D concentrations and % body fat in collegiate male wrestlers, regardless of seasonal changes. Recently, in addition to body fat tissue, circulating 25(OH)D concentrations have been negatively associated with ectopic lipid content in adults in the general population [23–25]. Although excessive ectopic fat accumulation, for example, increased skeletal muscle intramyocellular lipid (IMCL), has been independently associated with insulin resistance in obese and type 2 diabetes subjects, athletes who are highly insulin sensitive have similar muscle fat content to that observed in insulin-resistant subjects [26]. This metabolic conundrum has been termed “athletes’ paradox”. However, to date, the relationships between 25(OH)D concentrations and ectopic lipid are still unclear in collegiate football athletes.

Therefore, the primary aims of this study were to evaluate whether body fat or ectopic fat indicators are closely related to serum 25(OH)D concentrations in overweight male collegiate football athletes.

2. Materials and Methods

Subjects

Thirty-four male collegiate athletes aged 21 (range, 20–21) years participated in the current study in summer (from June to July), which included 13 collegiate American football athletes (10 linemen, 3 backs) and 21 collegiate rugby football athletes (12 forwards, 9 backs). The levels of % body fat, SFA, VFA, and VO_2 peak of one subject in the rugby football group were unobtainable for personal reasons. One person’s dietary data was excluded from analysis because his total energy intake was more than 9000 kcal.

All athletes were members of the American and rugby football teams in the East Japan College Football Association division 1. All participants had >3 years of training experience playing their type of football and performed resistance exercise at least 5 times per week for about 2 h per day. All participants were in good health and free from chronic diseases. Athletes were excluded if they were regularly taking vitamin D supplements or used sunscreen. All participants provided written informed consent before study enrollment, and the Ethical Committee of Waseda University approved this study (ethical approval code: 2011-060; June 2013). The study was conducted in accordance with the Declaration of Helsinki.

3. Procedures

3.1. Anthropometric Measurements

Body weight was measured using an electronic scale (Inner Scan BC-600; Tanita Inc., Tokyo, Japan), whereas height was measured using a stadiometer (YL-65; YAGAMI Inc., Nagoya, Japan), with subjects wearing minimal clothing and no shoes. BMI was calculated using the body weight and height measurements. WC was measured at the umbilical region with an inelastic measuring tape at the end of normal expiration to the nearest 0.1 cm. Dual-energy X-ray absorptiometry (DXA) (Hologic QDT-4500, DXA Scanner; Hologic Inc., Waltham, MA, USA) was used to measure the % body fat. The subjects were asked to wear loose-fitting, light clothes without any metal objects and to lie in a supine position on the scanning table during the total body fat scan.

3.2. Magnetic Resonance Imaging and Spectroscopy

The VFA and SFA were measured by magnetic resonance imaging as described previously [12]. The imaging conditions included a T1-weighted spin-echo and axial plane sequence with a slice thickness of 10 mm, a repetition time (TR) of 140 milliseconds, and an echo time (TE) of 12.3 milliseconds. The cross-sectional areas of the VFA and SFA at the umbilical level were determined using image analysis software (Slice-o-matic 4.3 for Windows; Tomovision, Montreal, QC, Canada). The coefficient of variation for the cross-sectional area at the umbilical level was 0.4%.

Hepatic ^1H magnetic resonance spectroscopy (MRS) was performed using a 1.5-T whole-body scanner (Signa 1.5 T; General Electric, Inc., Milwaukee, WI, USA) with an 8-channel body array coil. A single voxel ($30 \times 30 \times 20 \text{ mm}^3$) for spectroscopy was selected in the right lobe of the liver while avoiding major blood vessels. Voxel shimming was performed to optimize the homogeneity of the magnetic field. The proton spectra of the liver were acquired using the point-resolved spectroscopy technique (repetition time = 2500 milliseconds; echo time = 35 milliseconds; 64 measurements; 1024 sample points). Acquisition was synchronized to the respiratory cycle and triggered at the end of expiration. MRS data were quantified using LCModel version 6.3 (Stephen Provencher, Oakville, ON, Canada). Intrahepatic lipid (IHCL) content was defined as the signal intensities at 1.3 and 1.6 ppm relative to the signal intensity of water at 4.7 ppm. The coefficient of variation between measurements was 7.5%.

Vastus lateralis intramyocellular lipid (IMCL) and extramyocellular lipid (EMCL) were assessed by ^1H -MRS with a flexible surface coil wrapped around the participant's upper leg. Care was taken to avoid vascular structures and adipose tissue deposits within the voxel. The T1 magnetic resonance images were acquired by using a point-resolved spectroscopy sequence with the following acquisition parameters: repetition time 2000 milliseconds; echo time 35 milliseconds; field of view, 200 mm. The voxel volume was ($20 \times 20 \times 20 \text{ mm}^3$). The water signal was suppressed by using chemically selective saturation.

3.3. Analysis of Blood Samples

Blood samples were collected in Venoject-II AutoSep tubes (gel, 9 mL; Terumo Corporation, Tokyo, Japan) from an antecubital vein of the forearm between 0830 and 0930 h after a 12-h overnight fast and then centrifuged at $3000 \times g$ for 15 min at 4°C . Serum samples were stored at -80°C until the time of analysis. Serum enzymatic activities of aspartate aminotransferase (AST), alanine aminotransferase (ALT) and γ -glutamyl transferase (γ -GTP) were determined using the ultraviolet method (BML Inc., Tokyo, Japan), respectively. Serum 25(OH)D concentrations were measured in duplicate with commercially available enzyme-linked immunosorbent assay kits (25(OH)D; Immundiagnostik AG, Bensheim, Germany) according to the manufacturer's instructions. The Control samples provided by the Immundiagnostik AG group were analyzed with each run for quality control. The intra-assay coefficient of variation was 8.9% for 25(OH)D.

3.4. Cardiorespiratory Fitness

Cardiorespiratory fitness was assessed using a cycle ergometer (Ergomedic 828E; Monark, Varberg, Sweden), and the peak oxygen uptake was quantified as VO_2peak . The graded cycle exercise began at a workload of 45–90 W and was increased by 15 W/min until the subject could no longer maintain the required pedaling frequency of 60 rpm. Heart rate and ratings of perceived exertion were monitored each minute during exercise. Additional details were published elsewhere [12]. To correct for fat mass, the peak oxygen uptake per kilogram of fat free mass ($\text{VO}_2\text{peak}_{\text{FFM}}$) was calculated.

3.5. Dietary Intake

Information on total energy intake, carbohydrate, protein, fat, vitamin D and calcium intakes were analyzed using the computerized nutritional analysis system of the Food Frequency Questionnaire Based on Food Groups (Kenpakusha, Japan) [27].

3.6. Statistical Analysis

All statistical analyses were performed with SPSS software, version 22.0 (SPSS, Inc., Chicago, IL, USA). The Kolmogorov-Smirnov test was performed to assess the normality of data distribution. The Student *t* test (for normally distributed variables) or Mann-Whitney *U* test (for non-normally distributed variables) was used to evaluate the differences in these variables according to American football and rugby football groups. Pearson correlation coefficients were computed between serum 25(OH)D and body indicators. Linear multiple regression analysis was performed to assess the association of serum 25(OH)D with the fat indicators, adjusted for group, position and total energy intake. All measurements and calculated values are presented as mean \pm standard deviation (for normally distributed variables) or median (interquartile range) (for non-normally distributed variables). The statistical significance level was set at $p < 0.05$.

4. Results

Subject characteristics according to study groups are presented in Table 1 and Figure 1. The prevalence of vitamin D deficiency (<50 nmol/L) and vitamin D insufficiency (50–75 nmol/L) were 15.4% and 53.8% in American football collegiate athletes, and 19.0% and 61.9% in rugby football collegiate athletes, respectively. American collegiate football athletes had higher values of weight, WC, % body fat, and SFA than rugby football athletes ($p < 0.05$). No difference was observed in $\text{VO}_2\text{peak}_{\text{FFM}}$, IHCL, IMCL, EMCL, AST, ALT, γ -GTP, 25(OH)D, total energy, and other nutrients intake.

Table 1. Participant characteristics according to groups.

Variable	Total (n = 34)	American Football (n = 13)	Rugby Football (n = 21)
Age (years)	21.0 (20.0–21.0)	21.0 (20.0–22.0)	21.0 (20.0–21.0)
Height (cm)	175.0 ± 4.8	176.1 ± 4.5	174.4 ± 5.0
Weight (kg)	90.4 ± 10.9	95.5 ± 9.3	87.3 ± 10.8 *
BMI (kg/m ²)	29.5 ± 3.0	30.7 ± 2.0	28.7 ± 3.3
WC (cm)	90.3 ± 8.0	94.8 ± 6.4	87.5 ± 7.8 *
% Body fat	16.6 ± 4.1	19.4 ± 2.6	14.8 ± 3.9 ^{a,*}
SFA (cm ²)	168.0 (89.5–222.0)	207.0 (170.0–275.0)	112.5 (72.3–183.0) ^{a,*}
VFA (cm ²)	82.8 ± 32.4	95.1 ± 31.6	74.8 ± 31.2 ^a
VO ₂ peak _{FFM} (mL/kg/min)	48.7 ± 5.6	46.5 ± 5.4	50.1 ± 5.4 ^a
IHCL	0.037 (0.017–0.082)	0.047 (0.023–0.177)	0.024 (0.008–0.064)
IMCL	0.013 ± 0.006	0.014 ± 0.008	0.012 ± 0.005
EMCL	0.017 (0.012–0.030)	0.019 (0.012–0.032)	0.015 (0.012–0.028)
AST (IU/L)	24.5 (21.8–36.0)	25.0 (22.5–34.0)	24.0 (21.0–36.5)
ALT (IU/L)	34.1 ± 16.4	37.4 ± 20.8	32.1 ± 13.1
γ-GTP (IU/L)	28.0 (24.5–38.0)	29.0 (24.0–35.0)	27.0 (24.0–43.0)
25(OH)D (nmol/L)	61.3 (54.2–74.7)	62.3 (53.9–79.0)	60.3 (52.8–73.0)
Total energy intake (kcal)	3895.8 (3534.5–4365.0)	4071.7 (3662.8–5089.1)	3840.7 (3476.1–4232.5) ^b
Carbohydrate intake (g)	511.6 (461.9–621.0)	522.6 (479.3–658.9)	501.0 (450.2–571.1) ^b
Fat intake (g)	130.9 ± 35.0	143.0 ± 32.0	123.1 ± 35.3 ^b
Protein intake (g)	143.2 ± 33.8	155.0 ± 37.5	135.5 ± 29.6 ^b
Vitamin D intake (µg/day)	10.4 ± 4.1	11.2 ± 4.9	9.7 ± 12.4 ^b
Calcium intake (mg/day)	1185.4 ± 442.0	1284.8 ± 402.3	1120.8 ± 464.3 ^b

Data are presented as mean ± SD or median (IQR) values. BMI, body mass index; WC, waist circumference; SFA, subcutaneous fat area; VFA, visceral fat area; VO₂peak, peak oxygen uptake; IHCL, intrahepatic lipid; IMCL, intramyocellular lipid; EMCL, extramyocellular lipid; AST, aspartate aminotransferase; ALT, alanine aminotransferase; γ-GTP, γ-glutamyl transferase; 25(OH)D, 25-hydroxyvitamin D. ^a n = 20; ^b n = 20, one person was excluded from the analysis because his total energy was >9000 kcal in rugby football; * p < 0.05 vs. American football players.




Figure 1. Prevalence of vitamin D deficiency and insufficiency.

The correlations between serum 25(OH)D concentrations and body fat, ectopic lipid variables and other factors are showed in Table 2 and Figure 2A–D. Serum 25(OH)D concentrations were significantly and negatively correlated with the IHCL values ($r = -0.372, p = 0.030$). No significant associations were observed among serum 25(OH)D concentrations with IMCL, EMCL, and other variables ($p > 0.05$). After adjusting by study group, position, and total energy intake, the significant relationship between serum 25(OH)D concentrations and IHCL was still observed ($r = -0.378, p = 0.047$), whereas the other measures became insignificant ($p > 0.05$). In the subgroup analysis, serum 25(OH)D was observed to be negatively correlated with γ -GTP ($r = -0.504, p = 0.033$) and IHCL ($r = -0.462, p = 0.053$) in rugby football players after adjusting by total energy intake; whereas, no relationships were found in American football players. In addition, multiple stepwise regression analysis with IHCL content as the dependent variable indicated that the 25(OH)D concentrations was a stronger predictor of the IHCL content ($\beta = -0.363, p = 0.030$), independent of the % body fat and $VO_{2peakFFM}$, and after adjustment for study group, position, and total energy intake.

Table 2. Correlations between 25(OH)D concentrations and other variables.

Variables	25(OH)D		25(OH)D *	
	<i>r</i>	<i>p</i> -Value	<i>r</i>	<i>p</i> -Value
BMI (kg/m ²)	-0.231	0.189	-0.012	0.952
WC (cm)	-0.279	0.110	-0.200	0.307
% Body fat	-0.036	0.841	0.164	0.405
SFA (cm ²)	-0.138	0.444	-0.026	0.895
VFA (cm ²)	-0.307	0.083	-0.218	0.265
IHCL	-0.372	0.030	-0.378	0.047
IMCL	0.032	0.860	-0.080	0.686
EMCL	0.019	0.913	0.125	0.526
ALT (IU/L)	-0.098	0.583	-0.048	0.806
AST (IU/L)	0.174	0.324	0.217	0.268
γ -GTP (IU/L)	-0.297	0.088	-0.317	0.100

Data are presented as Pearson’s coefficients. BMI, body mass index; WC, waist circumference; SFA, subcutaneous fat area; VFA, visceral fat area; IHCL, intrahepatic lipid; IMCL, intramyocellular lipid; EMCL, extramyocellular lipid; AST, aspartate aminotransferase; ALT, alanine aminotransferase; γ -GTP, γ -glutamyl transferase; 25(OH)D, 25-hydroxyvitamin D. 25(OH)D, IHCL, EMCL, AST, and γ -GTP were log-transformed; SFA was square root transformed for analysis. Bold font indicates significance ($p < 0.05$). * Data are adjusted by group, position, and total energy intake.




Figure 2. Associations among serum 25(OH)D with VFA (A), EMCL (B), IMCL (C), and IHCL (D). 25(OH)D, 25-hydroxyvitamin D; VFA, visceral fat area; IMCL, intramyocellular lipid; EMCL, extramyocellular lipid; IHCL, intrahepatic lipid. Closed and open circles represent data from American footballers and rugby footballers, respectively.

5. Discussion

We investigated the relationships between serum 25(OH)D concentrations and body fat and ectopic fat in male collegiate football athletes. We first found that the serum 25(OH)D concentrations were inversely correlated with IHCL rather than traditional body fat indicators, suggesting increasing vitamin D may modulate the activity of liver adipocytes.

The prevalence of vitamin D deficiency (<50 nmol/L) and insufficiency (50–75 nmol/L) were 17.6% and 58.5%, respectively, in the present study, which is higher than the 25(OH)D concentrations in our previous study conducted in 81 adults in the general population (mean 65.4 vs. 35.3 nmol/L) [28]. Zhou et al. [29] also found obese men (BMI > 28 kg/m²) have relatively lower serum 25(OH)D concentrations in July (mean 46.1 vs. 65.4 nmol/L). Recently, our previous studies found that both acute exercise and long-term exercise training could increase serum 25(OH)D concentrations or prevent its seasonal reduction [30,31]. Football athletes have a regular exercise habit that could, in part, explain the relatively higher 25(OH)D concentrations in our study.

Since athletes perform a large amount of exercise training, they usually have been regarded as a healthy model with a lower risk of cardiovascular disease. However, according to recent studies, the heavy athletes are usually at significantly higher risk of cardiovascular diseases [3,4]. To enhance athletic performance, American and rugby football athletes usually have the intent to consume more energy to gain more weight [32], which was suggested to lead to lipid accumulation in adiposity and ectopic tissues [6,7]. Excessive lipid accumulation predisposes one to the development of cardiovascular diseases [2] and is negatively related to serum 25(OH)D concentrations [9–12]. In vitro, it has been suggested that vitamin D modulates the activity of adipocytes and may have effects on the inhibition of lipid accumulation through VDRs or calcium ions [15,16]. Prior evidence indicated that low circulating 25(OH)D concentrations were associated with future weight gain [17], and vitamin D treatment could significantly decrease hepatic triglyceride accumulation in obese and hyperglycemic mice [18,19].

Previous studies have showed that serum 25(OH)D concentrations were negatively correlated with BMI and % body fat, and with abdominal fat indicators in adults in the general population [9–12]. However, the relationships between 25(OH)D and fat variables were unclear in collegiate football

athletes. Heller et al. [21] found that serum 25(OH)D concentrations were negatively associated with BMI and % body fat in 42 male and female athletes who participate in several kinds of sports. A similar result was also observed in collegiate male wrestlers, and the negative relationship between serum 25(OH)D concentrations and % body fat was observed regardless of seasonal changes [22]. In the present study, correlation analyses showed that serum 25(OH)D concentrations were not related to any of the body fat variables, including BMI, % body fat, WC, and SFA in collegiate football athletes, except there was a trend of a negative correlation between serum 25(OH)D and VFA ($r = -0.307$, $p = 0.083$). According to our knowledge, few research studies have focused on the relationships between 25(OH)D and central body fat indicators in Asian football athletes. Although circulating 25(OH)D concentrations were negatively correlated with BMI and VFA in non-Asian adults, our previous study found that serum 25(OH)D concentrations were negatively correlated with VFA levels rather than BMI in Japanese adults [12]; Hao et al. [33] also found that the levels of VFA were the strongest predictors of serum 25(OH)D in Chinese men. The discrepancy of the relationships between 25(OH)D and these body fat indicators may be attributed to the different fat distribution in Asians and non-Asians [34]. Moreover, the relatively small sample size and narrow body fat range could be another potential explanation for the discrepancy.

In addition to body fat in adipose tissues, the relationships between fat accumulation in ectopic tissue and the incidence of cardiovascular disease have received more attention recently [2]. Prior evidence has showed that there are negative relationships between 25(OH)D concentrations and liver and muscle fat in adults in the general population [23–25]. However, this relationship has not been investigated in young collegiate football athletes. Our study found that there is a negative relationship between serum 25(OH)D concentrations and IHCL rather than IMCL and EMCL even after adjusting for several factors in collegiate football players. These disparate results of the associations of serum 25(OH)D concentrations with IMCL and EMCL may be partly related to “athletes’ paradox” [26]. The lipid droplets in muscle are usually located adjacent to the muscle mitochondria, which indicates that they function as a substrate energy source during exercise in athletes [35]. Exercise-trained athletes may have similar levels of IMCL with insulin-resistant obesity and type 2 diabetes mellitus [36]. In addition, we found that there was an inverse relationship between serum 25(OH)D concentrations and γ -GTP concentrations ($r = -0.504$, $p = 0.033$) in rugby players in the subgroup analysis. These data, including the finding from recent mice [18,19], suggest that increasing serum 25(OH)D concentrations may lead to the inhibition of hepatic lipid accumulation in overweight athletes.

The present study first found that higher serum 25(OH)D concentrations were closely related to the content of IHCL rather than traditional body fat indicators in male collegiate football athletes. However, a small sample size could be a limitation. Additionally, it is still unclear whether the results observed in our study could be extrapolated to other kinds of athletes. Finally, the cross-sectional nature of the study does not allow for determination of cause and effect. Longitudinal studies conducted on other kinds of sports are necessary to confirm the associations between serum 25(OH)D concentrations and fat tissue and ectopic fat indicators.

6. Conclusions

In conclusion, we observed that the prevalence of vitamin D deficiency and insufficiency was higher, but relatively higher than in adults in the general population. In addition, serum 25(OH)D concentrations were more closely associated with the intrahepatic cellular fat content than any of the indicators traditionally used to assess adiposity in overweight collegiate athletes.

Our results suggest that maintaining higher circulating 25(OH)D concentrations may lead to lower IHCL content, with the potential for subsequent reductions in cardiovascular disease risks in overweight athletes, and perhaps in others. These possibilities warrant testing in adequately designed randomized controlled trials.

Author Contributions: Conceptualization, X.S. and Z.-B.C.; Methodology and Writing—Original Draft Preparation, X.S.; Data Curation, K.T., S.O., and M.H.; Writing—Reviewing and Editing, Z.-B.C. and M.H.; Conceptualization and Supervision, Z.-B.C. All authors read and approved the final manuscript.

Funding: This study was supported in part by a Waseda University Grant for Special Research Projects (2014B-419 to M.H.); a grant from the Ministry of Education, Culture, Sports, Science, and Technology-Supported Program for the Strategic Research Foundation at Private University (to M.H.); grant from National Natural Science Foundation of China (number (No.) 31571226 and 81703220); the Program for Professor of Special Appointment (Eastern Scholar) at Shanghai Institutions of Higher Learning (No. TP2014057) and Shanghai Key Lab of Human Performance (Shanghai University of Sport) (No. 11DZ2261100); and a grant from China Postdoctoral Science Foundation (No. 2017M613154).

Acknowledgments: The authors thank Katsuhiko Suzuki and Shizuo Sakamoto for their assistance. The authors are grateful to all subjects for their participation in this study.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Saydah, S.; Bullard, K.M.; Cheng, Y.; Ali, M.K.; Gregg, E.W.; Geiss, L.; Imperatore, G. Trends in cardiovascular disease risk factors by obesity level in adults in the United States, NHANES 1999–2010. *Obesity* **2014**, *22*, 1888–1895. [[CrossRef](#)] [[PubMed](#)]
2. Shulman, G.I. Ectopic fat in insulin resistance, dyslipidemia, and cardiometabolic disease. *N. Engl. J. Med.* **2014**, *371*, 2237–2238. [[CrossRef](#)] [[PubMed](#)]
3. Guo, J.; Zhang, X.; Wang, L.; Guo, Y.; Xie, M. Prevalence of metabolic syndrome and its components among Chinese professional athletes of strength sports with different body weight categories. *PLoS ONE* **2013**, *8*, E79758. [[CrossRef](#)] [[PubMed](#)]
4. Murata, H.; Oshima, S.; Torii, S.; Taguchi, M.; Higuchi, M. Characteristics of body composition and cardiometabolic risk of Japanese male heavyweight Judo athletes. *J. Physiol. Anthropol.* **2016**, *35*, 10. [[CrossRef](#)] [[PubMed](#)]
5. Selden, M.A.; Helzberg, J.H.; Waeckerle, J.F.; Browne, J.E.; Brewer, J.H.; Monaco, M.E.; Tang, F.; O’keefe, J.H. Cardiometabolic abnormalities in current National Football League players. *Am. J. Cardiol.* **2009**, *103*, 969–971. [[CrossRef](#)] [[PubMed](#)]
6. Sevastianova, K.; Santos, A.; Kotronen, A.; Hakkarainen, A.; Makkonen, J.; Silander, K.; Peltonen, M.; Romeo, S.; Lundbom, J.; Lundbom, N.; et al. Effect of short-term carbohydrate overfeeding and long-term weight loss on liver fat in overweight humans. *Am. J. Clin. Nutr.* **2012**, *96*, 727–734. [[CrossRef](#)] [[PubMed](#)]
7. Iggman, D.; Rosqvist, F.; Larsson, A.; Amlov, J.; Beckman, L.; Rudling, M.; Riserus, U. Role of dietary fats in modulating cardiometabolic risk during moderate weight gain: A randomized double-blind overfeeding trial (LIPOGAIN Study). *J. Am. Heart Assoc.* **2014**, *3*, e001095. [[CrossRef](#)] [[PubMed](#)]
8. Bikle, D. Nonclassic actions of vitamin D. *J. Clin. Endocrinol. Metab.* **2009**, *94*, 26–34. [[CrossRef](#)] [[PubMed](#)]
9. Dong, Y.; Pollock, N.; Stallmann-Jorgensen, I.S.; Gutin, B.; Lan, L.; Chen, T.C.; Keeton, D.; Petty, K.; Holick, M.F.; Zhu, H. Low 25-hydroxyvitamin D levels in adolescents: Race, season, adiposity, physical activity, and fitness. *Pediatrics* **2010**, *125*, 1104–1111. [[CrossRef](#)] [[PubMed](#)]
10. Kremer, R.; Campbell, P.P.; Reinhardt, T.; Gilsanz, V. Vitamin D status and its relationship to body fat, final height, and peak bone mass in young women. *J. Clin. Endocrinol. Metab.* **2009**, *94*, 67–73. [[CrossRef](#)] [[PubMed](#)]
11. Sulistyoningrum, D.C.; Green, T.J.; Lear, S.A.; Devlin, A.M. Ethnic-specific differences in vitamin D status is associated with adiposity. *PLoS ONE* **2012**, *7*, e43159. [[CrossRef](#)] [[PubMed](#)]
12. Sun, X.; Cao, Z.B.; Tanisawa, K.; Ito, T.; Oshima, S.; Ishimi, Y.; Tabata, I.; Higuchi, M. Associations between the serum 25(OH)D concentration and lipid profiles in Japanese men. *J. Atheroscler. Thromb.* **2015**, *22*, 355–362. [[CrossRef](#)] [[PubMed](#)]
13. Wortsman, J.; Matsuoka, L.Y.; Chen, T.C.; Lu, Z.; Holick, M.F. Decreased bioavailability of vitamin D in obesity. *Am. J. Clin. Nutr.* **2000**, *72*, 690–693. [[CrossRef](#)] [[PubMed](#)]
14. Vimalaswaran, K.S.; Vimalaswaran, K.S.; Berry, D.J.; Lu, C.; Tikkanen, E.; Pilz, S.; Hiraki, L.T.; Cooper, J.D.; Dastani, Z.; Li, R.; et al. Causal relationship between obesity and vitamin D status: Bi-directional Mendelian randomization analysis of multiple cohorts. *PLoS Med.* **2013**, *10*, e1001383. [[CrossRef](#)] [[PubMed](#)]

15. Song, Q.; Sergeev, I.N. Calcium and vitamin D in obesity. *Nutr. Res. Rev.* **2012**, *25*, 130–141. [[CrossRef](#)] [[PubMed](#)]
16. Sergeev, L.N.; Song, Q. High vitamin D and calcium intakes reduce diet-induced obesity in mice by increasing adipose tissue apoptosis. *Mol. Nutr. Food Res.* **2014**, *58*, 1342–1348. [[CrossRef](#)] [[PubMed](#)]
17. Mai, X.M.; Chen, Y.; Camargo, C.A.; Langhammer, A. Cross-sectional and prospective cohort study of serum 25-hydroxyvitamin D level and obesity in Adults. *Am. J. Epidemiol.* **2012**, *175*, 1029–1036. [[CrossRef](#)] [[PubMed](#)]
18. Cheng, S.; So, W.Y.; Zhang, D.; Cheng, Q.; Boucher, B.J.; Leung, P.S. Calcitriol reduces hepatic triglyceride accumulation and glucose output through Ca²⁺/CaMKK beta/AMPK activation under insulin resistant conditions in type 2 diabetes. *Curr. Mol. Med.* **2016**, *16*, 747–758. [[CrossRef](#)] [[PubMed](#)]
19. Borges, C.C.; Salles, A.F.; Brighenti, I.; Mandarim-De-Lacerda, C.A.; Aguila, M.B. Vitamin D deficiency increases lipogenesis and reduces Beta-oxidation in the liver of diet-induced obese mice. *J. Nutr. Sci. Vitaminol.* **2018**, *64*, 106–115. [[CrossRef](#)] [[PubMed](#)]
20. Cassity, E.P.; Redzic, M.; Teager, C.R.; Thomas, D.T. The effect of body composition and BMI on 25(OH)D response in vitamin D-supplemented athletes. *Eur. J. Sport Sci.* **2016**, *16*, 773–779. [[CrossRef](#)] [[PubMed](#)]
21. Heller, J.E.; Thomas, J.J.; Hollis, B.W.; Larson-Meyer, D.E. Relation between vitamin D status and body composition in collegiate athletes. *Int. J. Sport Nutr. Exerc. Metab.* **2015**, *25*, 128–135. [[CrossRef](#)] [[PubMed](#)]
22. Barcal, J.N.; Thomas, J.T.; Hollis, B.W.; Austin, K.J.; Alexander, B.M.; Larson-Meyer, D.E. Vitamin D and weight cycling: Impact on injury, illness, and inflammation in collegiate wrestlers. *Nutrients* **2016**, *8*, 775. [[CrossRef](#)] [[PubMed](#)]
23. Gilsanz, V.; Kremer, A.; Mo, A.O.; Wren, T.A.L.; Kremer, R. Vitamin D status and its relation to muscle mass and muscle fat in young women. *J. Clin. Endocr. Metab.* **2010**, *95*, 1595–1601. [[CrossRef](#)] [[PubMed](#)]
24. Scott, D.; Joham, A.; Teede, H.; Gibson-Helm, M.; Harrison, C.; Cassar, S.; Hutchison, S.; Ebeling, P.R.; Stepto, N. Associations of vitamin D with inter- and intra-muscular adipose tissue and insulin resistance in women with and without polycystic ovary syndrome. *Nutrients* **2016**, *8*, 774. [[CrossRef](#)] [[PubMed](#)]
25. Wang, D.; Lin, H.; Xia, M.; Aleteng, Q.; Li, X.; Ma, H.; Pan, B.; Gao, J.; Gao, X. Vitamin D levels are inversely associated with liver fat content and risk of non-alcoholic fatty liver disease in a Chinese middle-aged and elderly population: The Shanghai Changfeng Study. *PLoS ONE* **2016**, *11*, e0157515.
26. Dube, J.J.; Amati, F.; Stefanovic-Racic, M.; Toledo, F.G.S.; Sauers, S.E.; Goodpaster, B.H. Exercise-induced alterations in intramyocellular lipids and insulin resistance: The athlete's paradox revisited. *Am. J. Physiol. Endocrinol. Metab.* **2008**, *294*, E882–E888. [[CrossRef](#)] [[PubMed](#)]
27. Sunami, A.; Sasaki, K.; Suzuki, Y.; Oguma, N.; Ishihara, J.; Nakai, A.; Yasuda, J.; Yokoyama, Y.; Yoshizaki, T.; Tada, Y.; et al. Validity of a semi-quantitative food frequency questionnaire for collegiate athletes. *J. Epidemiol.* **2016**, *26*, 284–291. [[CrossRef](#)] [[PubMed](#)]
28. Sun, X.; Cao, Z.B.; Tanisawa, K.; Ito, T.; Oshima, S.; Higuchi, M. Vitamin D supplementation reduces insulin resistance in Japanese adults: A secondary analysis of a double-blind, randomized, placebo-controlled trial. *Nutr. Res.* **2016**, *36*, 1121–1129. [[CrossRef](#)] [[PubMed](#)]
29. Zhou, J.; Zhu, Y.; Guo, P.; Chen, Z.; Xie, F.; Liu, X.; He, S. Serum 25(OH)D and lipid levels in Chinese Obese and normal weight males before and after oral vitamin D supplementation. *Biomed. Environ. Sci.* **2013**, *26*, 801–807. [[PubMed](#)]
30. Sun, X.; Cao, Z.B.; Taniguchi, H.; Tanisawa, K.; Higuchi, M. Effect of an acute bout of endurance exercise on serum 25(OH)D concentrations in young adults. *J. Clin. Endocrinol. Metab.* **2017**, *102*, 3937–3944. [[CrossRef](#)] [[PubMed](#)]
31. Sun, X.; Cao, Z.B.; Tanisawa, K.; Taniguchi, H.; Kubo, T.; Higuchi, M. Effects of chronic endurance exercise training on serum 25(OH)D concentrations in elderly Japanese men. *Endocrine* **2018**, *59*, 330–337. [[CrossRef](#)] [[PubMed](#)]
32. Berglund, L.; Sundgot-Borgen, J.; Berglund, B. Adipositas athletica: A group of neglected conditions associated with medical risks. *Scand. J. Med. Sci. Sports* **2011**, *21*, 617–624. [[CrossRef](#)] [[PubMed](#)]
33. Hao, J.; Ma, X.; Shen, Y.; Ni, J.; Luo, Y.; Xiao, Y.; Bao, Y.; Jia, W. Associations of serum 25-hydroxyvitamin D₃ levels with visceral adipose tissue in Chinese men with normal glucose tolerance. *PLoS ONE* **2014**, *9*, e86773. [[CrossRef](#)] [[PubMed](#)]

34. Yoon, K.H.; Lee, J.H.; Kim, J.W.; Cho, J.H.; Choi, Y.H.; Ko, S.H.; Zimmet, P.; Son, H.Y. Epidemic obesity and type 2 diabetes in Asia. *Lancet* **2016**, *368*, 1681–1688. [[CrossRef](#)]
35. Bosma, M. Lipid droplet dynamics in skeletal muscle. *Exp. Cell Res.* **2016**, *340*, 180–186. [[CrossRef](#)] [[PubMed](#)]
36. Goodpaster, B.H.; He, J.; Watkins, S.; Kelley, D.E. Skeletal muscle lipid content and insulin resistance: Evidence for a paradox in endurance-trained athletes. *J. Clin. Endocr. Metab.* **2001**, *86*, 5755–5761. [[CrossRef](#)] [[PubMed](#)]



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).