



COVID-19-Related Mortality Risk in People With Severe Mental Illness: A Systematic and Critical Review

Marc De Hert^{1,2*}, Victor Mazereel¹, Marc Stroobants³, Livia De Picker^{4,5},
Kristof Van Assche^{2,6} and Johan Detraux⁷

¹ Department of Neurosciences, Center for Clinical Psychiatry, University Psychiatric Center, KU Leuven, Kortenberg, Belgium, ² Antwerp Health Law and Ethics Chair, University of Antwerp, Antwerp, Belgium, ³ Biomedical Library, University Psychiatric Center, KU Leuven, Kortenberg, Belgium, ⁴ Collaborative Antwerp Psychiatric Research Institute, University of Antwerp, Antwerp, Belgium, ⁵ University Psychiatric Hospital Campus Duffel, Duffel, Belgium, ⁶ Research Group Personal Rights and Property Rights, Faculty of Law, University of Antwerp, Antwerp, Belgium, ⁷ Department of Neurosciences, Public Health Psychiatry, University Psychiatric Center, KU Leuven, Kortenberg, Belgium

OPEN ACCESS

Edited by:

Andrea Fiorillo,
University of Campania Luigi
Vanvitelli, Italy

Reviewed by:

Daria Smirnova,
Samara State Medical
University, Russia
Martina Rojnic Kuzman,
University Hospital Centre
Zagreb, Croatia

*Correspondence:

Marc De Hert
marc.dehert@upckuleuven.be

Specialty section:

This article was submitted to
Psychosomatic Medicine,
a section of the journal
Frontiers in Psychiatry

Received: 20 October 2021

Accepted: 03 December 2021

Published: 13 January 2022

Citation:

De Hert M, Mazereel V, Stroobants M,
De Picker L, Van Assche K and
Detraux J (2022) COVID-19-Related
Mortality Risk in People With Severe
Mental Illness: A Systematic and
Critical Review.
Front. Psychiatry 12:798554.
doi: 10.3389/fpsy.2021.798554

Background: Increasing clinical evidence suggests that people with severe mental illness (SMI), including schizophrenia spectrum disorders, bipolar disorder (BD), and major depressive disorder (MDD), are at higher risk of dying from COVID-19. Several systematic reviews examining the association between psychiatric disorders and COVID-19-related mortality have recently been published. Although these reviews have been conducted thoroughly, certain methodological limitations may hinder the accuracy of their research findings.

Methods: A systematic literature search, using the PubMed, Embase, Web of Science, and Scopus databases (from inception to July 23, 2021), was conducted for observational studies assessing the risk of death associated with COVID-19 infection in adult patients with pre-existing schizophrenia spectrum disorders, BD, or MDD. Methodological quality of the included studies was assessed using the Newcastle-Ottawa Scale (NOS).

Results: Of 1,446 records screened, 13 articles investigating the rates of death in patients with pre-existing SMI were included in this systematic review. Quality assessment scores of the included studies ranged from moderate to high. Most results seem to indicate that patients with SMI, particularly patients with schizophrenia spectrum disorders, are at significantly higher risk of COVID-19-related mortality, as compared to patients without SMI. However, the extent of the variation in COVID-19-related mortality rates between studies including people with schizophrenia spectrum disorders was large because of a low level of precision of the estimated mortality outcome(s) in certain studies. Most studies on MDD and BD did not include specific information on the mood state or disease severity of patients. Due to a lack of data, it remains unknown to what extent patients with BD are at increased risk of COVID-19-related mortality. A variety of factors are likely to contribute to the increased mortality risk of COVID-19 in these patients. These include male sex, older age, somatic comorbidities (particularly cardiovascular diseases), as well as disease-specific characteristics.

Conclusion: Methodological limitations hamper the accuracy of COVID-19-related mortality estimates for the main categories of SMIs. Nevertheless, evidence suggests that SMI is associated with excess COVID-19 mortality. Policy makers therefore must consider these vulnerable individuals as a high-risk group that should be given particular attention. This means that targeted interventions to maximize vaccination uptake among these patients are required to address the higher burden of COVID-19 infection in this already disadvantaged group.

Keywords: severe mental illness, schizophrenia, bipolar disorder, major depressive disorder, mortality, COVID-19

BACKGROUND

People with severe mental illness (SMI), including schizophrenia (SZ) (lifetime prevalence: 0.7%), bipolar disorder (BD) (lifetime prevalence: 0.4–1.1%), and major depressive disorder (MDD) (lifetime prevalence: 15–18%) (1–3), have a two to three times higher mortality rate than the general population (4–7). This mortality gap translates into a 10–20 years shortened life expectancy (6, 8) and appears to be widening (9).

It is well-known that the majority of deaths in individuals with SMI are due to physical diseases, predominantly cardiovascular diseases (8, 10). Non-medical factors, including unhealthy lifestyles, disparities in physical health care, and stigmatizing attitudes toward people with SMI, contribute to the higher risk of death (11–13). Disease-related factors, such as unawareness of physical problems and challenges in appraising health information due to cognitive deficits, delusions, and, in general, lower educational attainment and health literacy (11, 14–18), as well as the use of psychotropic medication (antipsychotics, antidepressants, and mood stabilizers) (4, 10, 12) may further increase the risk of physical comorbidities.

Research has shown that several respiratory viruses, including human coronaviruses, can have neuroinvasive properties (19). The central nervous system (CNS) is also a potential target for the SARS-CoV-2 virus, because angiotensin-converting enzyme 2 (ACE 2) receptors, used by the virus to enter the cells, are equally expressed in glial cells and neurons in the brain (20–23). Although there is still no convincing evidence for direct neuropathogenic effects of SARS-CoV-2 (24, 25), COVID-19 infection can cause CNS damage (26–28). In a prospective autopsy cohort study, extensive inflammatory changes, affecting both white and gray matter, were detected in the brain of patients with lethal COVID-19. This inflammatory response was most pronounced in the olfactory bulbs and medulla oblongata. Results of this study suggest that CNS changes are due to a maladaptive immune response, rather than the consequence of a direct virus-induced effect, given that viral presence was low at late stages of COVID-19 (29). The first longitudinal imaging study (which had not been peer reviewed as of Oct 13, 2021), comparing structural and functional brain scans acquired from individuals before and after SARS-CoV-2 infection ($n = 401$) with scans from a well-matched control group ($n = 384$), demonstrated brain atrophy, mainly in the limbic regions with direct neuronal connectivity to the primary olfactory and

gustatory system, in addition to a more diffuse loss of gray matter. These authors suggested that the observed brain changes may be due to a direct virus effect, or to neuroinflammation, following viral infection and initiating chronic neuronal dysfunctions (30).

Different pathogenic pathways may be involved (31). One of the proposed mechanisms is that the SARS-CoV-2 virus enters the CNS through the neuronal retrograde route. In this case the virus infects neurons in the periphery and uses the olfactory nerve pathway to gain access to the CNS and cause infections of immune-functioning microglia or astrocytes in the CNS (19, 30). This, however, does not rule out a pathway from the nose to the brain by other mechanisms (such as the vascular route). A review by Uversky et al. (20) stated that there are at least seven candidate routes the SARS-CoV-2 virus can use to reach the CNS.

Nevertheless, because of these findings concerns have been raised regarding the neuroinvasive effect of COVID-19 infection in patients with pre-existing neuropsychiatric disorders, in particular in SMIs, which are often already characterized by a systemic pro-inflammatory state (32, 33). According to one review, 0.9–4% of individuals infected with SARS-CoV-2 develop psychotic spectrum disorders (34). A retrospective cohort study reported the following numbers (<6 months after acute infection): 0.9% in COVID-19 infection without hospitalization, 2.9% after hospitalization, and 7% after COVID-19-related encephalopathy (28). However, some of these cases probably are due to COVID-19-related psychosocial stress or treatment (e.g., steroid treatment in patients with COVID-19) (34, 35).

Recently, five systematic reviews and meta-analyses (36–40), assessing the risk of COVID-19-related mortality in patients with a psychiatric disorder, have demonstrated that people with SMI have a higher COVID-19-related mortality risk, compared to general population controls, and even compared to people with other psychiatric disorders. This risk remained high after adjustment for age, sex, and other confounders. Although these reviews have been thoroughly conducted, certain methodological limitations may hinder the accuracy of their research findings. Most of these reviews did not use a comprehensive search strategy for COVID-19 (36–39), or did not include the Embase database (36, 37, 40). Several of these reviews not only included laboratory-confirmed COVID-19 cases, but also patients where the SARS-CoV-2 infection was based on a clinical diagnosis made by physicians (36, 37, 39), or did not make this clear in their methodology section (40). Furthermore, reviews reporting

mortality data on mood disorders did not distinguish unipolar and BDs (although most of the included studies in these reviews relied on electronic medical records that do not allow a fine-grained analysis of clinical variables) and/or pre-existing from post-infection or comorbid disorders [e.g., (38)]. Finally, a thorough discussion part is also missing in most of these reviews.

AIM OF THE STUDY

To overcome the above-mentioned limitations a novel systematic literature search was conducted to assess the risk of death associated with COVID-19 infection in people with SMI (schizophrenia spectrum disorders, BD, and MDD), compared with patients without SMI, or without any psychiatric disorder, and a thorough discussion part was provided.

METHODS

Search Strategy

A comprehensive literature search, using the PubMed, Embase, Web of Science, and Scopus databases (from inception to July 23, 2021), was conducted without language restriction for studies reporting data on the risk of death associated with COVID-19 infection in adult patients with schizophrenia spectrum disorders, BD, or MDD, compared with controls (patients without a SMI or without a psychiatric disorder). Two of the authors (JD and MD) and two experienced biomedical information specialists worked closely together to construct effective search strings for the different databases. Full search strategies are available as **Supplementary Material 1**. Duplicates were removed using EndNote X9 and Rayyan QCRI (JD). After removing duplicates, titles and abstracts were screened by JD. Articles that were deemed potentially relevant according to the selection criteria were selected. JD and MD independently reviewed the full text of the selected articles and assessed their eligibility. They also attempted to identify additional studies through a systematic search of the reference lists of selected articles and of previously published systematic reviews/meta-analyses.

Selection Criteria

Inclusion criteria were:

- (i) Published, peer-reviewed, original studies,
- (ii) Population-based observational studies, including case-control, cohort, or cross-sectional studies,
- (iii) Studies including patients with laboratory-confirmed COVID-19 cases (i.e., a positive real-time reverse transcription-polymerase chain reaction test),
- (iv) Studies including patients with a clinically confirmed pre-existing SMI (i.e., schizophrenia spectrum disorders, BDs, and/or unipolar depression), using a widely-accepted standardized disease coding system, such as the Diagnostic and Statistical Manual of Mental Disorders (DSM) or the International Classification of Diseases (ICD), and
- (v) Studies reporting COVID-19-related mortality outcomes [i.e., odds ratio, risk ratio, hazard ratio (HR), or associated

metrics] and comparing COVID-19-related mortality risks of SMI patients with non-SMI patients, or patients without a psychiatric disorder.

Studies that did not include patients with pre-existing SMI (thus studies where COVID-19 may have been an antecedent to the onset of the SMI disorder), a control group, COVID-19-related mortality outcomes, original data, or were not peer-reviewed and published (preprints, conference papers), as well as case reports, reviews, meta-analyses, and studies where SMI mortality outcomes were grouped with those for non-SMIs, or studies where COVID-19 was not ascertained according to laboratory testing were excluded.

This systematic review adhered to the 2020 Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) guidelines (41). We did, however, not register our protocol prior to submitting the manuscript for publication.

Data Extraction

Data from the included studies were extracted from each article by JD to collect the following information:

- (i) Study characteristics: author information, publication year, country where the study was conducted, study design (cross-sectional, cohort or case-control, prospective vs. retrospective), time period studied during the COVID-19 pandemic,
- (ii) Patient characteristics: primary diagnoses included in the study, sample sizes of SMI patients and controls, mean/median age (years), and sex distribution,
- (iii) Outcome measure: adjusted and unadjusted mortality data (odds ratio, risk ratio, HR, or associated metrics with 95% confidence interval estimates), and
- (iv) Covariates in adjusted risk.

Assessment of Risk of Bias

The quality of each observational study was rated using the Newcastle-Ottawa Scale (NOS), whereby a higher score indicated higher methodological quality and lower risk of bias. The NOS was adapted for cross-sectional data.


RESULTS

Search Strategy

The original search in the PubMed ($n = 131$), Embase ($n = 1,133$), Web of Science ($n = 105$), and Scopus ($n = 77$) databases yielded a total of 1,446 reports. Of these, 265 duplicate reports were removed. Overall, 37 references of published studies were selected as potentially eligible, of which 12 original records met the inclusion criteria. One record, identified through cross-referencing and which fulfilled the inclusion criteria, was added, resulting in a total of 13 original records. The results of the study selection are shown in the PRISMA flow diagram (see **Figure 1**).

Study and Patient Characteristics

Study and patient characteristics, as well as mortality data and covariates are presented in **Table 1**. Median age of SMI patients across studies ranged from 40 to 66 years and was



not reported in nine studies. Two studies were carried out in Denmark, two in France, one in Israel, one in South Korea, two in Spain, one in the U.K., and four in the U.S. Severe mental illness was almost exclusively categorized according to ICD classification criteria. In the French and Israeli studies only patients with schizophrenia (spectrum disorders) were analyzed. The U.S. study of Egede et al. (46) analyzed exclusively data of BD patients.

Quality Assessment

Data on study quality are presented in **Supplementary Material 2**. Methodological quality was high in nine studies, and moderate in four studies.

Study Results

For studies analyzing mortality data separately for patients with schizophrenia spectrum disorders, BD and MDD, fully

adjusted risks (= adjusted for demographic factors and one or more comorbidities or other covariates) ranged from 1.30 to 4.36 for schizophrenia spectrum disorders, and from 2.04 to 2.68 for depression. Only two studies (42, 46) reported COVID-19-related mortality data separately for patients with BD. In the study of Egede et al. (46), including exclusively BD patients, a HR of 2.67 (95% CI: 1.07–6.67) was found, while Barcella et al. (42) did not find a significant HR [adjusted HR = 1.94 (0.97–3.90)] in the fully adjusted model. A variety of factors were likely to contribute to the increased mortality risk of COVID-19 in these patients. These included male sex, older age, and somatic comorbidities, as evidenced by the reduction of the unadjusted risk after adjusting for these demographic factors and comorbidities. However, as even after this adjustment the risk of COVID-19-related mortality was still increased, other factors also seem to play a primordial role (see **Table 1**).

TABLE 1 | Study characteristics and mortality data of the included studies (33, 42–53).

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Barcella et al. (42)	Denmark	Cohort study	February 27 2020–January 2 2021	Schizophrenia spectrum disorders (F20–29) BD (F30, 31, 38) Unipolar depression, including MDD (F32–34, F39) (ICD-8 and ICD-10) Controls = patients without psychiatric disorder	Schizophrenia spectrum disorders ($n = 984$) vs. controls ($n = 127,281$) BD ($n = 485$) vs. controls ($n = 127,281$) Unipolar depression ($n = 3,764$) vs. controls ($n = 127,281$)	40 (median) 48.8% 45.3 (median) 37.3% 44 (median) 30.8%	<p>Standardized average RR: 2.29 (1.36–3.22)</p> <p>Unadjusted HR:4.12 (2.64–6.43)</p> <p>Adjusted HR:2.41 (1.53–3.79)</p> <p>Standardized average RR: 1.87 (1.12–3.12)</p> <p>Unadjusted HR:3.34 (1.66–6.70)</p> <p>Adjusted HR:1.94 (0.97–3.90)</p> <p>Standardized average RR: 1.92 (1.39–2.44)</p> <p>Unadjusted HR:2.63 (1.97–3.52)</p> <p>Adjusted HR:2.04 (1.52–2.74)</p>	Age, sex, highest obtained education, income, ischemic heart disease, congestive heart failure, cerebrovascular disease, CKD, HP, peripheral artery disease, DM, COPD, asthma, substance abuse, and malignancy

(Continued)

TABLE 1 | Continued

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Nemani et al. (43) [†]	U.S.	Retrospective cohort study	March 3 2020–May 31 2020	Schizophrenia spectrum disorders (F20, 22, 23, 25, 28, 29) Mood disorders -Unipolar depression, including MDD F32, 33, 34, 39 -BD F30, 31 (ICD-10) Controls = patients without psychiatric disorders, with the exception of patients with organic mental disorders, mental disorders due to substance use, mental retardation, and disorders of psychological development.	Lifetime SZ vs. controls: 75/6,349 Lifetime mood disorders vs. controls: 564/6,349 Recent SZ vs. controls: 46/6,349 Recent mood disorders vs. controls: 374/6,349	N/A	<p>Unadjusted OR: 2.93 (1.75–4.92) Demographically adjusted OR: 2.87 (1.62–5.08)</p> <p>Fully adjusted OR: 2.67 (1.48–4.80) Unadjusted OR: 1.82 (1.45–2.29) Demographically adjusted OR: 1.25 (0.98–1.61) Fully adjusted OR: 1.14 (0.87–1.49) Unadjusted OR: 2.84 (1.47–5.52) Demographically adjusted OR: 3.13 (1.50–6.54)</p> <p>Fully adjusted OR: 2.67 (1.26–5.69) Unadjusted OR: 2.19 (1.69–2.84) Demographically adjusted OR: 1.52 (1.13–2.03)</p> <p>Fully adjusted OR: 1.27 (0.94–1.73)</p>	Demographically adjusted: age, sex, race Fully adjusted (demographically + medical risk factors): age, sex, race, smoking status, HP, heart failure, myocardial infarction, DM, CKD, COPD, and cancer

(Continued)

TABLE 1 | Continued

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Tzur Bitan et al. (44)	Israel	Retrospective cohort study	March 2020–October 2020	SZ (ICD-9 or ICD-10) (F20) Controls = people without schizophrenia randomly drawn from the general population	642 patients vs. 709 controls	51.51 (mean, SZ) 51.37 (mean, controls) 61%	Non-adjusted OR: 3.14 (1.34–7.36) Adjusted OR (95% CI): 3.27 (1.39–7.68)	Age, sex
Jeon et al. (45)	South Korea	Retrospective cohort study	1 December 2019–15 May 2020	Schizophrenia spectrum disorders (ICD-10) (F20–F29) Mood disorders (BD+ MDD) (ICD-10) (F30–F39) Controls = patients without a psychiatric disorder	159 patients vs. 628 controls 273 patients vs. 1,060 controls	N/A	Adjusted OR: 2.25 (0.36–14.03) Adjusted OR (95% CI): 2.33 (0.96–5.66)	Cohort matched by age, sex, and Charlson Comorbidity Index with up to four people without mental disorder and adjusted for type of insurance, medical history of DM and pneumonia, and use of β -blockers and anticonvulsants
Egede et al. (46)	U.S.	Cross-sectional analysis	March 20–July 10 2020	BD (ICD-9 and ICD-10) (F30.1–F30.4, F30.9, F31.1–F31.6, F31.73–F31.78, F31.9) Controls = patients without a psychiatric disorder	38 patients vs. 1,330 controls	52.3 (mean) 29.7%	Non-adjusted HR: 2.83 (1.15–6.96) Demographically adjusted HR: 2.63 (1.07–6.49) Fully adjusted HR: 2.67 (1.07–6.67)	Gender, age, race/ethnicity, location, and primary payor, tobacco use, and BMI.

(Continued)

TABLE 1 | Continued

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Fond et al. (47)	France	Case-control study	February 27 2020–May 4 2020	SZ (F20,22,25) (ICD-10) Controls = patients without a psychiatric disorder	15 patients vs. 1,077 controls	66 (median) 73.3%	<p>Non-adjusted OR: 3.80 (1.19–12.20)</p> <p>Adjusted OR: 4.36 (1.09–17.44)</p> <p>Adjusted OR: 4.28 (1.07–17.20)</p> <p>Adjusted OR: 4.33 (1.08–17.34)</p>	Age, sex, smoking status, obesity, Charlson Comorbidity Index
Fond et al. (48)	France	Cohort study	February 1 2020–June 9 2020	SZ (F20,22,25) (ICD-10) Controls = patients without a SMI	823 patients vs. 49,927 controls	48.8%	<p>Unadjusted OR: 1.25 (1.05–1.49)</p> <p>Adjusted OR: 1.30 (1.08–1.56)</p> <p>Significant interaction between SZ and age ($p = 0.0006$): SZ patients between 65 and 80 years have a significantly higher risk of death than controls of same age [+7.89%; Adj. OR: 1.62 (1.28–2.06)]</p>	Age, sex, social deprivation, smoking status, overweight and obesity, Charlson Comorbidity Index, origin of the patient, hospital category, number of hospital stays for COVID-19, geographical areas of hospitalization

(Continued)

TABLE 1 | Continued

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Reilev et al. (49)	Denmark	Cohort study	February 27 2020–May 19 2020	SMI=SZ (F20), schizoaffective disorder (F25), or BD (F30,31)(ICD-10) Controls = patients without a SMI	76 patients vs. 11,046 controls	N/A	Non-adjusted OR: 3.8 (2.1–7.0) Demographically adjusted OR: 2.5 (1.2–5.1) Fully adjusted OR: 1.9 (0.9–3.9)	Age, sex Age, sex, and number of comorbidities
Poblador-Plou et al. (50)	Spain	Retrospective cohort study	March 4 2020–May 17 2020	Mood disorders (ICD-9-CM) Controls = patients without a psychiatric disorder	202 patients vs. 569 controls	N/A	Adjusted OR: 1.38 (0.98–1.95) (men with mood disorder) Adjusted OR: 1.46 (1.12–1.91) (women with mood disorder)	Age Age
Yang et al. (33)	U.K.	Retrospective cohort study	January 31 2020–July 26 2020	MDD (ICD-9 or ICD-10) Psychotic disorders (ICD-10) Controls = non-psychiatric patients	22,352 patients vs. 398,662 controls 1,431 patients vs. 419,583 controls	N/A	Adjusted OR: 2.68 (2.03–3.54) Adjusted OR: 3.50 (1.70–7.17)	Adjusted for birth year, sex, race or ethnicity, Townsend deprivation index, educational attainment, annual household income, BMI, smoking status, and history of chronic cardiac disease, DM, COPD, CKD, and asthma.

(Continued)

TABLE 1 | Continued

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Castro et al. (51)	U.S.	Retrospective cohort study	February 15 2020–May 24 2020	Mood disorders (MDD and BD) (ICD-10) Controls = individuals without a psychiatric disorder	717 patients vs. 2,271 controls	N/A	Mortality risk \geq hospital <i>day 12</i> Unadjusted HR: 2.16 (1.54–3.02) Demographically adjusted HR: 2.00 (1.42–2.82) Fully adjusted HR: 1.54 (1.05–2.25) Mortality risk $<$ hospital <i>day 12</i> Unadjusted HR: 1.17 (0.91–1.50) Demographically adjusted HR: 1.08 (0.83–1.39) Fully adjusted HR: 0.88 (0.65–1.18)	Age, sex, race, ethnicity, admission site (academic medical center compared with community hospital), socioeconomic status, and Charlson comorbidity index
Diez-Quevedo et al. (52)	Spain	Retrospective cohort study	March 1 2020–November 17 2020	Mood disorders (ICD-10) Controls = patients without mood disorders	279 patients (mood disorder) vs. controls	N/A	Adjusted HR: 1.52 (1.13–2.06)	Sex, age, history of medical and psychiatric disorders

(Continued)

TABLE 1 | Continued

Study	Country	Study design	Time period	Diagnosis	Sample size	Mean/median age (years) Male gender	Risk (95% CI)	Covariates in adjusted risk
Wang et al. (53)	U.S.	Retrospective case-control study	Up to July 29 2020	Depressive disorder (code 3548900) (including MDD) (SNOMED-CT) Controls = patients without a psychiatric disorder	1,460 patients with a recent (past year, but prior to COVID-19) diagnosis of depression	N/A	Unadjusted OR: 1.56 (1.28–1.91)	Age, sex, ethnicity, and medical comorbidities (cancers, CVDs, type 2 DM, obesity, CKD, COPD, asthma, and SUDs)

BD, bipolar disorder; BMI, body mass index; CI, confidence interval; CKD, chronic kidney disease; COPD, chronic obstructive pulmonary disease; CVD, cardiovascular diseases; DM, diabetes mellitus; HF, hypertension; HR, hazard ratio; ICD-9-CM, International Classification of Diseases, Ninth Revision, Clinical Modification; ICD, International Classification of Diseases; MDD, major depressive disorder; N/A, not available; OR, odds ratio; RR, risk ratio; SMI, severe mental illness; SNOMED-CT, systematized nomenclature of medicine-clinical terms; SUDs, substance use disorders. Bold values indicate statistical significance between groups.
 † A secondary analysis was limited to patients with recently documented psychiatric diagnoses of interest recorded in an encounter between January 1, 2019, and March 3, 2020 (recent diagnoses).

DISCUSSION

Our systematic review shows that, after full adjustment for relevant confounders, the extent of the variation in COVID-19-related mortality rates between studies including people with schizophrenia spectrum disorders was large. COVID-19-related mortality risk was found to be 2- to 4-fold increased for patients with schizophrenia spectrum disorders, compared with controls. There are several reasons for this variation:

- (1) studies showing higher adjusted mortality estimates included rather small samples of patients (47), and/or presented mortality data with large confidence intervals (33, 45, 47), indicating a low level of precision of the estimated mortality outcome(s),
- (2) while some studies were strictly limited to patients with SZ (44), others included disorders covering more or less the whole spectrum of SZ-like disorders (42, 43, 45). In addition, there is (besides the lack of information about psychopharmacological treatments and psychiatric treatment settings of patients) a lack of specific information about the severity and the status (first-episode vs. chronic) of the disease. These elements are important to consider since different forms of the disorder may have different risks of COVID-19-related mortality, and
- (3) the comparison group sometimes included non-SMI patients and at other times patients without a psychiatric disorder.

Although results are more stable for studies on MDD (COVID-19-related mortality risk in these patients seems to be 1.5- to 2-fold increased, compared with controls), these studies also in most cases did not include specific information on the mood state or disease severity of the patients. While some studies were strictly limited to patients with MDD (33, 51), others (42, 43) also included mild or moderate forms of the disease.

Finally, several studies (45, 49, 50, 52) involved a mixed population and did not make a distinction between the three main categories of SMIs. Studies on BD patients clearly are lacking.

Recently, several systematic reviews and meta-analyses (36, 37, 39, 40), showed that pre-existing mental disorders were associated with an increased COVID-19-related mortality risk, compared to controls, even after adjustment for age, sex, and other confounders. In the meta-analysis of Fond et al. (37), patients with SMI (schizophrenia spectrum disorders and/or BD) were found to have the highest risk of COVID-19-related mortality (adjusted OR = 1.67; 95% CI: 1.02–2.73). Vai et al. (36) also observed that the most robust associations were found for psychotic disorders (adjusted OR = 1.68; 95% CI: 1.29–2.18) and mood disorders (adjusted OR = 1.43; 95% CI: 1.15–1.79), after adjustment for age, sex, and other confounders, with a statistically significant difference ($p = 0.0047$) identified between adjusted estimates for SMI patients (adjusted OR = 1.55; 95% CI: 1.30–1.85) and patients with non-SMI psychiatric disorders (adjusted OR = 1.09; 95% CI: 0.92–1.29). A very recently published cohort study confirmed the previously published evidence (54). These results thus show that patients with SMI have a statistically significantly higher risk of death than patients with

non-SMI disorders. Moreover, analyses stratified by the number of redeemed psychotropic medications indicated that COVID-19-related mortality risk increases with higher psychotropic medication use (42). All these results therefore suggest an association between mental illness severity and COVID-19-related mortality risk.

A retrospective cohort study in patients with mood disorders has found that COVID-19-related mortality risk in these patients seems to be particularly elevated 2 weeks after admission, while there seems to be little difference in mortality risk with controls during early hospitalization (51). However, the meta-analysis of Vai et al. (36) found no evidence of increased in-hospital mortality in patients with psychiatric disorders vs. those without. Moreover, COVID-19-related mortality risk was significantly higher among psychiatric patients who were not admitted to the hospital than among hospitalized patients.

Factors Underlying the Association Between SMI and COVID-19-Related Mortality

Several demographic factors and somatic comorbidities have been identified that contribute to the higher observed mortality estimates associated with COVID-19 infection in people with SMI (42–44, 49, 50, 53, 55, 56) and without SMI (57–63). These include older age (≥ 65 –70 years old), male gender, low socioeconomic status (SES) and educational level, and physical diseases (cardiovascular disease, hypertension, chronic obstructive pulmonary disease, chronic kidney disease, diabetes). These identified factors may have a more profound impact on people with a SMI.

Firstly, as in the general population, particularly older individuals (≥ 60 years) with SMI are vulnerable to COVID-19. However, there seems to be an excess mortality due to COVID-19 among patients of this age group. Deaths due to COVID-19 were found to be 4 times higher for those with SMI, compared to individuals without SMI within the same age group [people with SMI: 0.01% (40–59 years), 0.12% (60–69 years), 0.46% (70–79 years), vs. individuals without SMI: 0.01% (40–59 years), 0.03% (60–69 years), 0.11% (70–79 years)] (55). Fond et al. (48) found that patients with SZ between 65 and 80 years had a higher risk of death [7.69% (< 55 years) and 30.29% (≥ 65 < 80 years)], compared to individuals without a SMI of the same age [4.04% (< 55 years) and 22.4% (≥ 65 < 80 years)] [Adj. OR (95% CI): 1.62 (1.27–2.06, $p = 0.0002$)]. These results can be linked to the accelerated biological aging hypothesis, one of the major causes of the higher premature mortality rates that are observed in people with SMI (64, 65). This means that aging of both body and brain, and, in particular for COVID-19, the senescence of immune cells (66), might be more rapid in these people. One study found that molecular brain age (i.e., biological age of the brain) was 2–6 years higher than the chronological age in individuals with SZ, and 4.7–7.5 years higher in subjects with BD. No increase in brain aging was noted in subjects with MDD (67).

The impact of SES, which has been found to be associated with health care access (68), on the risk of COVID-19-related mortality remains to be elucidated, particularly in

individuals with SMI (55). Due to the complexity of SES and its metrics (such as the Townsend Deprivation Index or the Distressed Communities Index), it stays unclear which individual components are associated with COVID-19-related mortality. While the impact of certain socioeconomic aspects (e.g., lower education and race) on COVID-19-related mortality has been shown (69, 70), other components (such as poverty and unemployment) were found to be protective against COVID-19-related mortality (69). Moreover, these measurements remain indirect indices of health care access and are assessed only once at baseline. Therefore, misclassification due to the absence of repeated measurements might exist (33). The same applies to the impact of the type of care/facility on the treatment outcomes for persons with SMI with COVID-19, particularly during the first wave of the COVID-19 pandemic. At the beginning of this pandemic, in the U.S. (71) and in Europe (72) many inpatient psychiatric facilities created psychiatric COVID-19-positive units (PCU). When tested positive, psychiatric patients were transferred to these units, where they were treated medically for their COVID-19 illness by internists and medical nurse practitioners. Only if patients showed signs of respiratory distress, such as shortness of breath or chest pain, they were transferred to a medical emergency room for further evaluation (71). The organization of PCU, however, varied considerably across countries and over time (72). Therefore, it remains unknown what the effect is of the type of care/facility on COVID-19-related mortality figures in these persons.

Secondly, a higher somatic comorbidity burden in patients with SMI, compared to non-psychiatric patients, may also partly explain the increased COVID-19-related mortality risk. However, with the exception of cardiovascular diseases, results for other somatic comorbidities are sometimes inconsistent. In SMI and non-psychiatric populations hypertension, diabetes, and chronic obstructive pulmonary disease have been found to correlate with an increased risk of fatality in most (43, 49, 73–75), but not all studies (55, 76). Nevertheless, it is well-recognized that individuals with these comorbidities are at an increased risk for a severe course of COVID-19 (77). In addition, somatic comorbidities seem in general to be an important driver of the observed increased COVID-19-related infection and mortality estimates in patients with SMI (33, 43, 49, 53). The reasons why certain somatic comorbidities are associated with more severe COVID-19 illness in people with SMI are not yet fully understood (78).

Although results are inconsistent, the use of psychotropic medications may be another important risk factor. A recent meta-analysis (36) showed that, after adjustment for age, sex, and other confounders, COVID-19-related mortality was associated with exposure to antipsychotics (initiated before contracting COVID-19) (adjusted OR = 2.43, 95% CI: 1.81–3.25), but not to antidepressants (adjusted OR = 1.18, 95% CI: 0.93–1.50). However, a very recently published retrospective cohort study did not observe an association between antipsychotic use and COVID-19-related mortality (79). An important obstacle in finding an answer to the question whether there exists an association between the use of psychotropic medications and

COVID-19-related mortality is that specific data on the use of psychotropic medication and psychiatric status (acute phase vs. stabilization phase) across studies is lacking.

Some antipsychotic medications (particularly clozapine) seem to increase susceptibility to pneumonia and pneumonia-related mortality risk in individuals with SMI due to sedation, impaired swallowing and hypersalivation (80–82). Clozapine can also suppress immune function (78). In animal models, short- to intermediate-term exposure to clinically relevant levels of risperidone has been shown to induce inflammatory and adaptive immune process dysregulation, possibly affecting susceptibility to respiratory infections, including COVID-19 (83). Nevertheless, some studies found a protective effect for COVID-19 infection in patients treated with antipsychotics (84, 85). Based on preclinical findings, the antipsychotics chlorpromazine and haloperidol have been suggested to offer protection against SARS-CoV-2, possibly through their interactions with sigma-1 receptors, inducing anti-inflammatory effects by inhibiting cytokine production (86–90). However, initial observational clinical studies did not confirm that these agents offer protection against COVID-19 infection or COVID-19-related mortality (84, 91). However, the results of these studies have to be interpreted with caution, because of possible confounding factors. Moreover, it is important to make a distinction between acute and long-term treatment effects of antipsychotic treatment.

Valproate, a mood stabilizer, also may be associated with an increased risk of respiratory infections (92). By contrast, lithium, another mood stabilizer, seems to be associated with a decreased risk of respiratory infections and demonstrated potential antiviral properties at a preclinical level (92, 93). Lithium has even been proposed as a candidate treatment for COVID-19. It can suppress NOD-like receptor family pyrin domain containing-3 (NLRP3) inflammasome activity (which is implicated in the release of pro-inflammatory cytokines during the cytokine storm), inhibits cell death (resulting in a decrease in lung parenchymal damage), and is characterized by immune-regulatory mechanisms (preventing the harmful effects of immune hyperactivation) (94). However, its antiviral properties, as well as its safety as a potential antiviral agent (due to its narrow therapeutic index and high risk of toxicity), remain to be confirmed in clinical settings (93, 95).

More promising is the association that has been observed between FIASMA (Functional Inhibitors of Acid Sphingomyelinase) treatments, including certain SSRI and non-SSRI antidepressants such as fluvoxamine and amitriptyline, and a reduction in clinical deterioration and mortality risk in patients with COVID-19. Acid sphingomyelinase (ASM) is an important lipid-metabolizing enzyme catalyzing the hydrolysis of sphingomyelin into ceramide and phosphorylcholine (91, 96). SARS-CoV-2 probably activates this ASM-ceramide system, facilitating viral entry and infection of human nasal epithelial cells by clustering ACE-2 receptors (97, 98). Functional Inhibitors of Acid Sphingomyelinase antidepressants are thought to impair SARS-CoV-2 entry into epithelial cells by functional inhibition of the ASM-ceramide system (99). Several retrospective (100–102) and prospective (103) observational studies, as well as a small double-blind randomized trial (104) showed that taking a FIASMA treatment was associated with a lower risk of clinical

deterioration or death in both non-psychiatric and psychiatric patients with COVID-19. The anti-inflammatory properties of certain antidepressants, probably due to their high affinity for sigma-1 receptors (105), may have additional value in managing COVID-19. Nevertheless, more large-scale double-blind controlled randomized clinical trials of these medications in patients with COVID-19 are needed (99, 106, 107).

Polypharmacy has been found to be associated with a higher risk of developing COVID-19 (108). Psychotropic polypharmacy is quite common in patients with SMI (12). According to a Swedish study, 25% of patients dispensed antipsychotic drugs receive a combination of two or more antipsychotic drugs. These patients also did more often take anxiolytics and sedatives than those prescribed antipsychotic monotherapy (109). Psychotropic polypharmacy, particularly during the treatment of elderly people with SMI, seems to be associated with greater adverse effects on most physical diseases, compared to monotherapy (4), as it carries the risk of adverse drug reactions. As the possible contribution of antipsychotic polypharmacy to the general excess mortality in people with SMI remains unclear (110, 111), further meta-analyses are needed analyzing mortality outcomes based on specific antipsychotic combinations rather than pooling data irrespective (111). The risks of adverse drug reactions due to psychotropic polypharmacy may be higher among certain regions in the world. Because of these aspects, the impact of psychotropic polypharmacy on COVID-19-related mortality therefore remains unknown.

Benzodiazepines (BZDs) and BZD-related medications (BZDRs) may also be of concern. A Swedish study in patients with SZ (112) showed that high exposure to BZDs is associated with an up to 70% higher mortality risk, compared with no users of BZDs. This is an important observation, knowing that the use of these medications by patients with SMI probably is even more common in other developed countries as the U.S (112). Although the risk of respiratory impairment associated with BZD use in the general population remains debated, in several studies current or recent exposure to certain BZDs or BZDRs has been found to be associated with an increased pneumonia risk (113), particularly in critically ill patients in intensive care units (114) or elderly (115, 116), whose immune system is vulnerable. BZDs and BZDRs, taken by 30–60% of individuals with SZ or BD (55), illnesses already characterized by a systemic pro-inflammatory state (see further), therefore may increase the risk for COVID-19-related mortality in these persons.

Finally, clinically relevant drug interactions between psychotropic medications and antiviral COVID-19 therapies may exist. The co-administration of protease inhibitors (blocking the protease enzyme that the virus needs to replicate), with certain antipsychotics (e.g., haloperidol and quetiapine), the mood stabilizer carbamazepine, or the BZDs midazolam or triazolam should be avoided because of increased toxicity and possible life-threatening events (117, 118).

Results of the studies included in our review indicate that COVID-19-related mortality risk, even after adjustment for all above mentioned factors, remains high in these patients (see **Table 1**). This indicates that SMI-related issues (i.e., immunological disturbances) may further increase the risk of

COVID-19-related mortality. In one of the included studies [i.e., (43)] the high risk of mortality associated with schizophrenia spectrum disorders ranked second behind age in strength of an association among all known demographic and medical risk factors examined.

Research has shown that disease-related immune dysregulation may provide some explanation for the higher susceptibility of people with SMI for severe clinical outcomes of COVID-19 (42, 83, 119, 120). Hyperactivation of the immune system, leading to excess release of pro-inflammatory cytokines (hypercytokinemia) or a “cytokine storm” (cytokine release syndrome), seems to play a major role in the process of disease aggravation in patients with COVID-19 infection (121–127). Circulating levels of inflammatory biomarkers, including interleukin-1 (IL-1), interleukin-6 (IL-6), tumor necrosis factor alpha (TNF- α), and C-reactive protein (CRP), are often excessively elevated during severe SARS-CoV-2 infection. This disproportionate release of cytokines beyond that of a controlled immune response has been associated with poor outcomes and an increased risk of mortality (22, 122, 124, 126, 128, 129). As SMIs already are often characterized by a systemic pro-inflammatory state or overproduction of pro-inflammatory cytokines, which may persist even after patients' symptoms have improved (32, 33), the systemic hyperinflammation triggered by SARS-CoV-2 infection may be more pronounced in these individuals, leading to excess tissue damage, multi-organ failure, and death. A reduction in lymphocyte natural killer cell activity (a common finding in severe COVID-19) in some patients with SMI may further explain why COVID-19-related mortality rates are higher in these individuals (119, 120, 130–132). However, these hypotheses remain to be tested more rigorously.

Particularly in older patients, perturbations in gut microbiome composition, which seem to be related to elevated concentrations of inflammatory cytokines, may exacerbate COVID-19-related severity (133). This observation may be important knowing that people with SMI present with various alterations of the gut microbiome (134).

Prevention Strategies and Possible Therapeutic Options

Given the strong association between COVID-19-related mortality and SMI, it is paramount that COVID-19 vaccination and equitable access to COVID-19 vaccines for people with SMI should be a matter of priority (135–137). This should be even more obvious knowing that the management of physical diseases (including comorbid conditions causing more severe COVID-19 illness) in people with SMI is already suboptimal, due to non-medical factors such as stigmatization and disparities in physical health care (10, 11, 13). It is therefore astonishing to note that some governments within and outside the European Union (e.g., India) are still doubting whether these individuals should be prioritized for COVID-19 vaccination (135).

However, only granting priority access to people with SMI in national vaccination strategies will not be sufficient, as a significant COVID-19 vaccination gap seems to exist between these individuals and the general population, despite having

been granted early universal or priority access to SARS-CoV-2 vaccination (138–140). Targeted interventions to maximize vaccination uptake among these patients will be needed (139, 140). There are ways mental health professionals and agencies can address barriers to COVID-19 vaccination, based on the Increasing Vaccination Model (IVM). Identifying and addressing internal conflicts (by using motivational interviewing), social network interventions (making clinician recommendations build on interpersonal trust), and direct behavior change interventions (including reminders and primes, automatic appointments, and presumptive healthcare professional communication) can be helpful in this regard (141). Developing an intentional vaccine delivery strategy in conjunction with experts, utilizing multiple communication channels, and expanding vaccine delivery outside of the hospital to reach patients can be another strategy (142). Finally, the involvement of peers, family, or volunteers to support people with SMI in making healthcare choices may also be helpful. These types of actions can pay off. Our research group has shown that vaccine willingness among patients with psychiatric disorders in our university psychiatric hospital with a targeted prevention program was just as high as in the general population: 93% or 1,070 of 1,151 patients who were offered COVID-19 vaccination accepted this vaccination (143). Other studies confirmed that vaccination willingness among these patients is at least almost as high as in the general population (144, 145).

Several therapeutic options for this vulnerable population may exist to reduce the increased mortality rate: the use of medications that target specific inflammatory markers, the use of a cytokine filter targeting multiple different cytokines at a larger scale, natural killer cell-based immunotherapies, and the use of nicotine, nicotinic receptor agonists, or positive modulators of these receptors (activation of these receptors, particularly $\alpha 7$ nAChR, can suppress production of pro-inflammatory cytokines as these receptors are abundantly expressed in a variety of immune cells) (122, 127, 130, 131). Early prediction of a cytokine storm is made possible by several biochemical and hematological markers (128). In addition to reducing pathogen exposure, individual immunity in this vulnerable population can be enhanced by promoting a healthy lifestyle, regular exercise, balanced nutrition, and quality sleep (125). However, more research on these therapeutic options is urgently needed.

Patients with SMI often have lower vitamin D levels (146). Several meta-analyses (147, 148) and publications (146, 149–153) suggest that vitamin D supplementation may be potentially effective in preventing COVID-19 infection and mitigating the clinical course of the disease. Study results, however, remain difficult to interpret due to possible confounding factors (149, 154).

STRENGTHS AND LIMITATIONS

Our review has particular strengths. Compared to other available systematic reviews, we developed a more comprehensive search strategy for the retrieval of reports of controlled trials. With the exception of Ceban et al. (38), our review is the only one that also

included the Embase database to obtain a more comprehensive coverage of the existing US and EU literature. Finally, a thorough and critical discussion of this issue is presented in this paper.

Nevertheless, our systematic and critical review has its limitations that are inherent to the nature of the available evidence, and in that respect comparable to the previously published reviews. An important limitation is that all included studies are observational and mostly retrospective, and therefore causal relationships cannot be inferred. Most of the included studies in our systematic review were carried out during the first wave of the COVID-19 pandemic, which lasted from February/March until May/June 2020. During this period, in most countries testing was largely restricted to individuals exhibiting symptoms or to certain risk groups, due to a limited PCR-testing capacity at that time. This biased deployment of testing can distort true estimates of COVID-19-related mortality rate in people with SMI. Most of the studies on mood disorders involved mixed populations and did not make a distinction as to whether an individual with a mood disorder had MDD or BD. The two largest studies on COVID-19-related mortality risk in patients with BD to date (42, 155) only included around 500 patients. This is an important limitation given the possible differential impact of COVID-19-related mortality risk across SMIs. Therefore, more studies are needed that present separate data for patients with BD and MDD to gain a better estimate of true COVID-19-related mortality risks associated with these groups of people with SMI. Control groups also varied across studies: some control groups excluded patients with psychiatric disorders, while others included non-SMI disorders. Finally, most studies did not provide detailed patient sociodemographic characteristics (i.e., mean/median age, gender), or other detailed information on clinical/psychiatric variables (such as severity and status of the disease), and smoking histories. Future studies therefore should further explore these issues to better understand which specific patients are at an increased risk of COVID-19-related mortality.

CONCLUSION

Even without taking COVID-19 into account, people with SMI already have a two to three times higher mortality rate

than the general population, largely attributable to somatic comorbidities. Our review has shown that individuals with SMI, particularly patients with schizophrenia spectrum disorders, are at significantly higher risk of COVID-19-related mortality, not only due to higher somatic comorbidity rates and the possible use of psychotropic medication, but also to unknown factors at the moment that will have to be explored in future research. Severe mental illness therefore should be identified as a separate, independent risk factor for a more severe clinical course when infected with COVID-19 and targeted as a high-risk population. Consequently, targeted interventions to maximize vaccination uptake among these people should be prioritized in health policy worldwide.

DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**, further inquiries can be directed to the corresponding author/s.

AUTHOR CONTRIBUTIONS

JD: search and analysis. All authors contributed to the article and approved the submitted version.

ACKNOWLEDGMENTS

The authors wish to thank Krizia Tuand and Thomas Vandendriessche, the biomedical reference librarians of the KU Leuven Libraries—2Bergen—Learning Centre Désiré Collen (Leuven, Belgium), for their help in conducting the systematic literature search.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fpsy.2021.798554/full#supplementary-material>

REFERENCES

- McIntyre RS, Berk M, Brietzke E, Goldstein BI, López-Jaramillo C, Kessing LV, et al. Bipolar disorders. *Lancet*. (2020) 396:1841–56. doi: 10.1016/S0140-6736(20)31544-0
- Malhi GS, Mann JJ. Depression. *Lancet*. (2018) 392:2299–312. doi: 10.1016/S0140-6736(18)31948-2
- Owen MJ, Sawa A, Mortensen PB. Schizophrenia. *Lancet*. (2016) 388:86–97. doi: 10.1016/S0140-6736(15)01121-6
- Correll CU, Detraux J, De Lepeleire J, De Hert M. Effects of antipsychotics, antidepressants and mood stabilizers on risk for physical diseases in people with schizophrenia, depression and bipolar disorder. *World Psychiatry*. (2015) 14:119–36. doi: 10.1002/wps.20204
- Hayes JE, Miles J, Walters K, King M, Osborn DP. A systematic and meta-analysis of premature mortality in bipolar affective disorder. *Acta Psychiatr Scand*. (2015) 131:417–25. doi: 10.1111/acps.12408
- Laursen TM, Musliner KL, Benros ME, Vestergaard M, Munk-Olsen Mortality and life expectancy in persons with severe unipolar depression. *J Affect Disord*. (2016) 193:203–7. doi: 10.1016/j.jad.2015.12.067
- Walker ER, McGee RE, Druss BG. Mortality in mental disorders and global disease burden implications: a systematic review and meta-analysis. *JAMA Psychiatry*. (2015) 72:334–41. doi: 10.1001/jamapsychiatry.2014.2502
- Liu NH, Daumit GL, Dua T, Aquila R, Charlson F, Cuijpers P, et al. Excess mortality in persons with severe mental disorders: a multilevel intervention framework and priorities for clinical practice, policy and research agendas. *World Psychiatry*. (2017) 16:30–40. doi: 10.1002/wps.20384

9. De Hert M, Detraux J. Reversing the downward spiral for people with severe mental illness through educational innovations. *World Psychiatry*. (2017) 16:41–2. doi: 10.1002/wps.20377
10. De Hert M, Correll CU, Bobes J, Cetkovich-Bakmas M, Cohen D, Asai I, et al. Physical illness in patients with severe mental disorders. I. Prevalence, impact of medications and disparities in health care. *World Psychiatry*. (2011) 10:52–77. doi: 10.1002/j.2051-5545.2011.tb00014.x
11. Kohn L, Christiaens W, Detraux J, De Lepeleire J, De Hert M, Gillain B, et al. Barriers to somatic health care for persons with severe mental illness in Belgium: a qualitative study of patients' and healthcare professionals' perspectives. *Front Psychiatry*. (2021). doi: 10.3389/fpsy.2021.798530. [Epub ahead of print].
12. De Hert M, Detraux J, Vancampfort D. Impact of medication effects on physical health. In: Corrigan PW, Ballentine SL, editors. *Health and Wellness in People Living with Serious Mental Illness*. Washington, DC: American Psychiatric Association Publishing (2021). p. 103–29.
13. De Hert M, Cohen D, Bobes J, Cetkovich-Bakmas M, Leucht S, Ndeti DM, et al. Physical illness in patients with severe mental disorders. II. Barriers to care, monitoring and treatment guidelines, plus recommendations at the system and individual level. *World Psychiatry*. (2011) 10:138–51. doi: 10.1002/j.2051-5545.2011.tb00036.x
14. Kim SW, Park WY, Jhon M, Kim M, Lee JY, Kim SY, et al. Physical health literacy and health related behaviors in patients with psychosis. *Clin Psychopharmacol Neurosci*. (2019) 17:279–87. doi: 10.9758/cpn.2019.17.2.279
15. Sheffield JM, Karcher NR, Barch DM. Cognitive deficits in psychotic disorders: a lifespan perspective. *Neuropsychol Rev*. (2018) 28:509–33. doi: 10.1007/s11065-018-9388-2
16. Tempelaar WM, Termorshuizen F, MacCabe JH, Boks MPM, Kahn RS. Educational achievement in psychiatric patients and their siblings: a register-based study in 30 000 individuals in the Netherlands. *Psychol Med*. (2017) 47:776–84. doi: 10.1017/S0033291716002877
17. Dickerson FB, Kreyenbuhl J, Goldberg RW, Fang L, Medoff D, Brown CH, et al. A 5-year follow-up of diabetes knowledge in persons with serious mental illness and type 2 diabetes. *J Clin Psychiatry*. (2009) 70:1057–8. doi: 10.4088/jcp.08l04602
18. Isohanni I, Jones PB, Järvelin MR, Nieminen P, Rantakallio P, Jokelainen J, et al. Educational consequences of mental disorders treated in hospital. A 31-year follow-up of the Northern Finland 1966 Birth Cohort. *Psychol Med*. (2001) 31:339–49. doi: 10.1017/s003329170100304x
19. Desforges M, Le Coupanec A, Dubeau P, Bourgouin A, Lajoie L, Dubé M, et al. Human coronaviruses and other respiratory viruses: underestimated opportunistic pathogens of the central nervous system? *Viruses*. (2019) 12:14. doi: 10.3390/v12010014
20. Uversky VN, Elrashdy F, Aljadawi A, Ali SM, Khan RH, Redwan EM. Severe acute respiratory syndrome coronavirus 2 infection reaches the human nervous system: how? *J Neurosci Res*. (2021) 99:750–77. doi: 10.1002/jnr.24752
21. Yassin A, Nawaiseh M, Shaban A, Alsherbini K, El-Salem K, Soudah O, et al. Neurological manifestations and complications of coronavirus disease 2019 (COVID-19): a systematic review and meta-analysis. *BMC Neurol*. (2021) 21:138. doi: 10.1186/s12883-021-02161-4
22. Liu C, Martins AJ, Lau WW, Rachmaninoff N, Chen J, Imberti L, et al. Time-resolved systems immunology reveals a late juncture linked to fatal COVID-19. *Cell*. (2021) 184:1836.e22–57.e22. doi: 10.1016/j.cell.2021.02.018
23. Ni W, Yang X, Yang D, Bao J, Li R, Xiao Y, et al. Role of angiotensin-converting enzyme 2 (ACE2) in COVID-19. *Crit Care*. (2020) 24:422. doi: 10.1186/s13054-020-03120-0
24. Maury A, Lyoubi A, Peiffer-Smadja N, de Broucker T, Meppiel E. Neurological manifestations associated with SARS-CoV-2 and other coronaviruses: a narrative review for clinicians. *Rev Neurol (Paris)*. (2021) 177:51–64. doi: 10.1016/j.neurol.2020.10.001
25. Schirinzi T, Landi D, Liguori C. COVID-19: dealing with a potential risk factor for chronic neurological disorders. *J Neurol*. (2021) 268:1171–8. doi: 10.1007/s00415-020-10131-y
26. Cagnazzo F, Arquizan C, Derraz I, Dargazanli C, Lefevre PH, Riquelme C, et al. Neurological manifestations of patients infected with the SARS-CoV-2: a systematic review of the literature. *J Neurol*. (2021) 268:2656–65. doi: 10.1007/s00415-020-10285-9
27. Collantes MEV, Espiritu AI, Sy MCC, Anlacan VMM, Jamora RDG. Neurological manifestations in COVID-19 infection: a systematic review and meta-analysis. *Can J Neurol Sci*. (2021) 48:66–76. doi: 10.1017/cjn.2020.146
28. Taquet M, Geddes JR, Husain M, Luciano S, Harrison PJ. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *Lancet Psychiatry*. (2021) 8:416–27. doi: 10.1016/S2215-0366(21)00084-5
29. Schurink B, Roos E, Radonic T, Barbe E, Bouman CSC, de Boer HH, et al. Viral presence and immunopathology in patients with lethal COVID-19: a prospective autopsy cohort study. *Lancet Microbe*. (2020) 1:e290–9. doi: 10.1016/S2666-5247(20)30144-0
30. Douaud G, Lee S, Alfaro-Almagro F, Arthofer C, Wang C, Lange F, et al. Brain imaging before and after COVID-19 in UK Biobank. *medRxiv [Preprint]*. (2021) 2021.06.11.21258690. doi: 10.1101/2021.06.11.21258690
31. Roy D, Ghosh R, Dubey S, Dubey MJ, Benito-León J, Kanti Ray B. Neurological and neuropsychiatric impacts of COVID-19 pandemic. *Can J Neurol Sci*. (2021) 48:9–24. doi: 10.1017/cjn.2020.173
32. Rajkumar RP. Cross-national variations in COVID-19 mortality: the role of diet, obesity and depression. *Diseases*. (2021) 9:36. doi: 10.3390/diseases9020036
33. Yang H, Chen W, Hu Y, Chen Y, Zeng Y, Sun Y, et al. Pre-pandemic psychiatric disorders and risk of COVID-19: a UK Biobank cohort analysis. *Lancet Healthy Longev*. (2020) 1:e69–79. doi: 10.1016/S2666-7568(20)30013-1
34. Brown E, Gray R, Lo Monaco S, O'Donoghue B, Nelson B, Thompson A, et al. The potential impact of COVID-19 on psychosis: a rapid review of contemporary epidemic and pandemic research. *Schizophr Res*. (2020) 222:79–87. doi: 10.1016/j.schres.2020.05.005
35. Dinakaran D, Manjunatha N, Naveen Kumar C, Suresh BM. Neuropsychiatric aspects of COVID-19 pandemic: a selective review. *Asian J Psychiatr*. (2020) 53:102188. doi: 10.1016/j.ajp.2020.102188
36. Vai B, Mazza MG, Delli Colli C, Foiselle M, Allen B, Benedetti F, et al. Mental disorders and risk of COVID-19-related mortality, hospitalisation, and intensive care unit admission: a systematic review and meta-analysis. *Lancet Psychiatry*. (2021) 8:797–812. doi: 10.1016/S2215-0366(21)0232-7
37. Fond G, Nemani K, Etchecopar-Etchart D, Loundou A, Goff DC, Lee SW, et al. Association between mental health disorders and mortality among patients with covid-19 in 7 countries: a systematic review and meta-analysis. *JAMA Psychiatry*. (2021) 78:1208–17. doi: 10.1001/jamapsychiatry.2021.2274
38. Ceban F, Nogo D, Carvalho IP, Lee Y, Nasri F, Xiong J, et al. Association between mood disorders and risk of COVID-19 infection, hospitalization, and death: a systematic review and meta-analysis. *JAMA Psychiatry*. (2021) 2021:e211818. doi: 10.1001/jamapsychiatry.2021.1818
39. Liu L, Ni SY, Yan W, Lu QD, Zhao YM, Xu YY, et al. Mental and neurological disorders and risk of COVID-19 susceptibility, illness severity and mortality: a systematic review, meta-analysis and call for action. *EClinicalMedicine*. (2021) 40:101111. doi: 10.1016/j.eclinm.2021.101111
40. Toubasi AA, AbuAnzeh RB, Tawileh HBA, Aldebei RH, Alryalat SAS. A meta-analysis: the mortality and severity of COVID-19 among patients with mental disorders. *Psychiatry Res*. (2021) 299:113856. doi: 10.1016/j.psychres.2021.13856

41. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. (2021) 372:n71. doi: 10.1136/bmj.n71
42. Barcella CA, Polcwiartek C, Mohr GH, Hodges G, Søndergaard K, Niels Bang C, et al. Severe mental illness is associated with increased mortality and severe course of COVID-19. *Acta Psychiatr Scand*. (2021) 144:82–91. doi: 10.1111/acps.13309
43. Nemani K, Li C, Olfson M, Blessing EM, Razavian N, Chen J, et al. Association of psychiatric disorders with mortality among patients with COVID-19. *JAMA Psychiatry*. (2021) 78:380–6. doi: 10.1001/jamapsychiatry.2020.4442
44. Tzur Bitan D, Krieger I, Kridin K, Komantscher D, Scheinman Y, Weinstein O, et al. COVID-19 prevalence and mortality among schizophrenia patients: a large-scale retrospective cohort study. *Schizophr Bull*. (2021) 47:1211–7. doi: 10.1093/schbul/sbab012
45. Jeon HL, Kwon JS, Park SH, Shin JY. Association of mental disorders with SARS-CoV-2 infection and severe health outcomes: nationwide cohort study. *Br J Psychiatry*. (2021) 2021:1–8. doi: 10.1192/bjp.2020.20251
46. Egede C, Dawson AZ, Walker RJ, Garacci E, Campbell JA, Egede LE. Relationship between mental health diagnoses and COVID-19 test positivity, hospitalization, and mortality in Southeast Wisconsin. *Psychol Med*. (2021) 2021:1–269. doi: 10.1017/S0033291721002312
47. Fond G, Pauly V, Orleans V, Antonini F, Fabre C, Sanz M, et al. Increased in-hospital mortality from COVID-19 in patients with schizophrenia. *Encephale*. (2021) 47:89–95. doi: 10.1016/j.encep.2020.07.003
48. Fond G, Pauly V, Leone M, Llorca PM, Orleans V, Loundou A, et al. Disparities in intensive care unit admission and mortality among patients with schizophrenia and COVID-19: a national cohort study. *Schizophr Bull*. (2021) 47:624–34. doi: 10.1093/schbul/sbaa158
49. Reilev M, Kristensen KB, Pottegård A, Lund LC, Hallas J, Ernst MT, et al. Characteristics and predictors of hospitalization and death in the first 11 122 cases with a positive RT-PCR test for SARS-CoV-2 in Denmark: a nationwide cohort. *Int J Epidemiol*. (2020) 49:1468–81. doi: 10.1093/ije/dyaa140
50. Poblador-Plou B, Carmona-Pirez J, Ioakeim-Skoufa I, Poncel-Falcó A, Blik-Bueno K, Cano-Del Pozo M, et al. Baseline chronic comorbidity and mortality in laboratory-confirmed COVID-19 cases: results from the PRECOVID Study in Spain. *Int J Environ Res Public Health*. (2020) 17:5171. doi: 10.3390/ijerph17145171
51. Castro VM, Gunning FM, McCoy TH, Perlis RH. Mood disorders and outcomes of covid-19 hospitalizations. *Am J Psychiatry*. (2021) 178:541–7. doi: 10.1176/appi.ajp.2020.20060842
52. Diez-Quevedo C, Iglesias-González M, Giralto-López M, Rangil T, Sanagustin D, Moreira M, et al. Mental disorders, psychopharmacological treatments, and mortality in 2150 COVID-19 Spanish inpatients. *Acta Psychiatr Scand*. (2021) 143:526–34. doi: 10.1111/acps.13304
53. Wang Q, Xu R, Volkow ND. Increased risk of COVID-19 infection and mortality in people with mental disorders: analysis from electronic health records in the United States. *World Psychiatry*. (2021) 20:124–30. doi: 10.1002/wps.20806
54. Das-Munshi J, Chang KC, Bakolis I, Broadbent M, Dregan A, Hotopf M. All-cause and cause-specific mortality in people with mental disorders and intellectual disabilities, before and during the COVID-19 pandemic: cohort study. *Lancet Reg Health Eur*. (2021) 11:100228. doi: 10.1016/j.lanpe.2021.100228
55. Maripuu M, Bendix M, Öhlund L, Widerström M, Werneke U. Death associated with coronavirus (covid-19) infection in individuals with severe mental disorders in Sweden during the early months of the outbreak—an exploratory cross-sectional analysis of a population-based register study. *Front Psychiatry*. (2021) 11:609579. doi: 10.3389/fpsy.2020.609579
56. De Picker LJ. Closing COVID-19 mortality, vaccination, and evidence gaps for those with severe mental illness. *Lancet Psychiatry*. (2021) 8:854–5. doi: 10.1016/S2215-0366(21)00291-1
57. Pijls BG, Jolani S, Atherley A, Derckx RT, Dijkstra JIR, Franssen GH, et al. Demographic risk factors for COVID-19 infection, severity, ICU admission and death: a meta-analysis of 59 studies. *BMJ Open*. (2021) 11:e044640. doi: 10.1136/bmjopen-2020-044640
58. Bae S, Kim SR, Kim MN, Shim WJ, Park SM. Impact of cardiovascular disease and risk factors on fatal outcomes in patients with COVID-19 according to age: a systematic review and meta-analysis. *Heart*. (2021) 107:373–80. doi: 10.1136/heartjnl-2020-317901
59. Silverio A, Di Maio M, Citro R, Esposito L, Iuliano G, Bellino M, et al. Cardiovascular risk factors and mortality in hospitalized patients with COVID-19: systematic review and meta-analysis of 45 studies and 18,300 patients. *BMC Cardiovasc Disord*. (2021) 21:23. doi: 10.1186/s12872-020-01816-3
60. Liu Y, Lu H, Wang W, Liu Q, Zhu C. Clinical risk factors for mortality in patients with cancer and COVID-19: a systematic review and meta-analysis of recent observational studies. *Expert Rev Anticancer Ther*. (2021) 21:107–19. doi: 10.1080/14737140.2021.1837628
61. Zhang L, Hou J, Ma FZ, Li J, Xue S, Xu ZG. The common risk factors for progression and mortality in COVID-19 patients: a meta-analysis. *Arch Virol*. (2021) 166:2071–87. doi: 10.1007/s00705-021-05012-2
62. Corona G, Pizzocaro A, Vena W, Rastrelli G, Semeraro F, Isidori AM, et al. Diabetes is most important cause for mortality in COVID-19 hospitalized patients: systematic review and meta-analysis. *Rev Endocr Metab Disord*. (2021) 22:275–96. doi: 10.1007/s11154-021-09630-8
63. Jordan RE, Adab P, Cheng KK. Covid-19: risk factors for severe disease and death. *BMJ*. (2020) 368:m1198. doi: 10.1136/bmj.m1198
64. Laursen TM. Causes of premature mortality in schizophrenia: a review of literature published in 2018. *Curr Opin Psychiatry*. (2019) 32:388–93. doi: 10.1097/YCO.0000000000000530
65. Eyder LT, Jeste DV. Aging of the body and the brain in schizophrenia. *Schizophr Res*. (2018) 196:1–3. doi: 10.1016/j.schres.2017.11.036
66. De Picker LJ, Yolken R, Benedetti F, Borsini A, Branchi I, Fusar-Poli P, et al. ECNP Immuno-NeuroPsychiatry TWG. Viewpoint | European COVID-19 exit strategy for people with severe mental disorders: too little, but not yet too late. *Brain Behav Immun*. (2021) 94:15–7. doi: 10.1016/j.bbi.2021.01.008
67. Lin CW, Chang LC, Ma T, Oh H, French B, Puralowski R, et al. Older molecular brain age in severe mental illness. *Mol Psychiatry*. (2020) 10.1038/s41380-020-0834-1. doi: 10.1038/s41380-020-0834-1
68. Karanth S, Fowler ME, Mao X, Wilson LE, Huang B, Pisu M, et al. Race, socioeconomic status, and health-care access disparities in ovarian cancer treatment and mortality: systematic review and meta-analysis. *JNCI Cancer Spectr*. (2019) 3:pkz084. doi: 10.1093/jncics/pkz084
69. Hawkins RB, Charles EJ, Mehaffey JH. Socio-economic status and COVID-19-related cases and fatalities. *Public Health*. (2020) 189:129–34. doi: 10.1016/j.puhe.2020.09.016
70. Gaglioti AH, Li C, Douglas MD, Baltrus PT, Blount MA, Zahidi R, et al. Population-level disparities in COVID-19: measuring the independent association of the proportion of black population on COVID-19 cases and deaths in US counties. *J Public Health Manag Pract*. (2021) 27:268–77. doi: 10.1097/PHH.0000000000001354
71. Brody BD, Parish SJ, Kanellopoulos D, Russ MJ, A. COVID-19 testing and triage algorithm for psychiatric units: one hospital's response to the New York region's pandemic. *Psychiatry Res*. (2020) 291:113244. doi: 10.1016/j.psychres.2020.113244
72. Rojnic Kuzman M, Vahip S, Fiorillo A, Beezhold J, Pinto da Costa M, Skugarevsky O, et al. Mental health services during the first wave of the COVID-19 pandemic in Europe: results from the EPA Ambassadors Survey and implications for clinical practice. *Eur Psychiatry*. (2021) 64:e41. doi: 10.1192/j.eurpsy.2021.2215
73. Rabbani G, Shariful Islam SM, Rahman MA, Amin N, Marzan B, Robin RC, et al. Pre-existing COPD is associated with an increased risk of mortality and severity in COVID-19: a rapid systematic review and meta-analysis. *Expert Rev Respir Med*. (2021) 15:705–16. doi: 10.1080/17476348.2021.1866547
74. Alqahtani JS, Oyelade T, Aldhahir AM, Alghamdi SM, Almeahmadi M, Alqahtani AS, et al. Prevalence, severity and mortality associated

- with COPD and smoking in patients with COVID-19: a rapid systematic review and meta-analysis. *PLoS ONE*. (2020) 15:e0233147. doi: 10.1371/journal.pone.0233147
75. Rosenthal N, Cao Z, Gundrum J, Sianis J, Safo S. Risk factors associated with in-hospital mortality in a US national sample of patients with COVID-19. *JAMA Netw Open*. (2020) 3:e2029058. doi: 10.1001/jamanetworkopen.2020.29058
 76. Bajgain KT, Badal S, Bajgain BB, Santana MJ. Prevalence of comorbidities among individuals with COVID-19: a rapid review of current literature. *Am J Infect Control*. (2021) 49:238–46. doi: 10.1016/j.ajic.2020.06.213
 77. CDC COVID-19 Response Team. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019 - United States, February 12-March 28, 2020. *MMWR Morb Mortal Wkly Rep*. (2020) 69:382–6. doi: 10.15585/mmwr.mm6913e2
 78. Shinn AK, Viron M. Perspectives on the COVID-19 pandemic and individuals with serious mental illness. *J Clin Psychiatry*. (2020) 81:20com13412. doi: 10.4088/JCP.20com13412
 79. Nemani K, Conderino S, Marx J, Thorpe LE, Goff DC. Association between antipsychotic use and COVID-19 mortality among people with serious mental illness. *JAMA Psychiatry*. (2021) e212503. doi: 10.1001/jamapsychiatry.2021.2503
 80. Haga T, Ito K, Sakashita K, Iguchi M, Ono M, Tatsumi K. Risk factors for pneumonia in patients with schizophrenia. *Neuropsychopharmacol Rep*. (2018) 38:204–9. doi: 10.1002/npr2.12034
 81. Yang SY, Liao YT, Liu HC, Chen WJ, Chen CC, Kuo CJ. Antipsychotic drugs, mood stabilizers, and risk of pneumonia in bipolar disorder: a nationwide case-control study. *J Clin Psychiatry*. (2013) 74:e79–86. doi: 10.4088/JCP.12m07938
 82. Kuo CJ, Yang SY, Liao YT, Chen WJ, Lee WC, Shau WY, et al. Second-generation antipsychotic medications and risk of pneumonia in schizophrenia. *Schizophr Bull*. (2013) 39:648–57. doi: 10.1093/schbul/sbr202
 83. May M, Slitzky M, Rostama B, Barlow D, Houseknecht KL. Antipsychotic-induced immune dysfunction: a consideration for COVID-19 risk. *Brain Behav Immun Health*. (2020) 6:100097. doi: 10.1016/j.bbih.2020.100097
 84. Prokopez CR, Vallejos M, Lopredo LS, Sfriso LE, Chiappella LC, Arce C, et al. An analysis of the possible protective effect of antipsychotics for SARS-CoV-2 in patients under treatment for severe mental illnesses. *Schizophr Res*. (2021) 233:99–100. doi: 10.1016/j.schres.2021.06.019
 85. Moga S, Teodorescu A, Ifteni P, Gavris C, Petric PS. Inflammatory response in SARS-CoV-2 infection of patients with schizophrenia and long-term antipsychotic treatment. *Neuropsychiatr Dis Treat*. (2021) 17:3053–60. <https://doi.org/10.2147/NDT.S325062>
 86. Plaze M, Attali D, Prot M, Petit AC, Blatzer M, Vinckier F, et al. Inhibition of the replication of SARS-CoV-2 in human cells by the FDA-approved drug chlorpromazine. *Int J Antimicrob Agents*. (2021) 57:106274. doi: 10.1016/j.ijantimicag.2020.106274
 87. Plaze M, Attali D, Petit AC, Blatzer M, Simon-Lorier E, Vinckier F, et al. Repurposing chlorpromazine to treat COVID-19: the reCoVery study. *Encephale*. (2020) 46:169–72. doi: 10.1016/j.encep.2020.05.006
 88. Vaugeois J-M. Psychotropic drugs with cationic amphiphilic properties may afford some protection against SARS-CoV-2: a mechanistic hypothesis. *Psychiatry Res*. (2020) 291:113220. doi: 10.1016/j.psychres.2020.113220
 89. Brimson JM, Prasanth MI, Malar DS, Brimson S, Thitilertdecha P, Tencomnao T. Drugs that offer the potential to reduce hospitalization and mortality from SARS-CoV-2 infection: the possible role of the sigma-1 receptor and autophagy. *Expert Opin Ther Targets*. (2021) 25:435–49. doi: 10.1080/14728222.2021.1952987
 90. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, et al. A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature*. (2020) 583:459–68. doi: 10.1038/s41586-020-2286-9
 91. Hoertel N, Sánchez-Rico M, Vernet R, Jannot AS, Neuraz A, Blanco C, et al. AP-HP/Universities/INSERM Covid-19 research collaboration and AP-HP Covid CDR Initiative. Observational study of haloperidol in hospitalized patients with COVID-19. *PLoS ONE*. (2021) 16:e0247122. doi: 10.1371/journal.pone.0247122
 92. Landén M, Larsson H, Lichtenstein P, Westin J, Song J. Respiratory infections during lithium and valproate medication: a within-individual prospective study of 50,000 patients with bipolar disorder. *Int J Bipolar Disord*. (2021) 9:4. doi: 10.1186/s40345-020-00208-y
 93. Murru A, Manchia M, Hajek T, Nielsen RE, Rybakowski JK, Sani G, et al. International Group for The Study of Lithium Treated Patients (IGSLI). Lithium's antiviral effects: a potential drug for CoVID-19 disease?. *Int J Bipolar Disord*. (2020) 8:21. doi: 10.1186/s40345-020-00191-4
 94. Qaswal AB, Suleiman A, Guzu H, Harb T, Atiyat B. The potential role of lithium as an antiviral agent against SARS-CoV-2 via membrane depolarization: review and hypothesis. *Sci Pharm*. (2021) 89:11. doi: 10.3390/scipharm89010011
 95. Rajkumar RP. Lithium as a candidate treatment for COVID-19: promises and pitfalls. *Drug Dev Res*. (2020) 81:782–5. doi: 10.1002/ddr.21701
 96. Kornhuber J, Rhein C, Müller CP, Mühle C. Secretory sphingomyelinase in health and disease. *Biol Chem*. (2015) 396:707–36. doi: 10.1515/hsz-2015-0109
 97. Carpinteiro A, Edwards MJ, Hoffmann M, Kochs G, Gripp B, Weigang S, et al. Pharmacological inhibition of acid sphingomyelinase prevents uptake of SARS-CoV-2 by epithelial cells. *Cell Rep Med*. (2020) 1:100142. doi: 10.1016/j.xcrm.2020.100142
 98. Carpinteiro A, Gripp B, Hoffmann M, Pöhlmann S, Hoertel N, Edwards MJ, et al. Inhibition of acid sphingomyelinase by ambroxol prevents SARS-CoV-2 entry into epithelial cells. *J Biol Chem*. (2021) 296:100701. doi: 10.1016/j.jbc.2021.100701
 99. Hoertel N, Sánchez-Rico M, Cougoule C, Gulbins E, Kornhuber J, Carpinteiro A, et al. Repurposing antidepressants inhibiting the sphingomyelinase acid/ceramide system against COVID-19: current evidence and potential mechanisms. *Mol Psychiatry*. (2021) 2021:1–2. doi: 10.1038/s41380-021-01254-3
 100. Hoertel N, Sánchez-Rico M, Gulbins E, Kornhuber J, Carpinteiro A, Lenze EJ, et al. AP-HP/Université de Paris/INSERM COVID-19 research collaboration, AP-HP COVID CDR Initiative, “Entrepôt de Données de Santé” AP-HP Consortium. Association between FIASMA and reduced risk of intubation or death in individuals hospitalized for severe COVID-19: an observational multicenter study. *Clin Pharmacol Ther*. (2021) 110:1498–1511. doi: 10.1002/cpt.2317
 101. Hoertel N, Sánchez-Rico M, Vernet R, Beeker N, Jannot AS, Neuraz A, et al. AP-HP/Universities/INSERM COVID-19 Research Collaboration and AP-HP COVID CDR Initiative. Association between antidepressant use and reduced risk of intubation or death in hospitalized patients with COVID-19: results from an observational study. *Mol Psychiatry*. (2021) 26:5199–5212. doi: 10.1038/s41380-021-01021-4
 102. Darquennes G, Le Corre P, Le Moine O, Loas G. Association between functional inhibitors of acid sphingomyelinase (FIASMA) and reduced risk of death in COVID-19 patients: a retrospective cohort study. *Pharmaceuticals (Basel)*. (2021) 14:226. doi: 10.3390/ph14030226
 103. Seftel D, Boulware DR. Prospective cohort of fluvoxamine for early treatment of coronavirus disease 19. *Open Forum Infect Dis*. (2021) 8:ofab050. doi: 10.1093/ofid/ofab050
 104. Lenze EJ, Mattar C, Zorumski CF, Stevens A, Schweiger J, Nicol GE, et al. Fluvoxamine vs placebo and clinical deterioration in outpatients with symptomatic COVID-19: a randomized clinical trial. *JAMA*. (2020) 324:2292–300. doi: 10.1001/jama.2020.22760
 105. Sukhatme VP, Reiersen AM, Vayttaden SJ, Sukhatme VV. Fluvoxamine: a review of its mechanism of action and its role in COVID-19. *Front Pharmacol*. (2021) 12:652688. doi: 10.3389/fphar.2021.652688
 106. Salles J, Briand-Mésange F, Trudel S, Ausseil J, Salles JP, Chap H. Can antidepressants unlock prescription of rimonabant in the fight against COVID-19? *Mol Psychiatry*. (2021) 2021:1–2. doi: 10.1038/s41380-021-01221-y
 107. Stip E, Arnone D, Abdel Aziz K, Javaid SF. Diversity of mechanism of action of psychotropic drugs in their anti-COVID-19 properties. *Mol Psychiatry*. (2021) 2021:1–5. doi: 10.1038/s41380-021-01222-x
 108. McQueenie R, Foster HME, Jani BD, Katikireddi SV, Sattar N, Pell JB, et al. Multimorbidity, polypharmacy, and COVID-19 infection within the UK Biobank cohort. *PLoS ONE*. (2020) 15:e0238091. doi: 10.1371/journal.pone.0238091

109. Bergendal A, Schiöler H, Wettermark B, Björkstén KS. Concomitant use of two or more antipsychotic drugs is common in Sweden. *Ther Adv Psychopharmacol.* (2015) 5:224–31. doi: 10.1177/2045125315588647
110. Kadra G, Stewart R, Shetty H, MacCabe JH, Chang CK, Taylor D, et al. Long-term antipsychotic polypharmacy prescribing in secondary mental health care and the risk of mortality. *Acta Psychiatr Scand.* (2018) 138:123–32. doi: 10.1111/acps.12906
111. Buhagiar K, Templeton G, Blyth H, Dey M, Giacco D. Mortality risk from long-term treatment with antipsychotic polypharmacy vs monotherapy among adults with serious mental illness: a systematic review and meta-analysis of observational studies. *Schizophr Res.* (2020) S223:18–28. doi: 10.1016/j.schres.2020.08.026
112. Tiihonen J, Mittendorfer-Rutz E, Torniainen M, Alexanderson K, Tanskanen A. Mortality and cumulative exposure to antipsychotics, antidepressants, and benzodiazepines in patients with schizophrenia: an observational follow-up study. *Am J Psychiatry.* (2016) 173:600–6. doi: 10.1176/appi.ajp.2015.15050618
113. Sun GQ, Zhang L, Zhang LN, Wu Z, Hu DF. Benzodiazepines or related drugs and risk of pneumonia: a systematic review and meta-analysis. *Int J Geriatr Psychiatry.* (2019) 34:513–21. doi: 10.1002/gps.5048
114. Caroff DA, Szumita PM, Klompas M. The relationship between sedatives, sedative strategy, and healthcare-associated infection: a systematic review. *Infect Control Hosp Epidemiol.* (2016) 37:1234–42. doi: 10.1017/ice.2016.129
115. Huang KH, Tai CJ, Kuan YH, Chang YC, Tsai TH, Lee CY. Pneumonia risk associated with the use of individual benzodiazepines and benzodiazepine related drugs among the elderly with Parkinson's Disease. *Int J Environ Res Public Health.* (2021) 18:9410. doi: 10.3390/ijerph18179410
116. Taipale H, Tolppanen AM, Koponen M, Tanskanen A, Lavikainen P, Sund R, et al. Risk of pneumonia associated with incident benzodiazepine use among community-dwelling adults with Alzheimer disease. *CMAJ.* (2017) 189:E519–29. doi: 10.1503/cmaj.160126
117. Gatti M, De Ponti F, Pea F. Clinically significant drug interactions between psychotropic agents and repurposed COVID-19 therapies. *CNS Drugs.* (2021) 35:345–84. doi: 10.1007/s40263-021-00811-2
118. University of Liverpool. *COVID-19 Drug Interactions.* Available online at: <https://www.covid19-druginteractions.org/checker>. (accessed December 18, 2021).
119. Momtazmanesh S, Zare-Shahabadi A, Rezaei N. Cytokine alterations in schizophrenia: an updated review. *Front Psychiatry.* (2019) 10:892. doi: 10.3389/fpsy.2019.00892
120. Mohan M, Perry BI, Saravanan P, Singh SP. COVID-19 in people with schizophrenia: potential mechanisms linking schizophrenia to poor prognosis. *Front Psychiatry.* (2021) 12:666067. doi: 10.3389/fpsy.2021.666067
121. Mukaetova-Ladinska EB, Kronenberg G. Psychological and neuropsychiatric implications of COVID-19. *Eur Arch Psychiatry Clin Neurosci.* (2021) 271:235–48. doi: 10.1007/s00406-020-01210-2
122. Rizvi S, Danic M, Silver M, LaBond V. Cytosorb filter: an adjunct for survival in the COVID-19 patient in cytokine storm? A case report. *Heart Lung.* (2021) 50:44–50. doi: 10.1016/j.hrtng.2020.09.007
123. Ye Q, Wang B, Mao J. The pathogenesis and treatment of the 'Cytokine Storm' in COVID-19. *J Infect.* (2020) 80:607–13. doi: 10.1016/j.jinf.2020.03.037
124. Kar SK, Verma P, Verma SK, Saxena SK. Mental illnesses among COVID-19 patients: possible immunological underpinnings. *Asian J Psychiatr.* (2020) 53:102376. doi: 10.1016/j.ajp.2020.102376
125. Kim SW, Su KP. Using psychoneuroimmunity against COVID-19. *Brain Behav Immun.* (2020) 87:4–5. doi: 10.1016/j.bbi.2020.03.025
126. Liu F, Li L, Xu M, Wu J, Luo D, Zhu Y, et al. Prognostic value of interleukin-6, C-reactive protein, and procalcitonin in patients with COVID-19. *J Clin Virol.* (2020) 127:104370. doi: 10.1016/j.jcv.2020.104370
127. Tizabi Y, Getachew B, Copeland RL, Aschner M. Nicotine and the nicotinic cholinergic system in COVID-19. *FEBS J.* (2020) 287:3656–63. doi: 10.1111/febs.15521
128. Rowaiye AB, Okpalefe OA, Onuh Adejoke O, Ogidigo JO, Hannah Oladipo O, Ogu AC, et al. Attenuating the effects of novel COVID-19 (SARS-CoV-2) infection-induced cytokine storm and the implications. *J Inflamm Res.* (2021) 14:1487–510. doi: 10.2147/JIR.S301784
129. Moradian N, Gouravani M, Salehi MA, Heidari A, Shafeghat M, Hamblin MR, et al. Cytokine release syndrome: inhibition of pro-inflammatory cytokines as a solution for reducing COVID-19 mortality. *Eur Cytokine Netw.* (2020) 31:81–93. doi: 10.1684/ec.2020.0451
130. Bao C, Tao X, Cui W, Hao Y, Zheng S, Yi B, et al. Natural killer cells associated with SARS-CoV-2 viral RNA shedding, antibody response and mortality in COVID-19 patients. *Exp Hematol Oncol.* (2021) 10:5. doi: 10.1186/s40164-021-00199-1
131. Market M, Angka L, Martel AB, Bastin D, Olanubi O, Tennakoon G, et al. Flattening the COVID-19 curve with natural killer cell based immunotherapies. *Front Immunol.* (2020) 11:1512. doi: 10.3389/fimmu.2020.01512
132. Manickam C, Sugawara S, Reeves RK. Friends or foes? The knowns and unknowns of natural killer cell biology in COVID-19 and other coronaviruses in July 2020. *PLoS Pathog.* (2020) 16:e1008820. doi: 10.1371/journal.ppat.1008820
133. Vignesh R, Swathirajan CR, Tun ZH, Rameshkumar MR, Solomon SS, Balakrishnan P. Could perturbation of gut microbiota possibly exacerbate the severity of COVID-19 via cytokine storm? *Front Immunol.* (2021) 11:607734. doi: 10.3389/fimmu.2020.607734
134. Nguyen TT, Kosciolk T, Eyler LT, Knight R, Jeste DV. Overview and systematic review of studies of microbiome in schizophrenia and bipolar disorder. *J Psychiatr Res.* (2018) 99:50–61. doi: 10.1016/j.jpsychires.2018.01.013
135. The Lancet Psychiatry. A matter of priority: COVID-19 vaccination and mental health. *Lancet Psychiatry.* (2021) 8:551. doi: 10.1016/S2215-0366(21)00209-1
136. De Hert M, Mazereel V, Detraux J, Van Assche K. Prioritizing COVID-19 vaccination for people with severe mental illness. *World Psychiatry.* (2021) 20:54–5. doi: 10.1002/wps.20826
137. Mazereel V, Van Assche K, Detraux J, De Hert M. COVID-19 vaccination for people with severe mental illness: why, what, and how? *Lancet Psychiatry.* (2021) 8:444–50. doi: 10.1016/S2215-0366(20)30564-2
138. Tzur Bitan D, Kridin K, Cohen AD, Weinstein O. COVID-19 hospitalisation, mortality, vaccination, and postvaccination trends among people with schizophrenia in Israel: a longitudinal cohort study. *Lancet Psychiatry.* (2021) 8:901–8. doi: 10.1016/S2215-0366(21)00256-X
139. Curtis HJ, Inglesby P, Morton CE, MacKenna B, Walker AJ, Morley J, et al. Trends and clinical characteristics of COVID-19 vaccine recipients: a federated analysis of 57.9 million patients' primary care records *in situ* using OpenSAFELY. *MedRxiv.* (2021) 01.25.21250356. doi: 10.1101/2021.01.25.21250356
140. De Picker LJ, Dias MC, Benros ME, Vai B, Branchi I, Benedetti F, et al. Severe mental illness and European COVID-19 vaccination strategies. *Lancet Psychiatry.* (2021) 8:356–9. doi: 10.1016/S2215-0366(21)00046-8
141. Brewer NT, Abad N. Ways that mental health professionals can encourage COVID-19 vaccination. *JAMA Psychiatry.* (2021). doi: 10.1001/jamapsychiatry.2021.2951
142. Beste LA, Chen A, Geyer J, Wilson M, Schuttner L, Wheat C, et al. Best practices for an equitable Covid-19 vaccination program. *NEJM Catal Innov Care Deliv.* (2021) 2:238. doi: 10.1056/CAT.21.0238
143. Mazereel V, Vanbrabant T, Desplenter F, De Hert M. COVID-19 vaccine uptake in patients with psychiatric disorders admitted to or residing in a university psychiatric hospital. *Lancet Psychiatry.* (2021) 8:860–1. doi: 10.1016/S2215-0366(21)00301-1
144. Jepsen OH, Kølbaek P, Gil Y, Speed M, Dinesen PT, Sønderkov KM, et al. COVID-19 vaccine willingness amongst patients with mental illness compared with the general population. *Acta Neuropsychiatr.* (2021) 33:273–6. doi: 10.1017/neu.2021.15
145. Danenberg R, Shemesh S, Tzur Bitan D, Maoz H, Saker T, Dror C, et al. Attitudes of patients with severe mental illness towards COVID-19 vaccinations: a preliminary report from a public psychiatric hospital. *J Psychiatr Res.* (2021) 143:16–20. doi: 10.1016/j.jpsychires.2021.08.020
146. Werneke U, Gaughran F, Taylor DM. Vitamin D in the time of the coronavirus (COVID-19) pandemic - a clinical review from a public health and public mental health perspective. *Ther Adv Psychopharmacol.* (2021) 11:20451253211027699. doi: 10.1177/20451253211027699

147. Liu N, Sun J, Wang X, Zhang T, Zhao M, Li H. Low vitamin D status is associated with coronavirus disease 2019 outcomes: a systematic review and meta-analysis. *Int J Infect Dis.* (2021) 104:58–64. doi: 10.1016/j.ijid.2020.12.077
148. Teshome A, Adane A, Girma B, Mekonnen ZA. The impact of vitamin D Level on COVID-19 infection: systematic review and meta-analysis. *Front Public Health.* (2021) 9:624559. doi: 10.3389/fpubh.2021.624559
149. Ferrari D, Locatelli M, Briguglio M, Lombardi G. Is there a link between vitamin D status, SARS-CoV-2 infection risk and COVID-19 severity? *Cell Biochem Funct.* (2021) 39:35–47. doi: 10.1002/cbf.3597
150. Faniyi AA, Lugg ST, Faustini SE, Webster C, Duffy JE, Hewison M, et al. Vitamin D status and seroconversion for COVID-19 in UK healthcare workers. *Eur Respir J.* (2021) 57:2004234. doi: 10.1183/13993003.04234-2020
151. Viani-Walsh D, Kennedy-Williams S, Taylor D, Gaughran F, Lally J. Vitamin D deficiency in schizophrenia implications for COVID-19 infection. *Ir J Psychol Med.* (2020) 2020:1–10. doi: 10.1017/ipm.2020.107
152. Grant WB, Lahore H, McDonnell SL, Baggerly CA, French CB, Aliano JL, et al. Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients.* (2020) 12:988. doi: 10.3390/nu12040988
153. Fond G, Masson M, Richieri R, Korchia T, Etchecopar-Etchart D, Sunhary de Verville PL, et al. The Covid-19 infection: an opportunity to develop systematic vitamin D supplementation in psychiatry. *Encephale.* (2021) 2021:S0013-7006(21)00057-9. doi: 10.1016/j.encep.2021.02.002
154. Shah K, Saxena D, Mavalankar D. Vitamin D supplementation, COVID-19 and disease severity: a meta-analysis. *QJM.* (2021) 114:175–81. doi: 10.1093/qjmed/hcab009
155. Fond G, Pauly V, Leone M, Orleans V, Garosi A, Lancon C, et al. Mortality among inpatients with bipolar disorders and COVID-19: a propensity score matching analysis in a national French cohort study. *Psychol Med.* (2021) 2021:1–10. doi: 10.1017/S0033291721003676

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 De Hert, Mazereel, Stroobants, De Picker, Van Assche and Detraux. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.