



The use of vocal coordination in male African elephant group departures: evidence of active leadership and consensus

Caitlin E. O’Connell-Rodwell^{1,2,3,4,5}, Jodie L. Berezin^{3,4}, Alessio Pignatelli⁶ and Timothy C. Rodwell^{3,7}

¹ Center for Conservation Biology, Stanford University, Stanford, CA, United States of America

² Center for the Environment, Harvard University, Cambridge, MA, United States of America

³ Utopia Scientific, San Diego, CA, United States of America

⁴ Eaton-Peabody Laboratories, Massachusetts Eye and Ear, Harvard Medical School, Boston, MA, United States of America

⁵ Department of Otolaryngology, Head & Neck Surgery, Harvard Medical School, Boston, MA, United States of America

⁶ Department of Biostatistics, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD, United States of America

⁷ Department of Medicine, University of California, San Diego, School of Medicine, La Jolla, CA, United States of America

ABSTRACT

Group-living animals engage in coordinated vocalizations to depart from a location as a group, and often, to come to a consensus about the direction of movement. Here, we document for the first time, the use of coordinated vocalizations, the “let’s go” rumble, in wild male African elephant group departures from a waterhole. We recorded vocalizations and collected behavioral data as known individuals engaged in these vocal bouts during June-July field seasons in 2005, 2007, 2011, and 2017 at Mushara waterhole within Etosha National Park, Namibia. During departure events, we documented which individuals were involved in the calls, the signature structure of each individual’s calls, as well as the ordering of callers, the social status of the callers, and those who initiated departure. The “let’s go” rumble was previously described in tight-knit family groups to keep the family together during coordinated departures. Male elephants are described as living in loose social groups, making this finding particularly striking. We found that this vocal coordination occurs in groups of closely associated, highly bonded individuals and rarely occurs between looser associates. The three individuals most likely to initiate the “let’s go” rumble bouts were all highly socially integrated, and one of these individuals was also the most dominant overall. This finding suggests that more socially integrated individuals might be more likely to initiate, or lead, a close group of associates in the context of leaving the waterhole, just as a high-ranking female would do in a family group. The fact that many individuals were involved in the vocal bouts, and that departure periods could be shorter, longer, or the same amount of time as pre-departure periods, all suggest that there is consensus with regard to the act of leaving, even though the event was triggered by a lead individual.

Submitted 17 May 2024

Accepted 26 June 2024

Published 22 July 2024

Corresponding author

Caitlin E. O’Connell-Rodwell,

ceoconnell@stanford.edu,

caitlin_oconnell@meei.harvard.edu

Academic editor

Cédric Sueur

Additional Information and
Declarations can be found on
page 21

DOI 10.7717/peerj.17767

© Copyright

2024 O’Connell-Rodwell et al.

Distributed under

Creative Commons CC-BY-NC 4.0

OPEN ACCESS

Subjects Animal Behavior, Ecology, Zoology

Keywords Vocal coordination, Active leadership, Male African elephant, Socially integrated, Antiphony, Signature rumbles, Consensus, Dominance, *Loxodonta africana*

INTRODUCTION

Group-living animals rely on vocalizations to identify and communicate with individuals at a distance, assess reproductive status, facilitate social interactions, and coordinate movement (Bousquet, Sumpter & Manser, 2011; O'Connell-Rodwell et al., 2012; Poole et al., 1988; Stewart & Harcourt, 1994; Walker et al., 2017). Coordinating movement confers advantages, such as not getting separated from the rest of the group (Boinski & Campbell, 1995; Walker et al., 2017), ensuring group members have met their physiological needs (e.g., food and water) (Sueur et al., 2010), and conserving energy by moving in relative synchrony, minimizing localization effort if separated (Black, 1988; Boinski, 1991). Mountain gorillas and redfronted lemurs have pre-departure vocalizations called “grunts” (Sperber et al., 2017; Stewart & Harcourt, 1994) and white-faced capuchins make pre-departure “trills,” (Boinski & Campbell, 1995), that cause the entire group to get ready and then move from an area. In wild dog packs, the incidence of sneezing increases prior to departure, acting as a quorum to confirm the group is ready to depart (Walker et al., 2017).

Elephant vocalizations contain information about sex (Baotic & Stoeger, 2017), age and body size (Stoeger & Baotic, 2016), condition, and social and ovulation status (Poole et al., 1988; Soltis, Leong & Savage, 2005). Information encoded within calls makes it possible to identify individuals (McComb et al., 2003; Stoeger & Baotic, 2016; Wierucka, Henley & Mumby, 2021) as well as address one another with unique calls (Pardo et al., 2024). Elephant vocalizations are also used to coordinate action within family groups, often initiated by either the matriarch or another dominant female within the family (O'Connell-Rodwell et al., 2012; Poole, 2011; Poole et al., 1988).

In elephant family groups, matriarchs have been described as leaders (Lee & Moss, 2012) because they make decisions for their family and act as knowledge repositories based on their experiences (Mutinda, Poole & Moss, 2011). Matriarchs assess predator threats to determine when to act (McComb et al., 2011), and make foraging decisions and initiate movement (Mutinda, Poole & Moss, 2011), such as when to leave the waterhole (O'Connell-Rodwell et al., 2012).

While male elephants are not considered group living animals, many individuals appear to spend a lot of time in all-male groups (Chiyo et al., 2011; Evans & Harris, 2008; Goldenberg et al., 2014; Lee et al., 2011; Poole & Moss, 1989). However, little research has been conducted to assess the potential of male elephant coordination or active leadership. While male elephants have weaker associations within all-male groups than females do within their families (Archie, Moss & Alberts, 2006; Chiyo et al., 2011), their social lives are very complex. Male elephants establish dominance hierarchies within social networks (O'Connell-Rodwell et al., 2011) and gather in large groups where males of all ages prefer to associate with older, mature males (Evans & Harris, 2008). Preference for older males is likely attributed to older males taking on similar roles as matriarchs (i.e., mentor or

leadership role): older males aid in maintaining social cohesion (*Chiyo et al., 2011*), mediate aggressive behaviors (*Allen, Croft & Brent, 2021; Slotow et al., 2000*), and provide ecological information about resource location and effective navigation through the environment (*Allen et al., 2020*).

In general, individuals within bonded social groups coordinate their behavior and activities, which serves to maintain social stability through the use of physical interactions and vocalizations (*Seltmann et al., 2013*). While the evidence presented from photographs appears to support passive leadership, *i.e.*, younger individuals following older individuals (*Allen et al., 2020*), we propose that some highly associated individuals, and especially the highest-ranking male within an extended social network, may engage in active leadership tactics by initiating group departures vocally.

In this study, we document the use of “let’s go” rumble (LGR) vocalizations within bonded groups of male African elephants. We also show that these LGR events are mostly initiated by the most socially integrated individual. The initial LGR vocalization within a waterhole visit event triggers a series of highly synchronized and coordinated vocalizations within repeated bouts, a patterning that *Poole (2011)* refers to as cadenced rumbles, as the dynamic resembles, and likely is a form of conversation to reach consensus. We refer to bouts in this context as LGR cadenced rumble bouts, as they occur in bouts with often long periods of silence between them in the context of departure. This phenomenon was previously described only in the context of family groups preparing for departure, whereby a dominant female stops drinking, orients in the departure direction, and emits the LGR accompanied with slow ear flapping (*O’Connell-Rodwell et al., 2012; Poole, 2011; Poole et al., 1988*) and for the first time, we report that male elephants display exactly the same behavior. We discuss the value of having such a vocal tool to trigger action and coordinate movement of a group of associates, as well as highlighting the evidence for, and implications of, active leadership of highly socially integrated individuals within male elephant groups.

MATERIALS & METHODS

Field site and elephant identification

Data were collected during June–July field seasons in 2005, 2007, 2011, and 2017 at Mushara waterhole (hereafter referred to as Mushara) in Etosha National Park, Namibia. Mushara is located within a 0.22 km² clearing. Data were collected from an 8-meter-tall research tower, located approximately 80 m from the waterhole. The waterhole is fed by a permanent, artisanal spring, and is the only stable source of water within 10 km², making it an important resource during the dry season. For additional details about the field site, see recent publications (*Berezin et al., 2023; O’Connell-Rodwell et al., 2022a; O’Connell-Rodwell et al., 2022*). Namibian Ministry of Environment and Tourism, permit codes: #877/2005 for 1 February 2005 to 31 January 2006; #1141/2007 for 7 March 2007 to 28 February 2008; #1489/2011 for 12 June 2011 to 30 May 2012; #2188/2016 for 1 June 2016 to 30 June 2017.

Elephants have been individually identified at Mushara since 2004 using unique, recognizable morphological characteristics such as ear tear patterns, tail hair configurations,

tusk size and shape, and scarring. Elephants were assigned to age classes based on overall body size, shoulder height, hindfoot length, and skull and face morphometrics (Moss, 1996; O'Connell-Rodwell et al., 2022a; O'Connell-Rodwell et al., 2022). Age classes include: one-quarter (1Q), 10–14 years old; two-quarter (2Q), 15–24 years old; three-quarter (3Q), 25–34 years old; full, 35–49 years old; and elder, 50 years and older.

Keystone individual (the most socially integrated and dominant individual in a population) identification using social network and dominance hierarchy analyses was described recently; portions of this text were previously published as part of a preprint (O'Connell-Rodwell et al., 2024a; O'Connell-Rodwell et al., 2024b) and will be summarized in brief. For the social network analysis, we constructed association networks based on co-presence at the waterhole during field seasons. Weighted matrices of dyad-level association indices were built based on the Simple Ratio Index of association, ranging from 0–1, with higher indices representing individuals who are closely associated (Cairns & Schwager, 1987; Whitehead, 2008).

For the dominance hierarchy, we used dyad-level displacement (when an individual forces another to change his position; (O'Connell-Rodwell et al., 2011), to construct an ordinal hierarchy using the normalized David's Score (David, 1987; de Vries, Stevens & Vervaecke, 2006; Gammell et al., 2003). David's Score is calculated using the proportion of wins or losses across all dyads an individual is present in, while also considering the total number of dominance interactions observed. The highest values are associated with those who most consistently win contests. One individual (#22) had the highest average eigenvector centrality (most socially-integrated) and the highest dominance rank of all individuals included in the analysis across five years (2007 to 2011).

Data acquisition

We recorded LGR vocalization events in the context of male elephants leaving Mushara waterhole. For each LGR event, we quantified the temporal spacing of the event, the onset of the departure period, the characteristics and individuality of LGR rumbles, the level of association between individuals that engaged in the bouts, and the behavior patterns within events, as well as bout initiation and serial participation of known individuals within the bouts.

Behavioral data and vocalization recordings were collected opportunistically during the evening and night (approximately 5:00 p.m. to 2:00 a.m.) when ambient sound and wind shear was low enough to record extremely low-frequency male vocalizations made in the range of 11 Hz. After dark, light-enhancing technology was attached to a standard HD video recorder and 3x magnification was used to visually identify individuals and document their behavior. In the new moon period, an infrared spotlight was also attached to the recorder to enhance visibility of tusks, ear tears and tail hair for individual identification.

Vocalizations were recorded using a Neumann Km131 microphone (Berlin, Germany) at a sampling rate of 48 kHz and placed 20 m from the waterhole, powered remotely *via* a 12-volt battery in the field tower. Vocalization data collected in 2005–2011 was recorded using a TEAC DAT digital recorder, and in 2017, a Sound Devices solid-state digital recorder (Reedsburg, WI, USA) was used. All vocalizations recorded were logged by date,

time, type, and social context, including all individuals involved, the locations of callers, and those participating in the vocal bouts when known. Calls were flagged when it wasn't possible to tell who the caller was, due to an obstruction (another elephant, the tower, or too far away to distinguish which individual was ear-flapping), or overlap with another caller, and were labeled as unknown.

Events were described as a period when a group of male elephants entered the clearing (from the forest) to the time when they departed the clearing. The criteria used to select events was as follows: (1) audio recordings were captured for the full event (from arrival to departure), (2) males arrived and departed together, and (3) females were not present during any time of the event, nor any other behaviorally impactful disturbances. Events were divided into pre-departure and departure periods following protocols described in [O'Connell-Rodwell et al. \(2012\)](#): pre-departure began when the elephants entered the clearing and was defined by greetings between males and drinking water, and ended when the departure period began. Departure began when a known male initiated the behavior associated with the LGR (and could be heard in almost all cases, due to the proximity of the microphone to the caller at the waterhole, as well as low-frequency sounds being more easily detectible after dark, given the low wind shear and quiet background) and ended when all elephants left the clearing. The microphone was monitored remotely using headphones plugged into the recorder in the tower.

Behaviorally, LGRs were identified when a known male stepped away from the waterhole, stood still and rumbled, most often while flapping his ears, and positioned facing away from the waterhole ([O'Connell-Rodwell et al., 2012](#); [Poole et al., 1988](#)). This first rumble marked the onset of the departure period.

After the initial rumble was emitted, the individual repeated the vocalization, while remaining stationary, or while walking away from the waterhole. This initial LGR call, or sequence of repeated single LGR calls sometimes over the course of several minutes, then triggered a bout of coordinated responses from the rest of the bonded group, a pattern that [Poole \(2011\)](#) refers to as cadenced rumbles. Each caller within the coordinated interactive bout was noted by ear-flapping behavior, while standing stationary or walking out to follow the initiator. If there was no ear-flapping, the males were spaced far enough apart to tell where the call was coming from. If the males were close together and there was no ear flapping, both males were noted and the call was ascribed to the two possible callers. These bouts were recorded until the group hit the edge of the clearing.

Acoustic analysis

Rumbles were analyzed using Raven Pro 1.6 (Cornell Lab of Ornithology, Ithaca, NY, USA) with a Hann window size of 65,536, a hop size of 32,768, with 50% overlap. The window size is larger than previous publications ([Stoeger & Baotic, 2016](#); [Wierucka, Henley & Mumby, 2021](#)) to precisely identify the fundamental frequency and harmonics. However, this extremely precise frequency resolution comes at the cost of a lower time resolution. "Let's go" cadenced rumble bouts have slightly overlapping rumbles and any calls made within two seconds were considered within a single bout. We chose this two-second window because there were no other vocalizations that occurred outside the cadenced

rumble bouts, and since the bouts might occur a minute or even five or so minutes apart, it seemed appropriate to include a vocalization that occurred within two seconds as being part of the bout. This differed from the 1.5 s window that was used for the female bouts. Within family groups, there are so many vocalizations and individuals involved in the LGR cadenced rumble bouts, that the response time is quicker, and many more bouts within the departure period to consider as unique, whereas for the males, there are long periods of silence, and as such, it seemed appropriate to extend the window to two seconds. For non-overlapping rumbles, the full rumble was selected. For rumbles that did overlap, only the non-overlapping section is selected. For this study, only slightly overlapping bouts were considered as part of the LGR bout, or departure conversation. Individual rumbles outside of the 2 s window were assumed to not be part of the LGR bout sequence. For bouts with more than three rumbles, only the first three rumbles of each bout were considered in the acoustic analysis.

A combination of parameters were used to identify individuals: (1) field notes, detailing the behavioral observation noting the time of “let’s go” rumble behaviors and the corresponding times on the audio recorder; noting the callers by ear flapping and or location, if they were far enough away from any others to designate a caller, and (2) the rule of non-consecutive rumble criteria (*O’Connell-Rodwell et al., 2012*), where it is assumed that overlapping rumbles cannot be produced by the same individual (but could be caller #1 and #3).

Where it was difficult to behaviorally discern between two individuals, principal components analysis (PCA) visualizations of rumble characteristics were used to identify unique individuals. The same parameters used in the PERMANOVA (described below) were used in the PCA. Since we always knew which elephants were actually present at the waterhole during each event (which was never more than six elephants vocalizing, [Table 1](#)), we could plot all known individuals as well as the unknowns ([Fig. 1](#)). The PCA grouped the known individuals clearly, and even if two individuals were close together on the PCA axes we could still distinguish them. For example, #61/#132 means that #61 and #132 were both present at the waterhole along with the known callers (event 7). We were able to visualize the unknown calls (in addition to those of other elephants) into two distinct groups of calls: one for #61 and one for #132 ([Fig. 1](#)). However, we do not have specific notes on when #61 and/or #132 is vocalizing, therefore we cannot assign one group of calls to #61 and the other to #132. Hence, the #61/#132 (1) and #61/#132 (2). The same was true for the pair #105/#69 (1) and #105/#69 (2) (event 2, [Fig. 1](#)).

Following the methodology of *Wierucka, Henley & Mumby (2021)*, we measured five key acoustic parameters: frequency 5% (frequency that divides the rumble into two frequency intervals containing 5% and 95% of the energy), Frequency 95% (frequency that divides the rumble into two frequency intervals containing 95% and 5% of the energy), Bandwidth 90% (the difference between the 5% and 95% frequencies), Center frequency (divides the rumble into the two frequency intervals of equal energy), and Duration 90% (the differences between the 5% and 95% times) (abbreviated definitions reproduced from *Charif, Waack & Strickman (2010)* and *Wierucka, Henley & Mumby (2021)*). We also chose to measure the fundamental frequency (*Stoeger & Baotic, 2016*); both the fundamental

Table 1 Description of “let’s go” rumble (LGR) events used in this study. Year, event number, number of bouts, number of rumbles per event, the duration of the event, number of males engaged in LGRs versus the total number within the group, as well as who initiated and the social context (presence of keystone male). Events with missing information refer to the events included for associations analyses of individuals involved in vocal coordination but were not used in acoustic analysis (see ‘Methods’ for more details).

LGR event	Year	# Bouts	# Rumbles Recorded (analyzed)	Total Event time (minutes)	# Elephants vocalizing	# Elephants present	Initiator of event	Keystone male present?
1	2005	8	22(20)	55.0	4	4	#22	Yes
2	2007	10	26(24)	98.2	6	6	#22	Yes
3	2007	1	3(3)	50.0	3	5	#67	No
4	2007	3	9(8)	37.3	3	4	#22	Yes
5	2011	5	17(13)	31.9	3	3	#84	No
6	2011	11	29(26)	57.0	5	8	#46	No
7	2017	10	32(28)	56.0	4	5	#46	No
8	2007	–	–	–	–	4	–	No
9	2007	–	–	–	–	4	–	Yes
10	2007	–	–	–	–	6	–	Yes
11	2007	–	–	–	–	5	–	No
12	2007	–	–	–	–	7	–	No

frequency and duration 90% were measured using a rectangular selection box around the entire call.

Statistical analysis

To evaluate whether the initial “let’s go” rumble triggered departure, we used a paired Wilcoxon Signed Rank test to assess whether the pre-departure time was significantly longer than the departure time, using the function `wilcox.test` in the R “stats” package (*R Core Team, 2023*). Similarly, a Wilcoxon Signed Rank test was also used to evaluate whether the number of rumbles significantly increased in the departure period (compared to the pre-departure period). Since longer events would be expected to have more rumbles, we calculated the rate of rumbles as the number of rumbles per minute in each period.

Next, we wanted to assess whether each LGR emitted contained a unique signature distinctive to each known individual, reproducing the methodology of *Wierucka, Henley & Mumby (2021)*. Acoustic parameter data was normalized on a scale of 0 to 1 due to the different variable types, mean values of each variable, and disparate standard deviations. We used a permutational multivariate analysis of variance (PERMANOVA) test using the `adonis` function in the “vegan” package (*Oksanen et al., 2022*) with a Euclidean distance matrix of the frequency parameters. To test the assumption of homogeneity of the variance-covariance matrix, we first used the `betadisper` function in the “vegan” package to calculate the average distance of an individual’s calls to their calculated centroid and then used an ANOVA test to assess whether the mean distance to centroid differed significantly among individuals (suggesting that individuals had large variances in their acoustic parameters). We then performed a pair-wise comparison of the mean distances among individuals




Figure 1 Principal components analysis (PCA) visualizations of the seven events and 19 unique individuals. To facilitate visual comparisons across events, the PCA analysis was performed using data from all events so that each event could be visualized using the same axes (PC1 and PC2). See associated code and methods section for additional details.

Full-size DOI: [10.7717/peerj.17767/fig-1](https://doi.org/10.7717/peerj.17767/fig-1)

using a Wilk's test and false discovery rate method for p -value adjustment using the `pairwise.perm.manova` function in the "RVAideMemoire" package (Herve, 2023).

Lastly, we assessed whether the males involved in "let's go" events had significantly higher associations than those not involved in the "let's go" events. Only "let's go" events and association data from the 2007 field season were used, due to the large number of dyads observed, with data available for all individuals included in "let's go" events. To increase the sample size of dyadic-relationships within "let's go" events, we included five additional groups of individuals that were observed and acoustically recorded in a "let's go" rumble event (Table 1). These events could not be included in acoustic analysis, due to the lack of clear arrival times and audio recording of the entire event. Of the 25 individuals that came to the waterhole at least three times in 2007, there were a total of 223 unique dyads observed of the 300 possible combinations. Of these unique dyads, 64 dyads involved 17 individuals who were both taking part in LGR events, while 159 of these dyads involved at least one individual who was not involved in LGR events. We used a Mann–Whitney U -test to assess for significant differences between the two groups of individuals (those observed in LGR events and those who are not), using the `wilcox.test` function in the "stats" package.

All statistical analyses and visualizations were performed using R statistical software (version 4.3.1) (R Core Team, 2023), with significance set at an alpha level of $\alpha = 0.05$.

RESULTS

LGR events and temporal spacing

The final acoustic analysis included data from 7 LGR events, with a total of 48 bouts and 122 analyzed rumbles (Table 1). A total of 19 individuals were recorded across the 7 LGR events (Table 2), with a mean group size was 4.9 individuals, with a range of 3 to 8 individuals (with only 7 individuals present across LGR events who did not vocalize). 16 individuals involved in the LGR events were in the 3Q, full, or elder age classes (25+ years old), with only 3 individuals in the 1Q (10–14 years old) and 2Q (15–24 years old) age classes (Table 2).

LGR events were defined by the pre-departure period, which was the arrival of a group of male elephants at the waterhole where they drank and socialized, followed by the departure period which was initiated by the onset of a "let's go" rumble. Three rumble types were observed during these events, namely the first single call by the initiator (Fig. 2A) which triggered the highly synchronized and coordinated bouts that contained slightly overlapping rumbles emitted within bouts (Fig. 2C), by some or all of the individuals within the group at the waterhole. Sometimes, the initial vocalization was followed by an overlapping, "duet" call by the initiator and a close associate (Fig. 2B). A spectrogram of an excerpt from event 2 depicts vocalizations in real time (Fig. 3). Sometimes, the initiator emitted a call, but did not get an immediate response, and proceeded to call several more times and even started walking away from the waterhole, before others responded (Fig. 3). In the example depicted in Fig. 3, of a subset of rumbles, the keystone male, #22, emitted two LGR before triggering several bouts of rumbles that he almost always led (Fig. 4). The repeated vocal bouts resulted in the act of leaving the waterhole, most often as a group,

Table 2 Summary data for individual elephants ($n = 19$). Individually-known elephants and their corresponding age class, including the years that rumble vocalizations were recorded, along with a description of the number of rumbles recorded and rumble characteristics. For individuals who changed age classes over the course of the study, an age class is provided for each observation year.

Individual	Years recorded	Age class	Rumble count	Mean frequency 5% (Hz)	Mean fundamental frequency (Hz)	Mean center frequency (Hz)	Mean Frequency 95% (Hz)	Mean Bandwidth 90 (Hz)	Mean Duration (s)
#18	2005, 2007, 2017	Full	15	11.04	13.29	16.89	70.95	59.91	4.57
#22	2005, 2007	Full	23	12.93	13.99	28.80	78.02	65.09	4.04
#25	2007	Full	6	14.16	14.29	31.25	81.01	66.85	4.22
#46	2007, 2011, 2017	3Q, Full, Full	23	10.89	12.99	19.52	82.22	71.33	4.55
#48	2007	Elder	2	8.06	13.78	28.56	106.20	98.15	2.49
#61/#132 (1)	2017	Full/Elder	8	9.52	13.07	13.73	45.04	35.52	3.94
#61/#132 (2)	2017	Full/Elder	1	13.18	13.28	67.38	96.68	83.50	4.16
#65	2007, 2011	2Q, 3Q	7	11.72	13.09	20.30	64.45	52.73	4.25
#67	2007	3Q	1	7.32	21.75	24.90	146.48	139.16	2.53
#76	2011	Full	7	13.18	13.64	25.78	77.01	63.83	5.96
#83	2011	3Q	4	10.25	12.17	19.41	76.54	66.28	2.44
#84	2011	Full	5	11.43	12.49	24.90	87.31	75.88	4.32
#105/#69 (1)	2007	Full/3Q	1	16.11	16.50	74.71	96.68	80.57	3.24
#105/#69 (2)	2007	Full/3Q	6	9.52	13.87	23.68	87.40	77.88	3.01
#118	2011	Full	1	17.58	16.56	36.62	92.29	74.71	5.58
#119	2011	Full	5	11.13	13.63	20.22	85.55	74.41	3.89
#146	2011	1Q	4	10.99	12.71	24.17	82.76	71.78	3.01
Unk1	2005	Unknown	2	10.99	11.48	17.58	24.90	13.92	2.94
Unk2	2005	Unknown	1	10.25	13.97	14.65	89.36	79.10	4.13

though sometimes there were stragglers that returned to the water for one more drink before following the rest of the group out of the clearing.

The duration of the pre-departure period was longer than the departure period for four of the seven LGR events (Fig. 5). The median pre-departure time (30.0 ± 9.68 min, range = 15.67, 42.50) was longer than the departure time (21.67 ± 16.5 min, range = 4.91, 55.97) but was not significant (Wilcoxon Signed Rank test $p = 0.469$, effect size $r = 0.32$). Event 2 was unique in that there was an initial bout, then 43 min passed before a series of 9 bouts occurred in quick succession. During the 43 min between the first bout and the series, 14 individual rumbles were vocalized by the keystone individual (#22). When tested without event 2, the median times were still not significantly different (Wilcoxon Signed Rank test $p = 0.313$, effect size $r = 0.47$).

By construction in this study, but for comparison with a previous study (O'Connell-Rodwell et al., 2012), the rate of rumbles in the departure period was significantly higher than the pre-departure period (Wilcoxon Signed Rank test $p = 0.016$, effect size $r = 0.89$). For all events, pre-departure periods were silent, with no vocalizations recorded. The




Figure 2 Three spectrograms depicting three different male elephant coordinated departure rumble vocalizations. (A) The “let’s go” rumble (LGR) is a single call from the initiator that triggers the highly synchronized and coordinated interactive LGR bouts, or cadence calls, in response to the initial LGR. This trigger was initiated by the most dominant or most socially integrated individuals. (B) The LGR duet between the initiator and a close associate sometimes followed the single trigger vocalization. (C) LGR bouts, or cadence calls, are emitted by the rest of the group of close associates in response to the LGR trigger. Duration of calls and mean frequency were measured on the fundamental frequency. The nonoverlapping portions of the cadence calls were used to analyze individual identity and confirmed by field notes.

Full-size DOI: [10.7717/peerj.17767/fig-2](https://doi.org/10.7717/peerj.17767/fig-2)

median (\pm SD) rate of rumbles per minute in the departure period was 0.84 ± 1.12 (range = 0.26, 3.46; mean = 1.25).

Across all events, the mean (\pm SD) number of bouts per departure period was 6.86 ± 3.89 with a range of 1 to 11. The mean (\pm SD) number of rumbles was 19.71 ± 10.67 with a range of 3 to 32, while the mean (\pm SD) number of rumbles per bout was 2.88 ± 0.96 with a range of 2 to 6. The mean (\pm SD) duration of bouts was 10.54 ± 3.81 s with a range of 3.77 and 19.51 s. The average time between bouts was 156.55 ± 405.40 s (2.61 ± 6.76 min) with a range of 2.80 and 3,624.23 s (0.047 to 60.40 min).

Rumble characteristics and individual differences

The mean (\pm SD) duration of rumbles was 4.15 ± 1.42 s and the mean Frequency 5% was 11.53 Hz ± 2.31 . Additional rumble characteristics are presented in Table 3. We




Figure 3 This spectrogram depicts a portion of a male elephant “let’s go” rumble (LGR) event within the departure period. The initiator emits a single LGR call, and the call sometimes gets repeated several times (in this event, the last two single calls in a series of repeated single calls are depicted, prior to triggering the LGR bout, or cadence responses from the rest of the group). After emitting a LGR upon leaving the waterhole, the initiator did not get a response (at 1:24:30), the lower intensity dB level depicting that the initiator is heading away from the group and repeats the call again about a minute later (1:26:10). When he does not get a response the second time, he returns to the group at the waterhole and emits another LGR, this time triggering a series of LGR bouts emitted by the rest of the initiator’s close associates in response. The vocal bouts result in the act of leaving the waterhole, most often as a group, though sometimes there are stragglers that return to the water for one more drink before following the rest of the group out of the clearing.

Full-size DOI: 10.7717/peerj.17767/fig-3

Bout	First caller	Second caller	Third caller
1	#22	#65	-
2	#22	#25	#105/#69 (1)
3	#22	#25	#22
4	#48	#105/#69 (2)	-
5	#22	#25	#65
6	#48	#105/#69 (2)	-
7	#105/#69 (2)	#22	-
8	#22	#105/#69 (2)	-
9	#65	#22	#105/#69 (2)
10	#22	#105/#69 (2)	-

Figure 4 Example of the order of callers (rumbles) in each cadenced rumble bout for event 2. The key-stone male (#22) initiates most of the bouts in this event, while the other initiators were all mid-ranking.

Full-size DOI: 10.7717/peerj.17767/fig-4

found significant individual differences in the five acoustic parameters among the 19 individuals included in the study (PERMANOVA $R^2 = 0.522$, $p = 0.001$). The assumption of homogeneity of variance was not violated, ($F = 1.34$, $DF = 18$, $p = 0.182$), suggesting that individuals have similar variation and co-variation across their rumble characteristics.




Figure 5 Total time (in minutes) the pre-departure and departure periods for the seven LGR events. Dashed lines connecting two solid black circles represent an event's corresponding pre-departure and departure time, to visualize the relationship between the amount of time groups were present in each period. Thick horizontal bars represent the median, while the white diamond shape within each plot represents the mean. The vertical length of the box represents the interquartile range and the vertical lines are the minimum and maximum values. The median pre-departure time was longer than departure time, but the periods did not differ significantly (Wilcoxon Signed Rank test $p = 0.469$). For four events, the pre-departure time was longer than the departure time, while the other three events had longer departure periods.

Full-size DOI: [10.7717/peerj.17767/fig-5](https://doi.org/10.7717/peerj.17767/fig-5)

Pairwise comparisons showed that many of the individuals significantly differed in their acoustic characteristics (Table 4).

Associations, dominance, and the keystone individual

Of all the frequent visitors to Mushara in 2007, individuals within LGR groups had a mix of association levels amongst its members, where some individuals had high association strengths, and others had low. Dyads involving two individuals within LGR events (highlighted in yellow; Fig. 6) had significantly higher association indices than dyads in which at least one individual was not involved in LGR events (highlighted in blue; Fig. 6) (Mann–Whitney U test $p = 0.0001$, median difference = 0.05, effect size = 0.26). The median (\pm SD) index for those involved in an LGR event was 0.16 ± 0.17 (mean = 0.21, range = 0.04 to 0.92), while the median for those not observed in an LGR group was 0.11 ± 0.07 (mean = 0.12, range = 0.04 to 0.36).

Table 3 Rumble parameter characteristics.

Parameter	Mean	Median	SD	Min	Max
Duration (s)	4.15	4.13	1.42	1.51	8.89
Bandwidth (Hz)	65.49	67.38	17.74	11.72	139.16
Center Frequency (Hz)	23.26	24.17	9.01	11.72	74.71
Frequency 5% (Hz)	11.53	11.72	2.31	5.86	17.58
Frequency 95% (Hz)	77.02	79.10	17.65	21.97	146.48
Fundamental Frequency (Hz)	13.47	13.05	1.84	10.19	22.92

Notes.

SD, standard deviation; Min, minimum frequency; Max, maximum frequency.

For the three events where he was present, the keystone male (#22) initiated the departure of the group by emitting the first LGR and was also the first caller in LGR cadenced rumble bouts 61.9% (13/21) of the time. When the keystone male was present, six (of nine) other individuals in his groups initiated cadenced rumble bouts, but only 1 or 2 times each, making the keystone male 1.6 times more likely to initiate these bouts than any other individual within groups where he was present. Across all events (when #22 was present and when he was not), 12 of the 19 individuals initiated cadenced rumble bouts. When the keystone male was not present, one individual (#46) initiated 54.5% (12/22) of the bouts in the three events he was present in. All other individuals initiated bouts five times or fewer (for example, see Fig. 4).

Across the seven events, four males (#22, #46, #67, and #84) initiated the departure period by emitting an LGR. Males #22, #46, and #67 had high centrality rankings of 1, 6, and 8, respectively, out of 25 individuals evaluated (data was not available for male #84, the departure initiator of event 5). Of these three individuals, only male #22 was the highest ranked in the dominance hierarchy overall, while males #46 and #67 were mid-ranking overall and not the highest ranked members in their respective LGR groups (Fig. 7).

DISCUSSION

Since male elephants have been described as living in loose groups of associates (Archie, Moss & Alberts, 2011; Chiyo et al., 2011), it is surprising to document them engaging in highly coordinated vocal behavior, used to departure from the waterhole as a group of associates, just as group-living animals do. And even more surprising, is that they do so with vocal patterning and synchrony (Figs. 2 and 3) previously only described in females living within family groups (O'Connell-Rodwell et al., 2012; Poole et al., 1988) as part of a departure conversation, or cadenced rumbling (Poole, 2011). To add to these surprising findings is the fact that this vocal coordination during departure only occurs within male groups that have strong associations and are much rarer between loose associates (Fig. 6).

This solicitous behavior suggests much deeper relationships than random meet ups at a waterhole while drinking, whereby individuals might engage in social interactions with bonded associates, and from there, perhaps passively follow a dominant or socially integrated individual upon departure. Similar vocal coordination among associates was also

Table 4 Pairwise comparisons of an individual's mean distance to their centroid. Adjusted p -values are shown, with significant results in bold. Numbers represent individual elephants. Where there is a dashed line, both individuals had only one rumble analyzed; with only one rumble, a mean distance to the centroid could not be calculated as the centroid was represented by the single rumble.

	18	48	76	65	84	105/69 (1)	105/69 (2)	22	119	46	67	118	146	83	25	61/132 (1)	61/132 (2)	Unk1	
48	0.045																		
76	0.012	0.093																	
65	0.358	0.086	0.024																
84	0.049	0.137	0.039	0.012															
105/69 (1)	0.137	0.368	0.213	0.213	0.227														
105/69 (2)	0.029	0.167	0.012	0.090	0.221	0.220													
22	0.012	0.067	0.042	0.092	0.252	0.103	0.042												
119	0.138	0.178	0.037	0.091	0.557	0.227	0.352	0.167											
46	0.214	0.074	0.032	0.103	0.586	0.095	0.109	0.012	0.727										
67	0.153	0.368	0.220	0.221	0.227	–	0.220	0.111	0.227	0.126									
118	0.116	0.368	0.329	0.213	0.227	–	0.221	0.234	0.227	0.116	–								
146	0.064	0.138	0.032	0.085	0.227	0.252	0.585	0.209	0.576	0.221	0.252	0.252	0.460						
83	0.039	0.138	0.024	0.057	0.116	0.252	0.440	0.093	0.182	0.103	0.252	0.252	0.460						
25	0.012	0.111	0.116	0.024	0.103	0.221	0.068	0.536	0.088	0.024	0.227	0.344	0.136	0.093					
61/132 (1)	0.012	0.096	0.012	0.012	0.020	0.209	0.012	0.012	0.012	0.012	0.209	0.183	0.012	0.029	0.020				
61/132 (2)	0.153	0.368	0.220	0.220	0.227	–	0.221	0.149	0.227	0.116	–	–	0.252	0.252	0.220	0.215			
Unk1	0.068	0.368	0.091	0.091	0.116	0.368	0.103	0.039	0.116	0.032	0.368	0.368	0.138	0.138	0.103	0.133	0.368		
Unk2	0.399	0.368	0.421	0.307	0.368	–	1.000	0.344	1.000	0.931	–	–	0.615	0.432	0.227	0.205	–	0.368	




Figure 6 Association indices for individuals involved in LGR events (yellow) and those who are not (blue). (A) Social network for 2007 for $n = 25$ individuals. The lines between individuals represent the strength of association, binned into four levels, ranging from 0 to 1. Circles represent individuals and the size of the circle represents their centrality within the social network, binned into four levels. Three of the four LGR event initiators are marked with asterisks next to their ID numbers. (B) Violin plot of the association indices for dyads involving at least one individual not observed in an LGR group ($n = 159$ dyads) and for dyads in which both individuals were observed in LGR an group (but not necessarily together) ($n = 64$ dyads). The shape of the violin plot is a visual representation of the distribution of the data. The box plots provide further details about the data distribution where the thick black line represents the median and the black diamond represents the mean. The length of the box is the interquartile range, with the vertical lines representing the minimum and maximum values, and filled circles represent outliers.

Full-size [DOI: 10.7717/peerj.17767/fig-6](https://doi.org/10.7717/peerj.17767/fig-6)

found in bonobos, whereby more bonded individuals were more effective at coordinating group action (Levrero *et al.*, 2019), and adult male Barbary macaques most frequently recruited those with whom they had affiliative relationships (Seltmann *et al.*, 2013). Although the level of dyadic associations varied in some male elephant groups—some individuals having low associations—each individual had a stronger association with at least one other individual in the group. This lends further evidence to the idea that these vocal bouts, or conversations, expedited departure amongst bonded individuals.

The most intriguing aspect of these findings is that three of the departure initiators (males #22, #46, and #67) were highly socially integrated (central) within the association network (Fig. 6), and only one of those individuals was also very high ranking overall (male #22; Fig. 7), all three being nearly fully mature (>25 years old; #67) or fully mature adults (>35 years old; #22 and #46) (O’Connell-Rodwell *et al.*, 2022a; O’Connell-Rodwell *et al.*, 2022). Species social structure is thought to impact the coordination of movement (Seltmann *et al.*, 2013), but results have been inconclusive as to who has the most social




Figure 7 Dominance hierarchy for 2007, highlighting the three departure initiators for which data was available (in red). A total of 25 non-musth individuals were included in the analysis. Male #22 (the key-stone individual) was ranked 1, while males #67 and #46 were ranked 13 and 14, respectively.

Full-size DOI: [10.7717/peerj.17767/fig-7](https://doi.org/10.7717/peerj.17767/fig-7)

influence (*Petit & Bon, 2010*). For example, social integration and maturity were important for coordinated movement in cattle (*Šárová et al., 2013; Sueur et al., 2018*). Being an adult, high-ranking male was important for Barbary macaques (*Seltnann et al., 2013*). And, lastly, dominance rank was the most important for successive rallying and departure for African wild dogs (*Walker et al., 2017*).

Highly socially integrated individuals were the departure initiators. This data suggests that network centrality is critical with regard to taking initiative to coordinate the group. Since two of the four initiators were mid-ranking and one was the highest ranking, the results of this small dataset suggests the possibility that dominance might not be as important as centrality with regard to leadership within groups of male elephants. In a follow up study, we plan to compare the importance of dominance status *versus* social

integration as they relate to leadership. Socially integrated individuals are thought to act as sources of social information (King & Sueur, 2011), due to the quantity of connections within their network. Central individuals might also have greater access to information (Palacios-Romo, Castellanos & Ramos-Fernandez, 2019), making them more attractive as companions than less socially integrated individuals. For example, in male elephants, we constructed the dominance hierarchy based on displacements at the waterhole, thus, being a dominant male often does not necessarily convey to others that an individual has knowledge about the social or physical environment.

Socially integrated individuals were the most likely to initiate the departure period, but several other individuals initiated cadenced rumble bouts within the events (Table 1, Fig. 4). Additionally, a majority of the individuals in the group participated in the bouts (Table 1), suggesting that the final decision of when to depart is shared in a consensus (Sueur & Petit, 2008). Collective decision-making is thought to be more accurate than a decision made with a lack of consensus, since it's based on the knowledge of many individuals (Conradt & Roper, 2005). For our male groups, the individuals who participated in the vocal bouts were all at least 25 years old (3Q age class; with the exception of individual #65; Table 2), all of whom would have decades of shared knowledge. Further, even the individuals who did not participate in the vocalizations (many of whom were mature adults) are considered to be part of the decision-making process just by following and "agreeing" non-vocally to the decision being made by the other individuals in the group (Conradt & Roper, 2005).

The pre-departure and departure periods did not significantly differ in duration, but three of the seven events had longer departure times than pre-departure (Fig. 5). In contrast to family groups where the matriarch has the most knowledge of the environment (McComb et al., 2001; McComb et al., 2011; Mutinda, Poole & Moss, 2011), the adult male elephants in our LGR groups likely all have similar repositories of environmental knowledge and are independent adults. As such, the initiators of the departure likely have less "control" than a matriarch might have over her family group and might require the males to have longer periods of decision-making, contributing to our observed longer departure periods. Additional variables, such as group size, rumble rate, or level of bondedness, might also impact departure duration and warrant further investigation.

In accordance with our definition, we found a significant increase in the rate of rumbles made within LGR cadenced rumble bouts in the departure *versus* the pre-departure period, where all events had zero rumbles in the pre-departure period. These results contrast with previous findings in female elephants where there were considerably more vocalizations in general made in the pre-departure period (prior to the onset of the first LGR) (O'Connell-Rodwell et al., 2012) than we observed in the male groups. Male elephants are described as being less vocal overall than females (reviewed in Morris-Drake & Mumby (2017)), which likely explains why there were no vocalizations in the pre-departure period. Additionally, between males and females, no matter how many are in the group, there always tended to be three callers on average per LGR cadenced rumble bout in response to the LGR. This suggests that male and female groups may have similar organizational principles of leadership and consensus.

These results offer the first evidence of active leadership in male African elephants, whereby socially integrated and/or dominant individuals, actively determine the departure time for the group, just as matriarchs or high-ranking females do. A leader, or active leader, is defined as one who solicits those to follow them and exerts social influence over a group by means of their dominance rank, social position, experience, or a specific behavior (King, Johnson & Van Vugt, 2009; Pyritz et al., 2011). In contrast, passive leadership occurs when an individual might be unintentionally leading (King, Johnson & Van Vugt, 2009; Pyritz et al., 2011), such as what was previously described in male elephants where younger individuals followed mature males (Allen et al., 2020).

This coordination among males within highly associated groups begs the question of what advantage individuals might have in maintaining a group's integrity over time and space. Maintaining bonds within groups strengthens group cohesion (de Waal, 1986), which for social males, could facilitate coalition behavior, thus providing a competitive advantage over resources, such as scarce waterpoints in an arid environment. This competitive edge over adversaries might outweigh having to share resources with associates (Conradt & Roper, 2000) and also reduces competition over scarce waterpoints (O'Connell-Rodwell et al., 2011). Additionally, this behavior might benefit genetically related individuals involved in coordinated vocal departures, whereby shared social and environmental knowledge could serve to enhance reproductive benefits. Future relatedness studies on associates may shed light on this possibility and would reflect previous findings on females being able to discern paternity (Archie et al., 2007). Lastly, the repetition and synchronized nature of only overlapping right at the end of a call creates a much longer call that could be heard at a greater distance, which might alert conspecifics to their departure.

We found some significant differences in rumble characteristics amongst individuals, supporting previous findings using similar methodologies (Stoeger & Baotic, 2016; Wierucka, Henley & Mumby, 2021). Our frequency 5% was extremely similar to Wierucka, Henley & Mumby (2021) and also fit within the range of the fundamental frequency previously reported (Baotic & Stoeger, 2017; Poole et al., 1988; Stoeger & Baotic, 2016). Further, our center frequency, duration, bandwidth, and frequency 95% fall within the range of those of Wierucka, Henley & Mumby (2021) (Table 3). These quantifiable differences in call structure between at least some individuals (Table 4) are likely distinguishable by others within the cohort and could be used to keep track of who is calling at what distances to facilitate coordination while leaving the area.

LGRs have sex-based differences, where the male rumbles tend to be relatively monotonic, like the females, but often with less frequency modulation (Fig. 2A) than female LGR calls measured at the same field site (Fig. 1 in O'Connell-Rodwell et al. (2012)). This may be due to the fact that the females can become very insistent within a dispute about a particular departure direction, thus modulation increases (O'Connell-Rodwell et al., 2012). When individuals do not respond to an LGR, the frequency modulation of the call tends to increase, often with an increase in dB as well, which is true for both males and females. In addition, the mean duration of the male LGR was four seconds (± 1.4), one second longer than the average female LGR at this field site (O'Connell-Rodwell et al., 2012). Both LGR and rumbles within a bout of cadenced rumbling were considerably

longer (median 5.2 and 5.1 s respectively) in Amboseli (Poole, 2011) than Etosha male or female rumbles in LGR cadenced rumble bouts. The mean fundamental frequency for the males was 13.6 Hz (\pm 1.6 Hz), which is similar to the findings of Baotic & Stoeger (2017) where the females were slightly higher in frequency by 2–6 Hz.

The harmonic structure differs in the male LGR and cadenced rumble bouts from those found in females at this site, as well as sites in Amboseli, in that the dB level is consistent between the fundamental frequency (F0) and first two harmonics, and only slightly lower at the third and fourth harmonics, and then markedly lower only starting at the fifth harmonic. In the female LGR and cadenced rumbles, F0 and the first harmonic are consistent but the second and third harmonics are markedly weaker, with the fourth, fifth, and sixth harmonics being higher in amplitude.

Finally, female LGR structure differs in Etosha as compared with Amboseli, where F0 and the first and second harmonics are dominant, the third and fourth harmonics have markedly lower amplitude, and the fifth, sixth, and seventh harmonics have lower but visible amplitude for Amboseli females (Poole, 2011), versus Etosha females, where F0 and the first harmonics are dominant, the second and third harmonics almost absent, with the fourth, fifth, and sixth harmonics present but weaker than F0 and the first harmonic (O'Connell-Rodwell et al., 2012). The dB patterning is also different between the females at both study sites and would be interesting to compare in a future analysis for the possibility of a dialect between the two populations.

CONCLUSIONS

This study reports the first evidence of the use of vocal coordination in the departures of closely associated, male African elephants. We also provide the first evidence of active leadership in male elephants, whereby socially integrated individuals begin the departure period by actively recruiting their associate's company during departure, using a "let's go" rumble vocalization. Most of the other group members participate in the decision making process concerning the time of the departure, similar to the negotiation of family groups (O'Connell-Rodwell et al., 2012), contrasting previous findings of passive leadership in males, where older males appeared to be unintentionally leading subordinates to resources (Allen et al., 2020).

These findings provide further support that mature males, and perhaps certain individuals such as those leading the LGR events here, are important for male elephant society (Allen et al., 2020; Allen, Croft & Brent, 2021; Chiyo et al., 2011; Goldenberg et al., 2014; Lee et al., 2011; Slotow et al., 2000). Further studies are needed to understand the underlying advantages of such surprisingly coordinated vocal bouts within groups of male African elephants, the level of coordination and vocal manipulation, as well as conditions that evoke such behavior that has not yet been documented in other populations.

ACKNOWLEDGEMENTS

The authors thank the Namibian Ministry of Environment and Etosha Ecological Institute for their support of this research. We also thank the contributing volunteers of Utopia

Scientific for supporting field work. We are also grateful to Jason D. Wood for early help with acoustic analysis and training with Raven Pro. We thank the reviewers for their constructive feedback on the manuscript.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

A grant from the National Geographic Society, the USFWS African Elephant Conservation Fund, and The Oakland Zoo Conservation Fund supported the field work and equipment in 2005 and 2007. The field work for this study was funded by the Stanford University Vice Provost Office for Undergraduate Education (VPUE) Faculty and Student Grants in 2011 and 2017 and the contributing volunteers of Utopia Scientific. Data analysis was supported by Harvard University Center for the Environment (HUCE) Summer Undergraduate Research Fund for an undergraduate student summer stipend, Utopia Scientific Donor Volunteers, The Elephant Sanctuary, and the National Institute of Health (NIH) grant number K01 DC017812 of the NIDCD. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

National Geographic Society.

USFWS African Elephant Conservation Fund.

The Oakland Zoo Conservation Fund.

Stanford University Vice Provost Office for Undergraduate Education (VPUE) Faculty and Student Grants.

Harvard University Center for the Environment (HUCE) Summer Undergraduate Research Fund.

Utopia Scientific Donor Volunteers, The Elephant Sanctuary.

National Institute of Health (NIH) grant number: K01 DC017812.

Competing Interests

Caitlin E O'Connell-Rodwell is a non-salaried founder and Chief Executive Officer of Utopia Scientific, a U.S. 501(c)3 non-profit public organization based in San Diego, and is currently working in Namibia and other countries focused on elephant research and conservation as well as science education and public health messaging. Timothy C Rodwell is a non-salaried founder and treasurer of Utopia Scientific. Jodie L Berezin is a scientist for Utopia Scientific and receives salary compensation for work with Utopia Scientific. Otherwise, the authors declare no competing interests.

Author Contributions

- Caitlin E. O'Connell-Rodwell conceived and designed the experiments, performed the experiments, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.

- Jodie L. Berezin analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the article, and approved the final draft.
- Alessio Pignatelli analyzed the data, prepared figures and/or tables, and approved the final draft.
- Timothy C. Rodwell conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the article, and approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

This study was conducted according to the guidelines of the Declaration of Helsinki and research grants approved by the Namibian Ministry of Environment, Forestry, and Tourism. This study was purely observational.

Data Availability

The following information was supplied regarding data availability:

The raw data is available in the [Supplemental Files](#).

The code used to analyze the data is available at Zenodo: Utopia Scientific Lab. (2024). Utopia-Sci-Lab/Male-Elephant-Vocal-Coordination: zenodo release (zenodo). Zenodo. <https://doi.org/10.5281/zenodo.11593367>.

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.17767#supplemental-information>.

REFERENCES

- Allen CRB, Brent LJN, Motsentwa T, Weiss MN, Croft DP. 2020.** Importance of old bulls: leaders and followers in collective movements of all-male groups in African savannah elephants (*Loxodonta africana*). *Scientific Reports* **10**:13996 DOI [10.1038/s41598-020-70682-y](https://doi.org/10.1038/s41598-020-70682-y).
- Allen CRB, Croft DP, Brent LJN. 2021.** Reduced older male presence linked to increased rates of aggression to non-conspecific targets in male elephants. *Proceedings of the Royal Society B: Biological Sciences* **288**:20211374 DOI [10.1098/rspb.2021.1374](https://doi.org/10.1098/rspb.2021.1374).
- Archie EA, Hollister-Smith JA, Poole JH, Lee PC, Moss CJ, Maldonado JE, Fleischer RC, Alberts SC. 2007.** Behavioural inbreeding avoidance in wild African elephants. *Molecular Ecology* **16**:4138–4148 DOI [10.1111/j.1365-294X.2007.03483.x](https://doi.org/10.1111/j.1365-294X.2007.03483.x).
- Archie EA, Moss CJ, Alberts SC. 2006.** The ties that bind: genetic relatedness predicts the fission and fusion of social groups in wild African elephants. *Proceedings of the Royal Society B: Biological Sciences* **273**:513–522 DOI [10.1098/rspb.2005.3361](https://doi.org/10.1098/rspb.2005.3361).
- Archie EA, Moss CJ, Alberts SC. 2011.** Friends and relations: kinship and the nature of female elephant social relationships. In: Moss CJ, Croze HJ, Lee PC, eds. *The Amboseli elephants: a long-term perspective on a long-lived mammal*. Chicago: University of Chicago Press, 238–245.

- Baotic A, Stoeger AS. 2017.** Sexual dimorphism in African elephant social rumbles. *PLOS ONE* 12:e0177411 DOI 10.1371/journal.pone.0177411.
- Berezin JL, Odom AJ, Hayssen V, O'Connell-Rodwell CE. 2023.** A snapshot into the lives of elephants: camera traps and conservation in Etosha National Park, Namibia. *Diversity* 15:1146 DOI 10.3390/d15111146.
- Black JM. 1988.** Preflight signalling in swans: a mechanism for group cohesion and flock formation. *Ethology* 79:143–157 DOI 10.1111/j.1439-0310.1988.tb00707.x.
- Boinski S. 1991.** The coordination of spatial position: a field study of the vocal behaviour of adult female squirrel monkeys. *Animal Behaviour* 41:89–102 DOI 10.1016/S0003-3472(05)80505-6.
- Boinski S, Campbell AF. 1995.** Use of trill vocalizations to coordinate troop movement among white-faced Capuchins: a second field test. *Behaviour* 132:875–901 DOI 10.1163/156853995X00054.
- Bousquet CA, Sumpter DJ, Manser MB. 2011.** Moving calls: a vocal mechanism underlying quorum decisions in cohesive groups. *Proceedings of the Royal Society B: Biological Sciences* 278:1482–1488 DOI 10.1098/rspb.2010.1739.
- Cairns SJ, Schwager SJ. 1987.** A comparison of association indices. *Animal Behaviour* 35:1454–1469 DOI 10.1016/S0003-3472(87)80018-0.
- Charif RA, Waack AM, Strickman LM. 2010.** *Raven Pro 1.4 user's manual*. Ithaca: Cornell Lab of Ornithology.
- Chiyo PI, Archie EA, Hollister-Smith JA, Lee PC, Poole JH, Moss CJ, Alberts SC. 2011.** Association patterns of African elephants in all-male groups: the role of age and genetic relatedness. *Animal Behaviour* 81:1093–1099 DOI 10.1016/j.anbehav.2011.02.013.
- Conradt L, Roper TJ. 2000.** Activity synchrony and social cohesion: a fission–fusion model. *Proceedings of the Royal Society B: Biological Sciences* 267:2213–2218 DOI 10.1098/rspb.2000.1271.
- Conradt L, Roper TJ. 2005.** Consensus decision making in animals. *Trends in Ecology & Evolution* 20:449–456 DOI 10.1016/j.tree.2005.05.008.
- David HA. 1987.** Ranking from unbalanced paired-comparison data. *Biometrika* 74:432–436 DOI 10.1093/biomet/74.2.432.
- de Waal FB. 1986.** The integration of dominance and social bonding in primates. *The Quarterly Review of Biology* 61:459–479 DOI 10.1086/415144.
- de Vries H, Stevens JMG, Vervaecke H. 2006.** Measuring and testing the steepness of dominance hierarchies. *Animal Behaviour* 71:585–592 DOI 10.1016/j.anbehav.2005.05.015.
- Evans KE, Harris S. 2008.** Adolescence in male African elephants, *Loxodonta africana*, and the importance of sociality. *Animal Behaviour* 76:779–787 DOI 10.1016/j.anbehav.2008.03.019.
- Gammell MP, de Vries H, Jennings DJ, Carlin CoM, TJ Hayden. 2003.** David's score: a more appropriate dominance ranking method than Clutton-Brock et al's index. *Animal Behaviour* 66:601–605 DOI 10.1006/anbe.2003.2226.
- Goldenberg SZ, de Silva S, Rasmussen HB, Douglas-Hamilton I, Wittemyer G. 2014.** Controlling for behavioural state reveals social dynamics among

- male African elephants, *Loxodonta africana*. *Animal Behaviour* **95**:111–119
[DOI 10.1016/j.anbehav.2014.07.002](https://doi.org/10.1016/j.anbehav.2014.07.002).
- Herve M. 2023.** RVAideMemoire: testing and plotting procedures for biostatistics. R Package Version 0.9-83-7.
- King AJ, Johnson DD, Van Vugt M. 2009.** The origins and evolution of leadership. *Current Biology* **19**:R911–R916 [DOI 10.1016/j.cub.2009.07.027](https://doi.org/10.1016/j.cub.2009.07.027).
- King AJ, Sueur C. 2011.** Where next? Group coordination and collective decision making by primates. *International Journal of Primatology* **32**(6):1245–1267
[DOI 10.1007/s10764-011-9526-7](https://doi.org/10.1007/s10764-011-9526-7).
- Lee PC, Moss CJ. 2012.** Wild female African elephants (*Loxodonta africana*) exhibit personality traits of leadership and social integration. *Journal of Comparative Psychology* **126**:224–232 [DOI 10.1037/a0026566](https://doi.org/10.1037/a0026566).
- Lee PC, Poole J, Njiraini NW, Sayialel CN, Moss C. 2011.** Male social dynamics: independence and beyond. In: *The Amboseli elephants: a long-term perspective on a long-lived mammal*. Chicago: University of Chicago Press.
- Levrero F, Touitou S, Fredet J, Nairaud B, Guery JP, Lemasson A. 2019.** Social bonding drives vocal exchanges in Bonobos. *Scientific Reports* **9**:711
[DOI 10.1038/s41598-018-36024-9](https://doi.org/10.1038/s41598-018-36024-9).
- McComb K, Moss C, Durant SM, Baker L, Sayialel S. 2001.** Matriarchs as repositories of social knowledge in African elephants. *Science* **292**:491–494
[DOI 10.1126/science.1057895](https://doi.org/10.1126/science.1057895).
- McComb K, Reby D, Baker L, Moss C, Sayialel S. 2003.** Long-distance communication of acoustic cues to social identity in African elephants. *Animal Behaviour* **65**:317–329 [DOI 10.1006/anbe.2003.2047](https://doi.org/10.1006/anbe.2003.2047).
- McComb K, Shannon G, Durant SM, Sayialel K, Slotow R, Poole J, Moss C. 2011.** Leadership in elephants: the adaptive value of age. *Proceedings of the Royal Society B: Biological Sciences* **278**:3270–3276 [DOI 10.1098/rspb.2011.0168](https://doi.org/10.1098/rspb.2011.0168).
- Morris-Drake A, Mumby HS. 2017.** Social associations and vocal communication in wild and captive male savannah elephants *Loxodonta africana*. *Mammal Review* **48**:24–36
[DOI 10.1111/mam.12106](https://doi.org/10.1111/mam.12106).
- Moss CJ. 1996.** Getting to know a population. In: Kangwana K, ed. *Studying elephants*. Nairobi: African Wildlife Foundation, 58–74.
- Mutinda H, Poole JH, Moss CJ. 2011.** Decision making and leadership in using the ecosystem. In: Moss CJ, Croze HJ, Lee PC, eds. *The Amboseli elephants: a long-term perspective on a long-lived mammal*. Chicago: The University of Chicago Press, 246–259.
- O’Connell-Rodwell CE, Berezin JL, Kinzley C, Freeman PT, Sandri MN, Kieschnick D, Abarca M, Hayssen V. 2024a.** To be unique or blend in: dynamics of male African elephant character durability across time and social contexts. *bioRxiv*.
[DOI 10.1101/2024.05.24.595367](https://doi.org/10.1101/2024.05.24.595367).
- O’Connell-Rodwell CE, Berezin JL, Pignatelli A, Rodwell TC. 2024b.** The use of vocal coordination in male African elephant group departures: evidence of active leadership and consensus. *bioRxiv* [DOI 10.1101/2024.05.31.596833](https://doi.org/10.1101/2024.05.31.596833).

- O'Connell-Rodwell CE, Freeman PT, Kinzley C, Sandri MN, Berezin JL, Wiśniewska M, Jessup K, Rodwell TC. 2022. A novel technique for aging male African elephants (*Loxodonta africana*) using craniofacial photogrammetry and geometric morphometrics. *Mammalian Biology* **102b**:591–613 DOI [10.1007/s42991-022-00238-2](https://doi.org/10.1007/s42991-022-00238-2).
- O'Connell-Rodwell CE, Sandri MN, Berezin JL, Munevar JM, Kinzley C, Wood JD, Wisniewska M, Kilian JW. 2022a. Male African elephant (*Loxodonta africana*) behavioral responses to estrous call playbacks may inform conservation management tools. *Animals* **12**:1162 DOI [10.3390/ani12091162](https://doi.org/10.3390/ani12091162).
- O'Connell-Rodwell CE, Wood JD, Kinzley C, Rodwell TC, Alarcon C, Wasser SK, Sapolsky R. 2011. Male African elephants (*Loxodonta africana*) queue when the stakes are high. *Ethology Ecology & Evolution* **23**:388–397 DOI [10.1080/03949370.2011.598569](https://doi.org/10.1080/03949370.2011.598569).
- O'Connell-Rodwell CE, Wood JD, Wyman M, Redfield S, Puria S, Hart LA. 2012. Antiphonal vocal bouts associated with departures in free-ranging African elephant family groups (*Loxodonta africana*). *Bioacoustics* **21**:215–224 DOI [10.1080/09524622.2012.686166](https://doi.org/10.1080/09524622.2012.686166).
- Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P, Stevens M, Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G, Chirico M, De Caceres M, Durand S, Evangelista H, Fitz John R, Friendly M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlenn D, Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Braak CTer, Weedon J. 2022. vegan: community ecology package. R Package Version 2.6-4.
- Palacios-Romo TM, Castellanos F, Ramos-Fernandez G. 2019. Uncovering the decision rules behind collective foraging in spider monkeys. *Animal Behaviour* **149**:121–133 DOI [10.1016/j.anbehav.2019.01.011](https://doi.org/10.1016/j.anbehav.2019.01.011).
- Pardo MA, Fristrup K, Lolchuragi DS, Poole JH, Granli P, Moss C, Douglas-Hamilton I, Wittemyer G. 2024. African elephants address one another with individually specific name-like calls. *Nature Ecology & Evolution* **8**:1353–1364 DOI [10.1038/s41559-024-02420-w](https://doi.org/10.1038/s41559-024-02420-w).
- Petit O, Bon R. 2010. Decision-making processes: the case of collective movements. *Behav Processes* **84**:635–647 DOI [10.1016/j.beproc.2010.04.009](https://doi.org/10.1016/j.beproc.2010.04.009).
- Poole J. 2011. Behavioral contexts of elephant acoustic communication. In: Moss CJ, Croze HJ, Lee PC, eds. *The Amboseli elephants: a long-term perspective on a long-lived mammal*. Chicago: University of Chicago Press, 125–161.
- Poole J, Moss C. 1989. Elephant mate searching: group dynamics and vocal and olfactory communication. In: *symposia of the Zoological Society of London*, 61.
- Poole JH, Payne K, Langbauer WR, Moss CJ. 1988. The social contexts of some very low frequency calls of African elephants. *Behavioral Ecology and Sociobiology* **22**:385–392 DOI [10.1007/BF00294975](https://doi.org/10.1007/BF00294975).
- Pyritz LW, King AJ, Sueur C, Fichtel C. 2011. Reaching a consensus: terminology and concepts used in coordination and decision-making research. *International Journal of Primatology* **32**:1268–1278 DOI [10.1007/s10764-011-9524-9](https://doi.org/10.1007/s10764-011-9524-9).

- R Core Team.** 2023. R: a language and environment for statistical computing. Version 4.3.1. Vienna: R Foundation for Statistical Computing. Available at <https://www.r-project.org>.
- Seltmann A, Majolo B, Schulke O, Ostner J.** 2013. The organization of collective group movements in Wild Barbary Macaques (*Macaca sylvanus*): social structure drives processes of group coordination in macaques. *PLOS ONE* **8**:e67285 DOI [10.1371/journal.pone.0067285](https://doi.org/10.1371/journal.pone.0067285).
- Slotow R, Van Dyk G, Poole J, Page B, Klocke A.** 2000. Older bull elephants control young males. *Nature* **408**:425–426 DOI [10.1038/35044191](https://doi.org/10.1038/35044191).
- Soltis J, Leong K, Savage A.** 2005. African elephant vocal communication II: rumble variation reflects the individual identity and emotional state of callers. *Animal Behaviour* **70**:589–599 DOI [10.1016/j.anbehav.2004.11.016](https://doi.org/10.1016/j.anbehav.2004.11.016).
- Sperber AL, Werner LM, Kappeler PM, Fichtel C, Wright J.** 2017. Grunt to go—Vocal coordination of group movements in redfronted lemurs. *Ethology* **123**:894–905 DOI [10.1111/eth.12663](https://doi.org/10.1111/eth.12663).
- Stewart KJ, Harcourt AH.** 1994. Gorillas' vocalizations during rest periods: signals of impending departure? *Behaviour* **130**:29–40 DOI [10.1163/156853994X00127](https://doi.org/10.1163/156853994X00127).
- Stoeger AS, Baotic A.** 2016. Information content and acoustic structure of male African elephant social rumbles. *Scientific Reports* **6**:27585 DOI [10.1038/srep27585](https://doi.org/10.1038/srep27585).
- Sueur C, Deneubourg JL, Petit O, Couzin ID.** 2010. Differences in nutrient requirements imply a non-linear emergence of leaders in animal groups. *PLOS Computational Biology* **6**:e1000917 DOI [10.1371/journal.pcbi.1000917](https://doi.org/10.1371/journal.pcbi.1000917).
- Sueur C, Kuntz C, Debergue E, Keller B, Robic F, Siegwalt-Baudin F, Richer C, Ramos A, Pelé M.** 2018. Leadership linked to group composition in Highland cattle (*Bos taurus*): Implications for livestock management. *Applied Animal Behaviour Science* **198**:9–18 DOI [10.1016/j.applanim.2017.09.014](https://doi.org/10.1016/j.applanim.2017.09.014).
- Sueur C, Petit O.** 2008. Organization of group members at departure is driven by social structure in Macaca. *International Journal of Primatology* **29**:1085–1098 DOI [10.1007/s10764-008-9262-9](https://doi.org/10.1007/s10764-008-9262-9).
- Šárová R, Špinka M, Stěhulová I, Ceacero F, Šimečková M, Kotrba R.** 2013. Pay respect to the elders: age, more than body mass, determines dominance in female beef cattle. *Animal Behaviour* **86**:1315–1323 DOI [10.1016/j.anbehav.2013.10.002](https://doi.org/10.1016/j.anbehav.2013.10.002).
- Walker RH, King AJ, McNutt JW, Jordan NR.** 2017. Sneeze to leave: African wild dogs (*Lycaon pictus*) use variable quorum thresholds facilitated by sneezes in collective decisions. *Proceedings of the Royal Society B: Biological Sciences* **284**(1862):20170347 DOI [10.1098/rspb.2017.0347](https://doi.org/10.1098/rspb.2017.0347).
- Whitehead H.** 2008. *Analyzing animal societies: quantitative methods for vertebrate social analysis*. Chicago, IL: University of Chicago Press.
- Wierucka K, Henley MD, Mumby HS.** 2021. Acoustic cues to individuality in wild male adult African savannah elephants (*Loxodonta africana*). *PeerJ* **9**:e10736 DOI [10.7717/peerj.10736](https://doi.org/10.7717/peerj.10736).